Novel Mutations in Human Luteinizing Hormone Beta Subunit Related to Polycystic Ovary Syndrome among Sudanese Women

Nidal Essa, Suliman Osman, Salah Jemaah, Mohamed A. Hassan, Rashid Eltayeb.

Introduction: Polycystic ovary syndrome (PCOS) is a common disorder that is not fully understood. Multiple hormonal and metabolic factors impact on disease pathophysiology resulting in various phenotypic characteristics among the PCOS population. Luteinizing hormone beta subunit (LHB) (protein ID P01229) is mapped on (chr19p13.3) and consists of three exons. Luteinizing hormone (LH) has a central role in stimulation ovarian steroidogenesis, in particular androgen production, and the promotion of ovulation.

Objectives: To determine if genetic variations of LHB are associated with PCOS among Sudanese families.

Methods: A prospective laboratory based cross-sectional study to examine genetic mutations in LHB that associate with PCOS in families (cases; n=35 families, 90 females and controls; n=11 families, 30 females) in Khartoum State, Sudan. Quantitative enzyme linked immuno-sorbent assay (ELISA) and polymerase chain reaction (PCR) with Sanger sequencing were used to analyze biochemical parameters and detect polymorphisms. Protein structure and function bioinformatics analysis was conducted using standard software.

Results: PCOS cases had significantly different biochemical parameters from the controls (LH: $p<0.001$; testosterone: $p<0.001$; fasting glucose: $p=0.02$; insulin: $p=0.01$; triglycerides: $p=0.03$; total cholesterol: $p<0.001$; high density lipoprotein (HDL): $p=0.012$; low density lipoprotein (LDL): $p<0.001$). There were no differences in follicle stimulating hormone (FSH) ($p=0.984$) or prolactin ($p=0.068$). Sanger sequencing revealed 5 single nucleotide polymorphisms (rs5030775, A18T; rs746167425, R22K; rs1800447, W28R; rs35270001, H30R; and rs34349826, I35T) located on (exon 2) of *LHB* gene that were statistically correlated with serum LH, Testosterone and insulin levels among PCOS families.

Conclusion: This is the first molecular family-based study in Sudan exploring the genetics of the *LHB* gene in women manifesting PCOS. These novel mutations give further information about the role of genetic inheritance and may explain some of the altered ovarian function and responses in women with PCOS.

Keywords: PCOS, Luteinizing hormone, SNPs, infertility, family studies

Introduction:

Polycystic ovary syndrome (PCOS) is a disorder which is associated with a hormone imbalance, anovulation and multiple paused follicles in the ovaries that is linked to disparate metabolic disorders.^{1,2} It has a complex etiology with genetic, epigenetic and environmental factors. Adherence to the three common diagnostic criteria of increased androgens, polycystic ovaries on ultrasound scan and anovulation (1990 National Institutes of Health's (NIH), the Rotterdam criteria and the Androgen Excess Society), which cover the different phenotypic features of PCOS, suggests a prevalence of 1.6-18% among women of reproductive age.²⁻⁴

There are epigenetic factors that can promote PCOS. In various animal models, and in women, increased exposure of the developing fetus to increased androgen concentrations before birth can result in developmental programming of the features of PCOS. Prenatal androgens result in dynamic changes in gene expression, resulting hyperinsulinemia, adipocyte dysfunction, visceral obesity, and PCOS in offspring.^{9, 10} In addition, there are environmental factors involved in PCOS. An increase of weight can unmask a PCOS phenotype in adulthood.

However, there are genetic elements of PCOS, although the genes involved remain poorly characterized. Various familial aggregation studies suggest that PCOS is an autosomal dominant syndrome and there is an association in both sisters and brothers of women with PCOS with infertility problems. In addition monozygotic and dizygotic twin studies suggest a genetic contribution to PCOS. Recent genome-wide association studies (GWAS) in Han Chinese women with PCOS identified 11 genetic loci that related to PCOS. These loci were in genes involved in steroidogenesis, steroid hormone action, gonadotrophin action, regulation of internal secretion, energy balance, chronic inflammation, and other pathways relevant to PCOS.¹¹⁻¹⁶

PCOS is associated with an exaggerated LH pulse frequency and amplitude that enhances androgen secretion. In addition, there are alterations in the androgen synthetic pathway that associate with enhanced ovarian and adrenal androgen production¹⁷⁻¹⁹. Moreover, excessive androgen secretion results on many types of hormonal and metabolic

abnormalities, which are characterized by hirsutism, anovulation, clinical hyperandrogenism, hyperinsulinemia, insulin resistance (IR), and obesity, which are all more common in PCOS¹⁻³. Therefore, assessment of hormonal profile among PCOS population using high-quality techniques such as liquid chromatography tandem mass spectrometry (LC-MS/MS), plays a key diagnostic marker of PCOS.

As androgens are driven by LH, we focused on the *LHB* gene. We aimed to examine the relationship between single nucleotide polymorphisms (SNPs) of the *LHB* gene with the clinical features of PCOS among Sudanese women within the reproductive age.

Methodology:

This study was approved by the national ethics committee - Federal Ministry of Health (3-12-19). The inclusion criteria for PCOS families were women diagnosed with PCOS, according to the Rotterdam criteria that had one or two sisters diagnosed as PCOS within the reproductive age (20-40) years. In control families neither the woman nor any sisters had any features of PCOS. Any women below the reproductive age or with family history of adrenal hyperplasia, hyperprolactinemia, Cushing syndrome, thyroid disorders, ovarian cancer or breast cancer and causes of infertility other than PCOS were excluded from the study.

This prospective laboratory based cross-sectional study involved blood sampling carried out in different fertility centers as well as different localities (families' homes) in Khartoum State. Hormonal profile estimations were carried out using validated tests within the clinical service laboratory with quantitative enzyme linked immunosorbent assay (ELISA) and enzymatic methods for measurement of fasting blood glucose and lipid profile. DNA extraction from whole blood was carried out using a commercial kit according to the manufacturer's instructions. Primer 3 was used to design the *LHB* primers (Table 1). Routine polymerase chain reaction (PCR) and gel electrophoresis was used for band amplification and visualization. The Sanger dioxynucleotide sequencing method was used for genotyping and the bioinformatics software (BLAST, USCF Chimera, Project Hope and Raptor X) was used for protein structure, function and prediction.

Descriptive and biological information about study groups were analyzed by SPSS version 23 using descriptive analysis for age, BMI, and biochemical analysis. T-tests were used for statistical significance after testing for normality, Chi square test were used for proportion analysis and Person correlation were used for statistical association. Statistical significance was $p < 0.05$.

Table 1: The *LHB* gene primers for PCR protocol.

Gene	Forward	Length	Reverse	Length	Product length
<i>LH-B</i>	GGAGGCCTCTTTCTGGAGGG	20	GGAAGCCCTCTGTTCCTGC	19	571

Results:

The patient characteristics and biochemical parameters are shown in (Table 2). There was no difference in age and BMI, serum levels of FSH and PRL assessment between the groups. However, there were clear increases in LH and testosterone concentrations ($p < 0.001$) in the PCOS group. This meant that the LH: FSH ratio was significantly elevated in women with PCOS ($P < 0.001$). In addition, there was evidence of insulin resistance and dyslipidaemia in the women with PCOS (Table 2). Women with PCOS suffered from signs of hyperandrogenism with hirsutism (42.9%), acne (54.5%) and seborrhea (30%) (Table 3) and testosterone concentrations were significantly linked to menstrual irregularity (Table 4). This highlights the validity of the cohort of PCOS patients when compared to controls.

DNA was extracted and sequenced from the cases and controls and the chemical structure of the LH protein determined (Figure 1). Within the PCOS families there were five SNPs that were identified within exon (2) of $LH\beta$ (Figure 2, Table 5). These were significantly associated with serum levels of LH, testosterone and menstrual irregularity (Tables 6, 7).

Table (2) Characteristics of PCOS and controls cohorts:

Parameters	Study groups	Mean± SD	<i>p-value</i>
Age. yrs.	Case	35.3±13,9	0.283
	Control	37±18,1	
BMI	Case	28.6±112,9	0.529
	Control	26.6±59,3	
FBG/mmol/l	Case(n=90)	6.2 ±1.8	0.020*
	Control(n=30)	5.4±0.9	
LH (ng/dl)	Case (n=90)	17.7±12.3	0.000*
	Control(n=30)	9.7±5.4	
FSH (ng/dl)	Case(n=90)	5.4±4.6	0.984
	Control(n=30)	5.4±2.9	
PRL (ng/dl)	Case(n=90)	12.4±8.2	0.068
	Control(n=30)	8.0±6.2	
LH/FSH ratio	Case(n=90)	5.5±6.1	0.000*
	Control(n=30)	1.9±0.8	
TG (mmol/l)	Case(n=90)	1.5±0.6	0.030*
	Control(n=30)	1.1±0.3	
T.Chol (mmol/l)	Case(n=90)	4.1±1.5	0.000*
	Control(n=30)	2.2±0.4	
HDL(mmol/l)	Case(n=90)	1.2±0.2	0.012*
	Control(n=30)	1.4±0.2	
LDL(mmol/l)	Case(n=90)	2.3±1. 6	0.000*
	Control(n=30)	0.27±0.5	
Insulin (ng/dl)	Case(n=90)	15.2±8.4	0.010*
	Control(n=30)	7.7±6.2	
Testosterone (ng/dl)	Case(n=90)	33.6±18.0	0.000*
	Control(n=30)	10.5±7.0	

* The mean difference is significant at the 0.050 level

Table (3): The association between clinical hyperandrogenism and menstrual cycle status among study groups.

Clinical Hyperandrogenism		Status of MC		<i>P-value</i>
		Regular	Irregular	
Hirsutism	Yes	15%	24%	0.000*
	No	42%	17%	
Acne	Yes	25%	27%	0.040*
	No	32.2%	15%	
Alopecia	Yes	17%	18	0.040*
	No	42.2%	22.2%	
Seborrhea	Yes	17%	17.7%	0.130
	No	42.2%	24.4%	

* The mean difference is significant at the 0.05 level.

Table (4): The association between hormonal profile and menstrual cycle status among study groups.

Parameters	Status of MC		<i>P-value</i>
	Regular ($\bar{x} \pm SD$)	Irregular ($\bar{x} \pm SD$)	
LH (ng/dl)	16.9 $\pm$ 127,9	16,079 $\pm$ 106,9	0.741
FSH (ng/dl)	5,077 $\pm$ 41,2	5,750 $\pm$ 47,4	0.476
PRL (ng/dl)	11,708 $\pm$ 75,4	11.932 $\pm$ 88,8	0.900
Insulin(ng/dl)	12.90 $\pm$ 8,3	15.84 $\pm$ 8,6	0.108
Testosterone (ng/dl)	26.00 $\pm$ 17,8	36.18 $\pm$ 18,5	0.011*

* The mean difference is significant at the 0.05 level.

Figure (1): (A)The normal chemical structure of LH protein, (B) The chromatogram graph for the sequenced protein, and (C) The gel electrophoresis showing the LH band.

Figure (2) Example of normal (A) and mutated amino acids (red color) on sequenced exon (2) LH β gene (B).

Table (5) Distribution of LH β genotypes and allele frequencies in healthy controls and patients with PCOS

SNP ID	Genotype	Cases	Controls
rs5030775 (A18T)	Homozygous A	100%	0%
rs746167425 (R22k)	Homozygous A	100%	0%
rs1800447 (W28R)	Homozygous C	100%	0%
rs35270001 (H30R)	Heterozygous A/G	94%	0%
rs34349826 (I35T)	Homozygous C	100%	0%

Table (6): The association of LH β polymorphisms and menstrual irregularity among study groups:

Status of MC	Mutations		p-value
	Present	Absent	
Regular	45.5%	12.2%	0.034*
Irregular	40%	0.02%	

* The mean difference is significant at the 0.05 level.

Table (7): The association of LH β polymorphisms and hormonal profile among study groups:

Parameters	Mutation		95% CI		p-value
	Present	Absent			
LH (ng/dl)	17.732 $\pm$ 23.2	9.6 $\pm$ 53.5	11.395	149.870	.023*
FSH (ng/dl)	5.3 $\pm$ 46.11	5.3 $\pm$ 28.61	-26.038	26.567	.984
Prolactin (ng/dl)	12.442 $\pm$ 82.2	8.0 $\pm$ 61.8	-.3297	91.821	.068
Insulin (ng/dl)	15.3 $\pm$ 8.4	7.6 $\pm$ 6.1	2.673	12.410	.003*
Testosterone (ng/dl)	33.6 $\pm$ 18.0	10.7 $\pm$ 7.0	17.369	28.827	.000*

4. Discussion:

In this study we investigated if different biochemical parameters were associated with polymorphisms of the *LHB* gene among Sudanese families who had more than one PCOS sufferer and compared this to control families without women with PCOS. The biochemical parameters within the PCOS group were consistent with PCOS and showed that they were different to the control group. We identified five polymorphisms in the *LHB* gene in the families with PCOS in Sudan. There were no polymorphisms in the control cases.

In this study the biochemical parameters were within the normal ranges but significantly different from the control group of the study, except BMI, serum levels of FSH and PRL. The cases and controls were clearly different symptomatically. Women with PCOS were more likely to suffer from menstrual irregularity (46.8%) and signs of clinical hyperandrogenism; facial hirsutism (42.9%), Acne (54.5%), Seborrhea (33.3%) and Alopecia (35.6%) (Esmaeilzadeh et al., 2014; Patel, 2018) than controls. They also showed different biochemical profiles with consensus to the international hallmarks of PCOS of hormonal imbalance (Act, 2018; Azziz, 2016; Escobar-Morreale, 2018; Harris et al., 2017).

In agreement with many previous studies the PCOS cohort also showed dyslipidemia (Esmaeilzadeh et al., 2015; Shoaib et al., 2015; Bennal et al., 2014). In contrast, Sadeghi et al., 2014 suggested that this was obesity dependent. They showed that there is no association between serum lipid profile, total cholesterol level, LDL cholesterol and HDL cholesterol (p-value =0.86, 0.21, 0.76 respectively) and PCOS except in the term of TG (p-value =0.04) in PCOS women with BMI > 25 kg/m². While only obese PCOS women showed significant difference on serum TC as well as LDL-C (p=0.000) when compared with overweight/normal PCOS indicating that obesity, rather than PCOS per se, makes women more susceptible to dyslipidemia and cardiovascular disease (Castelo-Branco et al., ; Spałkowska et al., 2018). In our study there was no difference in BMI, and the mean BMI was only slightly overweight, but women with PCOS demonstrated dyslipidaemia.

In this cohort the higher glucose and insulin suggests the PCOS women are more predisposed to T2DM on their future life similar. This finding is similar to Forslund et al., (2020) and Rashidi (2018) who showed increased risk across different BMI groups. Obesity is a risk factor for impaired glucose tolerance as is PCOS with increased androgen concentrations. There is an adverse effects of obesity on serum testosterone level which is an inverse correlation among PCOS women with BMI ≥ 30 ($p=0.014$) while lean PCOS women had positive correlation ($p=0.009$). This suggests that the interaction of obesity and hyperandrogenism in the development of impaired glucose tolerance is complex (Spałkowska et al., 2018; Saxena et al., 2013). Hyperandrogenemia is responsible for accumulation of the adipose tissue in the upper body (android fat distribution pattern; Good et al.). Interestingly Aydin et al., concluded that no differences in body composition between lean women with PCOS and healthy controls. Overall the influences of PCOS and obesity and their interaction on fat distribution and function requires further study.

We report that sisters of affected women have irregular menstrual cycles and this agrees with the Turkish study that showed that sisters had menstrual irregularity and larger ovaries with higher serum androstenedione and dehydroepiandrosterone sulfate levels than sisters without PCO, suggesting a spectrum of clinical phenotypes in PCO families. This parallels what was reported in a multicenter study with (1466) subjects, of whom (363) had PCOS and (79) polycystic ovaries (PCO) without other symptoms of PCOS. This is further evidence for PCOS clustering in families.

We examined the LHB gene as LH-stimulated androgen synthesis is fundamental to PCOS. We found that the LHB polymorphisms (rs1800447 (W28R) and rs34349826 (I35T) homozygotes (CC) had an association with menstrual irregularities with infertility, and elevation of serum testosterone level (Furui et al., 1994), whereas the Finnish subjects whom were homo- or heterozygous for the polymorphic *LHB* allele were largely healthy. The first report of this mutation among the male population was a patient with hypogonadotropic hypogonadism provided clinical and experimental evidence that this novel mutation causes selective LH deficiency. Other studies showed significantly higher serum LH concentrations in V-LH carriers among the young Baltic

men and Estonian male infertility patients. This suggests polymorphism interaction with the genetic background.

Herein we described the presence of 5 single nucleotide polymorphisms of *LHB* gene, that shared between the investigated cases were with homozygous (AA) inheritance mode. They are located on the signal peptide and the homologue glycoprotein parts and thus could have significant influence on serum level of the hormone without inactivating or damaging its biological function. The most reported mutations of this research findings were (rs1800447 (W28R) and rs34349826 (I35T)) which are associated with protein glycosylation and have already been linked to response to controlled ovarian stimulation, male infertility and PCOS.

Protein glycosylation is one of the most frequent and relevant post-translational modifications and plays a key role in the modulation of protein properties particularly with regard to the half-life of the hormone, heterodimer stability and the intrinsic bioactivity. The carbohydrate content of LH is approximately 15%(N50) while that of FSH is 20%(N25,N42), and that of hCG is 30%(N33,N50) and the circulatory half-lives of the gonadotropins are about 20 mins for LH, 2 hours for FSH, and 12–24 hours for hCG. Molecular alterations of V-LHB, due to rs1800447 (Trp28Arg) and Ile35Thr (rs34349826) create an extra glycosylation signal (Asn-X-Thr) into the *LHB* chain, which introduces an oligosaccharide side chain into Asn13, similar to that present in HCG. This might increase the affinity binding between the peptide portion of the hormone and the receptor and also increase the lifespan of the hormone (Shafaghiet *et al.*, 2019).

Findings using serum variant homozygotes showed that V-LH is more active than WT-LH in bioassays in vitro, but it has a shorter half-life in circulation. The shorter half-life of V-LH is possibly compensated for by about (40%) higher promoter activity of the gene, explained by 8 SNPs in the (50-flanking regions) of *V-LHB*. They were reported as having a modulatory effect on the hyperandrogenemia of PCOS. More recently, Alviggi *et al.* have confirmed that *v-LHB* polymorphism produces a less active form of the hormone which is not able to support satisfactorily FSH activity during controlled ovarian stimulation, and that the *v-LHβ* carriers experience a considerable

decrease of the number of transferred embryos. The presence of this mutation statistically associated with high serum levels of LH and testosterone and irregular menstrual cycles highlighting the essential role of LH/FSH cooperation in the latest stages of follicle maturation.

The polymorphism rs5030775 (A18T) is associated with more potent stimulation of the IP3 response than wild type LH mapped on the signal peptide position, which is a short peptide located in the N-terminal of proteins which is responsible for protein recognition and secretion and it is often cleaved to generate the mature protein. Therefore, presence of this mutation within the leader sequence, which has been reported as potential stimulator of IP3 in contrast to the cAMP pathway, could cause an alteration of GnRH pulse frequency due to overstimulation of IP3. This in turn would increase *LHB* gene expression and might influence abnormal recognition of the prohormone within the endoplasmic reticulum when binding with SRP occurs. This would affect its interaction with the LHCGR receptor. Interestingly, a point mutation associated with autosomal recessive familial isolated hypoparathyroidism caused improper cleavage by signal peptidase at the normal position. Remarkably, this mutation ((historically reported as Thr-3A) A18T) was also found in 3% of the Rwanda population with a heterogeneous pattern that might be a sign of the evolutionarily background origin of PCOS and support the idea that a high level of serum *LHB* is associated with increased pulse frequencies because of high GnRH pulse frequencies that favor *LHB* gene expression, whereas low GnRH pulse frequencies favor *FSHB* gene expression.

We are not aware of any literature in regarding to rs746167425 (R22K) and rs35270001 (H30R). With regards to their physiochemical properties Arginine mutated into a Lysine at position 22, which is smaller, would not be expected to affect protein stability. However, Histidine, located very close to a residue that makes a cysteine bond, forms a hydrogen bond with Tryptophan on position 28. When mutated to Arginine there is positive charge and a reduction in size. This creates an extra glycosylation site on the mutated form that might cause loss of hydrophobic interactions with other molecules on the surface of the protein.

In conclusion, The diagnosis of PCOS makes the patient aware of possible fertility concerns, dysfunctional bleeding, endometrial cancer, obesity, diabetes, dyslipidemia, hypertension, and theoretical increased risk of cardiovascular disease. Since PCOS has genetic elements, knowledge of polymorphisms that affect LH action may bring awareness of PCOS to family members and future children. Given that insulin resistance is heavily associated with PCOS, with early genetic identification these individuals could be directed to increased screening to help reduce adverse diabetic and cardiovascular outcomes that will likely have better long-term outcomes. Interventions may include early lifestyle advice, as well as insulin sensitizing medication, such as metformin, screening for diabetes and hyperlipidemia. Overall, in Sudan there seem to be genetic markers that highlight the risk of PCOS that can be detected before PCOS is manifest.

References:

1. Franks, S., College, I., Kingdom, U., Hospital, H., & Kingdom, U. (2018). Polycystic Ovary Syndrome. In *Encyclopedia of Endocrine Diseases, 2nd Edition* (2nd ed., Issue February). Elsevier Inc. <https://doi.org/10.1016/B978-0-12-801238-3.03910-6>.
2. Crespo, R. P., Bachega, T. A. S. S., Mendonça, B. B., & Gomes, L. G. (2018). An update of genetic basis of PCOS pathogenesis. *Archives of Endocrinology and Metabolism*, 62(3), 352–361. <https://doi.org/10.20945/2359-3997000000049>
3. Day, F., Karaderi, T., Jones, M. R., Meun, C., He, C., Drong, A., Kraft, P., Lin, N., Huang, H., Broer, L., Magi, R., Saxena, R., Laisk, T., Urbanek, M., Hayes, M. G., Thorleifsson, G., Fernandez-Tajes, J., Mahajan, A., Mullin, B. H., ... Welt, C. K. (2018). Large-scale genome-wide meta-analysis of polycystic ovary syndrome suggests shared genetic architecture for different diagnosis criteria. *PLoS Genetics*, 14(12), 1–20. <https://doi.org/10.1371/journal.pgen.1007813>.
4. Deswal, R., Nanda, S., & Dang, A. S. (2019). Systems Biology in Reproductive Medicine Association of Luteinizing hormone and LH receptor gene polymorphism with susceptibility of Polycystic ovary syndrome Association of Luteinizing hormone and LH receptor gene polymorphism with. *Systems Biology in Reproductive Medicine*, 0(0), 1–9. <https://doi.org/10.1080/19396368.2019.1595217>
5. Janjic, M. M., Prévide, R. M., Patrick, A. F., Sherman, A., Smi, K., Daniel, A., Bje, I., & Stoji, S. S. (2019). *Divergent expression patterns of pituitary gonadotropin subunit and GnRH receptor genes to continuous GnRH in vitro and in vivo*. 1–14. <https://doi.org/10.1038/s41598-019-56480-1>
6. Shafaghi, M., Akbar, A., & Minuchehr, Z. (2019). Rational design of hyper-glycosylated human luteinizing hormone analogs (a bioinformatics approach). *Computational Biology and Chemistry*, 79(January), 16–23. <https://doi.org/10.1016/j.compbiolchem.2019.01.002>

7. Pan, J., Tan, Y., Wang, F., Hou, N., Xiang, Y., Zhang, J., & Liu, Y. (2018). *Aberrant expression and DNA methylation of lipid metabolism genes in PCOS: a new insight into its pathogenesis*. 1–12. <https://doi.org/10.1186/s13148-018-0442-y>
8. Jin, P., & Xie, Y. (2018). Treatment strategies for women with polycystic ovary syndrome. *Gynecological Endocrinology*, 34(4), 272–277. <https://doi.org/10.1080/09513590.2017.1395841>
9. Pasquali, R. (2018). Contemporary approaches to the management of polycystic ovary syndrome. *Therapeutic Advances in Endocrinology and Metabolism*, 9(4), 123–134. <https://doi.org/10.1177/2042018818756790>
10. Nataraja, S., Sriraman, V., & Palmer, S. (2018). *Allosteric Regulation of the Follicle-Stimulating Hormone Receptor*. 159(May), 2704–2716. <https://doi.org/10.1210/en.2018-00317>
11. Shaaban, Z., Khoradmehr, A., Reza, M., Shirazi, J., & Tamadon, A. (2018). *Pathophysiological mechanisms of gonadotropins – and steroid hormones – related genes in etiology of polycystic ovary syndrome*. <https://doi.org/10.22038/ijbms.2018.31776.7646>
12. Zou, J., Wu, D., Liu, Y., & Tan, S. (2018). Association of luteinizing hormone / choriogonadotropin receptor gene polymorphisms with polycystic ovary syndrome risk: a meta- analysis. *Gynecological Endocrinology*, 0(0), 1–5. <https://doi.org/10.1080/09513590.2018.1498834>
13. Id, U. B., Id, K. M., & Wijesundera, S. (2018). *Identification of selected genetic polymorphisms in polycystic ovary syndrome in Sri Lankan women using low cost genotyping techniques*. 1, 1–19.
14. Casarini, L., Santi, D., & Emilia, R. (2018). LH (Luteinizing Hormone). In *Encyclopedia of Endocrine Diseases, 2nd Edition* (2nd ed., Issue March). Elsevier Inc. <https://doi.org/10.1016/B978-0-12-801238-3.64299-X>
15. Escobar-Morreale, H. F. (2018). Polycystic ovary syndrome: Definition, aetiology, diagnosis and treatment. *Nature Reviews Endocrinology*, 14(5), 270–284. <https://doi.org/10.1038/nrendo.2018.24>
16. Wolf, W. M., Wattick, R. A., Kinkade, O. N., & Olfert, M. D. (2018). Geographical prevalence of polycystic ovary syndrome as determined by region and race/ethnicity. *International Journal of Environmental Research and Public Health*, 15(11), 1–13. <https://doi.org/10.3390/ijerph15112589>

17. Le, A. R. T. I. C. (2018). *Delayed Diagnosis and a Lack of Information Associated With Dissatisfaction in Women With Polycystic Ovary Syndrome*. 102(July), 604–612. <https://doi.org/10.1210/jc.2016-2963>
18. Dadachanji, R., Shaikh, N., & Mukherjee, S. (2018). Genetic Variants Associated with Hyperandrogenemia in PCOS Pathophysiology. *Genetics Research International*, 2018. <https://doi.org/10.1155/2018/7624932>
19. Patel, S. (2018). Polycystic ovary syndrome (PCOS), an inflammatory, systemic, lifestyle endocrinopathy. *Journal of Steroid Biochemistry and Molecular Biology*, 182(February), 27–36. <https://doi.org/10.1016/j.jsbmb.2018.04.008>
20. Sun, X., Wu, X., Duan, Y., Liu, G., Yu, X., & Zhang, W. (2017). *Family-Based Association Study of rs17300539 and rs12495941 Polymorphism in Adiponectin Gene and Polycystic Ovary Syndrome in a Chinese Population*. 78–84. <https://doi.org/10.12659/MSM.901944>
21. Mohammad, M. B., & Seghinsara, A. M. (2017). Polycystic ovary syndrome (PCOS), diagnostic criteria, and AMH. *Asian Pacific Journal of Cancer Prevention*, 18(1), 17–21. <https://doi.org/10.22034/APJCP.2017.18.1.17>
22. Karakas, S. E. (2017). Clinica Chimica Acta New biomarkers for diagnosis and management of polycystic ovary syndrome. *Clinica Chimica Acta*, 471(June), 248–253. <https://doi.org/10.1016/j.cca.2017.06.009>
23. Sales, M. F., Rodrigues, K. F., Martins, S. R., Carvalho, L. M. L., Ferreira, C. N., S, M. O., Candido, A. L., Reis, F. M., Silva, I. F. O., Campos, F. M. F., & Gomes, K. B. (2017). *Molecular and Cellular Endocrinology Microparticles*: Inflammatory and haemostatic biomarkers in Polycystic Ovary Syndrome. 443. <https://doi.org/10.1016/j.mce.2017.01.017>
24. El Hayek, S., Bitar, L., Hamdar, L. H., Mirza, F. G., & Daoud, G. (2016). Poly Cystic Ovarian Syndrome: An updated overview. *Frontiers in Physiology*, 7(APR), 1–15. <https://doi.org/10.3389/fphys.2016.00124>
25. Merkin, S. S., Phy, J. L., Sites, C. K., & Yang, D. (2016). Environmental determinants of polycystic ovary syndrome. *Fertility and Sterility*, 106(1), 16–24. <https://doi.org/10.1016/j.fertnstert.2016.05.011>

- 26.D, D. M. T. F. P., D, B. N. M., & D, R. A. M. (2016). Evolutionary determinants of polycystic ovary syndrome□: part 2. *Fertility and Sterility*, May, 1–6. <https://doi.org/10.1016/j.fertnstert.2016.05.016>
- 27.Sezgin, E., & Yildiz, B. O. (2016). *Evolutionary determinants of polycystic ovary syndrome — part 1.* May. <https://doi.org/10.1016/j.fertnstert.2016.05.010>
- 28.Atwood, C. S., & Vadakkadath, S. (2016). Molecular and Cellular Endocrinology The spatiotemporal hormonal orchestration of human folliculogenesis, early embryogenesis and blastocyst implantation. *Molecular and Cellular Endocrinology*, 430, 33–48. <https://doi.org/10.1016/j.mce.2016.03.039>
- 29.Hayes, M. G., Urbanek, M., Ehrmann, D. A., Armstrong, L. L., Lee, J. Y., Sisk, R., Karaderi, T., Barber, T. M., Mccarthy, M. I., Franks, S., Lindgren, C. M., Welt, C. K., Diamanti-kandarakis, E., Panidis, D., Goodarzi, M. O., Azziz, R., Zhang, Y., & James, R. G. (2016). *Corrigendum□: Genome-wide association of polycystic ovary syndrome implicates alterations in gonadotropin secretion in European ancestry populations.* 10762. <https://doi.org/10.1038/ncomms10762>
- 30.Rohrbasser, L. J., Alsaffar, H., & Blair, J. (2016). *The Hypothalamus – Pituitary Axis.* <https://doi.org/10.1007/978-3-319-27318-1>
- 31.Zhu, R., Wong, Y., & Yong, E. (2016). Best Practice & Research Clinical Obstetrics and Gynaecology Sonographic evaluation of polycystic ovaries. *Best Practice & Research Clinical Obstetrics & Gynaecology*, 37, 25–37. <https://doi.org/10.1016/j.bpobgyn.2016.02.005>
- 32.Kanamarlapudi, V., Gordon, U. D., & López, A. (2016). Luteinizing hormone / chorionic gonadotrophin receptor overexpressed in granulosa cells from polycystic ovary syndrome ovaries is functionally active. *Reproductive BioMedicine Online*, 32(6), 635–641. <https://doi.org/10.1016/j.rbmo.2016.03.003>
- 33.Huang-doran, I., Franks, S., Morton, N. M., & Huang-doran, I. (2016). *Genetic Rodent Models of Obesity- Associated Ovarian Dysfunction and Subfertility□: insights into Polycystic Ovary Syndrome.* 7(June). <https://doi.org/10.3389/fendo.2016.00053>
- 34.Rosenfield, R. L., Ehrmann, D. A., & Biochemical, A. (2016). *The Pathogenesis of Polycystic Ovary Syndrome (PCOS): The Hypothesis of PCOS as Functional Ovarian.* 37(October), 467–520. <https://doi.org/10.1210/er.2015-1104>

35. Dhar, N., Mohan, A., Thakur, C., Chandra, N. R., & Dighe, R. R. (2016). Molecular and Cellular Endocrinology Dissecting the structural and functional features of the Luteinizing hormone receptor using receptor specific single chain fragment variables. *Molecular and Cellular Endocrinology*, 1–12. <https://doi.org/10.1016/j.mce.2016.02.022>
36. Pincas, H., González-maeso, J., Ruf-zamojski, F., & Sealson, S. C. (2016). *G Protein-Coupled Receptors*. <https://doi.org/10.1007/978-3-319-27318-1>
37. Azziz, R. (2016). Introduction: Determinants of polycystic ovary syndrome. *Fertility and Sterility*, 106(1). <https://doi.org/10.1016/j.fertnstert.2016.05.009>
38. Mcallister, J. M., Legro, R. S., Modi, B. P., & State, P. (2016). *HHS Public Access*. 26(3), 118–124. <https://doi.org/10.1016/j.tem.2014.12.004>. Functional
39. El-Shal, A. S., Zidan, H. E., Rashad, N. M., Abdelaziz, A. M., & Harira, M. M. (2016). Association between genes encoding components of the Leutinizing hormone/Luteinizing hormone-choriogonadotrophin receptor pathway and polycystic ovary syndrome in Egyptian women. *IUBMB Life*, 68(1), 23–36. <https://doi.org/10.1002/iub.1457>
40. Lan, C., Chen, M., Tai, K., Yu, D. C. W., Yang, Y. Y., Jan, P., Yang, Y. Y., Chen, H., Ho, H., Thathapudi, S., Kodati, V., Rutz, C., Klein, W., Sch, R., Bui, H. N., Sluss, P. M., Hayes, F. J., Blincko, S., Knol, D. L., ... Mukherjee, S. (2016). Molecular and Cellular Endocrinology Microparticles□: Inflammatory and haemostatic biomarkers in Polycystic Ovary Syndrome. *Fertility and Sterility*, 7(1), 1–12. <https://doi.org/10.1007/978-3-319-27318-1>
41. Dunaif, A. (2016). *Perspectives in Polycystic Ovary Syndrome□: From Hair to Eternity*. 101(March), 759–768. <https://doi.org/10.1210/jc.2015-3780>.
42. Irving F. Stein, Michael L. Leventhal, Amenorrhea associated with bilateral polycystic ovaries, American Journal of Obstetrics and Gynecology, Volume 29, Issue 2, 1935, Pages 181-191, ISSN 0002-9378, [https://doi.org/10.1016/S0002-9378\(15\)30642-6](https://doi.org/10.1016/S0002-9378(15)30642-6).
43. Verhoeven, M. O., & Lambalk, C. B. (2016). *Ovarian Physiology*. <https://doi.org/10.1007/978-3-319-27318-1>
44. Lee, H., Oh, J., Sung, Y., Chung, H., Kim, H., Kim, G. S., Cho, Y. S., & Kim, J. T. (2015). *Genome-wide association study identified new*

- susceptibility loci for polycystic ovary syndrome*. 30(3), 723–731. <https://doi.org/10.1093/humrep/deu352>
45. Rutz, C., Klein, W., & Sch, R. (2015). *N-Terminal Signal Peptides of G Protein-Coupled Receptors*: Significance for Receptor Biosynthesis , Trafficking , and Signal Transduction. 1–21. <https://doi.org/10.1016/bs.pmbts.2015.03.003>
46. Thathapudi, S., & Kodati, V. (2015). *Association of Luteinizing Hormone Chorionic*. 19(3), 128–132. <https://doi.org/10.1089/gtmb.2014.0249>
47. Almawi, W. Y., Hubail, B., & Arekat, D. Z. (2015). *Leutinizing hormone / choriogonadotropin receptor and follicle stimulating hormone receptor gene variants in polycystic ovary syndrome*. 607–614. <https://doi.org/10.1007/s10815-015-0427-0>
48. Bui, H. N., Sluss, P. M., Hayes, F. J., Blincko, S., Knol, D. L., Blankenstein, M. A., & Heijboer, A. C. (2015). *Clinica Chimica Acta Testosterone , free testosterone , and free androgen index in women*: Reference intervals , biological variation , and diagnostic value in polycystic ovary syndrome. *Clinica Chimica Acta*, 450, 227–232. <https://doi.org/10.1016/j.cca.2015.08.019>
49. Day, F. R., Hinds, D. A., Tung, J. Y., Stolk, L., Styrkarsdottir, U., Saxena, R., Bjornes, A., Broer, L., Dunger, D. B., Halldorsson, B. V., Lawlor, D. A., Laval, G., Mathieson, I., Mccardle, W. L., Louwers, Y., Meun, C., Ring, S., Uitterlinden, G., Wareham, N. J., ... Perry, J. R. B. (2015). *ovary syndrome*. 1–7. <https://doi.org/10.1038/ncomms9464>
50. Salilew-wondim, D., Wang, Q., Tesfaye, D., Schellander, K., Hoelker, M., Hossain, M., & Tsang, B. K. (2015). *Polycystic ovarian syndrome is accompanied by repression of gene signatures associated with biosynthesis and metabolism of steroids , cholesterol and lipids*. ???, 1–14. <https://doi.org/10.1186/s13048-015-0151-5>.

Revelation of Novel Mutations in Human Luteinizing Hormone Beta Subunit Related to Polycystic Ovary Syndrome among Sudanese Women

Nidal Essa, Suliman Osman, Salah Jemaah, Mohamed A. Hassan, Rashid Eltayeb.

Figure (1): The frequencies of presence of Acne, hirsutism, and status of MC between study groups; (54, 5) but 57.1% of them had no facial hirsutism similar to 77% of control group. 84.6% of our control had regular menstrual cycle and only 15.3 had irregular menstrual cycle, while the case group had the higher distribution than control group.

Figure (2) (A)The normal chemical structure of LH protein, (B) The chromatogram graph for the sequenced protein, (C) The gel electrophoresis showing the PCR product band (571bp).

Figure (3) illustrated both normal and mutated amino acids on sequenced exon (2) LH β gene