

Extraction of Viral Nucleic Acids with Carbon

Nanotubes Increases SARS-CoV-2 RT-qPCR

Detection Sensitivity

*Sanghwa Jeong¹, Eduardo G. Grandio¹, Nicole Navarro², Rebecca L. Pinals¹, Francis Ledesma¹, Darwin Yang¹, and Markita P. Landry^{*1,3-5}*

¹Department of Chemical and Biomolecular Engineering, University of California, Berkeley, California 94720, United States

²Department of Chemistry, University of California, Berkeley, California 94720, United States

³Innovative Genomics Institute (IGI), Berkeley, California 94720, United States

⁴California Institute for Quantitative Biosciences, QB3, University of California, Berkeley, California 94720, United States

⁵Chan-Zuckerberg Biohub, San Francisco, California 94158, United States

Name	Target Sequence in SARS-CoV-2	Target Region in SARS-CoV-2
Target1	AAAAAAAAAAAAAAAAAAAAA (poly A tail)	29861-29893
Target2	GCUGCUGAGGCUUCAAG	29024-29041
Target3	UUACAAACAUUGGCCGCA	29164-29181
Target4	AAAUUCAAGAUCAAGU	29304-29322
Target5	ACUGUGACUCUUCUCCU	29444-29461
Target6	UUUUCGCUUUUCCGUUA	29574-29591
Target7	GAACUUCUCCUGCUAGAA	28884-28901
Target8	GUGCUACAACUCCUCAA	28745-28762
Target9	ACUACCUAGGAACUGGGC	28605-28622
Target10	UAAAUUCCUCGAGGACA	28465-28482

Table S1. Sequences targeted in the SARS-CoV-2 viral genome (variant MT007544.1) with 10 target DNA sequences. Base location in the genome for each target is shown. All the target sequences, except Target 1, aim to bind N gene region in the SARS-CoV-2 RNA genome.

SARSCoV-2_N1	SARS-CoV-2_N1-F	GAC CCC AAA ATC AGC GAA AT
	SARS-CoV-2_N1-R	TCT GGT TAC TGC CAG TTG AAT CTG
	SARS-CoV-2_N1Probe	FAM-ACC CCG CAT TAC GTT TGG TGG ACC BHQ1

Table S2. Primer and probes used for RT-qPCR detection of SARS-CoV-2 viral RNA. All reagents come from the 2019-nCoV RUO Kit (Integrated DNA technologies).