Discernment of Mediator and Outcome Measurement in the PACE trial

Ewan Carr¹, Silia Vitoratou¹, Trudie Chalder^{2,3}, and Kimberley Goldsmith¹

¹ Department of Biostatistics and Health Informatics, Institute of Psychiatry, Psychology and Neuroscience, King's College London, London, United Kingdom.

² Academic Department of Psychological Medicine, Institute of Psychiatry, Psychology and Neuroscience, Weston Education Centre, King's College London, London, United Kingdom.

³ South London and Maudsley NHS Foundation Trust, London, United Kingdom.

Corresponding author: Dr Kimberley Goldsmith, Department of Biostatistics and Health Informatics, Institute of Psychiatry, Psychology & Neuroscience, King's College London, London, SE5 8AF, United Kingdom. +44 (0) 20 7848 0059; kimberley.goldsmith@kcl.ac.uk

Words: 3970; Tables: 3; Figures: 2

Abstract

Background: When measuring latent traits, such as those used in psychology and psychiatry, it can be unclear whether the instruments used are measuring different concepts. This issue is particularly important in context of mediation analysis, since for a sound mediation hypothesis the mediator and outcome should be distinct. We sought to assess the extent of measurement overlap between mediators and outcomes in the PACE trial (n=640).

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

Methods: Potential measurement overlap was assessed using generalised linear latent variable models where confirmatory factor models quantified the extent to which the addition of cross-loading items resulted in significant improvements in model fit.

Results: Out of 26 mediator-outcome pairs considered, only six showed evidence of cross-loading items, supporting the suggestion that mediator and outcome constructs in the PACE trial were conceptually distinct.

Conclusions: This study highlights the importance of assessing measurement overlap in mediation analyses with latent traits to ensure mediator and outcome instruments are distinct.

Keywords: Mediation, confirmatory factor analysis, construct overlap, latent trait.

Background

Studies in psychiatry and psychology are often interested in unobserved constructs such as behaviour or cognition that are captured using multiple items from a questionnaire. These constructs can be measured within the generalised linear latent variable models framework (Bartholomew et al., 2011; Skrondal & Rabe-Hesketh, 2004), consisting of two parts: the measurement part and the structural part of the model. The measurement part refers to the observed items which designate ('load' onto) a latent variable (the estimated magnitudes of these relationships are referred to as the 'factor loadings'); the structural part refers to the latent variables and their relationships.

The measurement part of the model is typically specified based on theory or prior empirical research. The assignment of observed items to latent constructs may be straightforward if, for example, the items are drawn from an established psychological

scale, or there is some other strong theoretical motivation for including, on a given instrument, some items over others. However, this is not always the case and there may be some ambiguity about which items to include on a given instrument. A single observed item might be related to two or more latent constructs, represented in the model by two or more statistically significant factor loadings. This arrangement would produce ‘cross-loadings’, where a single item loads initially onto the intended latent construct (the ‘primary loading’) as well as onto a second latent construct (the ‘cross-loading’).

In this study we are concerned with what happens when two measurement instruments share several cross-loading items and how our interpretations should be adjusted accordingly. While overlap may exist in the structural part of the model (e.g. Schaufeli & Bakker, 2004), our focus here is on overlap in the measurement part of the model: do the items measuring one construct also serve as indicators of another? We are particularly interested in how measurement overlap affects the interpretation of mediational hypotheses. Mediation is where a ‘third variable’ transmits the effect of one variable to another and is thus “intermediate in the causal sequence relating an independent variable to a dependent variable” (p. 1, MacKinnon, 2008). The single mediator model in a clinical trial setting postulates that the effect of a randomly-assigned treatment (R) on a continuous outcome (Y) is mediated by a third variable (M) (MacKinnon, 2008). Since Baron and Kenny’s widely-cited paper describing mediation and moderation in psychology (1986), methods for mediation analysis have grown in sophistication (Emsley et al., 2010; K. Goldsmith et al., 2018; K. A. Goldsmith et al., 2018; Preacher, 2015; VanderWeele et al., 2014, 2012). It is now recognised that single mediator model makes several inferential assumptions (MacKinnon, 2008) including temporal precedence (R occurs before M which occurs before Y), that the temporal

ordering is appropriate for the hypothesised mediational chain, and that there are no omitted influences (i.e. other variables influencing R , M , or Y , either observed or unobserved). Here we consider the additional (and often untested) assumption that M and Y represent distinct theoretical constructs. If the mediator represents an earlier assessment of the outcome then the model interpretation changes; the mediator no longer represents a ‘third variable’ in the mediational chain.

Psychological studies often involve latent constructs that necessarily share common themes and have the potential to overlap. Kaufman et al. (2005), for example, considered potential mediators of CBT for adolescents with depressive symptoms. Their potential mediators included the Automatic Thoughts Questionnaire (ATQ) (Hollon & Kendall, 1980) which shares similar themes and question wordings as their outcome measure, the Beck Depression Inventory (BDI-II) (Beck et al., 1996). In particular, regarding notions of worthlessness, self-dislike, and self-criticalness. This raises the question of whether these two constructs are truly distinct. Identifying potential overlap is not always straightforward. Given the difficulties of distinguishing and measuring some latent traits, cross-loading may occur even among items that do not appear, on face value, to be conceptually similar. As we describe below, many of the mediators and outcomes studied in this paper do not share common items or wordings but nonetheless could be seen to share similar themes.

This paper was motivated by an existing mediation analysis that tested whether unhelpful cognitions and behaviours mediated the effect of cognitive behavioural therapy (CBT) in treating patients with chronic fatigue syndrome (Chalder et al., 2015). Drawing on data from the PACE trial (White et al., 2011), this analysis considered whether the effect of treatment allocation (R ; 0 weeks) on fatigue (Y ; 52 weeks) was

93 mediated by variables measured at 12 weeks using the Cognitive Behavioural
 94 Responses Questionnaire (CBRQ; Ryan et al. (2017)). For example, illustrated in Figure
 95 1, 34% of the effect of CBT on fatigue was transmitted via ‘fear avoidance’ beliefs.

Figure 1. Standardised effects in mediation models, reproduced from Chalder et al. 2015 (p. 150).

Notes.

CBT = cognitive behaviour therapy; APT = adaptive pacing therapy. Besides treatment, models included: centre, Standardised Clinical Interview for DSM-IV (SCID) depression status, London criteria for myalgic encephalomyelitis status, International Chronic Fatigue Syndrome (CFS) criteria, baseline measures of both outcome variables, baseline Work and Social Adjustment Scale, SCID anxiety disorder status, age, sex, CFS group membership, receipt of benefits, benefits in dispute, physical illness attribution, fibromyalgia status, illness duration, Jenkins Sleep Score, employment status, body-mass index, and physical symptoms (PHQ-15) score.

96
 97 The mediators and outcomes used in the PACE trial were purposefully chosen to
 98 represent distinct theoretical constructs. The questionnaires used to measure these
 99 constructs had previously been psychometrically evaluated and found to be reliable and
 100 valid (Cella et al., 2011; Chalder et al., 1993; McHorney et al., 1993; Ryan et al., 2017).
 101 This first step is critical: careful selection of appropriate constructs can help avoid

ambiguity about which items correspond to which constructs and minimise potential overlap at the outset. However, despite these efforts, there remains the possibility of overlap between mediator and outcome constructs and the possibility that the identified mechanisms could in part be explained by measurement overlap (i.e. the mediator and outcome ‘measuring the same thing’). For example, items measuring the mediators include “I stay in bed to control my symptoms” and “When I experience symptoms, I rest” while items measuring the fatigue outcome include “Do you need to rest more?” and “Do you have problems with tiredness?”

This paper presents a procedure to assess the extent of measurement overlap between the mediators and outcomes used in the PACE trial to address the question: were the chosen mediators and outcomes ‘measuring the same thing?’ To our knowledge, no previous papers have addressed measurement overlap in the context of mediation analysis. We expect to find minimal overlap since the included instruments were chosen to represent distinct theoretical constructs and have previously been shown to be psychometrically reliable and valid. We note that our aim is to assess measurement overlap and not to re-evaluate the structure of these constructs, which has been addressed in previous studies (Cella et al., 2011; Chalder et al., 1993; McHorney et al., 1993; Ryan et al., 2017).

Methods

Data

The PACE trial was a four-arm randomised trial designed to assess the effectiveness of rehabilitative treatments for chronic fatigue syndrome (White et al., 2011). 640 patients were recruited from six specialist chronic fatigue syndrome clinics in the UK National Health Service between March 2005 and November 2008. Eligibility criteria, trial

methods and results are described elsewhere (White et al., 2011). For this analysis we used mediator data gathered at 12 weeks post-randomisation and outcome data gathered at 52 weeks post-randomisation, consistent with the published mediation analysis of PACE (Chalder et al., 2015).

Measures

We considered seven mediating factors consistent with those included in the PACE mediation analysis (Chalder et al., 2015), derived from questions on the Cognitive Behavioural Responses Questionnaire (Ryan et al., 2017). The five *cognitive* mediators were ‘fear avoidance,’ ‘catastrophising,’ ‘damage,’ ‘embarrassment avoidance,’ and ‘symptom focusing.’ Two *behavioural* mediators were ‘all-or-nothing behaviour’ and ‘behavioural avoidance.’ For this analysis, we included both the original versions of these factors as well as shortened versions (3-item) described in Ryan et al. (2017), with the exception of ‘catastrophising’ for which a short version was not available.

Two outcomes were considered, following the original PACE mediation analysis, both of which have previously been psychometrically validated. Fatigue was measured by the 11-item Chalder Fatigue Questionnaire (CFQ) (Cella et al., 2011; Chalder et al., 1993). Physical function was measured by the physical function subscale (PF, 10 items) of the Medical Outcomes Study Short Form Health Survey (SF-36) (McHorney et al., 1993). The factor structure and question wordings for each mediator and outcome are presented in Supplementary Table 1.

Statistical analyses

We assessed overlap between the mediators and outcomes by fitting three sets of confirmatory factor models, described below. There were 13 mediating factors (7

factors plus 6 short versions) and two outcomes giving a total of 26 mediator-outcome pairs. Overall model fit was assessed based on the comparative fit index (CFI, with preferred values higher than 0.90) (Bentler, 1990), the root mean square error of approximation (RMSEA, with preferred values less than 0.08) (Browne & Cudeck, 1993), and the relative χ^2 (ratio of χ^2 to model degrees of freedom). Close fit was indicated by $CFI \geq 0.95$, $RMSEA \leq 0.05$ and relative χ^2 close to 2 (Kaplan, 2008). Changes in model fit – for example, after adding a cross-loading to a model – were assessed using the DIFFTEST procedure in Mplus (Asparouhov & Muthén, 2006). This is a procedure for testing nested models with mean and variance adjusted χ^2 statistics. In our example, this involved fitting two models: the unrestricted model (H_1) where the cross-loading was freely estimated; and the restricted model (H_0) where the cross-loading parameter was constrained to 0. All models were fitted in Mplus 7.3 (Muthén & Muthén, 2015) using R version 4.0.3 (R Core Team, 2017) and the MplusAutomation package (Hallquist & Wiley, 2018). All models were estimated using the robust weighted least squares (WLSMV) estimator.

Before investigating overlap between constructs, we made some assumptions about the individual factors. Firstly, as these factors had previously been psychometrically evaluated, we assumed that each factor would independently achieve good overall model fit. Secondly, since our aim was to assess construct overlap, not the structure of individual scales, we assumed the psychometric structure of each factor was known and appropriate at the outset.

In Step 1, we estimated a single factor confirmatory factor model (uni-dimensional CFA) for each mediator-outcome pair (Figure 2, Step 1). This was a model where a single continuous latent variable was measured by the observed items for *both* the

mediator and outcome. For this model we hypothesised that overall model fit would be poor since these latent variables were expected to be measuring distinct constructs. Good model fit would have indicated that the observed items for the mediator and outcome could be adequately explained by a single latent variable and that the purported two-factor structure was not required, which was hypothesised to be unlikely.

In Step 2, we estimated a two-factor CFA model, with continuous latent variables for the mediator and outcome measured by their respective observed items (Figure 2, Step 2). Since we expected the mediator and outcome to be related at the structural level, following Chalder et al. (2015), we included a parameter modelling the covariance between each pair of latent variables. Poor fit for the Step 2 model would suggest possible overlap between the mediator and outcome. However, this was tested formally in the next step.

In Step 3, we added cross-loadings to the two-factor model based on the modification indices and improvements in model fit. A cross-loading here refers to a path from the latent variable representing the mediator to an observed item representing the outcome, or vice versa. Overlap was indicated to the extent that the addition of each additional cross-loading item resulted in a statistically significant improvement in overall model fit (tested using the DIFFTEST procedure in Mplus, described above). Starting with the two-factor model from Step 2, above:

- i. From the modification indices, select the cross-loading with the highest χ^2 value and add this to the model.
- ii. Use the DIFFTEST procedure to test whether the addition of this cross-loading item produces a statistically significant improvement in overall model fit ($p \leq 0.05$).

197 The above steps were repeated until the additional cross-loading did not result in a
198 statistically significant improvement in model fit, the modification indices did not
199 contain any cross-loadings, or the model failed to converge. For models that failed to
200 converge (which would indicate a poorly-defined model) the previous converging
201 iteration was considered as the final model. In this way, we identified the number of
202 cross-loading items for each mediator-outcome pair (Figure 2, Step 3).

203

Figure 2. Illustration of model-fitting procedure

204 Cross-loading items were classified according to the strength with which the item
 205 loaded onto the primary and secondary factors (Table 1). The primary factor was the
 206 factor that the item was theoretically intended to load onto, based on prior psychometric
 207 analyses; the secondary factor would represent a cross-loading. Factor loadings were
 208 classified as strong ($0.6 > \lambda \leq 1.0$), moderate ($0.3 > \lambda \leq 0.6$), weak ($0.2 > \lambda \leq 0.3$) or
 209 non-salient ($0.0 > \lambda \leq 0.2$) (Hair et al. 2010). A cross-loading was thus classified as
 210 *switched* if at Step 2 it loaded onto the primary factor with a strong or moderate loading,
 211 but after Step 3, ‘switched’ to load more strongly onto the secondary factor. An item
 212 was *shared* if after Step 3 it loaded with equal strength onto both the primary and
 213 secondary factors. A *strong cross-loading* was an item that, after Step 3, continued to
 214 load strongly onto the primary factor but also loaded with at least moderate strength
 215 onto the secondary factor. Finally, for each mediator-outcome pair we calculated the
 216 number of cross-loadings (total, and of each type) and the percentage of cross-loading
 217 items (i.e. the total number of cross-loadings after the Step 3, relative to the number of
 218 factor loadings at Step 2).

Table 1: Classification of cross-loading items, based on factor loading for primary and secondary factors.

Primary [†]	Secondary [†]	Type for pair
Non-salient	Non-salient	Non-salient
	Weak	Switched
	Moderate	Switched
	Strong	Switched
Weak	Non-salient	No cross-loading
	Weak	Shared
	Moderate	Switched
	Strong	Switched

Moderate	Non-salient	No cross-loading
	Weak	No cross-loading
	Moderate	Shared
	Strong	Strong cross-loading (switched)
Strong	Non-salient	No cross-loading
	Weak	No cross-loading
	Moderate	Strong cross-loading (switched)
	Strong	Shared

† Strong = $0.6 > \lambda \leq 1.0$; Moderate = $0.3 > \lambda \leq 0.6$; Weak = $0.2 > \lambda \leq 0.3$; Non-salient = $0.0 > \lambda \leq 0.2$

Results

In Step 1, almost all single-factor CFA models fitted poorly indicating, as expected, that there were few mediator-outcome combinations where a single latent variable underlying the observed items was consistent with the data. The lowest values of RMSEA and relative χ^2 were 0.18 and 21.41, respectively, thus indicating unacceptable fit. One pair had a CFI above 0.95 (for CFQ and the short version of ‘Damage’), however its RMSEA was 0.20. For almost all mediator-outcome pairs CFI was below 0.90 (results available upon request).

Between Steps 2 and 3, cross-loadings were added to most mediator-outcome pairs based on the procedure described above. There were only 4/26 pairs for which no cross-loadings were added (Supplementary Table 2). This was either because the first cross-loading to be tested did not result in a statistically significant improvement in model fit, or because the modification indices contained no cross-loadings. However, most of the cross-loadings added at Step 3 were non-salient. This means they produced a statistically significant improvement in model fit (based on DIFFTEST) but had non-salient factor loadings (below 0.2) onto the secondary factor. Excluding those that were

235 non-salient, only six mediator-outcome pairs exhibited cross-loading, involving a total
 236 of 18 cross-loading items. Of these, three were ‘shared’ and 15 were ‘strong’ cross-
 237 loadings (Table 2).

Table 2: Salient cross-loadings added during the iterative procedure

Mediator	Outcome [‡]	Number of factor loadings		Number of factor loadings added	Type of cross-loading			Percentage of salient cross-loadings [†]
		Before iterative procedure	After iterative procedure		Non-salient	Shared	Strong cross-loading	
Fear avoidance (short)	CFQ	14	22	8	4	0	4	29%
Avoidance/resting behaviour (short)	CFQ	14	20	6	2	1	3	29%
Damage (short)	CFQ	14	20	6	2	2	2	29%
Fear avoidance	CFQ	17	24	7	5	0	2	12%
Symptom focusing	CFQ	17	23	6	3	0	3	18%
Fear avoidance (short)	PF	13	17	4	3	0	1	8%

[†]This is defined as the number of salient cross-loadings relative to the number of factor loadings before carrying out the iterative procedure. [‡] CFQ = Chalder Fatigue Questionnaire; PF = Physical function subscale of the SF-36.

238

239 Table 3 presents factor loadings and question wordings for cross-loading items from the
 240 six mediator-outcome pairs. In total, there were only 18 instances of cross-loading.
 241 Most instances of overlap (17/18) involved items 8 to 11 from the fatigue questionnaire
 242 (CFQ). These were questions that assessed whether participants “feel weak” (item 8),
 243 “have difficulty concentrating” (item 9), “have problems thinking clearly” (item 10), or

“make slips of the tongue when speaking” (item 11). They cross-loaded mostly with the mediators: avoidance/resting behaviour (4 items), damage (4 items), fear avoidance (7 items), and symptom focusing (3 items). The other pair where there was a strong cross-loading was the SF-36 physical function item “Walking several hundred yards” and fear avoidance. Items exhibiting overlap, such as these, indicated that in addition to loading on the latent variable measuring fatigue, combinations of these items also loaded onto the mediator constructs listed above (e.g. avoidance/resting behaviour, damage, and fear avoidance). More precisely, the addition of each cross-loading to the two-factor model resulted in a statistically significant improvement in model fit. The responses to these items, therefore, are influenced by both the outcome and mediator, and thus represent a point of overlap between the two.

Table 3: Factor loadings and question wordings for cross-loading items

Pair		Item	Question wording	Factor loadings		Type of cross-loading
Outcome [‡]	Mediator			Outcome	Mediator	
CFQ	Avoidance/resting behaviour (short version)	CFQ_10	Do you have problems thinking clearly?	0.81	0.57	Strong cross-loading
		CFQ_11	Do you make slips of the tongue when speaking?	0.79	0.38	Strong cross-loading
		CFQ_8	Do you feel weak?	0.87	0.32	Strong cross-loading
		CFQ_9	Do you have difficulty concentrating?	0.81	0.61	Shared
CFQ	Damage (short version)	CFQ_10	Do you have problems thinking clearly?	0.71	0.63	Shared
		CFQ_11	Do you make slips of the tongue when speaking?	0.73	0.41	Strong cross-loading
		CFQ_8	Do you feel weak?	0.83	0.34	Strong cross-loading
		CFQ_9	Do you have difficulty concentrating?	0.71	0.61	Shared
CFQ	Fear avoidance	CFQ_10	Do you have problems thinking clearly?	0.99	-0.31	Strong cross-loading
		CFQ_9	Do you have difficulty concentrating?	1.00	-0.34	Strong cross-loading
CFQ	Fear avoidance (short version)	CFQ_10	Do you have problems thinking clearly?	0.68	0.59	Strong cross-loading
		CFQ_11	Do you make slips of the tongue when speaking?	0.70	0.39	Strong cross-loading
		CFQ_8	Do you feel weak?	0.81	0.31	Strong cross-loading
		CFQ_9	Do you have difficulty concentrating?	0.69	0.59	Strong cross-loading
CFQ	Symptom focusing	CFQ_10	Do you have problems thinking clearly?	0.90	-0.45	Strong cross-loading
		CFQ_11	Do you make slips of the tongue when speaking?	0.82	-0.32	Strong cross-loading
		CFQ_9	Do you have difficulty concentrating?	0.91	-0.43	Strong cross-loading
PF	Fear avoidance (short version)	PF_8	Walking several hundred yards [†]	0.83	-0.31	Strong cross-loading

‡ CFQ = Chalder Fatigue Questionnaire; PF = Physical function subscale of the SF-36.

† Does your health limit you in these activities?

255

256

257 Discussion

258 In this paper we assessed overlap between latent constructs used to quantify mediators
259 and outcomes in the PACE randomised controlled trial of rehabilitative treatments for
260 chronic fatigue syndrome (White et al., 2011). The purpose of examining overlap was to
261 verify a key assumption of the theoretical mediation model, namely, that these latent
262 constructs were ‘measuring different things.’ Otherwise, rather than evaluating a
263 mediational process, what we would be doing would be more akin to erroneously
264 assessing relationships between repeated measures of our outcome. Adopting a latent
265 variable modelling framework, we assessed whether there were cross-loadings between
266 latent variables representing the mediators and outcomes. Ideally, the observed items for
267 a given mediator should be largely unique and should not simultaneously load onto the
268 outcome. Conversely, the items measuring the outcome should not also serve as
269 indicators of the mediator.

270 We identified low levels of measurement overlap among the mediators and outcomes in
271 the PACE trial. The single-factor models fitted poorly, suggesting that the items
272 quantifying each mediator-outcome pair could not be explained by a single latent
273 variable. Our analysis suggested many potential cross-loadings, but most of these were
274 non-salient. Out of 26 mediator-outcome pairs, just six indicated cross-loading items,
275 and nearly all of these involved the same four items from a single factor (CFQ). This

suggests that the measures used to quantify the mediators and outcomes in PACE were largely distinct.

To our knowledge, no previous studies have examined item overlap in the context of mediation analysis, although several studies have assessed measurement overlap in more general settings. Parker et al. (1991) used exploratory factor analysis (EFA) to assess construct overlap between measures of alexithymia and depression, concluding that “alexithymia was a construct that is distinct and separate from depression” (p. 387). Bagby and Rector (1998) similarly used EFA to assess overlap in personality constructs linked to depression. They found some constructs to overlap (self-criticism and neuroticism) whereas others appeared distinct (dependency and neuroticism). Other studies have used latent variable techniques to assess overlap – e.g. latent class analysis (Mezuk et al., 2013) or item response theory (Hoertel et al., 2015) – but these have adopted differing interpretations of ‘overlap’ and thus are not directly comparable with our approach. For example, Mezuk et al. (2013) defined overlap as the extent to which two sets of observed items measuring depression and frailty identified similar subgroups in their sample. Hoertel et al. (2015) considered whether depressive symptoms (e.g. sadness, fatigue) ‘overlap’ based on differing response profiles in groups of women at different stages of pregnancy. Perhaps closest to our study is the work of Gunzler and Morris (2015) who adopted a latent variable framework to assess overlap between variables measuring depression and variables measuring fatigue and cognitive decline related to multiple sclerosis. Their procedure used MIMIC models (“Multiple Indicator, Multiple Cause”) to test for differential item functioning (DIF), whereas we used confirmatory factor analysis (CFA). Moreover, they took what could be termed a backwards stepwise approach in that they started with a model including all theoretically relevant cross-loadings and *removed* them one-by-one. We started with a

model without any cross-loadings and *added* them one-by-one. Both approaches appear valid methods for assessing measurement overlap. Their approach requires that potential cross-loadings are identified a priori based on theory or clinical experience, whereas our approach tests all possible cross-loadings. This exhaustive approach seems preferable when trying to rule out potential overlap, or when lacking strong theoretical expectations as to which precisely items will overlap – as is the case here, given that the chosen constructs were designed to be theoretically distinct.

When assessing mediation it is important to ensure that the mediator and outcome represent distinct constructs. Extensive item overlap would be problematic insofar as it changes the interpretation of the mediation model. In the most extreme case the mediator may no longer represent a distinct third variable. While some studies have sought to reduce overlap by removing similar items (e.g. Segerstrom et al., 2000), this is not always considered or appropriate (e.g. if the psychometric properties of a scale have been validated based on the complete set of items (Bartholomew et al., 2011) then removal of individual items may invalidate the scale).

We found little evidence of mediator-outcome overlap in the PACE trial data, with just 6/26 pairs suggesting overlap, and for each pair no more than 29% of items were affected. It is challenging to define precise thresholds as to what degree of overlap would be considered ‘problematic.’ The proportion of items that exhibit cross-loading is one possible criterion, but the types of cross-loadings also seem important (e.g. the number of ‘shared’ vs. ‘switched’ cross-loadings). It also seems important to consider the theoretical interpretation of the cross-loading items. Are most instances of cross-loading attributable to a single item, or is there widespread cross-loading across many items?

Where substantial overlap is detected – for example, more than half of the items measuring the mediator are simultaneously cross-loading onto the outcome – our interpretation of the mediation analysis should be revised accordingly. In this case, it might be argued that the mediator better represents an earlier measurement of the outcome rather than a distinct construct, so we cannot say with confidence that a *third* variable transmits the effect of our treatment to our outcome. Instead, the situation may be closer to the treatment (T_0) changing our outcome (T_2) via a change in an earlier measurement of our outcome (at T_1). This is an important distinction when it comes to understanding the nature of the processes we are studying – i.e. whether the process is mediational or more akin to repeated measurement of an outcome, which will affect the analysis and interpretation. We recommend that researchers planning mediational analysis using latent traits assess mediator/outcome overlap, and if it is appreciable, either change the measures used, or at least interpret accordingly.

Our analysis used assessments of the mediators at 12 weeks and outcomes at 52 weeks post-randomisation. These time points were chosen to correspond with the measures used in the published mediation analysis (Chalder et al., 2015). An alternative approach, however, might have been to examine construct overlap at baseline (0 weeks). This may be useful when a mediation analysis is planned but follow-up data are not yet available. Another advantage of using baseline measures to assess overlap, in a trial setting, is that they are unaffected by the treatment intervention. However, evidence of a lack of overlap at baseline does not imply absence of overlap at later time points. We suggest then that researchers should either examine overlap based on follow-up assessments that they intend to use for their mediation analysis (once such data become available) or should establish measurement invariance between baseline and follow-up measures.

Steps to establish longitudinal measurement invariance are described elsewhere (Fokkema et al., 2013).

Strengths and limitations

To our knowledge, no studies have addressed the issue of measurement overlap in the context of mediation analysis where the variables of interest are latent (measured via questionnaire). Our study benefited from a large sample size and examined mediators and outcomes that were pre-specified, had been measured on multiple occasions, and had previously been psychometrically validated (Cella et al., 2011; Chalder et al., 1993; McHorney et al., 1993; Ryan et al., 2017). In terms of limitations, the iterative procedure to identify potential cross-loadings made repeated use of the DIFFTEST command in Mplus but no adjustment for multiple comparisons was made. Hoertel et al. (2015), for example, used the Benjamini–Hochberg (1995) procedure to adjust *P*-values produced by DIFFTEST. However, whereas they were attempting to minimize the number of false positives, our analysis was more concerned with false negatives. We wanted to increase the likelihood of detecting cross-loading items, and were thus less concerned by inflated false positive rates due to multiplicity. Also, our approach was somewhat data-driven, in that we used the modification indices to select potential cross-loadings. Importantly, however, the modification indices were not used to select the ‘best’ or ‘correct’ model. The constructs and indicators used to measure mediators and outcomes in PACE were theoretically motivated and purposefully chosen to represent distinct theoretical constructs. We did not use the modification indices to alter these constructs. Rather, the modification indices were used as a computational tool to identify, out of several equally probable cross-loadings, the most influential cross-loading to test first. Ideally, researchers would have a sense about which items will be

potentially overlapping. However, in many applications this will not be the case, and the potentially cross-loading items may not be obvious, suggesting the need for a more exploratory approach. This approach is similar to how modification indices are used in the context of a Multiple Indicators Multiple causes (MIMIC) model to identify measurement invariance (e.g. Woods, 2009). An alternative approach might have been to select cross-loadings theoretically, as suggested by Gunzler and Morris. For example, the researcher might select a list of theoretically plausible cross-loadings, and *these* are tested during the iterative procedure, based on their position in the modification indices.

Conclusions

In conclusion, we found little evidence of construct overlap between mediators and outcomes measured in the PACE trial. This is an important result, supporting the findings of the published mediation analysis (Chalder et al., 2015). Where researchers find more extensive overlap between constructs measuring mediators and outcomes their interpretation of the mediation analysis should be adjusted accordingly, since the mediator can likely no longer be considered a distinct ‘third variable.’ Finally, we reiterate the importance of selecting appropriate and distinct constructs (and questionnaire items) at the point of study design and before data collection. Drawing on relevant theory to inform the selection of distinct constructs will help minimise construct overlap from the outset, allowing studies to obtain more robust conclusions about mediation hypotheses.

References

Asparouhov T, Muthén B. Robust chi square difference testing with mean and variance adjusted test statistics. Mplus Web Notes. 2006;No. 10:1–6.

396 Bagby RM, Rector NA. Self-criticism, dependency and the five factor model of
397 personality in depression: Assessing construct overlap. *Personality and Individual*
398 *Differences*. 1998 Jun;24(6):895–7.

399 Baron RM, Kenny DA. The moderator-mediator variable distinction in social
400 psychological research: Conceptual, strategic, and statistical considerations. *Journal of*
401 *personality and social psychology*. 1986;51(6):1173.

402 Bartholomew DJ, Knott M, Moustaki I. Latent variable models and factor analysis: A
403 unified approach. John Wiley & Sons; 2011.

404 Beck A, Steer R, Brown G. Manual for Beck Depression Inventory-II. San Antonio, TX:
405 Psychological Corporation; 1996.

406 Benjamini Y, Hochberg Y. Controlling the False Discovery Rate - a Practical and
407 Powerful Approach. *Journal of the Royal Statistical Society Series B-Methodological*.
408 1995;57(1):289–300.

409 Bentler PM. Comparative Fit Indexes in Structural Models. *Psychological Bulletin*.
410 1990;107:238–46.

411 Browne M, Cudeck R. Alternative ways of assessing model fit. In: Bollen K, Long J,
412 editors. *Testing Structural Equation Models*. Newbury Park: Sage; 1993.

413 Cella M, Sharpe M, Chalder T. Measuring disability in patients with chronic fatigue
414 syndrome: Reliability and validity of the Work and Social Adjustment Scale. *Journal of*
415 *Psychosomatic Research*. 2011 Sep;71(3):124–8.

416 Chalder T, Berelowitz G, Pawlikowska T, Watts L, Wessely S, Wright D, et al.
 417 Development of a fatigue scale. *Journal of Psychosomatic Research*. 1993
 418 Feb;37(2):147–53.

419 Chalder T, Goldsmith K, White P, Sharpe M, Pickles A. Rehabilitative therapies for
 420 chronic fatigue syndrome: A secondary mediation analysis of the PACE trial. *The*
 421 *Lancet Psychiatry*. 2015 Feb;2(2):141–52.

422 Emsley R, Dunn G, White I. Mediation and moderation of treatment effects in
 423 randomised controlled trials of complex interventions. *Statistical Methods in Medical*
 424 *Research*. 2010 Jun;19(3):237–70.

425 Fokkema M, Smits N, Kelderman H, Cuijpers P. Response shifts in mental health
 426 interventions: An illustration of longitudinal measurement invariance. *Psychological*
 427 *Assessment*. 2013;25(2):520–31.

428 Goldsmith KA, MacKinnon DP, Chalder T, White PD, Sharpe M, Pickles A. Tutorial:
 429 The practical application of longitudinal structural equation mediation models in
 430 clinical trials. *Psychological Methods*. 2018 Jun a;23(2):191–207.

431 Goldsmith K, Chalder T, White P, Sharpe M, Pickles A. Measurement error, time lag,
 432 unmeasured confounding: Considerations for longitudinal estimation of the effect of a
 433 mediator in randomised clinical trials. *Statistical Methods in Medical Research*. 2018
 434 Jun b;27(6):1615–33.

435 Gunzler DD, Morris N. A Tutorial on Structural Equation Modeling for Analysis of
 436 Overlapping Symptoms in Co-occurring Conditions Using MPlus. *Statistics in*
 437 *medicine*. 2015 Oct;34(24):3246–80.

438 Hallquist MN, Wiley JF. MplusAutomation: An R Package for Facilitating Large-Scale
 439 Latent Variable Analyses in Mplus. Structural Equation Modeling: A Multidisciplinary
 440 Journal. 2018 Jan;0(0):1–8.

441 Hoertel N, López S, Peyre H, Wall MM, González-Pinto A, Limosin F, et al. Are
 442 Symptom Features of Depression During Pregnancy, the Postpartum Period and Outside
 443 the Peripartum Period Distinct? Results from a Nationally Representative Sample Using
 444 Item Response Theory (irt). Depression and Anxiety. 2015 Feb;32(2):129–40.

445 Hollon SD, Kendall PC. Cognitive self-statements in depression: Development of an
 446 automatic thoughts questionnaire. Cognitive Therapy and Research. 1980
 447 Dec;4(4):383–95.

448 Kaplan D. Structural Equation Modelling: Foundations and Extensions. Second.
 449 London: SAGE Publications; 2008.

450 Kaufman NK, Rohde P, Seeley JR, Clarke GN, Stice E. Potential Mediators of
 451 Cognitive-Behavioral Therapy for Adolescents With Comorbid Major Depression and
 452 Conduct Disorder. Journal of Consulting and Clinical Psychology. 2005;73(1):38–46.

453 MacKinnon D. Introduction to Statistical Mediation Analysis. New York: Taylor &
 454 Francis Group; 2008.

455 McHorney CA, Ware JE, Raczek AE. The MOS 36-Item Short-Form Health Survey
 456 (SF-36): II. Psychometric and clinical tests of validity in measuring physical and mental
 457 health constructs. Medical Care. 1993 Mar;31(3):247–63.

458 Mezuk B, Lohman M, Dumenci L, Lapane KL. Are Depression and Frailty Overlapping
459 Syndromes in Mid- and Late-life? A Latent Variables Analysis. *The American Journal of*
460 *Geriatric Psychiatry*; Washington. 2013 Jun;21(6):560–9.

461 Muthén L, Muthén B. *Mplus User's Guide*. Seventh. Los Angeles, CA: Muthén &
462 Muthén; 2015.

463 Parker JDA, Bagby RM, Taylor GJ. Alexithymia and depression: Distinct or
464 overlapping constructs? *Comprehensive Psychiatry*. 1991 Sep;32(5):387–94.

465 Preacher KJ. Advances in Mediation Analysis: A Survey and Synthesis of New
466 Developments. *Annual Review of Psychology*. 2015;66(1):825–52.

467 R Core Team. *R: A Language and Environment for Statistical Computing*. Vienna,
468 Austria: R Foundation for Statistical Computing; 2017.

469 Ryan EG, Vitoratou S, Goldsmith KA, Chalder T. Psychometric properties and factor
470 structure of a shortened version of the Cognitive Behavioural Responses Questionnaire
471 (CBRQ). *Psychosomatic Medicine*. 2017 Oct

472 Schaufeli WB, Bakker AB. Job demands, job resources, and their relationship with
473 burnout and engagement: A multi-sample study. *Journal of Organizational Behavior*.
474 2004 May;25(3):293–315.

475 Segerstrom SC, Tsao JCI, Alden LE, Craske MG. Worry and Rumination: Repetitive
476 Thought as a Concomitant and Predictor of Negative Mood. *Cognitive Therapy and*
477 *Research*. 2000 Dec;24(6):671–88.

- 478 Skrandal A, Rabe-Hesketh S. Generalized latent variable modeling: Multilevel,
479 longitudinal, and structural equation models. Chapman and Hall/CRC; 2004.
- 480 VanderWeele TJ, Valeri L, Ogburn EL. The role of measurement error and
481 misclassification in mediation analysis. Epidemiology (Cambridge, Mass.). 2012
482 Jul;23(4):561–4.
- 483 VanderWeele TJ, Vansteelandt S, Robins JM. Effect decomposition in the presence of an
484 exposure-induced mediator-outcome confounder. Epidemiology (Cambridge, Mass.).
485 2014 Mar;25(2):300–6.
- 486 White P, Goldsmith K, Johnson A, Potts L, Walwyn R, DeCesare J, et al. Comparison of
487 adaptive pacing therapy, cognitive behaviour therapy, graded exercise therapy, and
488 specialist medical care for chronic fatigue syndrome (PACE): A randomised trial. The
489 Lancet. 2011 Mar;377(9768):823–36.
- 490 Woods CM. Empirical Selection of Anchors for Tests of Differential Item Functioning.
491 Applied Psychological Measurement. 2009 Jan;33(1):42–57.
- 492 **List of abbreviations**
- 493 BDI = Beck Depression Inventory; CBT = Cognitive behavioural therapy; CFA =
494 Confirmatory Factor Model; CFI = Comparative Fit Index; CFQ = Chalder Fatigue
495 Questionnaire; DIF = Differential Item Functioning; EFA = Exploratory Factor
496 Analysis HAM-D = Hamilton Rating Scale for Depression; MIMIC = Multiple
497 Indicators Multiple Causes PF = Physical function; RMSEA = Root Mean Square Error
498 of Approximation; RSS = Rumination on Sadness Scale; SF-36 = Medical Outcomes
499 Study Short Form Health Survey; WLSMV = Robust Weighted Least Squares.

Declarations

Ethics approval and consent to participate

This study uses data that have previously been published (Chalder et al. 2015; Ryan et al. 2017). The original PACE study was approved by the West Midlands Multicentre Research Ethics Committee (MREC 02/7/89).

Consent for publication

Not applicable.

Availability of data and materials

The dataset analysed in the current study are available in the Vivli repository, <https://search.vivli.org>. Some measures analysed are not publicly available because the authors are still analysing/publishing papers with them.

Competing interests

TC has received royalties from Sheldon Press and Constable and Robinson. Funding for the PACE trial was provided by the Medical Research Council, Department for Health for England, The Scottish Chief Scientist Office, and the Department for Work and Pensions.

The authors declare no other conflicts of interest.

Funding

This paper represents independent research part funded by the National Institute for Health Research (NIHR) Biomedical Research Centre (BRC) at South London and Maudsley NHS Foundation Trust and King's College London and Applied Research Collaboration South London (NIHR ARC South London) at King's College Hospital

522 NHS Foundation Trust. The views expressed are those of the author(s) and not
523 necessarily those of the NHS, the NIHR or the Department of Health and Social Care.
524 EC and SV are fully funded by the NIHR BRC; TC is part-funded by the NIHR BRC;
525 KG is part-funded by the BRC and ARC.

526 *Authors' contributions*

527 KG and TC came up with the concept for the study; all authors contributed to the design
528 of the study. EC carried out the data analysis. All authors contributed to the
529 interpretation of results. EC wrote the first draft of the manuscript. All authors
530 contributed to writing the manuscript. All authors gave critical revision of the article
531 and approved of the final version.

532 *Acknowledgements*

533 The authors acknowledge the other investigators on the PACE trial, Peter D. White and
534 Michael Sharpe, and the help of the PACE Trial Management Group.

Supplementary Table 1

Factor structure and question wordings for PACE mediators and outcomes.

	Factor	Items	Question wording
Outcomes	1. Chalder Fatigue Questionnaire (CFQ) ^a	CFQ_1	Do you have problems with tiredness?
		CFQ_2	Do you need to rest more?
		CFQ_3	Do you feel sleepy or drowsy?
		CFQ_4	Do you have problems starting things?
		CFQ_5	Do you start things without difficulty but get weak as you go on?
		CFQ_6	Are you lacking in energy?
		CFQ_7	Do you have less strength in your muscles?
		CFQ_8	Do you feel weak?
		CFQ_9	Do you have difficulty concentrating?
		CFQ_10	Do you have problems thinking clearly?
		CFQ_11	Do you make slips of the tongue when speaking?
	2. SF-36 Physical Functioning subscale (PF) ^b	PF_1	Vigorous activities, such as running, lifting heavy objects, vacuum cleaner, bowling, or playing golf
		PF_2	Moderate activities, such as moving a table, pushing a vacuum cleaner, bowling, or playing golf
		PF_3	Lifting or carrying groceries
		PF_4	Climbing several flights of stairs
		PF_5	Climbing one flight of stairs
		PF_6	Bending, kneeling, or stooping
		PF_7	Walking more than a mile
		PF_8	Walking several blocks
		PF_9	Walking one block
		PF_10	Bathing or dressing yourself
Mediators	1. Fear Avoidance ^d	SIQ_1	I am afraid that I will make my symptoms worse if I exercise.

	SIQ_2R	My symptoms would be relieved if I were to exercise (reversed).
	SIQ_3	Avoiding unnecessary activities is the safest thing I can do to prevent my symptoms from worsening.
	SIQ_7	Physical activity makes my symptoms worse.
	SIQ_8	Doing less helps symptoms.
	SIQ_11	I should avoid exercise when I have symptoms.
2. Fear Avoidance (short version) ^d	SIQ_1	(as above)
	SIQ_2R	(as above)
	SIQ_7	(as above)
3. Catastrophising ^d	SIQ_12	I worry that I may become permanently bedridden because of my problems.
	SIQ_13	I think that if my symptoms get too severe they may never decrease.
	SIQ_15	My illness is awful and I feel that it overwhelms me.
	SIQ_17	I will never feel right again.
4. Damage ^d	SIQ_4	The severity of my symptoms must mean there is something serious going on in my body.
	SIQ_5R	Even though I experience symptoms, I don't think they are actually harming me.
	SIQ_6	When I experience symptoms, my body is telling me that there is something seriously wrong.
	SIQ_9	Symptoms are a signal that I am damaging myself.
	SIQ_10	I am afraid I will have more symptoms if I am not careful.
5. Damage (short version) ^d	SIQ_4	(as above)
	SIQ_9	(as above)
	SIQ_10	(as above)
6. Symptom focusing ^d	SIQ_18	When I experience symptoms, I think about them constantly.
	SIQ_19	I worry when I am experiencing symptoms.
	SIQ_20	When I am experiencing symptoms it is difficult for me to think of anything else.
	SIQ_21	I think a great deal about my symptoms.
	SIQ_22	My symptoms are always at the back of my mind.
	SIQ_23	I spend a lot of time thinking about my illness.
7. Symptom focusing (short version) ^d	SIQ_21	(as above)

	version) ^d	SIQ_22	(as above)
		SIQ_23	(as above)
8.	Embarrassment avoidance ^d	SIQ_24	I am embarrassed about my symptoms.
		SIQ_25	I worry that people will think badly of me because of my symptoms.
		SIQ_26	The embarrassing nature of my symptoms prevents me from doing things.
		SIQ_27	I avoid social situations because I am scared my symptoms will get out of control.
		SIQ_28	I am ashamed of my symptoms.
		SIQ_29	My symptoms have the potential to make me look foolish in front of other people.
9.	Embarrassment avoidance (short version) ^d	SIQ_24	(as above)
		SIQ_25	(as above)
		SIQ_26	(as above)
10	All-or-nothing behaviour ^d	SIQ_34	I tend to overdo things when I feel energetic.
		SIQ_35	I find myself rushing to get things done before I crash
		SIQ_36	I tend to overdo things and then rest up for a while
		SIQ_37	I tend to do a lot on a good day and rest on a bad day
		SIQ_41	I'm a bit all or nothing when it comes to doing things
11	All-or-nothing behaviour (short version) ^d	SIQ_34	(as above)
		SIQ_35	(as above)
		SIQ_36	(as above)
12	Avoidance behaviour ^d	SIQ_30	I stay in bed to control my symptoms.
		SIQ_31	When I experience symptoms, I rest.
		SIQ_32	I tend to avoid activities that make my symptoms worse.
		SIQ_33	I tend to nap during the day to control my symptoms.
		SIQ_38	I sleep when I'm tired in order to control my symptoms.
		SIQ_39	I avoid making social arrangements in case I'm not up to it.
		SIQ_40	I avoid exerting myself in order to control my symptoms.
		SIQ_42	I avoid stressful situations.

13	Avoidance behaviour	SIQ_30	(as above)
.	(short version) ^d	SIQ_33	(as above)
		SIQ_38	(as above)

Item scoring

- a Each item has 4 response options ranging from ‘less than usual’ (score 0), ‘no more than usual’ (score 1), ‘more than usual’ (score 2), and ‘much more than usual’ (score 3).
- b There are three responses to each of the 10 items; ‘limited a lot’ (score 0), ‘limited a little’ (score 5) and ‘not limited at all’ (score 10).
- c Scores range from 0 (‘not at all impaired’) to 8 (‘very severely impaired’).
- d Scores on a 5-point Likert scale ranging from ‘strongly disagree’ (score 1) to ‘strongly agree’ (score 5).

Notes

CFQ = Chalder Fatigue Questionnaire; PF = Physical function subscale of the SF-36.

Supplementary Table 2

All cross-loadings added during the iterative procedure

Pairing		Number of factor loadings		Total cross-loadings added by iterative procedure	Type of cross-loadings		
Mediator	Outcome	Before adding cross-loadings	After adding cross-loadings		No cross- loading	Shared	Strong cross- loading
Fear avoidance (short)	CFQ	14	22	8	4	0	4
Fear avoidance	CFQ	17	24	7	5	0	2
Avoidance/resting behaviour (short)	CFQ	14	20	6	2	1	3
Damage (short)	CFQ	14	20	6	2	2	2
Symptom focusing	CFQ	17	23	6	3	0	3
Damage	PF	15	21	6	6	0	0
Damage (short)	PF	13	18	5	5	0	0
Fear avoidance (short)	PF	13	17	4	3	0	1
Fear avoidance	PF	16	20	4	4	0	0
Symptom focusing (short)	CFQ	14	17	3	3	0	0
Embarrassment avoidance	CFQ	17	20	3	3	0	0
All-or-nothing behaviour (short)	PF	13	15	2	2	0	0
Avoidance/resting behaviour (short)	PF	13	15	2	2	0	0
Symptom focusing (short)	PF	13	15	2	2	0	0
All-or-nothing behaviour	PF	15	17	2	2	0	0
Catastrophising	CFQ	15	17	2	2	0	0
Embarrassment avoidance	PF	16	18	2	2	0	0
Catastrophising	PF	14	15	1	1	0	0
Embarrassment avoidance (short)	PF	13	14	1	1	0	0
Symptom focusing	PF	16	17	1	1	0	0
Avoidance/resting behaviour	CFQ	19	20	1	1	0	0

Avoidance/resting behaviour	PF	18	19	1	1	0	0
All-or-nothing behaviour	CFQ	16	16	0	0	0	0
All-or-nothing behaviour (short)	CFQ	14	14	0	0	0	0
Damage	CFQ	16	16	0	0	0	0
Embarrassment avoidance (short)	CFQ	14	14	0	0	0	0

CFQ = Chalder Fatigue Questionnaire; PF = Physical function subscale of the SF-36; WSAS = Work and Social Adjustment Scale.