1 The E484K mutation in the SARS-CoV-2 spike protein reduces but does not abolish neutralizing
2 activity of human convalescent and post-vaccination sera.

3 Sonia Jangra, Ph.D.^{1,2,#}, Chengjin Ye, Ph.D.^{3,#}, Raveen Rathnasinghe, B.Sc.^{1,2,4}, Daniel Stadlbauer,
4 Ph.D.¹, PVI study group^{1,2}, Florian Krammer, Ph.D.¹, Viviana Simon, M.D., Ph.D.^{1,2,5}, Luis Martinez-
5 Sobrido, Ph.D.³, Adolfo García-Sastre, Ph.D.^{1,2,5,6,*}, Michael Schotsaert, Ph.D.^{1,2,*}

6 # Contributed equally

7 *Corresponding authors: Adolfo.Garcia-Sastre@mssm.edu, Michael.Schotsaert@mssm.edu

8 ¹Department of Microbiology, Icahn School of Medicine at Mount Sinai New York, NY, USA

9 ²Global Health and Emerging Pathogens Institute, Icahn School of Medicine at Mount Sinai New
10 York, NY, USA

11 ³Texas Biomedical Research Institute, San Antonio, TX, USA

12 ⁴Graduate School of Biomedical Sciences, Icahn School of Medicine at Mount Sinai, New York,
13 NY, USA

14 ⁵Department of Medicine, Division of Infectious Diseases, Icahn School of Medicine at Mount
15 Sinai New York, NY, USA

16 ⁶The Tisch Cancer Institute, Icahn School of Medicine at Mount Sinai New York, NY, USA

17

Abstract

One year in the coronavirus disease 2019 (COVID-19) pandemic, the first vaccines are being rolled out under emergency use authorizations. It is of great concern that newly emerging variants of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) can escape antibody-mediated protection induced by previous infection or vaccination through mutations in the spike protein. The glutamate (E) to Lysine (K) substitution at position 484 (E484K) in the receptor binding domain (RBD) of the spike protein is present in the rapidly spreading variants of concern belonging to the B.1.351 and P.1 lineages. We performed *in vitro* microneutralization assays with both the USA-WA1/2020 virus and a recombinant (r)SARS-CoV-2 virus that is identical to USA-WA1/2020 except for the E484K mutation introduced in the spike RBD. We selected 34 sera from study participants based on their SARS-CoV-2 spike ELISA antibody titer (negative [N=4] versus weak [N=8], moderate [N=11] or strong positive [N=11]). In addition, we included sera from five individuals who received two doses of the Pfizer SARS-CoV-2 vaccine BNT162b2. Serum neutralization efficiency was lower against the E484K rSARS-CoV-2 (vaccination samples: 3.4 fold; convalescent low IgG: 2.4 fold, moderate IgG: 4.2 fold and high IgG: 2.6 fold) compared to USA-WA1/2020. For some of the convalescent donor sera with low or moderate IgG against the SARS-CoV-2 spike, the drop in neutralization efficiency resulted in neutralization ID₅₀ values similar to negative control samples, with low or even absence of neutralization of the E484K rSARS-CoV-2. However, human sera with high neutralization titers against the USA-WA1/2020 strain were still able to neutralize the E484K rSARS-CoV-2. Therefore, it is important to aim for the highest titers possible induced by vaccination to enhance protection against newly emerging SARS-CoV-2 variants. Two vaccine doses may be needed for induction of high antibody titers against SARS-CoV-2. Postponing the second vaccination is suggested by some public health authorities in order to provide more individuals with a primer vaccination. Our data suggests that this may leave vaccinees less protected against newly emerging variants.

Introduction

One year in the coronavirus disease 2019 (COVID-19) pandemic, the first vaccines are being rolled out under emergency use authorizations. Recently, rapidly spreading variants of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), the virus that causes COVID-19, have been reported. It is of great concern that these newly emerging variants can escape neutralizing antibodies induced by previous infection and/or vaccination through mutations in the spike (S) protein, including the receptor binding domain (RBD), a target for neutralizing antibodies. We and others have previously reported that the asparagine (N) to tyrosine (Y) substitution at position 501 (N501Y), present in variants of concern belonging to the B.1.1.7, B.1.351 and P.1 lineages, does not seem to affect *in vitro* neutralization of SARS-CoV-2 viruses by human sera from convalescent or vaccinated human donors. However, there remains concern about additional substitutions like E484K present in B.1.351 and P.1 lineages allowing escape from neutralizing antibodies (1–4), thereby potentially rendering vaccine-induced immunity less protective.

In order to investigate the impact of the E484K mutation in the neutralizing activity of SARS-CoV-2 specific antisera, we performed *in vitro* microneutralization assays with both the USA-WA1/2020 virus and a recombinant (r)SARS-CoV-2 virus that is identical to USA-WA1/2020 except for the E484K mutation introduced in the spike RBD.

The E484K mutant rSARS-CoV-2 was generated using previously described reverse genetics based on the use of a bacterial artificial chromosome (BAC) (5–7). The USA-WA1/2020 reflects SARS-CoV-2 strains that circulated in the early phase of the COVID-19 pandemic. A total of 34 sera were selected from study participants based on their SARS-CoV-2 S enzyme linked immunosorbent

assay (ELISA) antibody titer (negative [N=4] versus weak [N=8], moderate [N=11] or strong positive [N=11]). In addition, we included sera from five individuals who received two doses of the Pfizer SARS-CoV-2 vaccine BNT162b2 (V1-V5). Demographics and available metadata for each participant is summarized in Supplementary Table 1. We performed all experiments in a blinded manner. The same sera have been tested for neutralization studies with a N501Y SARS-CoV-2 variant in our recent report (8).

Results

Sera from vaccinated donors gave high neutralization titers, similar to convalescent samples with the highest neutralization titers. Serum neutralization efficiency was lower against the E484K rSARS-CoV-2 (vaccination samples: 3.4 fold; convalescent low IgG: 2.4 fold, moderate IgG: 4.2 fold and high IgG: 2.6 fold based on geometric means) which was significantly different for the convalescent sera (see Figure 1), suggesting that the single E484K mutation in the RBD affects binding by serum polyclonal neutralizing antibodies from both convalescent and vaccinated donors. In the case of convalescent donor sera with low or moderate IgG against SARS-CoV-2 S protein, the drop in neutralization efficiency could result in neutralization ID₅₀ values similar to negative control samples, resulting in low or even absence of neutralization of the E484K recombinant virus by those sera.

Figure 1. Human convalescent and post-vaccination sera neutralize E484K rSARS-CoV-2 less efficient than USA-WA1/2020 in an *in vitro* microneutralization assay. Convalescent sera are subdivided in low, moderate and high IgG classes based on anti-spike ELISA titers. Two-sided Mann Whitney-U tests were performed to calculate statistical differences.

Conclusions

These data indicate that the E484K mutation present in circulating SARS-CoV-2 strains that belong to the B.1.351 and P.1 lineages reduces the neutralizing activity of human polyclonal sera induced in convalescent (infected with previous strains) and vaccinated individuals. The significant impact of a single point mutation in the neutralizing activity of polyclonal sera highlights the need for the rapid characterization of SARS-CoV-2 variants. However, human sera with high neutralization titers against the USA-WA1/2020 strain were still able to neutralize the

E484K rSARS-CoV-2. Therefore, it is important to aim for the highest titers possible induced by vaccination, as this should enhance the chances for protection even in the case of antigenic drift of circulating SARS-CoV-2 strains. Currently deployed SARS-CoV-2 vaccines are recommended as a prime-boost regimen. Because of vaccine shortage and relatively strong seroconversion being observed after a single dose, some public health authorities recommended to postpone the second booster vaccination in order to be able to provide more individuals with a first primer vaccination. This will result in lower neutralizing antibody titers. Our data show that this may be problematic in the context of newly emerging SARS-CoV-2 variants, as it may leave some vaccinees unprotected. It is currently unknown which neutralization titer correlates with (full) protection, and to what extent immune mechanisms beyond direct virus neutralization contribute to protection, especially for specific target groups with comorbidities that are currently being prioritized for vaccination.

Viruses that belong to the B.1.351 and P.1 lineages have originally been described in the Republic of South Africa and Brazil, but are now reported on multiple continents already. Therefore, while it is premature to update vaccines based on these lineages, it is important that the worldwide vaccination effort will aim at fully vaccinating as many people as possible using vaccination strategies that result in induction of high neutralizing antibody titers.

References

1. Weisblum Y, Schmidt F, Zhang F, DaSilva J, Poston D, Lorenzi JC, et al. Escape from neutralizing antibodies by SARS-CoV-2 spike protein variants. eLife [Internet]. [cited 2020 Dec 22];9. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7723407/>
2. Greaney AJ, Loes AN, Crawford KHD, Starr TN, Malone KD, Chu HY, et al. Comprehensive mapping of mutations to the SARS-CoV-2 receptor-binding domain that affect recognition by polyclonal human serum antibodies. bioRxiv. 2021 Jan 4;2020.12.31.425021.
3. Wang Z, Schmidt F, Weisblum Y, Muecksch F, Barnes CO, Finkin S, et al. mRNA vaccine-elicited antibodies to SARS-CoV-2 and circulating variants. bioRxiv. 2021 Jan 19;2021.01.15.426911.
4. Wu K, Werner AP, Moliva JJ, Koch M, Choi A, Stewart-Jones GBE, et al. mRNA-1273 vaccine induces neutralizing antibodies against spike mutants from global SARS-CoV-2 variants. bioRxiv. 2021 Jan 25;2021.01.25.427948.
5. Chiem K, Morales Vasquez D, Park J-G, Platt RN, Anderson T, Walter MR, et al. Generation and Characterization of recombinant SARS-CoV-2 expressing reporter genes. J Virol. 2021 Jan 11;
6. Chiem K, Ye C, Martinez-Sobrido L. Generation of Recombinant SARS-CoV-2 Using a Bacterial Artificial Chromosome. Curr Protoc Microbiol [Internet]. 2020 Dec [cited 2021 Jan 25];59(1). Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7646048/>

- 135 7. Ye C, Chiem K, Park J-G, Oladunni F, Platt RN, Anderson T, et al. Rescue of SARS-CoV-2 from
136 a Single Bacterial Artificial Chromosome. mBio [Internet]. 2020 Sep 25 [cited 2021 Jan
137 25];11(5). Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7520601/>
 - 138 8. Rathnasinghe R, Jangra S, Cupic A, Martínez-Romero C, Mulder LCF, Kehrer T, et al. The
139 N501Y mutation in SARS-CoV-2 spike leads to morbidity in obese and aged mice and is
140 neutralized by convalescent and post-vaccination human sera. medRxiv. 2021 Jan
141 20;2021.01.19.21249592.
- 142
- 143

Acknowledgements

PVI study group (in alphabetical order): H. Alshammary, A. Amoako, M. Awawda, K. Beach, C. M. Bermúdez-González, R. Chernet, L. Eaker, E. Ferreri, D. Floda, C. Gleason, Dr. G. Kleiner, D. Jurczynszak, J. Matthews, W. Mendez, Dr. LCF Mulder, K. Russo, A. Salimbangon, Dr. M. Saksena, A. Shin, L. Sominsky and K. Srivastava.

We thank all the study participants for their continued support of COVID19 research. This research was partly funded by CRIP (Center for Research for Influenza Pathogenesis), a NIAID supported Center of Excellence for Influenza Research and Surveillance (CEIRS, contract # HHSN272201400008C); by NCI grant U54CA260560; by the generous support of the JPB Foundation and the Open Philanthropy Project (research grant 2020-215611 (5384); and by anonymous donors to A.G-S. Work on SARS-CoV-2 in the Krammer and Simon laboratories was funded by the Collaborative Influenza Vaccine Innovation Centers (CIVIC) contract 75N93019C00051. Research in Martinez-Sobrido laboratory was partially funded by the New York Influenza Center of Excellence (NYICE), a member of the National Institute of Allergy and Infectious Diseases (NIAID), National Institutes of Health (NIH), Department of Health and Human Services, Centers of Excellence for Influenza Research and Surveillance (CEIRS) contract No. HHSN272201400005C (NYICE).

Conflicts of interest

The García-Sastre Laboratory has received research support from Pfizer, Senhwa Biosciences and 7Hills Pharma. Adolfo García-Sastre has consulting agreements for the following companies involving cash and/or stock: Vivaldi Biosciences, Contrafect, 7Hills Pharma, Avimex, Vaxalto, Accurius and Esperovax. The Icahn School of Medicine at Mount Sinai has filed patent

166 applications relating to SARS-CoV-2 serological assays and NDV-based SARS-CoV-2 vaccines
167 which list Florian Krammer as co-inventor. Daniel Stadlbauer and Viviana Simon are also listed
168 on the serological assay patent application as co-inventors. Florian Krammer has consulted for
169 Merck and Pfizer (before 2020), and is currently consulting for Seqirus and Avimex. The
170 Krammer laboratory is also collaborating with Pfizer on animal models of SARS-CoV-2.

Supplementary Table 1: Description of serum samples obtained from human subjects

SERUM					
Seropositive, vaccine	Spike IgG response	Sex	Age group (yrs)	Days post 1 vaccine dose (Pfizer)	
V1	Strong positive	F	>60	68	
V2	Strong positive	M	30-40	47	
V3	Strong positive	F	50-60	47	
V4	Strong positive	F	40-50	49	
V5	Strong positive	F	30-40	48	
Seropositive, infection	Spike IgG response	Sex	Age group (yrs)	Days post onset of symptoms	
P1	Weak positive	M	20-29	260	
P2	Weak positive	M	50-59	NA	
P3	Weak positive	F	30-39	111	
P4	Weak positive	F	30-39	221	
P5	Weak positive	F	30-39	254	
P6	Weak positive	F	20-29	247	
P7	Weak positive	M	30-39	220	
P8	Weak positive	F	20-29	Asymptomatic	
P9	Moderate positive	M	30-39	NA	
P10	Moderate positive	F	30-39	197	
P11	Moderate positive	F	50-59	Asymptomatic	
P12	Moderate positive	F	30-39	Asymptomatic	
P13	Moderate positive	M	30-39	234	
P14	Moderate positive	F	20-29	273	
P15	Moderate positive	M	30-39	Asymptomatic	
P16	Moderate positive	F	20-29	258	
P17	Moderate positive	F	20-29	246	
P18	Moderate positive	M	20-29	Asymptomatic	
P19	Moderate positive	F	50-59	204	
P20	Strong positive	F	50-59	NA	
P21	Strong positive	F	30-39	245	
P22	Strong positive	M	NA	170	
P23	Strong positive	F	>60	Asymptomatic	
P24	Strong positive	F	40-49	NA	
P25	Strong positive	F	50-59	191	
P26	Strong positive	F	30-39	NA	
P27	Strong positive	F	50-59	113	
P28	Strong positive	M	>60	Asymptomatic	
P29	Strong positive	M	18-19	218	

	P30	Strong positive	M	50-59	219
	Seronegative, post pandemic	Spike IgG response	Sex	Age group (yrs)	Days from last negative serology
	N1	Negative	F	40-50	23
	N2	Negative	F	20-29	24
	N3	Negative	F	20-29	23
	N4	Negative	F	30-35	22

Supplementary Methods section:

50% tissue culture infective dose (TCID₅₀) calculation and *in vitro* microneutralization assay:

To estimate the neutralizing efficiency of human sera, *in vitro* microneutralization assays were performed. Human sera were inactivated at 56°C for 30 min. Serum samples were serially diluted 3-fold starting from 1:30 dilution in Vero-E6-infection medium (DMEM+ 2% FBS+ 1% non-essential amino acids). The samples were incubated with 450 tissue culture infective dose 50 (TCID₅₀) of either USA-WA1/2020 or rSARS-CoV-2 E484K for 1 hour in an incubator at 37°C, 5% CO₂ followed by incubation with pre-seeded Vero-E6 at 37°C for 48 hours. The plates were fixed in 4% formaldehyde at 4°C overnight. For TCID₅₀ calculation, the virus stock was serially diluted 10-fold starting with 1:10 dilution and incubated on Vero-E6 cells for 48 hours followed by fixation in 4% Formaldehyde. The cells were washed with 1xPBS and permeabilized with 0.1% Triton X-100 in 1XPBS. The cells were washed again and blocked in 5% non-fat milk in 1xPBS+ 0.1% Tween-20 for 1 hour at room temperature. After blocking, the cells were incubated with anti-SARS-CoV-2 NP and anti-spike monoclonal antibodies, mixed in 1:1 ratio, for 1.5 hours at room temperature. The cells were washed in 1xPBS and incubated with 1:5000 diluted HRP-conjugated anti-mouse IgG secondary antibody for 1 hour at RT followed by a brief PBS wash. Finally, 100µl tetramethyl benzidine (TMB) substrate was added and incubated at RT until blue color appeared, and the reaction was terminated with 50µl 1M H₂SO₄. Absorbance was recorded at 450nm and 650nm and percentage reduction in infection was calculated as compared to negative control.

197 **Serum samples from human subjects:**

198 The study protocols for the collection of clinical specimens from individuals with and without
199 SARS-CoV-2 infection by the Personalized Virology Initiative were reviewed and approved by the
200 Mount Sinai Hospital Institutional Review Board (IRB-16-00791; IRB-20-03374). All participants
201 provided informed consent prior to collection of specimen and clinical information. All specimens
202 were coded prior to processing.

203

204