

1 *Table 1 Summary of results from the questionnaire survey*

2

		HCW, n=27 (%)	Pharm, n=27 (%)
<b>I</b>	<b>Demographics</b>		
	<i>Clients</i>		
<b>1</b>	Children under the age of 12		
	<10%	37	-
	Between 10-24%	22	-
	Between 25-50%	26	-
	>50%	15	-
<b>2</b>	Adults over the age of 60		
	<10%	44	-
	Between 10-24%	19	-
	Between 25-50%	33	-
	>50%	4	-
<b>3</b>	Age distribution, average age in years		
	Under 18	-	4
	18-29	-	52
	30-49	-	37
	50-75	-	7
<b>4</b>	Gender		
	Female	44	26
	Male	33	44
	Prefer not to say	22	30

		HCW, n=27 (%)	Pharm, n=27 (%)
<b>5</b>	IDPs, as proportion of clients		
	<10%	33	-
	Between 10-24%	48	-
	Between 25-50%	7	-
	>50%	11	-
	<i>Participants</i>		
<b>6</b>	Profession		
	Health Care Workers		
	Physicians	96	-
	Others (not stated)	4	-
	Pharmacists		
	Certified pharmacist	-	89
	Pharmacy worker	-	11
<b>7</b>	Training (pharmacists)		
	University	-	67
	Pharmacy	-	30
	Other	-	4
<b>8</b>	Years of practice		
	<1	8	-
	<5	44	22
	<10	30	48
	<15	19	15

		<b>HCW, n=27 (%)</b>	<b>Pharm, n=27 (%)</b>
<b>II</b>	<b>Perception of AMR</b>		
<b>9</b>	AMR is a problem in Yemen		
	Yes	96	96
	Unsure	4	4
	If yes, in what way?		
	Understudied and/or underreported	56 (15/27)	70 (19/27)
	High prevalence of AMR	19 (5/27)	33 (9/27)
	Both understudied/underreported and high prevalence	19 (5/27)	11 (3/27)
<b>10</b>	IDPs have a greater prevalence of AMR		
	No	52 (14/27)	30(8/27)
	Yes	48 (13/27)	63 (17/27)
<b>11</b>	Treatment failure incidences per month		
	<5	22	60
	<10	44	30
		<b>HCW, n=27 (%)</b>	<b>Pharm, n=27 (%)</b>
	<25	26	7
	<50	4	-
	>50	4	-
<b>12</b>	Source of information on AMR		
	Customers	-	37 (10/27)
	Friends	-	4 (1/27)
	Social media	-	4 (1/27)
	Training	-	44 (12/27)

		HCW, n=27 (%)	Pharm, n=27 (%)
<b>13</b>	<b>Tackling AMR</b>		
	Enforcement of prescription laws	37	15
	Increase awareness among patients	22	22
	Increased testing for susceptibility	11	4
	Increased training of pharmacists and physicians	30	59
	<b>Antimicrobial stewardship</b>		
<b>14</b>	Prescribing guide (Yes)	30	-
	If yes, what is the guide? (Local or Foreign, including online)	Foreign	-
<b>15</b>	Facility has a medicines formulary (Yes)	70	-
<b>16</b>	Dispensing guide		
	Yemen EML		0
	Other texts - including WHO, BNF, PNF		30 (8/27)
	Empirical (based on experience, or studies)		22 (6/27)
<b>17</b>	Drug and Therapeutic Committee (Yes)	19	-
<b>18</b>	Possibility of performing antimicrobial sensitivity tests (Yes)	67 (18/27)	-
	If yes (on-site)	56 (10/18)	-
<b>19</b>	AST performed before antibiotic selection (No)	81	-
	a. If not, reason for not performing:		
	Cost	76 (16/21)	-
	Availability	67 (14/21)	-
	Time	57 (12/21)	-
	b. If yes, how often is it used as a proportion of all cases		-
	<10	50 (6/12)	-
	Between 10-24	17 (2/12)	-
	Between 25-49	17 (2/12)	-

		<b>HCW, n=27 (%)</b>	<b>Pharm, n=27 (%)</b>
	75-99	17 (2/12)	-
<b>III</b>	<b>Practices</b>		
<b>20</b>	Pressure to prescribe broad-spectrum antibiotics (Yes)	74 (20/27)	
<b>21</b>	Reasons for choice of empirical treatment or antibiotic selection		
	Availability	52 (14/27)	37 (10/27)
	Broad-spectrum	52 (14/27)	44 (12/27)
	Convenience	11 (3/27)	7 (2/27)
	Cost	59 (16/27)	41 (11/27)
	Symptoms of patient	56 (15/27)	33 (9/27)
	Antibiotic susceptibility testing information	15 (4/27)	14 (4/27)
	Identification of bacterial infection	41 (11/27)	59 (16/27)
<b>22</b>	Antibiotics most commonly prescribed or dispensed		
	a. Spectrum		
	Broad-spectrum	-	78
	Narrow-spectrum	-	11
	Both equally	-	11
	b. Individual agents/ groups		
	Penicillin		
	Amoxicillin (and amoxicillin/clavulanic acid)	63 (17/27)	82 (22/27)
	Ampicillin	26 (7/27)	16 (5/27)
	Penicillin	22 (6/27)	30 (8/27)
	Cephalosporins		
	Ceftriaxone	22 (6/27)	22 (6/27)
	Cephalosporins (others, apart from ceftriaxone, including cefotaxime)	11 (3/27)	4 (1/27)
	Fluoroquinolones		

		<b>HCW, n=27 (%)</b>	<b>Pharm, n=27 (%)</b>
	Ciprofloxacin	52 (14/27)	74 (20/27)
	Quinolones (others apart from ciprofloxacin e.g., moxifloxacin, levofloxacin)	11 (3/27)	-
	Macrolides		
	Erythromycin	19 (5/27)	26 (7/27)
	Azithromycin	-	15 (4/27)
	Tetracyclines		
	Tetracycline	-	15 (4/27)
	Others, including carbapenems	11 (3/27)	7 (2/27)
<b>23</b>	Sale without prescriptions (Yes)	-	67
<b>24</b>	Counter-prescribing (Yes)	-	63
<b>25</b>	Conditions for which antibiotics demanded		
	Fever with other symptoms	-	11 (3/27)
	Inflammation, including pharyngitis, tonsillitis, otitis media	-	37 (10/27)
	Infection - UTI, RTI, including both upper and lower	-	33 (9/27)
<b>26</b>	Most important consideration for dispensing/counter- prescribing		
	Cost	-	15 (4/27)
	Disease	-	62 (16/27)
	Price patient can afford	-	22 (6/27)
<b>27</b>	Daily antibiotic prescribing incidence		
	<5	7	26
	<10	22	22
	<25	48	37
	<50	19	7
	>50	4	7
<b>28</b>	Daily antibiotic prescriptions received		

	<b>HCW, n=27 (%)</b>	<b>Pharm, n=27 (%)</b>
<5	-	15
<10	-	41
<25	-	41
≥50	-	8

- 3
- 4
- 5
- 6
- 7
- 8
- 9
- 10
- 11
- 12
- 13
- 14
- 15
- 16
- 17
- 18
- 19
- 20
- 21
- 22
- 23
- 24
- 25

26 **Appendix 1 Questionnaires**

27 A. Healthcare workers

28

29 1-Among the patients you see, approximately what percentage are children under 12?

- 30  less than 10%
- 31  between 10-24%
- 32  between 25-50%
- 33  Over 50%

34

35 2-Among the patients you see, approximately what percentage are over 60?

- 36  less than 10%
- 37  between 10-24%
- 38  between 25-50%
- 39  Over 50%

40

41 3-Do you see more male or female patients?

- 42  Male
- 43  Female
- 44  Prefer not to say

45

46 4-Among the patients you see, approximately what percentage of patients that you see are IDPs?

- 47  less than 10%
- 48  between 10-24%
- 49  between 25-50%
- 50  Over 50%

51

52 5-Do you think antibiotic resistance is a problem in Yemen?

- 53  Yes
- 54  No



55       ○ Unsure

56

57   5A. If yes, in what way? (Check all that apply)

58       ○ Understudied problem

59       ○ Underreported problem

60       ○ High levels of AMR are prevalent

61

62   6-Do you think IDPs have a higher risk of antibiotic resistance than population that has not been  
63 displaced?

64       ○ Yes

65       ○ No

66       ○

67   7-In your practice, is there a guide to prescribing antibiotics?

68       ○ Yes

69       ○ No

70   7A. If yes above, please describe the guide to prescribing antibiotics \_\_\_\_\_

71

72   8-Does the facility where you work have a Drug and Therapeutic Committee responsible for rational use  
73 of medicines?

74       ○ Yes

75       ○ No

76

77   9-Do you use Medicine formulary for your work in the health facility?

78       ○ Yes

79       ○ No

80

81   10-Is it possible to perform laboratory antibiotic susceptibility testing on-site or off-site?

82       ○ Yes

83       ○ No

84

85 10A. If yes, is it on-site or off-site?

86  On-site

87  Off-site

88

89 11-How often are antibiotics prescribed a day?

90  less than 5 incidences

91  less than 10 incidences

92  less than 25 incidences

93  less than 50 incidences

94  50 or more incidences

95

96 12-How often do you see antibiotic treatment failure within a month?

97  less than 5 incidences

98  less than 10 incidences

99  less than 25 incidences

100  less than 50 incidences

101  50 or more incidences

102

103 13-Are antibiotic susceptibility tests of bacteria performed before antibiotic selection?

104  Yes

105  No

106

107 13A. If not, why? (Check all that apply)

108  Cost

109  Availability

110  Time

111

112 13B. If Yes, how often is it used?

- 113       less than 10% of cases
- 114       10-24% of cases
- 115       25-49% of cases
- 116       50-74% of cases
- 117       75-99% of cases
- 118       Always (100%)

119      14- Which out of these antibiotics are most commonly prescribed? (Please choose up to three and  
120      specify for other)

- 121       Tetracycline
- 122       Ciprofloxacin
- 123       Erythromycin
- 124       Amoxicillin
- 125       Ampicillin
- 126       Penicillin
- 127       Other: \_\_\_\_\_

128

129      14A. If other, please list most commonly prescribed antibiotics \_\_\_\_\_

130

131      15-Why do you choose a particular antibiotic? (Check all that apply and specify for other)

- 132       Cost
- 133       Availability
- 134       Identification of bacterial infection
- 135       Antibiotic susceptibility testing information
- 136       Convenience
- 137       Broad-spectrum
- 138       Symptoms of patient
- 139       Other: \_\_\_\_\_

140

141 16- Do you feel pressured to prescribe broad-spectrum (tetracycline, ciprofloxacin, erythromycin,  
142 amoxicillin, ampicillin) antibiotics vs narrow-spectrum antibiotics (penicillins)?

143  Yes

144  No

145

146 17- How do you suggest the problem of antibiotic resistance can be tackled in Yemen? (Choose one)

147  Increase testing for susceptibility

148  Increase awareness among patients

149  Enforce prescription laws

150  Improved training of pharmacists and doctors

151  Other: \_\_\_\_\_

152

153 18-What is your profession? (e.g., doctor, nurse, healthcare worker, etc.).

154  Doctor

155  Nurse

156  Healthcare worker

157  Other: \_\_\_\_\_

158

159 19- How many years of professional experience do you have?

160  less than 1 year

161  less than 5 years

162  less than 10 years

163  less than 15 years

164  15 years or more

165

166

167

168

169

170 **B. Pharmacists**

171 1-Do you think antibiotic resistance is a problem in Yemen?

- 172  Yes
- 173  No
- 174  Unsure

175

176 1A. If yes, in what way? (Check all that apply)

- 177  Understudied problem
- 178  Underreported problem
- 179  High levels of AMR are prevalent

180

181 2-Do you think IDPs have a higher risk of antibiotic resistance compared to the population that has not  
182 been displaced?

- 183  Yes
- 184  No

185

186 3-How did you hear about antibiotic resistance?

- 187  Friends
- 188  Family
- 189  Customers
- 190  Training
- 191  Social Media

192 4-Do you sell over the counter antibiotics?

- 193  Yes
- 194  No

195

196 5-How often are prescriptions received for antibiotics a day?

- 197  less than 5 incidences
- 198  less than 10 incidences

199       less than 25 incidences

200       less than 50 incidences

201       50 or more incidences

202

203      6-How often are antibiotics dispensed a day?

204       less than 5 incidences

205       less than 10 incidences

206       less than 25 incidences

207       less than 50 incidences

208       50 or more incidences

209

210      7-How often do patients return due to treatment failure within a month?

211       less than 5 incidences

212       less than 10 incidences

213       less than 25 incidences

214       less than 50 incidences

215       50 or more incidences

216

217      8-Which antibiotics are most commonly dispensed? (Please choose up to three)

218       Tetracycline

219       Ciprofloxacin

220       Erythromycin

221       Amoxicillin

222       Ampicillin

223       Penicillin

224       Other: \_\_\_\_\_

225

226      If other, please list most commonly dispensed antibiotics \_\_\_\_\_

227

228 9-What conditions do you see the highest demand for antibiotics? \_\_\_\_\_

229

230 10-What is the average age of customers?

231  Under 18

232  18-29

233  30-49

234  50-75

235  Over 75

236 11-Are customers more often male or female?

237  Male

238  Female

239  Prefer not to say

240

241 12-Do you counter-prescribe antibiotics?

242  Yes

243  No

244

245 13-What guide do you use for counter-prescribing antibiotics? \_\_\_\_\_

246

247 14-Which antibiotics are most commonly counter-prescribed?

248  Tetracycline

249  Ciprofloxacin

250  Erythromycin

251  Amoxicillin

252  Ampicillin

253  Penicillin

254  Other: \_\_\_\_\_

255

256 If other, please list most commonly counter-prescribed antibiotics \_\_\_\_\_

- 257 15-What is the rationale for empiric antibiotic selection? (Check all that apply)
- 258     ○ Cost
- 259     ○ Availability
- 260     ○ Identification of bacterial infection
- 261     ○ Antibiotic susceptibility testing information
- 262     ○ Convenience
- 263     ○ Broad-spectrum
- 264     ○ Symptoms of patient
- 265     ○ Other: \_\_\_\_\_
- 266
- 267 16-What is the most important consideration for the dispensing/counter-prescribing of antibiotics?
- 268     ○ Cost
- 269     ○ Price patient can pay
- 270     ○ Availability of antibiotics
- 271     ○ Disease
- 272     ○ Other: \_\_\_\_\_
- 273
- 274 17-Which type of antibiotics are of higher demands; broad spectrum (tetracycline, ciprofloxacin,  
275 erythromycin, amoxicillin, ampicillin) or narrow spectrum antibiotics?
- 276     ○ Broad-spectrum
- 277     ○ Narrow-spectrum
- 278     ○ Equal
- 279
- 280 18-How do you suggest the problem of antibiotic resistance can be tackled in Yemen? (Choose one)
- 281     ○ Increased testing for susceptibility
- 282     ○ Increased training of pharmacists and doctors
- 283     ○ Increase awareness among patients
- 284     ○ Enforcement of prescription laws
- 285     ○ Other: \_\_\_\_\_



286

287 19-What is your profession?

288      Certified pharmacists

289      Worker in pharmacy

290 20-What is your professional training?

291      University degree

292      Training at a pharmacy

293      Other: \_\_\_\_\_

294

295 21-What is your profession and years of experience?

296      less than 1 year

297      less than 5 years

298      less than 10 years

299      less than 15 years

300      15 years or more

301

302

303

304

305

306

307

308

309

310

311

312

313

314

315 **Appendix 2 Demographics**

316 Participants: Physicians were almost all physicians, 96% (26/27) (there was no response from one  
317 participant). Pharmacists were mostly, 89% (24/27), certified pharmacists, with the rest 11% (3/27),  
318 being pharmacy workers. The majority had a University degree, 67% (18/27). 30% (8/27) reported being  
319 trained at a pharmacy, with one reporting “other” training.

320 Clients: Physicians reported seeing a different mix of children under the age of 12 years among clients  
321 that use their facilities. Most, 37% (10/27), reported that children comprised <10%, followed by 26%  
322 (7/27) who reported children as comprising between 25-50% of clients and 22% (6/27) where children  
323 comprised between 10-24%. In a few facilities, 15% (4/27), children comprised more than 50% of clients.  
324 This mix possibly reflecting participants’ facility type or specialization.

325 On the other hand, most pharmacists, 52% (14/27), reported that their clients were likely to be adults  
326 between the ages of 18-29 years, followed by 37% (10/27) who reported the age range to be between  
327 30-49 years. Clients under 18 and above 60 years comprised less than 10% respectively.

328 In terms of gender, when identified, most clients at hospitals were females (44%, 12/27); while most  
329 clients at pharmacies were males (44%, 12/27). More than one-fifths of participants chose not to  
330 indicate the gender of their clients: 22% (n=27) among physicians and 30% (n=27) for pharmacists.

331 IDPs as a proportion of clients at healthcare facilities varied, possibly with geographical location (north  
332 or south) as well as facility type, with all facilities seeing between less than 10% to more than 50%.  
333 Almost half of the health workers, 48% (13/27), reported IDPs to comprise between 10-24% of clients,  
334 followed by 33% (9/27) reporting this as less than 10%. An almost equal proportion saw either between  
335 25-50%, or more than 50%, respectively 7% (2/27) and 11% (3/27).

336

337