November 30, 2020

Epigenome-wide scan identifies differentially methylated regions for lung cancer using pre-diagnostic peripheral blood

Naisi Zhao¹, Mengyuan Ruan¹, Devin C. Koestler^{2,3}, Jiayun Lu⁴, Carmen J. Marsit⁶, Karl T. Kelsey^{7,8}, Elizabeth A. Platz^{4,5}, Dominique S. Michaud^{1,7}

Author Affiliations:

¹Department of Public Health & Community Medicine, Tufts University School of Medicine, Tufts University, Boston, MA

²Department of Biostatistics & Data Science, University of Kansas Medical Center, Kansas City, KS

³University of Kansas Cancer Center, University of Kansas Medical Center, Kansas City, KS

⁴Department of Epidemiology, Johns Hopkins Bloomberg School of Public Health, Baltimore, MD

⁵The Sidney Kimmel Comprehensive Cancer Center at Johns Hopkins, Baltimore, MD

⁶Department of Environmental Health and Department of Epidemiology, Rollins School of Public Health, Emory University, Atlanta, GA

⁷Department of Epidemiology, Brown University, Providence, RI

⁸Department of Pathology and Laboratory Medicine, Brown University, Providence, RI

Running Title: New genomic regions associated with lung cancer

Correspondence:

Dominique S. Michaud, ScD

Tufts University School of Medicine

136 Harrison Avenue

Boston, MA 02111

Tel: 617-636-0482

Email: Dominique.Michaud@tufts.edu

ORCID (if available)

Naisi Zhao: <https://orcid.org/0000-0002-0517-0709>

Mengyuan Ruan: <https://orcid.org/0000-0001-7555-8963>

Devin C. Koestler: <https://orcid.org/0000-0002-0598-0146>

Carmen J. Marsit: <https://orcid.org/0000-0003-4566-150X>

Karl T. Kelsey: <https://orcid.org/0000-0002-2302-1600>

Elizabeth A. Platz: <https://orcid.org/0000-0003-3676-8954>

Dominique S. Michaud: <https://orcid.org/0000-0002-8107-3011>

Abstract

Background: To reduce lung cancer burden in the US, a better understanding of biological mechanisms in early disease development could provide new opportunities for risk stratification.

Methods: In a nested case-control study, we measured blood leukocyte DNA methylation levels in pre-diagnostic samples collected from 430 men and women in the 1989 CLUE II cohort. Median time from blood drawn to diagnosis was 14 years for all participants. We compared DNA methylation levels by case/control status to identify novel genomic regions, both single CpG sites and differentially methylated regions (DMRs), while controlling for known DNA methylation changes associated with smoking using a previously described pack-years based smoking methylation score. Stratification analyses were conducted by time from blood draw to diagnosis, histology, and smoking status.

Results: We identified sixteen single CpG sites and forty DMRs significantly associated with lung cancer risk ($q < 0.05$). The identified genomic regions were associated with genes including *HI9*, *HOXA4*, *RUNX3*, *BRICD5*, *PLXNB2*, and *RP13*. For the single CpG sites, the strongest association was noted for cg09736286 in the *DIABLO* gene (OR [for 1 SD] = 2.99, 95% CI: 1.95-4.59, P-value = 4.81×10^{-7}). For the DMRs, we found that CpG sites in the *HOXA4* region were hypermethylated in cases compared to controls.

Conclusion: The single CpG sites and DMRs that we identified represented significant measurable differences in lung cancer risk, providing new insights into the biological processes of early lung cancer development and potential biomarkers for lung cancer risk stratification.

Keywords: Lung cancer, DNA methylation, Illumina MethylationEPIC array, epigenome-wide association study (EWAS), blood.

Introduction

Despite substantial reductions in lung cancer incidence and death rates over the past three decades (Siegel et al. 2020), lung cancer continues to be the leading cause of cancer death in the US and is projected to account for 135,720 deaths in 2020, approximately 22% of all cancer deaths (Siegel et al. 2020). To reduce the lung cancer burden in the US, cancer prevention and early detection remain a top priority (Moyer and Force 2014). However, while the conventional lung cancer screening method of using low-dose CT (LDCT) scans is effective in reducing lung cancer mortality (National Lung Screening Trial Research et al. 2011), it leads to high false positives rates (>95% of pulmonary nodules detected are benign) (Fabrikant et al. 2018), over-diagnosis (Heleno et al. 2018; Patz et al. 2014), radiation exposure (McCunney and Li 2014), and has had poor uptake (Quaife et al. 2016). A better understanding of biological mechanisms in early disease development, especially understanding gained through using DNA methylation data, may contribute new insights into lung cancer development and provide new opportunities for risk stratification, screening prioritization, and drug development.

Genome-wide DNA methylation profiling has provided new insight into risk factors, biological pathways, and disease processes. For instance, differences in blood leukocyte DNA methylation levels (proportion of CpGs methylated) have been associated with smoking (Baglietto et al. 2017; Shenker et al. 2013), elevated subclinical inflammation (Ahsan et al. 2017; Ligthart et al. 2016), obesity (Wahl et al. 2016; Xu et al. 2018), type II diabetes (Chambers et al. 2015), and heart disease (Agha et al. 2019; Huan et al. 2019). With respect to lung cancer, many of the alterations in blood leukocyte DNA methylation levels identified to date have been directly linked to smoking behaviors (Baglietto et al. 2017; Fasanelli et al. 2015; Shenker et al. 2013; Zhang et al. 2016b; Zhang et al. 2016c). One recent meta-analysis that included over 15,000 individuals identified 2,623 differentially methylated CpG sites that are related to cigarette smoking (Joehanes et al. 2016). Some of the identified smoking-related CpG sites have also been shown to mediate the effects of smoking on lung cancer (Baglietto et al. 2017; Battram et al. 2019; Fasanelli et al. 2015). Nevertheless, very few studies have investigated methylation changes associated with lung cancer risk that are not associated with smoking (Zhang et al. 2016a; Zhang et al. 2016b).

While smoking remains the strongest risk factor for lung cancer, identifying DNA methylation alterations that are not caused by active smoking could reveal important biological pathways. Among smokers, variation in methylation may indicate different genetic susceptibility to lung cancer, since individuals often differ in their ability to detoxify carcinogenic compounds or to repair induced DNA damage, for example. Alternatively, variation in methylation could be a result of other environmental risk factors, such as occupational exposures, changes in the immune response, radon, or secondhand smoke. Using blood samples collected from men and women without a cancer diagnosis in 1989 in the CLUE II cohort (Genkinger et al. 2004; Kakourou et al. 2015), we compared DNA methylation levels in individuals who later developed lung cancer with those who did not develop lung cancer in the same time frame. Specifically, we aimed to identify both single CpG sites, and differentially methylated regions, measured in pre-diagnostic blood samples of lung cancer cases and matched controls, that represent measurable differences in lung cancer risk independent of smoking exposures.

Materials and Methods

Study population – CLUE I/II cohort

Subjects for this study were selected from among participants of the CLUE II cohort who had also participated in the CLUE I cohort study (flowchart in Supplemental Figure 1 and additional details in Supplemental Methods). Both cohorts were based in Washington County, MD, and were initially established to identify serological precursors to cancer and other chronic diseases (Genkinger et al. 2004; Kakourou et al. 2015; Schober et al. 1987). CLUE II was conducted from May through October 1989, during which 32,894 individuals (25,076 were Washington County residents) provided a blood sample (Comstock et al. 1991). Among all participants, 98.3% were white, reflecting the population of this county at the time, and 59% were female. Participants provided health information at baseline, including the potential confounders attained education, cigarette smoking status, number of cigarettes smoked daily, cigar/pipe smoking status, and self-reported weight and height, from which body mass index (BMI) was calculated.

Lung cancer ascertainment

All incident lung cancer cases (ICD9 162 and ICD10 C34) were ascertained from linkage to the Washington County cancer registry (before 1992 to the present) and the Maryland Cancer Registry (since 1992 when it began to the present). We selected all 241 first primary incident lung cancer cases who had participated in CLUE I and were diagnosed after the day of blood draw in CLUE II through January 2018. Using incidence density sampling, we selected 1 control per case matching on age, sex, cigarette smoking status, number of cigarettes smoked daily, cigar/pipe smoking status, and date of CLUE II blood draw.

DNA methylation measurements

DNA extracted from buffy coat was bisulfite-treated using the EZ DNA Methylation Kit (Zymo) and DNA methylation was measured at specific CpG sites across the genome using the 850K Illumina Infinium MethylationEPIC BeadArray (Illumina, Inc, CA, USA) at the University of Minnesota Genomic Center (details in Supplemental Methods).

Statistical analysis

In the current nested case-control study, we aimed to identify novel genomic regions, both single CpG sites and differentially methylated regions where differences in DNA methylation levels are not explained by smoking exposures, by controlling for the known DNA methylation changes associated with smoking in the statistical analysis. To examine the association between CpG-specific DNA methylation and lung cancer risk, we conducted epigenome-wide association analysis using unconditional multivariable logistic regression to estimate odds ratios (OR) of lung cancer per 1 SD increase in methylation level at single CpG sites. To maximize power, we used unconditional logistic regression to include cases and controls without a matched pair, and included participants every time they were sampled. All models were adjusted for age at blood draw, sex, surrogate variables for batch effects (Leek et al. 2012; Leek et al. 2010), smoking status (never, former, current), pack-years based smoking methylation score (details in Supplemental Methods), BMI, and leukocyte cell composition (Houseman et al. 2012; Salas et al. 2018) (given the potential for confounding by cell composition) (Adalsteinsson et al. 2012). All p-values were

adjusted for multiple comparisons using the false discovery rate (FDR) method. Analyses of single CpGs with lung cancer were also stratified by smoking status and time from blood-draw to diagnosis, and separately by non-small cell (NSCLC) and small cell (SCLC) histology. All controls were included in these three types of stratification analyses. All statistical analyses were performed in R (version 3.5.0).

We used the DMRcate Bioconductor R package (Peters et al. 2015) to identify differentially methylated regions (DMRs) associated with lung cancer risk. Adjusting for the same covariates as in the single CpG analyses, DMRs were calculated using a parameter setting of $\lambda=1,000$ and kernel adjustment $C=2$ (default setting) (Peters et al. 2015). Statistically significant DMRs were required to have a minimum of two statistically significant single CpGs and to meet the multiple testing adjustment criteria of $FDR < 0.1$. Associations were also examined by time from blood-draw to diagnosis and lung cancer histology. Two of the most statistically significant regions were further evaluated for patterns by time to diagnosis.

Results

Population characteristics

Table 1 presents the characteristics of the 208 lung cancer cases and 222 controls that we included in this study. Over 99% of the majority of participants were white. The median time to lung cancer diagnosis was 14 years. The median age at blood draw in 1989 was 59 and 57 years in cases and controls, respectively. Overall, 55% of cases and controls were women and 11% were never smokers (Table 1).

Single CpG EWAS analysis

The EWAS analysis identified 16 differentially methylated CpGs that were statistically significant after multiple comparisons correction ($q < 0.05$). Results are presented in Table 2 (statistically significant CpGs were sorted by q-value) and Figure 1. Among the 16 CpGs, many were located in genomic regions that have been previously associated with lung cancer or other malignancies (*RUNX3* (Sato et al. 2005), *H19* (Huang et al. 2018), *BALP2L2* (Liu et al. 2020), *GPR132* (Chen et al. 2017), *CUEDC1* (Lopes et al. 2018), *SSBP4* (Guo et al. 2018), *AMPD2* (Gao et al. 2020), *ADAM11* (Sieuwerts et al. 2005), and *RTN4R* (He et al. 2020); the top 1000 CpGs based on q-value are presented in Supplemental Table 1).

CpGs previously reported to be associated with lung cancer risk, including cg05575921 in the *AHRR* gene (Fasanelli et al. 2015), cg03636183 in the *F2RL3* gene (Fasanelli et al. 2015), cg23387569 in the *AGAP2* gene (Baglietto et al. 2017), cg10151248 in the *PC* gene (Sandanger et al. 2018), and cg13482620 in the *B3GNTL1* gene (Sandanger et al. 2018), were not statistically significant in our analyses in which we adjusted for a pack-years methylation score. Adjusting for smoking status (never, former, current) but not for the packyears methylation score, two CpGs in the *AHRR* and *F2RL3* genes had similar-sized associations with lung cancer risk as previously reported (*AHRR*: OR [for 1 SD] = 0.43, 95% CI: 0.31-0.60, P-value = 6.76×10^{-7} vs. previously reported OR [for 1 SD] = 0.39, 95% CI: 0.24-0.61, P-value = 2.55×10^{-5} for cg05575921; *F2RL3*: OR for [1 SD] = 0.53, 95% CI: 0.40-0.70, P-value = 7.91×10^{-6} vs. previously reported OR [for 1 SD] = 0.51, 95% CI: 0.35-0.73, P-value = 4.19×10^{-4} for cg03636183) (Fasanelli et al. 2015). In addition, in the complete EWAS analysis conducted without adjusting for the packyears methylation score, only one CpG (cg14391737) was statistically significantly associated with lung cancer risk after adjusting for multiple comparisons. This CpG has been related to smoking in multiple studies (Joehanes et al. 2016) (top 1000 CpGs presented in Supplemental Table 2).

To further examine the associations of the significant CpGs with lung cancer risk, we stratified by time from blood draw to diagnosis (≤ 10 , >10 years). The magnitude of risk was similar in the two strata; small differences in risk were likely due to statistical variability (Table 2). For these 16 differentially methylated CpGs, the ORs of lung cancer were in general slightly higher for former smokers and for SCLC, than among current smokers or for NSCLC, respectively (Supplemental Table 3).

DMR analysis

Using the DMRcate package in R, we identified differentially methylated regions (DMRs) by case/control status (Peters et al. 2015). Instead of focusing on single CpG identification, the DMRcate method identifies regions of chromosomes that are differentially methylated by case-control status. After adjusting for both smoking status and packyears methylation score, forty DMRs were found to be statistically significantly associated with lung cancer risk (Table 3). Many of the top regions identified

included genes that have been linked to lung cancer in previous studies (*H19* (Huang et al. 2018), *HOXA4* (Xu et al. 2019), *PLXNB2* (Liu and Zhao 2019), *PRDM1* (Zhu et al. 2017), *TSPAN4* (Ying et al. 2019), *PHPT1* (Xu et al. 2010), *MSI2* (Kudinov et al. 2016), *CBX5* (Yu et al. 2012), *RCAN1* (Ma et al. 2017), *CCL5* (Huang et al. 2009), and *BRDT* (Grunwald et al. 2006)).

We conducted stratified analyses to examine whether our DMR results were modified by time to diagnosis or differed by histology. Among those with ≤ 10 years between blood draw and diagnosis, a region located on chromosome 20:36148699-36149271 (genes *NNAT* and *BLCAP*) was statistically significantly associated with lung cancer. By histology, one region located in *H19* gene and one region located in *MYEOV* gene were statistically significant DMRs for NSCLC and SCLC, respectively (Supplemental Table 4).

We conducted further analyses for two of the most statistically significant DMRs. These two regions, located on the chromosome 11 (*H19* gene) and 7 (*HOXA4* gene), consisted of 31 and 20 CpGs, respectively, that differed between the cases and controls (all were $q\text{-value} < 0.1$ FDR adjusted). For each of these two regions, we selected the CpGs with the strongest associations with lung cancer risk for comparison by time to cancer diagnosis (≤ 10 , > 10 years). Results for CpGs in these two regions are presented in Table 4 (top 10 statistically significant CpGs are included), with additional tables presented in Supplemental Table 5. For CpGs located in the *HOXA4* region, 18 out of 20 were statistically significant and all sites were hypermethylated in cases compared to controls. Results for these *HOXA4* region CpGs were similar in the ≤ 10 and > 10 year time to diagnosis groups. In the *H19* region, the strongest association was noted for cg00237904. This CpG was also identified in the top statistically significant CpGs in the single CpG EWAS (Table 2, $q\text{-value} < 0.05$).

Discussion

In this study, we identified both single CpGs and DMRs in lung cancer that are not primarily driven by smoking history, by using a DNA methylation-based packyears score to adjust for cumulative smoking. Using a prospective study design (with pre-diagnostic blood), we identified 16 single CpG sites and 40

DMRs regions that were associated with lung cancer risk; genes in these regions included *H19*, *HOXA4*, *RUNX3*, *BRICD5*, *RP13*, and *PLXNB2*.

Previous studies have used either case-control or nested case-control designs to study the association between methylation markers and lung cancer risk. Retrospective case-control studies on this topic are not comparable to our study because they either used a different methodology to measure blood leukocyte methylation (Wang et al. 2010), or measured methylation biomarkers in sputum as a classifier for lung cancer risk (Leng et al. 2017; Liu et al. 2017). Three nested case-control studies previously used pre-diagnostic, peripheral blood samples to examine DNA methylation levels associated with lung cancer risk, while adjusting for smoking using self-reported information (Baglietto et al. 2017; Fasanelli et al. 2015; Sandanger et al. 2018). Fasanelli 2015 (Fasanelli et al. 2015) and Baglietto 2016 (Baglietto et al. 2017) together identified six CpGs (cg05951221, cg21566642, cg05575921, cg06126421, cg23387569 and cg03636183) with significant ORs for lung cancer after adjusting for smoking using self-reported smoking status. Further stratification showed that five of the six CpGs had methylation levels strongly influenced by smoking. Sandanger 2018 found two additional CpGs, cg10151248 (*PC*) and cg13482620 (*B3GNTL1*), to be significantly associated with lung cancer risk after adjusting for smoking status, pack-years, and a comprehensive smoking index built using self-reported information (Sandanger et al. 2018). In our analyses, cg05575921 (*AHRR*), cg03636183 (*F2RL3*), and cg21566642 methylation levels were statistically significantly associated with lung cancer when we adjusted for self-reported smoking status, whereas cg10151248 (*PC*), cg13482620 (*B3GNTL1*), and cg23387569 (*AGAP2*) methylation levels did not significantly differ between the cases and controls regardless of whether we adjusted for the pack-years methylation score or not. cg05951221 and cg06126421 were not associated with lung cancer.

Our analyses identified novel genomic regions that are independent of smoking exposures. Many of the significant single CpGs we identified through EWAS were located in genomic regions that have been previously associated with lung cancer or other malignancies (*RUNX3* (Sato et al. 2005), *H19* (Huang et al. 2018), *BAIAP2L2* (Liu et al. 2020), *GPR132* (Chen et al. 2017), *CUEDCI* (Lopes et al. 2018), *SSBP4* (Guo et al. 2018), *AMPD2* (Gao et al. 2020), *ADAM11* (Sieuwerts et al. 2005), and *RTN4R* (He et al. 2020)). For

instance, the *RUNX3* region is a tumor suppressor gene that is implicated in lung cancer oncogenesis (Sato et al. 2005). Promoter hypermethylation of *RUNX3* has been associated with NSCLC survival (Yanagawa et al. 2007). Many of the top differentially methylated regions we identified using the DMRcate analysis included genes that have been previously linked to lung cancer in other studies (*H19* (Huang et al. 2018), *HOXA4* (Xu et al. 2019), *PLXNB2* (Liu and Zhao 2019), *PRDM1* (Zhu et al. 2017), *TSPAN4* (Ying et al. 2019), *PHPTI* (Xu et al. 2010), *MSI2* (Kudinov et al. 2016), *CBX5* (Yu et al. 2012), *RCAN1* (Ma et al. 2017), *CCL5* (Huang et al. 2009), and *BRDT* (Grunwald et al. 2006)), providing support for our findings. Of these eleven genetic regions already linked to lung cancer, many are shown to be connected to poor outcome in lung cancer. For instance, decreased expressed levels of *PLXNB2* (Liu and Zhao 2019) and *PRDM1* (Zhu et al. 2017) have been found to be correlated with poor prognosis in lung cancer, while *TSPAN4* (Ying et al. 2019), *PHPTI* (Xu et al. 2010), *MSI2* (Kudinov et al. 2016), and *CBX5* (Yu et al. 2012) are linked to metastasis. In addition, *HOXA4* (Xu et al. 2019), *RCAN1* (Ma et al. 2017), and *CCL5* (Huang et al. 2009) are involved in the growth, development, and migration of lung cancer cells. Other top regions identified in our study include genes that have been linked to breast cancer (*NNAT* (Nass et al. 2017), *RPL7A* (Zhu et al. 2001), and *HIST1H2BO* (Xie et al. 2019)), colorectal cancer (*RP11* (Sun et al. 2017; Wu et al. 2019)), endometrial cancer (*HELZ2* (Qiao et al. 2019)), pancreatic cancer (*LFNG* (Liu et al. 2016)), prostate cancer (*MAST3* (Dahlman et al. 2012)), renal cell carcinoma (*KCNJ1* (Guo et al. 2015)), and tumor progression (*ZC3H12D* (Huang et al. 2012)).

Of all the novel genomic regions that we identified as associated with lung cancer risk, the *H19* region was the only one that appeared in both of the single CpG EWAS and DMRcate results. The *H19* long noncoding RNA (lncRNA) has been previously implicated in lung cancer causation. Inhibition of lncRNA *H19* has been found to suppress the growth, migration, and invasion of NSCLC (Huang et al. 2018). In terms of disease development, loss of imprinting of the *H19* gene has been connected to a genome-wide loss of methylation, and associated with the transformation from normal to NSCLC (Anisowicz et al. 2008; Kondo et al. 1995; Langevin et al. 2015). In our analyses, we found that hypermethylation of many *H19* region CpGs were associated with lung cancer risk in CLUE II. The direction of this association was

unexpected since the overexpression of H19 LncRNA in lung tumor is often correlated with hypomethylation of the promoter region CpGs (Kondo et al. 1995). H19 LncRNA belongs to a highly conserved imprinted gene cluster that plays important roles in embryonal development and growth control (Gabory et al. 2010) and *H19* region methylation has been found to be influenced by early life exposures, including maternal factors during pregnancy (Miyaso et al. 2017), suggesting the possibility that external exposures could impact *H19* methylation. Since the blood samples were drawn years before cancer diagnosis in this study, the methylation patterns we observed could be regions that are modulated early on in lung cancer development. More research is needed to investigate the methylation pattern in blood prior to cancer diagnosis.

Use of the DMRcate analysis methods and the Illumina Infinium MethylationEPIC 850K BeadArray allowed us to investigate genome-wide regional methylation level differences between lung cancer cases and controls. Some of the regions we identified that have not previously been associated with lung cancer should be investigated in other populations. It is possible that some of single CpGs and DMRs that we identified after adjusting for the packyear methylation score could be related to risk factors unique to CLUE II. Further studies are needed to investigate pathways related to the novel genomic regions that we identified.

This study demonstrated the importance of carefully controlling for known DNA methylation changes associated with smoking to be able to identify novel genomic regions. We showed the potential for this approach to identify DMRs (i.e., not single CpG alterations) by case/control status using peripheral blood collected prior to lung cancer diagnosis. These findings suggest that methylation changes detectable years prior to cancer diagnosis could potentially influence lung cancer risk, providing new insights into the biological processes of early lung cancer development. Further work in other populations should be conducted to validate regions that we observed to be associated with lung cancer risk independent of smoking exposures, especially among different ethnic and racial groups.

Word Count: 3315

Acknowledgments

Cancer incidence data were provided by the Maryland Cancer Registry, Center for Cancer Surveillance and Control, Maryland Department of Health, 201 W. Preston Street, Room 400, Baltimore, MD 21201. We acknowledge the State of Maryland, the Maryland Cigarette Restitution Fund, and the National Program of Cancer Registries of the Centers for Disease Control and Prevention for the funds that helped support the availability of the cancer registry data.

Declarations

Competing interest

The authors report no conflicts of interest.

Funding

This work was supported by 2018 American Association for Cancer Research (AACR)-Johnson & Johnson Lung Cancer Innovation Science (18-90-52-MICH).

Note: The funders had no role in the design of the study; the collection, analysis, and interpretation of the data; the writing of the manuscript; and the decision to submit the manuscript for publication.

Authors' contributions

DSM., KTK and EAP designed the study, obtained funding and acquisition of data. JL assisted with preparation of dataset. DSM supervised all research activities. MR and NZ conducted the statistical analyses. DSM and NZ drafted the manuscript. DSM., EAP, KTK, DCK and CJM interpreted the data and provided critical revisions of the manuscript. All authors read and approved the final version of the manuscript.

Availability of data and material

The datasets generated during the current study are available from the corresponding author on reasonable request and will be deposited into dbGaP within 1 year.

Ethics approval

This study was approved by the Institutional Review Board at Johns Hopkins University Bloomberg School of Public Health and at Tufts University.

References

- Adalsteinsson BT et al. (2012) Heterogeneity in white blood cells has potential to confound DNA methylation measurements PLoS One 7:e46705 doi:10.1371/journal.pone.0046705
- Agha G et al. (2019) Blood Leukocyte DNA Methylation Predicts Risk of Future Myocardial Infarction and Coronary Heart Disease Circulation 140:645-657 doi:10.1161/CIRCULATIONAHA.118.039357
- Ahsan M et al. (2017) The relative contribution of DNA methylation and genetic variants on protein biomarkers for human diseases PLoS Genet 13:e1007005 doi:10.1371/journal.pgen.1007005
- Anisowicz A et al. (2008) A high-throughput and sensitive method to measure global DNA methylation: application in lung cancer BMC cancer 8:222
- Baglietto L et al. (2017) DNA methylation changes measured in pre-diagnostic peripheral blood samples are associated with smoking and lung cancer risk International journal of cancer Journal international du cancer 140:50-61 doi:10.1002/ijc.30431
- Batram T et al. (2019) Appraising the causal relevance of DNA methylation for risk of lung cancer International journal of epidemiology 48:1493-1504
- Chambers JC et al. (2015) Epigenome-wide association of DNA methylation markers in peripheral blood from Indian Asians and Europeans with incident type 2 diabetes: a nested case-control study The lancet Diabetes & endocrinology 3:526-534 doi:10.1016/S2213-8587(15)00127-8
- Chen P et al. (2017) Gpr132 sensing of lactate mediates tumor-macrophage interplay to promote breast cancer metastasis Proceedings of the National Academy of Sciences 114:580-585
- Comstock GW, Helzlsouer KJ, Bush TL (1991) Prediagnostic serum levels of carotenoids and vitamin E as related to subsequent cancer in Washington County, Maryland The American journal of clinical nutrition 53:260S-264S
- Dahlman KB et al. (2012) Modulators of prostate cancer cell proliferation and viability identified by short-hairpin RNA library screening PLoS One 7:e34414
- Fabrikant MS, Wisnivesky JP, Marron T, Taioli E, Veluswamy RR (2018) Benefits and Challenges of Lung Cancer Screening in Older Adults Clinical therapeutics doi:10.1016/j.clinthera.2018.03.003
- Fasanelli F et al. (2015) Hypomethylation of smoking-related genes is associated with future lung cancer in four prospective cohorts Nature communications 6:1-9
- Gabory A, Jammes H, Dandolo L (2010) The H19 locus: role of an imprinted non-coding RNA in growth and development Bioessays 32:473-480
- Gao Q-Z, Qin Y, Wang W-J, Fei B-J, Han W-F, Jin J-Q, Gao X (2020) Overexpression of AMPD2 indicates poor prognosis in colorectal cancer patients via the Notch3 signaling pathway World Journal of Clinical Cases 8:3197
- Genkinger JM, Platz EA, Hoffman SC, Comstock GW, Helzlsouer KJ (2004) Fruit, vegetable, and antioxidant intake and all-cause, cancer, and cardiovascular disease mortality in a community-dwelling population in Washington County, Maryland American journal of epidemiology 160:1223-1233 doi:10.1093/aje/
- Grunwald C et al. (2006) Expression of multiple epigenetically regulated cancer/germline genes in nonsmall cell lung cancer International journal of cancer 118:2522-2528
- Guo X et al. (2018) A comprehensive cis-eQTL analysis revealed target genes in breast cancer susceptibility loci identified in genome-wide association studies The American Journal of Human Genetics 102:890-903
- Guo Z et al. (2015) KCNJ1 inhibits tumor proliferation and metastasis and is a prognostic factor in clear cell renal cell carcinoma Tumor Biology 36:1251-1259
- He Y, Ji P, Li Y, Wang R, Ma H, Yuan H (2020) Genetic Variants Were Associated With the Prognosis of Head and Neck Squamous Carcinoma Frontiers in oncology 10:372

- Heleno B, Siersma V, Brodersen J (2018) Estimation of Overdiagnosis of Lung Cancer in Low-Dose Computed Tomography Screening: A Secondary Analysis of the Danish Lung Cancer Screening Trial *JAMA Intern Med* 178:1420-1422 doi:10.1001/jamainternmed.2018.3056
- Houseman EA et al. (2012) DNA methylation arrays as surrogate measures of cell mixture distribution *BMC bioinformatics* 13:86 doi:10.1186/1471-2105-13-86
- Huan T et al. (2019) Genome-wide identification of DNA methylation QTLs in whole blood highlights pathways for cardiovascular disease *Nat Commun* 10:4267 doi:10.1038/s41467-019-12228-z
- Huang C-Y, Fong Y-C, Lee C-Y, Chen M-Y, Tsai H-C, Hsu H-C, Tang C-H (2009) CCL5 increases lung cancer migration via PI3K, Akt and NF- κ B pathways *Biochemical pharmacology* 77:794-803
- Huang S et al. (2012) The putative tumor suppressor Zc3h12d modulates toll-like receptor signaling in macrophages *Cellular signalling* 24:569-576
- Huang Z, Lei W, Hu HB, Zhang H, Zhu Y (2018) H19 promotes non-small-cell lung cancer (NSCLC) development through STAT3 signaling via sponging miR-17 *Journal of cellular physiology* 233:6768-6776
- Joeanes R et al. (2016) Epigenetic signatures of cigarette smoking Circulation: cardiovascular genetics 9:436-447
- Kakourou A et al. (2015) Interleukin-6 and risk of colorectal cancer: results from the CLUE II cohort and a meta-analysis of prospective studies *Cancer Causes & Control* 26:1449-1460
- Kondo M, Suzuki H, Ueda R, Osada H, Takagi K, Takahashi T (1995) Frequent loss of imprinting of the H19 gene is often associated with its overexpression in human lung cancers *Oncogene* 10:1193-1198
- Kudinov AE et al. (2016) Musashi-2 (MSI2) supports TGF- β signaling and inhibits claudins to promote non-small cell lung cancer (NSCLC) metastasis *Proceedings of the National Academy of Sciences* 113:6955-6960
- Langevin SM, Kratzke RA, Kelsey KT (2015) Epigenetics of lung cancer *Translational Research* 165:74-90
- Leek JT, Johnson WE, Parker HS, Jaffe AE, Storey JD (2012) The sva package for removing batch effects and other unwanted variation in high-throughput experiments *Bioinformatics* 28:882-883
- Leek JT et al. (2010) Tackling the widespread and critical impact of batch effects in high-throughput data *Nature Reviews Genetics* 11:733-739
- Leng S et al. (2017) Gene methylation biomarkers in sputum as a classifier for lung cancer risk *Oncotarget* 8:63978
- Ligthart S et al. (2016) DNA methylation signatures of chronic low-grade inflammation are associated with complex diseases *Genome Biol* 17:255 doi:10.1186/s13059-016-1119-5
- Liu D et al. (2017) The indirect efficacy comparison of DNA methylation in sputum for early screening and auxiliary detection of lung cancer: a meta-analysis *International journal of environmental research and public health* 14:679
- Liu H, Zhao H (2019) Prognosis related miRNAs, DNA methylation, and epigenetic interactions in lung adenocarcinoma *Neoplasia* 66:487-493
- Liu J, Shangguan Y, Sun J, Cong W, Xie Y (2020) BAIAP2L2 promotes the progression of gastric cancer via AKT/mTOR and Wnt3a/ β -catenin signaling pathways *Biomedicine & Pharmacotherapy* 129:110414
- Liu P et al. (2016) Quantitative secretomic analysis of pancreatic cancer cells in serum-containing conditioned medium *Scientific reports* 6:37606
- Lopes R et al. (2018) CUEDC1 is a primary target of ER α essential for the growth of breast cancer cells *Cancer letters* 436:87-95
- Ma N, Shen W, Pang H, Zhang N, Shi H, Wang J, Zhang H (2017) The effect of RCAN1 on the biological behaviors of small cell lung cancer *Tumor Biology* 39:1010428317700405
- McCunney RJ, Li J (2014) Radiation risks in lung cancer screening programs *Chest* 145:618-624

- Miyaso H, Sakurai K, Takase S, Eguchi A, Watanabe M, Fukuoka H, Mori C (2017) The methylation levels of the H19 differentially methylated region in human umbilical cords reflect newborn parameters and changes by maternal environmental factors during early pregnancy *Environmental research* 157:1-8
 - Moyer VA, Force USPST (2014) Screening for lung cancer: U.S. Preventive Services Task Force recommendation statement *Ann Intern Med* 160:330-338 doi:10.7326/M13-2771
 - Nass N et al. (2017) High neuronatin (NNAT) expression is associated with poor outcome in breast cancer *Virchows Archiv* 471:23-30
 - National Lung Screening Trial Research T et al. (2011) Reduced lung-cancer mortality with low-dose computed tomographic screening *N Engl J Med* 365:395-409 doi:10.1056/NEJMoa1102873
 - Patz EF, Jr. et al. (2014) Overdiagnosis in low-dose computed tomography screening for lung cancer *JAMA Intern Med* 174:269-274 doi:10.1001/jamainternmed.2013.12738
 - Peters TJ et al. (2015) De novo identification of differentially methylated regions in the human genome *Epigenetics Chromatin* 8:6 doi:10.1186/1756-8935-8-6
 - Qiao Z, Jiang Y, Wang L, Wang L, Jiang J, Zhang J (2019) Mutations in KIAA1109, CACNA1C, BSN, AKAP13, CELSR2, and HELZ2 are associated with the prognosis in endometrial cancer *Frontiers in Genetics* 10:909
 - Quaife SL et al. (2016) The Lung Screen Uptake Trial (LSUT): protocol for a randomised controlled demonstration lung cancer screening pilot testing a targeted invitation strategy for high risk and 'hard-to-reach' patients *BMC cancer* 16:1-9
 - Salas LA, Koestler DC, Butler RA, Hansen HM, Wiencke JK, Kelsey KT, Christensen BC (2018) An optimized library for reference-based deconvolution of whole-blood biospecimens assayed using the Illumina HumanMethylationEPIC BeadArray *Genome Biol* 19:64 doi:10.1186/s13059-018-1448-7
 - Sandanger TM et al. (2018) DNA methylation and associated gene expression in blood prior to lung cancer diagnosis in the Norwegian Women and Cancer cohort *Scientific reports* 8:1-10
 - Sato K, Tomizawa Y, Iijima H, Saito R, Ishizuka T, Nakajima T, Mori M (2005) Epigenetic inactivation of the RUNX3 gene in lung cancer *Oncology reports* 15:129-135
 - Schober SE, Comstock GW, Helsing KJ, Salkeld RM, Morris JS, Rider AA, Brookmeyer R (1987) Serologic precursors of cancer: I. Prediagnostic serum nutrients and colon cancer risk *American journal of epidemiology* 126:1033-1041
 - Shenker NS et al. (2013) Epigenome-wide association study in the European Prospective Investigation into Cancer and Nutrition (EPIC-Turin) identifies novel genetic loci associated with smoking *Hum Mol Genet* 22:843-851 doi:10.1093/hmg/dd488
 - Siegel RL, Miller KD, Jemal A (2020) Cancer statistics, 2020 *CA Cancer J Clin* 70:7-30 doi:10.3322/caac.21590
 - Sieuwerts AM et al. (2005) How ADAM-9 and ADAM-11 differentially from estrogen receptor predict response to tamoxifen treatment in patients with recurrent breast cancer: a retrospective study *Clinical Cancer Research* 11:7311-7321
 - Sun L et al. (2017) Down-regulation of long non-coding RNA RP11-708H21. 4 is associated with poor prognosis for colorectal cancer and promotes tumorigenesis through regulating AKT/mTOR pathway *Oncotarget* 8:27929
 - Wahl S et al. (2016) Epigenome-wide association study of body mass index, and the adverse outcomes of adiposity *Nature* 541:81 doi:10.1038/nature20784
- <https://www.nature.com/articles/nature20784#supplementary-information>
- Wang L et al. (2010) Methylation markers for small cell lung cancer in peripheral blood leukocyte DNA *Journal of Thoracic Oncology* 5:778-785

- Wu Y et al. (2019) m 6 A-induced lncRNA RP11 triggers the dissemination of colorectal cancer cells via upregulation of Zeb1 *Molecular cancer* 18:1-16
- Xie W et al. (2019) Expression and potential prognostic value of histone family gene signature in breast cancer *Experimental and therapeutic medicine* 18:4893-4903
- Xu A-j, Xia X-h, Du S-t, Gu J-c (2010) Clinical significance of PHPT1 protein expression in lung cancer *Chinese medical journal* 123:3247-3251
- Xu K, Liu B, Ma Y (2019) The tumor suppressive roles of ARHGAP25 in lung cancer cells *OncoTargets and therapy* 12:6699
- Xu K, Zhang X, Wang Z, Hu Y, Sinha R (2018) Epigenome-wide association analysis revealed that SOCS3 methylation influences the effect of cumulative stress on obesity *Biol Psychol* 131:63-71 doi:10.1016/j.biopsycho.2016.11.001
- Yanagawa N, Tamura G, Oizumi H, Kanauchi N, Endoh M, Sadahiro M, Motoyama T (2007) Promoter hypermethylation of RASSF1A and RUNX3 genes as an independent prognostic prediction marker in surgically resected non-small cell lung cancers *Lung cancer* 58:131-138
- Ying X, Zhu J, Zhang Y (2019) Circular RNA circ-TSPAN4 promotes lung adenocarcinoma metastasis by upregulating ZEB1 via sponging miR-665 *Molecular genetics & genomic medicine* 7:e991
- Yu Y-H et al. (2012) Network biology of tumor stem-like cells identified a regulatory role of CBX5 in lung cancer *Scientific reports* 2:584
- Zhang X, Gao L, Liu Z-P, Jia S, Chen L (2016a) Uncovering driver DNA methylation events in nonsmoking early stage lung adenocarcinoma *BioMed research international* 2016
- Zhang Y, Breitling LP, Balavarca Y, Holleczer B, Schottker B, Brenner H (2016b) Comparison and combination of blood DNA methylation at smoking-associated genes and at lung cancer-related genes in prediction of lung cancer mortality *Int J Cancer* 139:2482-2492 doi:10.1002/ijc.30374
- Zhang Y, Elgizouli M, Schottker B, Holleczer B, Nieters A, Brenner H (2016c) Smoking-associated DNA methylation markers predict lung cancer incidence *Clin Epigenetics* 8:127 doi:10.1186/s13148-016-0292-4
- Zhu Y, Lin H, Li Z, Wang M, Luo J (2001) Modulation of expression of ribosomal protein L7a (rpL7a) by ethanol in human breast cancer cells *Breast cancer research and treatment* 69:29-38
- Zhu Z, Wang H, Wei Y, Meng F, Liu Z, Zhang Z (2017) Downregulation of PRDM1 promotes cellular invasion and lung cancer metastasis *Tumor Biology* 39:1010428317695929

Figure 1. Manhattan plot for significant DNA-methylation (CpG) probes. CpGs reaching statistical significance are annotated (located above the dotted red horizontal line [$p < 1 \times 10^{-6}$]).

Table 1. Pre-diagnostic characteristics of lung cancer cases and matched controls nested in the CLUE II cohort study

	Lung Cancer Cases	Controls
N	208	222
Median Age (Range, Years)	59 (30-83)	57 (28-81)
Sex		
Male	95 (46%) ^b	98 (44%)
Female	113 (54%)	124 (56%)
Cigarette Smoking Status		
Never Smoker	22 (11%)	22 (10%)
Ever Smoker	80 (38%)	88 (40%)
Current Smoker	106 (51%)	112 (50%)
Median Smoking Intensity (Range, Cig/Day) ^a	20 (0-80)	20 (0-80)
Cigar or Pipe Smoking		
Never	178 (86%)	189 (85%)
Ever	30 (14%)	33 (15%)
Median BMI, kg/m ² (Range)	25.6 (15.5-42.6)	25.9 (18.0-40.9)
Median Time before diagnosis (Range, Years)	14 (0-29)	--

^a 2 missing in cases and 2 missing in controls

^b Percentages were calculated among cases or among controls

Table 2. Association between single CpGs and lung cancer, CLUE II: statistically significant CpGs after multiple comparisons correction ($q < 0.05$)

CpG	Annotation			Methylation Beta Value				Odds Ratio per 1 SD increase (95% CI) ^a		
	Chr	Position	Gene	Mean in	Mean in	SD	Diff/SD	Overall	Time to diagnosis	
				Case	Control				≤ 10 years	> 10 years
cg06581459	chr19	6577668		0.606	0.585	0.055	0.383	2.26(1.70, 3.01)	2.27(1.55, 3.32)	2.15(1.54, 2.99)
cg21432334	chr16	2262293	BRICD5, PGP, C16orf79, RP11	0.819	0.809	0.034	0.312	2.28(1.69, 3.07)	2.37(1.56, 3.60)	2.10(1.48, 2.97)
cg11343375	chr20	61268342	RP13	0.673	0.657	0.047	0.343	2.42(1.74, 3.36)	2.27(1.44, 3.59)	2.44(1.66, 3.59)
cg08739693	chr1	110165726	AMPD2	0.785	0.776	0.036	0.256	2.17(1.62, 2.92)	2.28(1.52, 3.41)	2.21(1.56, 3.13)
cg11744966	chr22	38506771	BAIAP2L2	0.949	0.942	0.014	0.508	1.87(1.47, 2.37)	2.04(1.45, 2.88)	1.85(1.39, 2.46)
cg00237904	chr11	2020314	H19	0.519	0.489	0.076	0.401	1.94(1.50, 2.51)	1.78(1.27, 2.49)	1.87(1.40, 2.50)
cg12483301	chr1	25239107	RUNX3	0.697	0.678	0.058	0.333	2.81(1.88, 4.19)	3.05(1.70, 5.47)	2.72(1.72, 4.29)
cg11385536	chr1	227133801	CABC1, ADCK3	0.719	0.698	0.056	0.369	2.10(1.57, 2.80)	2.00(1.34, 2.98)	2.22(1.57, 3.12)
cg09736286	chr12	122712925	DIABLO	0.867	0.849	0.056	0.33	2.99(1.95, 4.59)	3.10(1.67, 5.74)	3.33(1.97, 5.64)
cg22032485	chr19	18531769	SSBP4	0.632	0.615	0.062	0.269	2.47(1.73, 3.52)	2.63(1.60, 4.30)	2.23(1.49, 3.33)
cg18332677	chr22	20239823	RTN4R	0.826	0.806	0.061	0.327	2.77(1.86, 4.13)	2.34(1.35, 4.07)	2.80(1.75, 4.48)
cg09447640	chr17	42854853	ADAM11	0.664	0.647	0.058	0.303	2.37(1.69, 3.34)	3.44(2.05, 5.79)	1.99(1.36, 2.92)
cg15979932	chr17	55980107	CUEDC1	0.812	0.796	0.04	0.389	2.01(1.53, 2.66)	2.36(1.54, 3.62)	1.84(1.34, 2.53)
cg02211449	chr10	110671609		0.126	0.143	0.046	-0.361	0.40(0.28, 0.58)	0.46(0.28, 0.77)	0.33(0.21, 0.51)
cg19907023	chr2	96803532	ASTL	0.839	0.823	0.043	0.379	2.10(1.56, 2.81)	1.76(1.18, 2.63)	2.23(1.57, 3.15)
cg12796320	chr14	105530592	GPR132	0.504	0.487	0.059	0.279	2.20(1.60, 3.01)	2.02(1.28, 3.18)	2.41(1.65, 3.52)

^a Model adjusted for age at blood draw, sex, smoking status (never, former, current), pack-years based methylation score, BMI, surrogate variables for batch effects, and cell proportions.

Table 3. Association between differential methylated regions (DMR) and lung cancer in CLUE II participants ^a

seqnames	start	end	width	no.cpgs	minfdr	maxbetafc	meanbetafc	overlapping.promoters
chr1	6646079	6646156	78	3	5.15E-06	0.00930873	0.00646122	ZBTB48-007, ZBTB48-002, ZBTB48-003
chr1	28261367	28261706	340	8	2.45E-06	-0.0078132	-0.0039881	SMPDL3B-002, SMPDL3B-001, SMPDL3B-003, SMPDL3B-004, SMPDL3B-006, SMPDL3B-007
chr1	92414221	92414985	765	11	2.03E-10	0.03582019	0.0135806	BRDT-008, BRDT-001, BRDT-002, BRDT-201, BRDT-202, BRDT-004, BRDT-005, BRDT-012, BRDT-003, BRDT-013, BRDT-007, BRDT-016, BRDT-017, BRDT-011, BRDT-009, BRDT-010, BRDT-015, BRDT-019, BRDT-020
chr1	173174650	173174920	271	6	6.58E-06	-0.0114329	-0.005826	TNFSF4-001, TNFSF4-002, TNFSF4-003
chr1	206306094	206306498	405	4	3.37E-06	-0.0156001	-0.0114794	RP11-38J22.6-001, RP11-38J22.6-003
chr2	164204752	164205188	437	5	5.72E-06	0.02526139	0.0170268	NA
chr3	6902824	6903067	244	6	7.67E-06	-0.0183154	-0.0088293	GRM7-001, GRM7-004, GRM7-003, GRM7-201, GRM7-202, GRM7-007, GRM7-002, GRM7-013, GRM7-015
chr3	71275993	71276631	639	5	2.44E-07	0.02400706	0.0153042	NA
chr5	134783229	134783581	353	3	1.91E-06	0.00988655	0.00539191	C5orf20-001
chr6	27859476	27859811	336	3	2.27E-07	0.01242064	0.01074769	HIST1H2BO-001, HIST1H3J-001, HIST1H2AM-001, HIST1H3J-002
chr6	74364495	74364700	206	4	2.63E-06	0.02515135	0.02166436	SLC17A5-001, SLC17A5-201
chr6	106546074	106546824	751	8	4.62E-14	0.02624875	0.01650889	PRDM1-002, PRDM1-007, PRDM1-003, RP1-134E15.3-001
chr6	149805995	149806659	665	11	1.81E-08	0.01785178	0.01265741	ZC3H12D-201, ZC3H12D-003, ZC3H12D-002, ZC3H12D-202, ZC3H12D-004, ZC3H12D-203
chr7	2556962	2557715	754	10	1.14E-08	0.01226153	0.00494892	LFNG-001, LFNG-201, LFNG-002, LFNG-004
chr7	27169674	27171213	1540	20	3.16E-15	0.0371663	0.0205328	HOXA4-001, HOXA4-201, HOXA4-004, HOXA-AS3-003
chr7	110730805	110731201	397	3	3.21E-06	-0.016941	-0.014336	LRRN3-003, LRRN3-001, LRRN3-002, LRRN3-201, LRRN3-004
chr7	153584038	153584748	711	10	1.28E-07	-0.020429	-0.00886	DPP6-001

chr8	101225181	101225902	722	6	2.40E-07	0.01812174	0.01370006	SPAG1-006
chr9	136216873	136217303	431	5	1.35E-07	0.01245556	0.00840205	RPL7A-005, RPL7A-009, MED22-002, MED22-006, SNORD36A-201, SNORD36B-201, MED22-003, SNORD24-201, RPL7A-004, MED22-005, MED22-010, RPL7A-001, RPL7A-003, MED22-004, MED22-009, RPL7A-006, RPL7A-007, RPL7A-008, MED22-011, RPL7A-002, SNORD36C-201
chr9	139745116	139745412	297	5	5.79E-08	0.01178954	0.00625226	PHPT1-001, MAMDC4-005, PHPT1-003, MAMDC4-201, PHPT1-005, PHPT1-004, MAMDC4-001, PHPT1-002, PHPT1-006, PHPT1-201
chr10	63808314	63809170	857	14	4.11E-08	0.0110823	0.00745464	ARID5B-002
chr10	104535792	104536052	261	8	6.65E-06	-0.0120183	-0.0071156	WBP1L-001
chr11	847042	847072	31	2	7.57E-06	0.01410401	0.01263935	TSPAN4-017, TSPAN4-005, TSPAN4-002, TSPAN4-012, TSPAN4-013, TSPAN4-018
chr11	2019436	2021103	1668	31	1.22E-28	0.03449739	0.00903111	H19-201, H19-008, H19-004, H19-002, H19-001, H19-005, H19-009, H19-012, H19-007, H19-013, H19-003, H19-006, H19-014
chr11	128736844	128737467	624	8	1.17E-07	-0.0292492	-0.009666	KCNJ1-002, KCNJ1-001, KCNJ1-005, KCNJ1-201, KCNJ1-003, KCNJ1-006
chr12	54653403	54653788	386	4	6.51E-07	-0.0179083	-0.0145844	CBX5-004, CBX5-003, CBX5-005
chr16	2261947	2262293	347	6	5.27E-11	0.01398617	0.00891788	BRICD5-001, RP11-304L19.8-001, PGP-003, BRICD5-002, BRICD5-003, BRICD5-004
chr17	17464853	17465682	830	8	2.21E-09	0.02334441	0.01559593	NA
chr17	34207332	34208012	681	6	4.60E-10	0.01767462	0.01343757	CCL5-001, CCL5-002, CCL5-003
chr17	55362520	55363072	553	4	5.30E-07	-0.0150749	-0.0124685	MSI2-005
chr19	17610751	17610785	35	2	2.91E-06	-0.0122955	-0.0107149	SLC27A1-009, SLC27A1-010
chr19	18209584	18209746	163	5	3.52E-06	-0.0104941	-0.0047016	MAST3-001, IL12RB1-005, IL12RB1-006, IL12RB1-002
chr20	36148133	36149750	1618	41	2.78E-17	0.02478338	0.0137516	NNAT-001, NNAT-002, BLCAP-009, BLCAP-010, BLCAP-005

chr20	62200050	62200711	662	6	6.94E-10	0.01497656	0.01241171	HELZ2-001, HELZ2-006
chr20	62688192	62688896	705	9	1.60E-06	0.01441642	0.01071442	TCEA2-008, RP13-152O15.5-001
chr21	35898717	35899468	752	8	5.29E-08	0.0200447	0.01593455	RCAN1-003, RCAN1-201, RCAN1-005, RCAN1-007
chr21	38362230	38363160	931	10	1.08E-08	0.04123433	0.01288671	HLCS-001, HLCS-004
chr22	31002362	31003283	922	14	1.55E-09	-0.0113028	-0.0050389	TCN2-001, PES1-003, PES1-002, TCN2-002, TCN2-003, TCN2-004, TCN2-005, PES1-012, PES1-007
chr22	50738160	50738890	731	3	2.49E-08	0.02463834	0.02309907	PLXNB2-012
chr22	51176117	51176734	618	7	1.57E-07	0.01094617	0.00757197	ACR-001, AC000036.4-001, ACR-003, ACR-002

^a Model adjusted for age at blood draw, sex, smoking status (never, former, current), pack-years based methylation score, BMI, surrogate variables for batch effects, and cell proportions.

Table 4. Odds ratio for top 10 CpGs (q-value<0.05) in two differentially methylated region (H19 and HOXA4) and lung cancer risk, and stratified by time to diagnosis in CLUE II participants

region	CpG	chr	pos	methylation beta value				Odds Ratio per 1 SD increase (95% CI) ^a		
				mean in cases	mean in controls	SD	Diff/SD	overall	by time to diagnosis	
									≤ 10 years	>10 years
Region 1 (gene: H19)	cg00237904	chr11	2020314	0.519	0.489	0.076	0.401	1.94(1.50, 2.51)	1.78(1.27, 2.49)	1.87(1.40, 2.50)
	cg27300742	chr11	2020129	0.569	0.550	0.067	0.288	1.86(1.43, 2.42)	1.57(1.10, 2.23)	1.97(1.45, 2.68)
	cg01895612	chr11	2020286	0.520	0.511	0.052	0.174	1.89(1.40, 2.56)	1.49(1.02, 2.17)	2.03(1.42, 2.91)
	cg24605090	chr11	2019943	0.562	0.549	0.060	0.216	1.86(1.39, 2.49)	1.65(1.09, 2.50)	2.02(1.43, 2.86)
	cg23476401	chr11	2020296	0.563	0.548	0.072	0.214	1.69(1.30, 2.19)	1.37(0.99, 1.90)	1.78(1.31, 2.41)
	cg25281616	chr11	2020279	0.588	0.580	0.056	0.151	1.75(1.30, 2.36)	1.41(0.96, 2.06)	1.84(1.30, 2.60)
	cg18362496	chr11	2019930	0.492	0.484	0.061	0.128	1.74(1.27, 2.38)	1.45(0.95, 2.21)	1.86(1.29, 2.69)
	cg03996735	chr11	2020104	0.744	0.729	0.055	0.274	1.42(1.14, 1.77)	1.22(0.92, 1.64)	1.58(1.21, 2.05)
	cg25579157	chr11	2020549	0.552	0.549	0.022	0.130	1.52(1.17, 1.98)	1.43(0.99, 2.06)	1.58(1.16, 2.16)
Region 2 (gene: HOXA4, HOXA-AS3, HOXA-AS2, HOXA3)	cg01539474	chr11	2019862	0.639	0.633	0.064	0.096	1.55(1.15, 2.09)	1.63(1.06, 2.50)	1.62(1.12, 2.35)
	cg11532431	chr7	27169674	0.810	0.781	0.100	0.282	1.45(1.17, 1.81)	1.70(1.22, 2.38)	1.38(1.07, 1.77)
	cg04317399	chr7	27170313	0.324	0.307	0.091	0.184	1.43(1.13, 1.82)	1.67(1.19, 2.36)	1.38(1.05, 1.82)
	cg17591595	chr7	27171051	0.732	0.720	0.074	0.167	1.39(1.12, 1.74)	1.33(0.98, 1.82)	1.51(1.16, 1.96)
	cg22997113	chr7	27170241	0.589	0.560	0.139	0.213	1.42(1.13, 1.78)	1.53(1.09, 2.13)	1.42(1.10, 1.84)
	cg07317062	chr7	27170388	0.393	0.371	0.123	0.182	1.40(1.11, 1.77)	1.50(1.08, 2.08)	1.39(1.06, 1.81)
	cg17457637	chr7	27170717	0.369	0.355	0.093	0.145	1.39(1.10, 1.76)	1.53(1.09, 2.15)	1.38(1.05, 1.80)
	cg00562553	chr7	27169740	0.875	0.862	0.073	0.184	1.34(1.08, 1.66)	1.47(1.07, 2.01)	1.34(1.05, 1.72)
	cg03724423	chr7	27170755	0.356	0.339	0.096	0.174	1.37(1.09, 1.73)	1.46(1.05, 2.04)	1.35(1.04, 1.76)
	cg23884241	chr7	27169957	0.562	0.556	0.066	0.091	1.40(1.10, 1.79)	1.53(1.07, 2.19)	1.33(1.02, 1.75)

cg24169822	chr7	27170994	0.434	0.418	0.105	0.146	1.37(1.09, 1.72)	1.49(1.07, 2.07)	1.35(1.04, 1.76)
------------	------	----------	-------	-------	-------	-------	------------------	------------------	------------------

^a Model adjusted for age at blood draw, sex, smoking status (never, former, current), pack-years based methylation score, BMI, surrogate variables for batch effects, and cell proportions.