Title: Cohort profile: COVID-19 in a cohort of pregnant women and their descendants, the MOACC-19 study

Authors: Javier Llorca[1,2], Carolina Lechosa-Muñiz[1,3], Pilar Gortázar[1,3], María Fernández-Ortiz[4], Yolanda Jubete[1,3], María J. Cabero[1,3,4] and the MOACC-19 group

Affiliations:

1. Universidad de Cantabria, Santander, Spain
2. CIBER de Epidemiología y Salud Pública (CIBERESP), Madrid, Spain
3. University Hospital Marques de Valdecilla, Santander, Spain
4. IDIVAL, Santander, Spain

Correspondence:

Javier Llorca
Facultad de Medicina
Universidad de Cantabria
Avda. Herrera Oria s/n
39011 Santander
Spain
Phone # 34-942 201 993
E-mail: javier.llorca@unican.es

Members of the MOACC-19 group:

Jéssica Alonso-Molero[1,4], Bárbara Arozamena[3], Laura Conde-Gil[3], Elsa Cornejo del Río[3], Rocío Cuesta-González[3], Trinidad Dierssen-Sotos[1,2,4], Pelayo Frank de Zulueta[3], Inés Gómez-Acebo[1,2,4], Coral Llano-Ruiz[3], Lorena Lasarte-Oria[3], Sonia López-Gómez[3], Sonia Mateo-Sota[3], Victoria Orallo[3], Rosa Pardo[3], Daniel Pérez González[3], María Sáez de Adana Herrero[3].

Supplementary Table 1. Characteristics of women tested positive for SARS-CoV-2 infection via RT-PCR, subcohorts 1 and 2

Variable	Category	N (%)
Study cohort	Sub-cohort 1 (Women delivering before 26/May/2020)	5
	Sub-cohort 2 (Women delivering from 26/May/2020 on)	9
Partner RT-PCR	Negative	7
	Positive	7
Age	<25 years	1 (7.1)
	25-29 years	3 (21.4)
	30-34 years	4 (28.6)
	35-39 years	4 (28.6)
	≥40 years	2 (14.3)
Education level	Primary studies	3 (21.4)
	Secondary studies	2 (14.3)
	Vocational training	2 (14.3)
	University studies	7 (50.0)
Work status	Unemployed / no active worker	5 (35.7)
	Employed	9 (64.3)
	Student	0
Occupation	Health care worker	4 (28.6)
	Working in restaurant or commerce	2 (14.3)
	Others	3 (21.4)
Fertilization type	Natural	12 (92.3)
	Artificial insemination	1 (7.7)
	In vitro fertilization (own ovule)	0
	In vitro fertilization (ovodonation)	0
	Missing	1 (7.7)
Smoker in pregnancy	No	14 (100.0)
	Yes	0
Alcohol consumption in pregnancy	No	14 (100.0)
	Yes	0
Parity	1	9 (64.3)
	2	1 /7.1)
	≥3	4 (28.6)
Type of delivery	Eutocic	8 (57.1)
	Caesarean rate	4 (28.6)
	Instrumental	2 (14.3)
Age (years)	mean±sd (min, max)	32.9±5.1 (23 – 40)
Gestational age leave work (weeks)	mean±sd (min, max)	23.4±12.7 (0 – 36)
Pregestational BMI (kg/m^2)	mean±sd (min, max)	23.3±4.0 (18.6 – 32.0)

Gestational weight gain (kg)	mean±sd (min, max)	11.2±4.1 (8 – 22)
Symptoms associated with COVID-19	Fever	3
	Chills	3
	Intense tiredness	2
	Sore throat	3
	Cough	3
	Short of breathing	2
	Headache	3
	Nausea	1
	Loss of sense	2

Supplementary Table 2. Characteristics of children born from women tested positive to SARS-CoV-2 infection using RT-PCR

Variable	Category	N = 7 (%)
Gestational age at birth	**<34 weeks**	0
	34-36 weeks	0
	≥37 weeks	14 (100.0
Birth weight	**<2500 g**	0
	2500-4000 g	14 (100.0)
	>4000 g	0
Gender	**Women**	4 (28.6)
	Men	10 (71.4)
Twin or singleton	**Singleton**	14 (100.0)
	Twin	0
Breastfeeding at hospital discharge	**Exclusive breastfeeding**	4 (28.6)
	Mixed	6 (42.9)
	Infant formula	4 (28.6)
Neonatology admission	**No**	11 (78.6)
	Yes	3 (21.4)
Birth size	**mean±sd (min, max)**	50.0±2.4 (47 – 53)
Birth cephalic perimeter	**mean±sd (min, max)**	34.7±1.5 (33 – 39)
Apgar 1'	**mean±sd (min, max)**	8.4±1.4 (4 – 9)
Apgar 5'	**mean±sd (min, max)**	9.2±0.9 (7 – 10)
pH at birth	**mean±sd (min, max)**	7.27±0.05 (7.20 – 7.36)
SARS-CoV-2 via RT-PCR		All negative