


Cancer* (ICD-10 prefix)	Cancer Cases		Fatalities Among		Mean HR	HR FDR
	Total	COVID-19+	COVID-19+ (%)	COVID-19- (%)		
Localized cancer	3688	235	19	2	9.85	5.9x10 ⁻³²
Lymphoma (C81–86,C884)	292	20	45	4.04	21.28	1.0x10 ⁻⁹
Melanoma (C43)	206	13	15	1.04	20.31	9.2x10 ⁻³
Leukemia (C91–95)	192	22	50	3.53	20.23	3.7x10 ⁻⁸
Plasma cell neoplasm (C90)	151	7	29	3.47	14.25	1.3x10 ⁻²
Colorectal cancer (C18–20)	431	25	20	1.72	12.04	6.2x10 ⁻⁵
Breast cancer (C50)	594	36	6	0.54	10.21	1.4x10 ⁻²
Kidney cancer (C64–65)	110	9	22	1.98	8.84	3.7x10 ⁻²
Lung cancer (C34)	198	12	33	5.91	6.57	3.5x10 ⁻³
Prostate cancer (C61)	816	59	7	1.59	4.20	1.5x10 ⁻²
Bladder cancer (C67)	399	26	8	2.14	4.07	7.8x10 ⁻²
Distant metastasis	918	53	38	9	5.08	9.6x10 ⁻¹⁰
Melanoma (C43)	42	2	100	8	49.37	5.7x10 ⁻³
Prostate cancer (C61)	136	8	62	4	22.11	6.8x10 ⁻⁶
Ovarian cancer (C56)	91	7	43	4	13.04	3.5x10 ⁻³
Lung cancer (C34)	105	6	50	15	5.95	1.3x10 ⁻²

Hazard Ratio of COVID-19 Fatalities



Notes:

- 1) All stratified analyses use COVID-19 plus covariates sex and age.
- 2) Localized cancer type stratified studies included those with at least 100 localized cases.
- 3) Distant metastasis cancer type stratified studies included those with at least 40 metastasized cases, except colorectal, breast, and pancreatic cancers, which had no COVID-19 associated deaths.