

Supplementary Table 1:


Combinatorial sample barcoding

Patient	Patient-specific antibodies	Timepoint-specific antibodies			
		Timepoint 1	Timepoint 2	Timepoint 3	Timepoint 4
2.6	Hashtag 1	anti-CD11a	anti-CD18	anti-CD45	
3.1	Hashtag 2	anti-CD11a	anti-CD18	anti-CD45	anti-CD20
3.12	anti-HLA-ABC	anti-CD11a	anti-CD18		
3.2	anti-HLA-DR	anti-CD11a	anti-CD18		
3.3	Hashtag 4	anti-CD11a	anti-CD18	anti-CD45	
3.4	Hashtag 5	anti-CD11a	anti-CD18	anti-CD45	


* Barcoded antibodies against human surface antigens with TotalSeq C oligonucleotides, specifically compatible with the Single Cell 5' workflow of 10X Genomics.

Supplementary Figure 1:

Binding of fluorescence-labeled BKPyV VLPs to B cells from healthy donor, seronegative KTx, seropositive KTx



Supplementary Figure 2- Viral load, neutralizing antibody titres, and summary clinical data of patients selected for scRNA seq experiment




Patient	Sex	Age at KTx	BKPyV genotype
3.1	F	60-70	Ib2
3.2	M	60-70	Ib2
3.3	M	60-70	IV
3.4	M	60-70	Ib2
2.6	F	50-60	Ib1
3.12	M	50-60	Ib2


Supplementary Figure 3:

Heavy and light chain v-gene usage in total and BKPyV-specific B-cells


Patient 3.1




Patient 3.1




Patient 3.4




Patient 3.4




Patient 2.6




Patient 2.6



Patient 3.2



Patient 3.2



■ BKPyV-specific B-cells ■ Total B-cells

Supplementary Figure 4:

Binding of fluorescence-labeled BKPyV VLPs to susceptible 293TT cells

