Do Assisted Living Facilities That Offer a Dementia Care Program Differ from Those That Do Not? A Population-Level Cross-Sectional Study in Ontario, Canada

Derek R. Manis, MHSc^{a-c}; Ahmad Rahim, MSc^c; Jeffrey W. Poss, PhD^d; Iwona A. Bielska, PhD^{a,b}; Susan E. Bronskill, PhD^{c,e-g}; Jean-Éric Tarride, PhD^{a,b}; Julia Abelson, PhD^{a,b}; and Andrew P. Costa, PhD^{a-c,h-j}

- a. Centre for Health Economics and Policy Analysis, McMaster University, Hamilton, Ontario, Canada
- b. Department of Health Research Methods, Evidence, and Impact, McMaster University, Hamilton, Ontario, Canada
- c. ICES, Toronto, Ontario, Canada
- d. School of Public Health and Health Systems, University of Waterloo, Waterloo, Ontario, Canada
- e. Institute of Health Policy, Management & Evaluation, University of Toronto, Toronto, Ontario, Canada
- f. Women's College Research Institute, Toronto, Ontario, Canada
- g. Sunnybrook Research Institute, Toronto, Ontario, Canada
- h. Centre for Integrated Care, St. Joseph's Health System, Hamilton, Ontario, Canada
- i. Schlegel Research Institute for Aging, Waterloo, Ontario, Canada
- j. Department of Medicine, McMaster University, Hamilton, Ontario, Canada

Corresponding Author: Derek R. Manis, McMaster University, 1280 Main St W, CRL-201, Hamilton, ON L8S 4K1, Canada. (manisd@mcmaster.ca) Twitter: [@DerekManis](https://twitter.com/DerekManis)

Running Title: "DEMENTIA CARE IN ASSISTED LIVING FACILITIES"

Keywords: Assisted Living Facilities; Retirement Homes; Dementia Care; Canada

Funding: This research did not receive any funding from agencies in the public, commercial, or not-for-profit sectors.

Word Count: 1,811 (Max: 2,000)

Abstract: 287 (Max: 300)

References: 30 (Max: 30)

Exhibitions: 2 Tables (Max: 3) (2 Supplemental Tables in Appendix)

BRIEF SUMMARY

Question: Do licensed assisted living facilities that provide a dementia care program differ from those that do not in Ontario, Canada?

Findings: Assisted living facilities that provide a dementia care program house many older adults, have many suites, and offer at least five of the other 12 regulated care services.

Meaning: Assisted living facilities that provide a dementia care program are larger and provide an array of care services.

Disclosures: We have no perceived or actual conflicts, financial or otherwise, to disclose.

Acknowledgments: This study was supported by ICES, which is funded by an annual grant from the Ontario Ministry of Health and Long-Term Care (MOHLTC). The opinions, results and conclusions reported in this paper are those of the authors and are independent from the funding sources. No endorsement by ICES or the Ontario MOHLTC is intended or should be inferred. We extend our gratitude to Paul Pham, Adriane Castellino, and Chloe Ma at the Retirement Homes Regulatory Authority for sharing the registry of licensed assisted living facilities. We also extend our thanks to Nathan M. Stall for his comments on an earlier draft of this manuscript.

Author Contributions: DRM and APC conceptualized the study. AR created the dataset, and DRM conducted the statistical analyses. DRM wrote the manuscript, and all authors critically read and contributed to it.

Data Availability: The dataset from this study is held securely in coded form at ICES. While data sharing agreements prohibit ICES from making the data set publicly available, access may be granted to those who meet pre-specified criteria for confidential access, available at www.ices.on.ca/DAS. The full data set creation plan and underlying analytic code are available from the corresponding author upon request, understanding that the programs may rely upon coding templates or macros that are unique to ICES.

ABSTRACT

Objective: To identify the characteristics of licensed assisted living facilities that provide a dementia care program compared to assisted living facilities that do not provide such a program.

Design, Setting, and Participants: Population-level cross-sectional study in Ontario, Canada on all licensed assisted living facilities in 2018 ($n = 738$).

Methods: Data on facility-level characteristics (e.g., resident and suite capacities, etc.) and the provision of other provincially regulated care services (e.g., pharmacist and medical services, skin and wound care, etc.) attributed to licensed assisted living facilities were examined.

Multivariable Poisson regression with robust standard errors was used to model the characteristics of assisted living facilities associated with the provision of a dementia care program.

Results: There were 123 (16.7%) assisted living facilities that provided a dementia care program. Nearly half of these facilities had a resident capacity exceeding 140 older adults (44.7%) and more than 115 suites (46.3%). All assisted living facilities that provided a dementia care program also provided nursing services, meals, assistance with bathing and hygiene, and administered medications. After adjusting for facility characteristics and other provincially regulated care services, the prevalence of a dementia care program was nearly three times greater in assisted living facilities that offered assistance with feeding than those that did not (Prevalence Ratio [PR] = 2.91, 95% Confidence Interval [CI] 1.98 to 4.29), and almost twice as great among assisted living facilities that provided medical services than those that did not (PR = 1.78, 95% CI 1.00 to 3.17).

Conclusions: A dementia care program was more prevalent in assisted living facilities that housed many older adults, had many suites, and provided at least five of the other 12 regulated

DEMENTIA CARE IN ASSISTED LIVING FACILITIES

4

- 24 care services. These findings deepen the understanding of specialized care for dementia in
- 25 assisted living facilities.

INTRODUCTION

Dementia affects more than half of all residents housed in assisted living facilities.¹⁻³ Older adults living with dementia are more likely to experience injuries requiring acute care, be diagnosed with pneumonia, and encounter difficulties with eating.^{4,5} Care for dementia is expensive and a widely cited reason for older adults requiring placement in a nursing home.⁶⁻⁹ Specialized care for older adults living with dementia, such as a dementia care program, has demonstrated reductions in acute health service use and long-stay transitions to a nursing home.^{8,10}

Assisted living facilities are referred to as retirement homes in the province of Ontario, Canada. Similar to assisted living facilities in the United States that are regulated at the state-level,¹¹ retirement homes are regulated by an independent, not-for-profit regulator (i.e., Retirement Homes Regulatory Authority [RHRA]) in Ontario since 2011.¹² The retirement home sector in Ontario has a resident capacity equivalent to that of its nursing home sector (i.e., more than 70,000 older adults),¹² which demonstrates that it is a sizeable, important sector providing supportive care to older adults. Unlike nursing homes, residency in a retirement home is exclusively financed through private, out-of-pocket payments by residents and/or their family caregivers.¹²⁻¹⁴

Much of the literature on dementia care in assisted living facilities addresses health service use among residents living with dementia, managing staff, and state-level regulations for dementia care.^{1,11,15-17} Studies explicitly investigating the characteristics of retirement homes or assisted living facilities that provide specialized care for dementia (i.e., a dementia care program), and how these characteristics compare to those that do not, are absent. The findings from such studies are important for identifying case mix and examining scope and breadth of

care for older adults with complex care needs. A growing proportion of residents of assisted living facilities live with dementia,¹⁸ and improving the understanding of dementia care programs in assisted living facilities contributes to informing the sector, community-based dementia care, and national dementia care strategies. The objective of this study is to identify the characteristics of licensed assisted living facilities that provide a dementia care program compared to assisted living facilities that do not provide such a program in Ontario, Canada.

METHODS

Study Design and Setting

A population-level cross-sectional study was conducted at ICES in Ontario, Canada. ICES is an independent, non-profit research institute funded by an annual grant from the Ontario Ministry of Health and Long-Term Care. As a prescribed entity under Ontario's privacy legislation, ICES is authorized to collect and use health care data for the purposes of health system analysis, evaluation, and decision support. Secure access to these data is governed by policies and procedures that are approved by the Information and Privacy Commissioner of Ontario. The REporting of studies Conducted using Observational Routinely-collected health Data (RECORD) statement guideline was followed (Appendix Table 2).¹⁹

Data

A list of all licensed assisted living facilities in Ontario in 2018 was obtained from the public register of the RHRA and imported to ICES ($n = 757$). The postal code of each assisted living facility was linked to Statistics Canada's Postal Code Conversion file, which is a specialized macro for use with health system administrative datasets containing postal codes. This macro is based on 2016 Census information, flags communities with a population less than 10,000 individuals as rural, and includes related data from Canada Post Corporation.²⁰ Nineteen

assisted living facilities ($n = 19$) were removed from the analysis because of missing facility-level and postal code data.

Exposures

The exposures of interest are facility-level characteristics (i.e., urban location, resident capacity, total suites, chain status, residual home status, and co-location with a nursing home) and the other 12 provincially regulated care services provided in an assisted living facility (Appendix Table 1).

Outcome

The primary outcome is whether the assisted living facility provided a dementia care program. Dementia care programs in assisted living facilities in Ontario are regulated to include communication strategies, mental stimulation activities, health and wellness monitoring and promotion, and identification of triggers for responsive behaviours.²¹ These programs must also be supervised by a regulated health care professional (e.g., registered nurse, physician, etc.), align with current evidence and best practices for dementia care, and be evaluated annually.²¹

Statistical Analysis

Counts, percentages, and standardized differences were calculated to describe the facility-level and care service characteristics of assisted living facilities that provided, and did not provide, a dementia care program.²² Multivariable Poisson regression with robust standard errors was used to model unadjusted and adjusted estimates with 95% confidence intervals to identify the characteristics of assisted living facilities associated with the provision of a dementia care program.^{23,24} Tests were two-tailed, and the level of statistical significance was set at $\alpha = 0.05$. The deviance goodness-of-fit test was calculated to assess whether the Poisson regression model was appropriate. Variance inflation factors were calculated to assess for multicollinearity.²⁵ Data

set processing was conducted in SAS Enterprise 9.4 (Cary, NC, USA) and statistical analyses were conducted in Stata MP 16.1 (College Station, TX, USA).

RESULTS

There were 738 licensed assisted living facilities in Ontario in 2018 ($n = 738$). Of these, 123 provided a dementia care program (16.7% versus 83.3% no dementia care program), and almost all were located in an urban area (92.7% versus 82.6% no dementia care program) (Table 1). Nearly half of these assisted living facilities had a resident capacity of 140 or more (44.7% versus 21.6% no dementia care program) and had more than 115 suites (46.3% versus 20.8% no dementia care program). All assisted living facilities that provided a dementia care program also provided nursing services, meals, assistance with bathing and hygiene, and administered medications ($n = 123$). Many of the standardized differences between assisted living facilities that provided a dementia care program and those that did not exceeded 10%, which indicated that assisted living facilities that provided a dementia care program were systematically different from those that did not.²²

Assistance with bathing and hygiene, provision of meals, administration of medication, and nursing services were removed from the adjusted model because of collinearity, and there was no evidence of multicollinearity in the adjusted model (i.e., variance inflation factors equal to or greater than a value of 10).²⁵ The deviance goodness-of-fit statistic was not statistically significant. After adjusting for facility characteristics and regulated care services, the prevalence of a dementia care program was almost three times greater in assisted living facilities with 115 or more suites (Prevalence Ratio [PR] = 2.78, 95% Confidence Interval [CI] 1.09 to 7.07) compared to assisted living facilities with 41 or fewer suites (Table 2). The prevalence of a dementia care program was nearly three times greater in assisted living facilities that provided assistance with

feeding (PR = 2.91, 95% CI 1.98 to 4.29), and the prevalence of a dementia care program was almost twice as great in assisted living facilities that provided medical services (PR = 1.78, 95% CI 1.00 to 3.17), compared to assisted living facilities that did not provide these care services.

DISCUSSION

Assisted living facilities that provided a dementia care program were systematically different from those that did not provide such a program. Specifically, assisted living facilities in Ontario that provided a dementia program had large resident capacities, many suites, and provided, at a minimum, nursing services, meals, assistance with bathing and hygiene, and administered medications. The prevalence of a dementia care program in an assisted living facility was greater in assisted living facilities where assistance with feeding and medical services were also provided.

More than 90% of assisted living facilities that provided a dementia care program were located in urban communities. Consistent with existing literature, this finding raises important equity considerations for older adults living with dementia in assisted living facilities located in rural and remote regions.²⁶ Rural assisted living facilities house fewer older adults and are more likely to have deficiencies in care provision than urban ones, including challenges with retaining appropriate care staff and resources to meet the needs of residents.²⁷ The use of videoconferencing and other information technology resources to provide dementia care should be considered to improve access to care for individuals living with dementia in rural and remote areas.²⁸

Most assisted living facilities that provided a dementia care program had capacity for more than 140 older adults and had more than 115 suites. Current practices for designing settings specifically for older adults living with dementia emphasize larger spaces that are not

characteristic of institutionalized congregate care,²⁹ and the presence and statistically significant association of many suites in assisted living facilities that provide a dementia care program aligns with the literature. In addition, this may indicate that many assisted living facilities that provide a dementia care program are large complexes, likely attributed to chains.

Given the challenges that older adults living with dementia face with respect to eating,⁵ it is expected that assistance with feeding would be a prevalent care service offered alongside a dementia care program in an assisted living facility. Moreover, the complex and intersecting care needs of older adults living with dementia, which includes polypharmacy,³⁰ underscores the need for a physician to monitor and evaluate the care plan. As such, the prevalence of medical services in assisted living facilities that offer a dementia care program is also expected.

As the assisted living sector is privately financed in Ontario, this study makes an important contribution to the literature to define the sector by modeling facility-level characteristics associated with the provision of a dementia care program. The findings are relevant to clinicians and policymakers actively considering dementia care options in communities to support older adults living with dementia and their caregivers. Family caregivers and consumers of assisted living services will also be interested in these findings to inform their decisions for care.

A limitation to this study is that the fees older adults residing in assisted living facilities pay each month and per care service could not be included in the adjusted model. This is due, in part, to the inability to capture this information in existing administrative health system data. Moreover, these data are not publicly available on the websites of assisted living facilities, through their member associations, or available to the RHRA through regulatory reporting requirements. Another limitation is that this study is descriptive; as such, no causal or temporal

claims can be made about the associations between the facility-level characteristics of assisted living facilities and the provision of a dementia care program. As with all secondary analyses of data, the data used in this study are susceptible to misclassification bias.

CONCLUSION

This study identified and compared facility-level characteristics of licensed assisted living facilities that provided a dementia care program to those that did not in Ontario, Canada in 2018. Assisted living facilities that offered a dementia care program housed more older adults and provided more care services. Future research might consider investigating the underlying differences in populations between residents of these facilities and their health outcomes attributed to care services provided in assisted living facilities.

REFERENCES

1. Thomas KS, Zhang W, Cornell PY, Smith L, Kaskie B, Carder PC. State Variability in the Prevalence and Healthcare Utilization of Assisted Living Residents with Dementia. *J Am Geriatr Soc.* 2020;68(7):1504-1511. doi:10.1111/jgs.16410
2. Thomas KS, Dosa D, Gozalo PL, et al. A Methodology to Identify a Cohort of Medicare Beneficiaries Residing in Large Assisted Living Facilities Using Administrative Data. *Med Care.* 2018;56(2):e10-e15. doi:10.1097/MLR.0000000000000659
3. Zimmerman S, Sloane PD, Reed D. Dementia prevalence and care in assisted living. *Health Aff (Millwood).* 2014;33(4):658-666. doi:10.1377/hlthaff.2013.1255
4. Rosa TD, Possin KL, Bernstein A, et al. Variations in Costs of a Collaborative Care Model for Dementia. *J Am Geriatr Soc.* 2019;67(12):2628-2633. doi:10.1111/jgs.16076
5. Mitchell SL, Teno JM, Kiely DK, et al. The Clinical Course of Advanced Dementia. *N Engl J Med.* 2009;361(16):1529-1538. doi:10.1056/NEJMoa0902234
6. Ng R, Lane N, Tanuseputro P, et al. Increasing Complexity of New Nursing Home Residents in Ontario, Canada: A Serial Cross-Sectional Study. *Journal of the American Geriatrics Society.* 2020;68(6):1293-1300. doi:10.1111/jgs.16394
7. Mondor L, Maxwell CJ, Hogan DB, et al. The Incremental Health Care Costs of Frailty Among Home Care Recipients With and Without Dementia in Ontario, Canada: A Cohort Study. *Medical Care.* 2019;57(7):512-520. doi:10.1097/MLR.0000000000001139
8. Jennings LA, Laffan AM, Schlissel AC, et al. Health Care Utilization and Cost Outcomes of a Comprehensive Dementia Care Program for Medicare Beneficiaries. *JAMA Intern Med.* 2019;179(2):161-166. doi:10.1001/jamainternmed.2018.5579
9. Morton-Chang F, Williams AP, Berta W, Laporte A. Towards a Community-Based Dementia Care Strategy: How do We Get There from Here? *World Health & Population.* 2019;18(1):6-39. doi:10.12927/whp.2019.26062
10. Jennings LA, Hollands S, Keeler E, Wenger NS, Reuben DB. The Effects of Dementia Care Co-Management on Acute Care, Hospice, and Long-Term Care Utilization. *J Am Geriatr Soc.* 2020;(7503062, h6v). doi:10.1111/jgs.16667
11. Nattinger MC, Kaskie B. Determinants of the Rigor of State Protection Policies for Persons With Dementia in Assisted Living. *J Aging Soc Policy.* 2017;29(2):123-142. doi:10.1080/08959420.2016.1236324
12. Roblin B, Deber R, Kuluski K, Silver MP. Ontario's Retirement Homes and Long-Term Care Homes: A Comparison of Care Services and Funding Regimes. *Can J Aging.* 2019;38(2):155-167. doi:10.1017/S0714980818000569

13. Silver BC, Grabowski DC, Gozalo PL, Dosa D, Thomas KS. Increasing Prevalence of Assisted Living as a Substitute for Private-Pay Long-Term Nursing Care. *Health Serv Res.* 2018;53(6):4906-4920. doi:10.1111/1475-6773.13021
14. Grabowski DC, Stevenson DG, Cornell PY. Assisted living expansion and the market for nursing home care. *Health Services Research.* 2012;47(6):2296-2315. doi:10.1111/j.1475-6773.2012.01425.x
15. Simmons SF, Coelho CS, Sandler A, Schnelle JF. A System for Managing Staff and Quality of Dementia Care in Assisted Living Facilities. *J Am Geriatr Soc.* 2018;66(8):1632-1637. doi:10.1111/jgs.15463
16. Simmons SF, Coelho CS, Sandler A, Shah AS, Schnelle JF. Managing Person-Centered Dementia Care in an Assisted Living Facility: Staffing and Time Considerations. *Gerontologist.* 2018;58(4):e251-e259. doi:10.1093/geront/gnx089
17. Carder PC. State Regulatory Approaches for Dementia Care in Residential Care and Assisted Living. *Gerontologist.* 2017;57(4):776-786. doi:10.1093/geront/gnw197
18. Cornell PY, Zhang W, Smith L, Fashaw S, Thomas KS. Developments in the Market for Assisted Living: Residential Care Availability in 2017. *J Am Med Dir Assoc.* 2020;21(11):1718-1723. doi:10.1016/j.jamda.2020.08.011
19. Benchimol EI, Smeeth L, Guttman A, et al. The REporting of studies Conducted using Observational Routinely-collected health Data (RECORD) Statement. *PLOS Medicine.* 2015;12(10):e1001885. doi:10.1371/journal.pmed.1001885
20. Government of Canada SC. Postal Code OM Conversion File Plus (PCCF+). Published March 6, 2017. Accessed November 24, 2020. <https://www150.statcan.gc.ca/n1/en/catalogue/82F0086X>
21. Government of Ontario. *O. Reg. 166/11: GENERAL.*; 2014. Accessed June 17, 2020. <https://www.ontario.ca/laws/regulation/110166#BK54>
22. Austin PC. Using the Standardized Difference to Compare the Prevalence of a Binary Variable Between Two Groups in Observational Research. *Communications in Statistics - Simulation and Computation.* 2009;38(6):1228-1234. doi:10.1080/03610910902859574
23. Barros AJ, Hirakata VN. Alternatives for logistic regression in cross-sectional studies: an empirical comparison of models that directly estimate the prevalence ratio. *BMC Med Res Methodol.* 2003;3(1):1-13. doi:10.1186/1471-2288-3-21
24. Tamhane AR, Westfall AO, Burkholder GA, Cutter GR. Prevalence Odds Ratio versus Prevalence Ratio: Choice Comes with Consequences. *Stat Med.* 2016;35(30):5730-5735. doi:10.1002/sim.7059
25. Kleinbaum DG, Kupper LL, Nizam A, Rosenberg ES. *Applied Regression Analysis and Other Multivariable Methods.* 5th ed. Cengage Learning; 2014.

26. Giebel C. “Current dementia care: what are the difficulties and how can we advance care globally?”. *BMC Health Serv Res*. 2020;20(1):414. doi:10.1186/s12913-020-05307-1
27. June JW, Meng H, Dobbs D, Hyer K. Using Deficiency Data to Measure Quality in Assisted Living Communities: A Florida Statewide Study. *J Aging Soc Policy*. 2020;32(2):125-140. doi:10.1080/08959420.2018.1563471
28. Wallack EM, Harris C, Ploughman M, Butler R. Telegerontology as a Novel Approach to Address Health and Safety by Supporting Community-Based Rural Dementia Care Triads: Randomized Controlled Trial Protocol. *JMIR Res Protoc*. 2018;7(2):e56. doi:10.2196/resprot.8744
29. Calkins MP. From Research to Application: Supportive and Therapeutic Environments for People Living With Dementia. *Gerontologist*. 2018;58(suppl_1):S114-S128. doi:10.1093/geront/gnx146
30. Mueller C, Molokhia M, Perera G, et al. Polypharmacy in people with dementia: Associations with adverse health outcomes. *Experimental Gerontology*. 2018;106:240-245. doi:10.1016/j.exger.2018.02.011

Table 1. Descriptive Characteristics of Licensed Assisted Living Facilities in 2018 ($n = 738$)

	Dementia Care Program		Standardized Difference
	Yes	No	
n (%)	123 (16.7)	615 (83.3)	
Facility Characteristics, n (%)			
Urban Location	114 (92.7)	508 (82.6)	0.309
Facility Capacity			0.583
6-49	24 (19.5)	155 (25.2)	
50-86	14 (11.4)	171 (27.8)	
87-139	30 (24.4)	156 (25.4)	
140+	55 (44.7)	133 (21.6)	
Total Suites			0.615
6-41	20 (16.3)	163 (26.5)	
42-70	16 (13.0)	168 (27.3)	
71-114	30 (24.4)	156 (25.4)	
115+	57 (46.3)	128 (20.8)	
Chain Status	74 (60.2)	281 (45.7)	0.293
Residential Home	8 (6.5)	71 (11.5)	0.176
Co-Located with Nursing Home	19 (15.4)	112 (18.2)	0.073
Care Services, n (%)			
Assistance with Bathing	123 (100.0)	581 (94.5)	0.342
Assistance with Hygiene	123 (100.0)	531 (86.3)	0.562
Assistance with Ambulation	117 to 123 (95.1 to 100.0) *	517 (84.1)	0.480
Assistance with Feeding	89 (72.4)	185 (30.1)	0.933
Assistance with Dressing	117 to 123 (95.1 to 100.0) *	532 (86.5)	0.507
Continence Care	117 to 123 (95.1 to 100.0) *	457 (74.3)	0.788
Skin and Wound Care	47 (38.2)	113 (18.4)	0.451
Provision of Meals	123 (100.0)	609 to 615 (99.0 to 100.0) *	0.057
Administration of Medications	123 (100.0)	609 to 615 (99.0 to 100.0) *	0.114
Pharmacist Services	117 to 123 (95.1 to 100.0) *	535 (87.0)	0.287
Nursing Services	123 (100.0)	574 (93.3)	0.377
Medical Services	107 (87.0)	401 (65.2)	0.528

* Small cell sizes (i.e., where six or fewer assisted living facilities have, or do not have, a characteristic) are suppressed due to privacy restrictions at ICES

Table 2. Associations with the Provision of a Dementia Care Program in Licensed Assisted Living Facilities

	Unadjusted PR (95% CI)	Adjusted PR (95% CI) †
Facility Characteristics		
Urban	2.26 (1.23 to 4.52) **	1.15 (0.61 to 2.17)
Facility Capacity		
6-49	1.00 (Reference)	1.00 (Reference)
50-86	0.56 (0.30 to 1.06)	0.34 (0.18 to 0.66) **
87-139	1.20 (0.73 to 1.98)	0.43 (0.20 to 0.93) *
140+	2.18 (1.41 to 3.37) ***	0.59 (0.25 to 1.42)
Total Suites		
6-41	1.00 (Reference)	1.00 (Reference)
42-70	0.80 (0.43 to 1.49)	1.40 (0.73 to 2.70)
71-114	1.48 (0.87 to 2.50)	2.28 (1.02 to 5.11) *
115+	2.82 (1.77 to 4.50) ***	2.78 (1.09 to 7.07) *
Chain Status	1.63 (1.17 to 2.27) **	1.21 (0.88 to 1.67)
Residential Home	0.58 (0.29 to 1.14)	0.75 (0.35 to 1.61)
Co-Located with a Nursing Home	0.85 (0.54 to 1.33)	1.21 (0.78 to 1.87)
Care Services		
Assistance with Ambulation	6.34 (2.05 to 19.57) **	0.96 (0.34 to 2.75)
Assistance with Feeding	4.43 (3.07 to 6.39) ***	2.91 (1.98 to 4.29) ***
Assistance with Dressing	15.67 (2.22 to 110.82) **	2.24 (0.26 to 18.96)
Continence Care	33.50 (4.71 to 238.20) ***	13.51 (1.64 to 111.67) *
Skin and Wound Care	2.23 (1.62 to 3.07) ***	1.18 (0.85 to 1.63)
Pharmacist Services	2.57 (1.17 to 5.66) *	0.91 (0.38 to 2.21)
Medical Services	3.03 (1.83 to 5.00) ***	1.78 (1.00 to 3.17) *

PR = Prevalence Ratio; CI = Confidence Interval

†Adjusted for all variables in the table

* $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$

APPENDIX

Appendix Table 1. Detailed Descriptions of Exposures

Variable	Description
Urban	Based on the PCCF flag, which is a specialized macro for use with health system administrative datasets containing postal codes. This macro is based on 2016 Census information, flags communities with a population less than 10,000 individuals as rural, and it includes related data from Canada Post Corporation.
Facility Capacity	Resident capacity of the assisted living facilities. This variable was transformed into quartiles for ease of interpretation.
Total Suites	Number of suites in the assisted living facility. This variable was transformed into quartiles for ease of interpretation.
Chain Status	Binary variable specifying whether the assisted living facility is part of a chain. Chains are defined as two or more assisted living facilities owned and operated by the same entity.
Residential Home	Binary variable specifying whether the assisted living facility is a residential home. Residential homes are not facilities and are located in residential areas.
Co-Located with a Nursing Home	Binary variable specifying whether the assisted living facility is co-located with a nursing home in the same building or land site.
Assistance with Bathing	Binary variable specifying whether the assisted living facility provides bathing respecting privacy and preferences.
Assistance with Hygiene	Binary variable specifying whether the assisted living facility provides grooming, oral care, foot care, and care for fingernails.
Assistance with Ambulation	Binary variable specifying whether the assisted living facility provides assistance with mobility devices and transferring and positioning techniques.
Assistance with Feeding	Binary variable specifying whether the assisted living facility provides assistance to residents to eat and drink safely, including assistance inserting and removing dentures.
Assistance with Dressing	Binary variable specifying whether the assisted living facility provides assistance with dressing, consistent with time of day and weather conditions.
Continence Care	Binary variable specifying whether the assisted living facility provides continence products and measures to prevent constipation.
Skin and Wound Care	Binary variable specifying whether the assisted living facility provides interventions and routine care to maintain the integrity of the resident's skin, prevent wounds and infections, and other preventive measures (e.g., physiotherapy, etc.).
Provision of Meals	Binary variable specifying whether the assisted living facility provides meals supporting good nutrition standards, food service workers, and individualized meals, if required.
Administration of Medications	Binary variable specifying whether the assisted living facility stores medications and has regulated health professionals providing medication to residents consistent with their needs and care plans.

Appendix Table 1. Detailed Descriptions of Exposures

Variable	Description
Pharmacist Services	Binary variable specifying whether the assisted living facility provides services from a pharmacist regulated by the Ontario College of Pharmacists.
Nursing Services	Binary variable specifying whether the assisted living facility provides services from a nurse regulated by the Ontario College of Nurses.
Medical Services	Binary variable specifying whether the assisted living facility provides services from a physician regulated by the Ontario College of Physicians and Surgeons.

Detailed descriptions of the care services are found in O. Reg. 166/11: GENERAL under *Retirement Homes Act, 2010*, S.O. 2010, c. 11. Available at: <https://www.ontario.ca/laws/regulation/110166#BK47>

Appendix Table 2. RECORD Checklist

	Item No.	STROBE items	Location in manuscript where items are reported	RECORD items	Location in manuscript where items are reported
Title and Abstract					
	1	(a) Indicate the study's design with a commonly used term in the title or the abstract (b) Provide in the abstract an informative and balanced summary of what was done and what was found	p. 1,3	<p>RECORD 1.1: The type of data used should be specified in the title or abstract. When possible, the name of the databases used should be included.</p> <p>RECORD 1.2: If applicable, the geographic region and timeframe within which the study took place should be reported in the title or abstract.</p> <p>RECORD 1.3: If linkage between databases was conducted for the study, this should be clearly stated in the title or abstract.</p>	p. 1,3
Introduction					
Background rationale	2	Explain the scientific background and rationale for the investigation being reported	p. 4		
Objectives	3	State specific objectives, including any prespecified hypotheses	p. 4		
Methods					

Appendix Table 2. RECORD Checklist

	Item No.	STROBE items	Location in manuscript where items are reported	RECORD items	Location in manuscript where items are reported
Study Design	4	Present key elements of study design early in the paper	p. 5		
Setting	5	Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection	p. 5		
Participants	6	<p>(a) <i>Cohort study</i> - Give the eligibility criteria, and the sources and methods of selection of participants. Describe methods of follow-up</p> <p><i>Case-control study</i> - Give the eligibility criteria, and the sources and methods of case ascertainment and control selection. Give the rationale for the choice of cases and controls</p> <p><i>Cross-sectional study</i> - Give the eligibility criteria, and the sources and methods of selection of participants</p>	p. 5	<p>RECORD 6.1: The methods of study population selection (such as codes or algorithms used to identify subjects) should be listed in detail. If this is not possible, an explanation should be provided.</p> <p>RECORD 6.2: Any validation studies of the codes or algorithms used to select the population should be referenced. If validation was conducted for this study and not published elsewhere, detailed methods and results should be provided.</p> <p>RECORD 6.3: If the study involved linkage of databases, consider use of a flow diagram or other graphical</p>	p. 5

Appendix Table 2. RECORD Checklist

	Item No.	STROBE items	Location in manuscript where items are reported	RECORD items	Location in manuscript where items are reported
		<p><i>(b) Cohort study</i> - For matched studies, give matching criteria and number of exposed and unexposed</p> <p><i>Case-control study</i> - For matched studies, give matching criteria and the number of controls per case</p>		display to demonstrate the data linkage process, including the number of individuals with linked data at each stage.	
Variables	7	Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable.	p. 5,6,16,17	RECORD 7.1: A complete list of codes and algorithms used to classify exposures, outcomes, confounders, and effect modifiers should be provided. If these cannot be reported, an explanation should be provided.	p. 5,6,16,17
Data sources/ measurement	8	For each variable of interest, give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group	p. 5,6,16,17		

Appendix Table 2. RECORD Checklist

	Item No.	STROBE items	Location in manuscript where items are reported	RECORD items	Location in manuscript where items are reported
Bias	9	Describe any efforts to address potential sources of bias	p. 5,6		
Study size	10	Explain how the study size was arrived at	p. 6		
Quantitative variables	11	Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen, and why	p. 6		
Statistical methods	12	(a) Describe all statistical methods, including those used to control for confounding (b) Describe any methods used to examine subgroups and interactions (c) Explain how missing data were addressed (d) <i>Cohort study</i> - If applicable, explain how loss to follow-up was addressed <i>Case-control study</i> - If applicable, explain how	p. 6		

Appendix Table 2. RECORD Checklist

	Item No.	STROBE items	Location in manuscript where items are reported	RECORD items	Location in manuscript where items are reported
		matching of cases and controls was addressed <i>Cross-sectional study</i> - If applicable, describe analytical methods taking account of sampling strategy (e) Describe any sensitivity analyses			
		..		RECORD 12.1: Authors should describe the extent to which the investigators had access to the database population used to create the study population.	p. 5
		..		RECORD 12.2: Authors should provide information on the data cleaning methods used in the study.	
		..		RECORD 12.3: State whether the study included person-level, institutional-level, or other data linkage across two or more databases. The methods of linkage and methods of linkage quality evaluation should be provided.	p. 5
Results					
Participants	13	(a) Report the numbers of individuals at each stage	p. 7	RECORD 13.1: Describe in detail the selection of the persons included	p. 7

Appendix Table 2. RECORD Checklist

	Item No.	STROBE items	Location in manuscript where items are reported	RECORD items	Location in manuscript where items are reported
		of the study (<i>e.g.</i> , numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analysed) (b) Give reasons for non-participation at each stage. (c) Consider use of a flow diagram		in the study (<i>i.e.</i> , study population selection) including filtering based on data quality, data availability and linkage. The selection of included persons can be described in the text and/or by means of the study flow diagram.	
Descriptive data	14	(a) Give characteristics of study participants (<i>e.g.</i> , demographic, clinical, social) and information on exposures and potential confounders (b) Indicate the number of participants with missing data for each variable of interest (c) <i>Cohort study</i> - summarise follow-up time (<i>e.g.</i> , average and total amount)	p. 7,13		
Outcome data	15	<i>Cohort study</i> - Report numbers of outcome	p. 7		

Appendix Table 2. RECORD Checklist

	Item No.	STROBE items	Location in manuscript where items are reported	RECORD items	Location in manuscript where items are reported
		events or summary measures over time <i>Case-control study</i> - Report numbers in each exposure category, or summary measures of exposure <i>Cross-sectional study</i> - Report numbers of outcome events or summary measures			
Main results	16	(a) Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (e.g., 95% confidence interval). Make clear which confounders were adjusted for and why they were included (b) Report category boundaries when continuous variables were categorized (c) If relevant, consider translating estimates of relative risk into absolute	p. 7,14		

Appendix Table 2. RECORD Checklist

	Item No.	STROBE items	Location in manuscript where items are reported	RECORD items	Location in manuscript where items are reported
		risk for a meaningful time period			
Other analyses	17	Report other analyses done—e.g., analyses of subgroups and interactions, and sensitivity analyses	N/A		
Discussion					
Key results	18	Summarise key results with reference to study objectives	p. 8		
Limitations	19	Discuss limitations of the study, taking into account sources of potential bias or imprecision. Discuss both direction and magnitude of any potential bias	p. 8	RECORD 19.1: Discuss the implications of using data that were not created or collected to answer the specific research question(s). Include discussion of misclassification bias, unmeasured confounding, missing data, and changing eligibility over time, as they pertain to the study being reported.	p. 9
Interpretation	20	Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence	p. 8		

Appendix Table 2. RECORD Checklist

	Item No.	STROBE items	Location in manuscript where items are reported	RECORD items	Location in manuscript where items are reported
Generalisability	21	Discuss the generalisability (external validity) of the study results			
Other Information					
Funding	22	Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based	p. 1		
Accessibility of protocol, raw data, and programming code		..		RECORD 22.1: Authors should provide information on how to access any supplemental information such as the study protocol, raw data, or programming code.	p. 2