

Supplementary Table 1. Demographic characteristics, healthy subjects.

	Total (n= 10)
Age	48.70 ± 5.48
Gender, female	6 (60.0%)
Weight	59.12 ± 6.34
BMI	22.81 ± 1.35
Hypertension	0 (0.0%)
Diabetes	0 (0.0%)
Alcohol consumption	2 (20.0%)
Galectin-3 (ng/mL)	9.65 [8.27 – 14.71]
Glucose (mg/dL)	89.8 ± 6.07
Triglycerides (mg/dL)	111.70 ± 46.31
Creatinine (mg/dL)	0.78 ± 0.14
ALT (U/L)	24.0 ± 8.47
AST (U/L)	25.6 ± 4.40

Data are reported as mean (\pm SD), median (IQR) and *n* (%). BMI= Body mass index, ALT= alanine transaminase, AST= aspartate transaminase.

Supplementary Table 2. Forward-stepwise multivariable logistic regression

	Variables	<i>p</i> value
Step 0	Age	0.260
	Gender, male	0.491
	Diabetes	0.244
	Hypertension	0.317
	Galectin-3 (binary)	< 0.0001
	INR	0.047
	Albumin	< 0.0001
	Triglycerides	< 0.01
	Ferritin	0.01
	D-Dimer	< 0.01
	CRP	< 0.0001
	Lymphocyte count	0.334
	Neutrophil count	<0.001
	NLR	<0.01

	Variables	OR (95% CI)	<i>p</i> value
Step 1	Galectin-3 (binary)	6.65 (2.82 – 15.70)	< 0.0001
Step 2	Galectin-3 (binary)	5.00 (2.00 – 12.45)	< 0.001
	Albumin	0.19 (0.07 – 0.51)	< 0.01
Step 3	Galectin-3 (binary)	3.89 (1.50 – 10.10)	< 0.01

	CRP	1.05 (1.00 – 1.11)	0.04
	Albumin	0.24 (0.09 – 0.65)	< 0.01

Variables included in every step of the forward-stepwise logistic regression. INR= International normalized ratio, CRP= C-reactive protein, NLR= Neutrophil to lymphocyte ratio.