Detection of Antibiotic-Resistant Bacteria, Resistance Determinants, and Mobile Elements in Surface Waters in Lebanon

Jennifer Moussa¹, Edmond Abboud² and Sima Tokajian^{1*}

¹Department of Natural Sciences, School of Arts and Sciences, Lebanese American University, Byblos, 1401, Lebanon

²Laboratory department, the Middle East Institute of Health University Center, Bsalim, Lebanon

¹ *Address for correspondence: Dr. Sima Tokajian, Department of Natural Sciences, School of Arts and Sciences, Lebanese American University, Byblos, 1401, Lebanon; Tel: +961-9-547254; Email: stokajian@lau.edu.lb

Abstract

The prevalence of antibiotic-resistant bacteria in surface water in Lebanon is a growing concern and understanding the mechanisms of the spread of resistance determinants is essential. We aimed at studying the occurrence of resistant organisms and determinants in surface water sources in Lebanon and understanding their mobilization and transmission. Water samples were collected from five major rivers in Lebanon. 91 isolates were recovered out of which 25 were multidrug-resistant (MDR) and accordingly were further characterized. *Escherichia coli* and *Klebsiella pneumoniae* were the most commonly identified MDR isolates. Conjugation assays coupled with *in silico* plasmid analysis were performed and validated using PCR-based replicon typing (PBRT) to identify and confirm incompatibility groups and the localization of β -lactamase encoding genes. *E. coli* EC23 carried a *bla*_{NDM-5} gene on a conjugative, multireplicon plasmid, while *bla*_{CTX-M-15} and *bla*_{TEM-1B} were detected in the majority of the MDR isolates. Different ST types were identified including the highly virulent *E. coli* ST131. Our results showed a common occurrence of bacterial contaminants in

25 surface water and an increase in the risk for the dissemination of resistance determinants
26 exacerbated with the ongoing intensified population mobility in Lebanon and the widespread
27 lack of wastewater treatment.

28 **Keywords:** Surface Water, Antibiotic Resistance, Population Mobility, *Escherichia coli*,
29 Mobile Elements

30 Introduction

31 The rapid spread of antibiotic resistance is a worldwide concern. Previously,
32 antimicrobial resistant bacteria were confined to hospitals and veterinary settings, but are
33 now widely disseminated in aquatic environments including rivers (Koczura *et al.* 2012),
34 sewage treatment plants (Ferreira Da Silva *et al.* 2007), hospital effluents (Spindler *et al.*
35 2012; Zhang *et al.* 2017), drinking (Walsh *et al.* 2011) and surface water (Pereira *et al.* 2013).
36 Aquatic environments constantly receive pathogenic and potentially pathogenic bacteria from
37 different sources including municipal, hospital, and agricultural waste and as such could be a
38 reservoir for multi-drug resistant organisms (Egervärn *et al.* 2017; Sanganyado and Gwenz
39 2019). The prevalence of antimicrobial resistant strains in such environments is a worldwide
40 concern due to the potential health hazards to the people exposed to such aquatic
41 environments through different activities (Baquero, Martínez and Cantón 2008).

42 Surface water is largely affected by natural processes, human activities and the fast
43 growth of the population which deteriorates water quality and threatens its use (Wilbers *et al.*
44 2014). In Lebanon, surface water sources are used as the main supply for agricultural
45 activities, electricity generation, leisure, and human consumption (Daou *et al.* 2018).
46 However, most water sources in Lebanon are contaminated with raw sewage and industrial
47 waste (Faour-Klingbeil *et al.* 2016).

Water supplies and sanitation have suffered a lot in Lebanon because of the civil war and later due to the influx of refugees living in informal settlements or under temporary living environments without access to safe water and sanitation (Daoud *et al.* 2018). In 1992, a study assessing water quality revealed that most of the water supply systems in Lebanon do not conform with the world health organization quality standards for community water supplies. In 2007, the study of coastal rivers showed high levels of fecal coliforms confirming a significant raw sewer water input (Hourri and El Jebrawi 2007). Diab *et al.* (2018) further validated the contamination of different water sources in Lebanon including spring and well waters, which are directly consumed without treatment, and estuaries which are used in watering crops and animals. Diab *et al.* (2018) also studied the distribution of multi-drug resistant bacteria in water resources. Similarly, Tokajian *et al.* (2018), examined the impact of population influx on the prevalence of ESBL-producing *E. coli* recovered from river effluents in Lebanon, and revealed the prevalence of drug resistant organisms and introduction of new resistance patterns into water systems.

Reports revealing the role of environmental factors and the environment, water and sanitation, in propagating resistance determinants are still limited in Lebanon. The purpose of this study was to estimate the occurrence and determine the molecular characteristics of antimicrobial resistant organisms and resistance determinants with the ultimate goal being to understand mobilization and transmission in surface water and mitigate where possible the spread.

Materials and Methods

Study Design and Sample Collection

A total of 15 water samples were collected between 2017 and 2018 from Al Qa'a refugee camp and four other major river effluents across Lebanon with possible sewage

contamination in the north, south, el beqaa, and Beirut (Figure 1). The collected volume was 5L using sterile containers, followed by 10x dilution before being inoculated on Blood and MacConkey agar. A total of 91 isolates were recovered and were named to reflect the type (genus and species) of the organism (EC= *E. coli*, EN= *Enterococcus* spp., KP= *Klebsiella pneumoniae*, SM= *Serratia marcescens*, SA= *Salmonella* spp., AB= *Acinetobacter baumannii*, DE= *Delftia* spp., HA= *Hafnia*, KL= *Kluyvera ascorbata*, SH= *Shewanella* spp., PR= *Providencia* spp., RA= *Raoultella* spp., CI= *Citrobacter* spp., EB= *Enterobacter* spp., AE= *Aeromonas* spp., PA= *Pseudomonas* spp.). Supplementary figure 5 shows a detailed description for all recovered isolates (Designation, location, date of isolation, and the type of the organism).

DNA extraction was performed using the NucleoSpin® Tissue DNA extraction kit (Macherey- Nagel, Germany) following the manufacturer's instructions and isolates were identified using 16S rRNA gene sequencing. *Escherichia coli* and *Klebsiella pneumoniae*, the two most commonly isolated organisms, were subjected to further characterization.

Antimicrobial Susceptibility Testing

All Gram-negative isolates were tested for resistance using the disk diffusion assay on Mueller-Hinton agar using 29 different antimicrobial agents belonging to ten classes (Table 1). Gram-positive isolates were tested against 11 different antimicrobial agents (Table 2). Isolates identified as *Pseudomonas* spp. were tested against 13 antibiotics (data not shown). E-test strips (AB BIODISK, Solna, Sweden) were used with one recovered carbapenem-resistant *E. coli* (EC23) to determine the minimal inhibitory concentration (MIC) of ertapenem, imipenem and meropenem. All results were interpreted according to the Clinical Laboratory Standards Institute guidelines (CLSI 2018).

96 **Multi-locus sequence typing (MLST)**

97 **1. *K. pneumoniae***

98 MLST was performed as described on the Institute Pasteur MLST database targeting
99 seven housekeeping genes (*rpoB*, *gapA*, *mdh*, *pgi*, *phoE*, *infB*, and *tonB*) using primers
100 with universal sequencing tails. Genes were sequenced using the universal oF and oR
101 primer pair. STs were assigned using the Institute Pasteur database.
102 (www.pasteur.fr/mlst).

103 **2. *E. coli***

104 The allelic profiles of the following seven housekeeping genes in *E. coli* were also
105 determined: *adk*, *fumC*, *gyrB*, *icd*, *mdh*, *purA* and *recA* using primers with universal
106 sequencing tails. Genes were sequenced using the same primers and STs were assigned using
107 the MLST Warwick database (www.enterobase.warwick.ac.uk). The Achtman multilocus
108 sequence typing (MLST) was also performed on all whole-genome sequenced isolates using
109 MLST 2.0 database available on the Center for Genomic Epidemiology (CGE)
110 (www.genomicepidemiology.org) (Larsen *et al.* 2012)

111 **PFGE fingerprinting**

112 Genomic DNA plugs for *E. coli* and *K. pneumoniae* were prepared according to the
113 standard PulseNet protocol (<http://www.pulsenetinternational.org>). Briefly, plugs were
114 digested using *Xba*I (EC 3.1.24.4) (ThermoScientific, Waltham, MA, USA) for 2 h at 37°C.
115 Electrophoresis was performed using the Bio-Rad laboratories CHEF DR-III system (Bio-
116 Rad Laboratories, Bio-Rad Laboratories Inc., Hercules, CA, USA). *Salmonella enterica*
117 *subsp. enterica* serovar *Braenderup* (ATCC® BAA664™) according to the standard
118 PulseNet protocol (<http://www.pulsenetinternational.org>). Gels were stained with ethidium
119 bromide. BioNumerics software version 7.6.1 (Applied Maths, St-Martens-Latem, Belgium)

was used to analyze the PFGE profiles and pulsotypes were clustered through dice correlation coefficients with 1.5% optimization and 1.5% tolerance.

Whole-genome sequencing

Based on the antimicrobial susceptibility test results, 25 isolates classified as being multi-drug resistant, acquired non-susceptibility to at least one agent in three or more antimicrobial categories (Magiorakos *et al.* 2012), or showed an interesting resistance pattern were chosen for in-depth molecular characterization through whole-genome sequencing. Library preparation was performed using the Illumina Nextera XT DNA Library preparation kit (Illumina, San Diego, CA, USA). Genomic DNA (gDNA) was used as input for library preparation. gDNA was subjected to end-repair, A-tailing, ligation of adaptors including sample-specific barcodes as per the manufacturer's recommendation. Qubit 2.0 fluorometer (Invitrogen, Carlsbad, CA, USA) was used to quantify the resulting library which was sequenced on an Illumina MiSeq (Illumina, San Diego, CA, USA) with paired-end 500 cycles protocol to read a length of 250 bp. Genome assembly was performed *de novo* using SPAdes Genome Assembler Version 3.13.0 along with read error correction (Nurk *et al.* 2017). Quality control checks on the raw sequence data was performed using FastQC version 0.11.9 (Andrews 2010). The assembled draft genomes were annotated using RAST (<http://rast.nmpdr.org/rast.cgi>) (Aziz *et al.* 2008).

Phylogenetic typing and Serotyping- *E. coli*

The phylogroup was determined as described previously (Clermont *et al.* 2013). The serotype was assigned using the SerotypeFinder 2.0. Results were further validated on GoSeqit (GoSeqit.com).

PCR amplification and sequencing of *wzi* gene.

To determine the capsular type, *wzi* gene typing was performed using *wzi*-TR and *wzi*-TF primers as previously described (Brisse *et al.* 2013). K-types were assigned using the Pasteur Institute database (<http://bigsd.b.pasteur.fr/klebsiella>).

Whole- genome based *in silico* typing and virulence and resistance profiling

In silico Plasmid and pMLST typing were performed on all whole-genome sequenced isolates using PlasmidFinder 1.3 and pMLST 2.0 by means of the Center for Genomic Epidemiology (CGE) online tools (Carattoli *et al.* 2014). Virulence genes were identified using VirulenceFinder 1.5 (Joensen *et al.* 2014). ResFinder v2.1 and the Comprehensive Antibiotic Resistance Database (CARD) were used to identify resistance genes (Zankari *et al.* 2012; Jia *et al.* 2017). Fimtyper 1.0 was used to identify the *E. coli* Fim type (Roer *et al.* 2017). Phage Search tool (PHASTER) was used for Phage identification (Arndt *et al.* 2016). IS-finder was used to identify insertion sequences (ISs) and IS-families (Siguier *et al.* 2006). plasmidSPAdes, version 3.12.0, was employed to generate separate plasmid contigs, and the output was visualized through Bandage and an assembly graph viewer. Mauve v2.4.0 was used for comparative genome alignment.

Pan-genome and recombination analysis

To further study the diversity within *E. coli* and *K. pneumoniae*, genomes were annotated using Prokka (version 1.13) with a similarity cutoff e-value 10^{-6} and minimum contig size of 200 bp (Seemann 2014). Annotated GFF3 files were piped into Roary (version 3.12). Choosing a minimum BLASTp identity of 95 and core gene prevalence in all (>99%) of the isolates (Page *et al.* 2015)

A maximum-likelihood phylogenetic tree based on the core genome alignment was constructed using FastTree2. Gubbins version 2.2.1 was used to assess recombination events in core genes and to construct a maximum-likelihood tree using RAxML. One reference genome was included in each tree. *E. coli* str. K-12 subtr. MG1665 (accession # U00096.3) was used as a reference for the maximum-likelihood phylogenetic tree for all sequenced *E. coli* isolates, *Klebsiella pneumoniae* subsp. *pneumoniae* HS11286 (accession # NC_016845.1) was used as a reference for the *K. pneumoniae* isolates and *Escherichia coli* ST131 strain EC958 (accession # HG941718.1) was used when comparing *E. coli* isolates belonging to ST131. The resulting phylogenetic trees, isolates metadata along with the pan-genome fingerprints of the isolates, core genome SNPs, and recombination hotspots were visualized using Phandango V1.3.0.

Plasmid typing and Conjugation assay

Plasmid incompatibility groups were identified by the PCR-based replicon-typing method using the DIATHEVA PCR-Based Replicon Typing (PBRT) kit (Diatheva, Fano, Italy).

Conjugation Assays

Conjugation-*bla*_{NDM-5}

Conjugal transfer of plasmid-borne *bla*_{NDM-5} from *E. coli* EC23 isolated from El Qa'a refugee camp was assessed by broth culture mating assay using *E. coli* J53 as a recipient as previously described by Inoue, Itoh and Mitsuhashi (1983) with some modifications. Exponential phase growing donor and recipient strains were mixed at a volume ratio of 1:3. The mating mixture was incubated for 2 h at 37°C and then plated on MacConkey agar supplemented with 100 µg mL⁻¹ sodium azide and 10 µg mL⁻¹ meropenem. Plasmid DNA

was extracted from the transconjugant using the plasmid Mini Kit (Qiagen GmbH, Hilden, Germany). PBRT and PCR for the *bla*_{NDM-5} were performed on the plasmid DNA

Conjugation-*bla*_{CTX-M-15}

Twelve isolates from different dates and locations were shown to carry *bla*_{CTX-M-15} (Figure 2). Conjugation experiments were conducted by the broth mating method using *bla*_{CTX-M-15} containing strains as donor and sodium azide-resistant *E. coli* J53 as the recipient, as previously described (Inoue, Itoh and Mitsuhashi 1983). The mating mixture was incubated for 2 h at 37°C and MacConkey agar supplemented with sodium azide (100 µg mL⁻¹) and cefixime (5 µg mL⁻¹) was used to select for transconjugants. Plasmids were typed according to their incompatibility group using PBRT (Carattoli 2009).

Additionally, a conjugation assay was conducted using *E. coli* EC7 as a recipient and *E. coli* EC36 as a donor isolate. Selection on trimethoprim (50 µg mL⁻¹) and cefixime (5 µg mL⁻¹) for *bla*_{CTX-M-15} positive transconjugants was used.

Results

Bacterial identification

A total of 91 organisms were isolated between 2017 and 2018 from four major sites: North (19/91, 20.9%), South (10/91, 10.9%), El Beqaa (17/91, 18.7%) and Beirut (17/91, 18.7%), in addition to El Qa'a refugee camp (28/91, 30.8%). Among the recovered isolates, 36/91 were identified as *E. coli* (39.6%), 12/91 *Enterococcus* spp. (13.2%), 11/91 *K. pneumoniae* (12.1%), the remaining 32/91 (35.1%) belonged to 13 different genera (Supplementary figure 1).

Antibiotic Resistance Profiles

Most of the recovered isolates (86.8%; n= 79/91) were Gram-negative. *Pseudomonas aeruginosa* was tested using a different panel of antimicrobial agents and as such data obtained was analyzed apart from the other recovered Gram-negative isolates. The disk diffusion assay revealed that 81% (62/77) of the isolates were resistant to amoxicillin, 64% (49/77) to ticarcillin, and 44% (34/77) to cefalotin. All isolates were sensitive to tigecycline. *E. coli* EC23 isolated from El Qa'a refugee camp in 2018 was the only isolate resistant to ertapenem, meropenem, and imipenem (Table 1)

In the Gram-positive isolates however, the majority (83.3%; n= 10/12) were resistant to trimethoprim/sulfamethoxazole, while 25% (3/12) showed resistance to erythromycin. All the isolates were sensitive to amoxicillin, vancomycin, teicoplanin and tigecycline (Table 2).

Twenty-five isolates, mainly *E. coli* (17/25) and *K. pneumoniae* (4/25), were identified as being MDR, showed resistance to at least one agent in three or more antimicrobial categories (Magiorakos *et al.* 2012), and subsequently were chosen for further in-depth molecular characterization using whole-genome sequencing (Table 3). In addition, two colistin resistant isolates were detected and they were identified as non-pigmented *Serratia marcescens* and one *Providencia alcalifaciens* which are intrinsically resistant to this antibiotic.

Antibiotic Resistance Determinants

The antibiotic susceptibility testing results were further validated *in silico* through the detection of resistance determinants. A total of 56 genes conferring resistance to one of the nine tested categories of antimicrobial agents (β -lactams, fluoroquinolones, fosfomycin, sulphonamides, trimethoprim, tetracycline, phenicol, macrolides and aminoglycosides) was detected.

We found 12 different variants of genes linked to β -lactam resistance. The *bla*_{TEM-1B} gene was the most prevalent within the MDR isolates, and was detected in 14 isolates (10 *E. coli*, 3 *K. pneumoniae*, and 1 *S. marcescens*). The *bla*_{CTX-M-15} found in 13 isolates (46%), out of which five were additionally positive for *bla*_{OXA-1}. Only one gene conferring resistance to carbapenems, namely *bla*_{NDM-5} was detected in this study from *E. coli* EC23. Fourteen different aminoglycoside resistance genes were also found. The most common were *aph*(3'')-Ib (8/25; 32%) and *aph*(6)-Id. We additionally, detected phenicol, macrolide, tetracycline, sulphonamide, fosfomycin and quinolone resistance determinants (Figure 2).

Plasmid typing

The combined results from PBRT and *in silico* replicon typing using PlasmidFinder 1.3 revealed 15 different replicons with the ones most prevalently detected Inc groups being IncFII (22.39%, 15/67), IncFIA (19.4%, 13/67), IncFIB (16.42%, 11/67) and IncFIB(K) (8.96%, 6/67). At least two different Inc groups were identified in the majority of the tested isolates (18/25, 72%). The highest number of Inc groups (n=7) was detected in EC18 isolated from El Qa'a refugee camp. The plasmidSPAdes version 3.12.0 was used to reconstruct the plasmids from paired-end reads. Based on that we showed the presence of a large multi-replicon plasmid having IncFIA, IncFIB, IncFII and IncI γ and carrying *bla*_{CMY-42} and *tetA* resistance determinants. Another smaller plasmid, missed due to Illumina short read sequencing and/or miss-assembly, carrying IncY, IncU, and IncI1 α replicons were also detected.

Molecular Typing

E. coli and *K. pneumoniae* were typed using PFGE, MLST, phylogrouping and serotyping to study genetic relatedness. The Achtman method

(<https://enterobase.warwick.ac.uk/species/ecoli>) was used to determine the MLST for *E. coli*, based on which we found that ST131 (16.7%, 6/36) and ST69 (11.1%, 4/36) were the most common. We also identified 17 other STs including ST405, ST394, and ST410. Four of the isolates did not match any allelic profile and as such were assigned new STs. Using Clermont *et al.* (2013) classification scheme, we were able to phylotype the 36 *E. coli* isolates and accordingly were distributed into five phylogroups: A (27.8%; n=10/36), B2 (36%; n=13/36), D (17%; n=6/36), C (11.1%; n=4/36), and E (5.6%; n=2/36) (Figure 3).

Serotypes of sequenced *E. coli* isolates were also determined *in silico* using SerotypeFinder 2.0. ST131 isolates (EC10, EC22, EC24, EC35, and EC36) belonged to O25b:H4 serotype except for EC7 which was typed as O16:H5b. Other serotypes were also found including: O102:H6 (ST405; EC8 and EC12). EC5 (ST2349), EC19 (ST746), and EC27 (ST10) however, had untypeable O type gene.

Using PFGE, a total of 34 pulsotypes were obtained. Isolates with the same ST did not cluster together; EC24, for example, typed as ST131 clustered away from the other five isolates from the same ST (Figure 3). It is noteworthy that EC3 (El Qa'a refugee camp) was untypeable.

With *K. pneumoniae*, seven different STs and K-types were identified while some were untypable. KP9 and KP6 belonged to ST76 and were linked to K-type 31. KP9 and KP6 additionally, shared a 96% similarity despite the difference observed in their susceptibility patterns. (Figure 4).

Genome Statistics and Comparative Analysis

Pan-genome analysis of all the MDR *E. coli* isolates revealed the presence of a total of 10,315 unique protein coding sequences. Genes identified as part of the core genome were detected in all and constituted 3,154 protein coding sequences. Genes present in 15-95% of

isolates were designated as being part of the shell genes and constituted 2,451 protein coding sequences, while those present in <15% constituted 4,710 protein coding sequences. Maximum-likelihood phylogenetic tree constructed based on single nucleotide polymorphisms (SNPs) placed the isolates in three different clusters. WgSNPs-based phylogenetic typing however, clustered the isolates based on differences in their STs, phylogroups, serotypes, and FimH. Again, three clusters were observed with one having all the isolates belonging to phylogroup B2 which included six isolates typed as ST131 and another as ST73. Isolates belonging to phylogroup D (ST405 and ST394), A, and, C also clustered together (Supplementary figure 2).

We also constructed a SNP-based maximum-likelihood phylogenetic tree for the isolates typed as ST131. We used a reference genome of the same MLST type, and of phylogroup B2 and serotype O25b:H4 (Accession # HG941718.1). EC7 with serotype O16:H5 clustered separately from the other isolates (Supplementary figure 3).

Using BLASTn search against the non-redundant nucleotide database we were able to associate the highly recombinant region to the *kpsFEDUCS* region 1 operon in *E. coli* ST131 isolates.

Conjugation assays

All CTX-M-15 bearing isolates were tested for transferability of the plasmid carrying the *bla*_{CTX-M-15} gene using *E. coli* J53. Transconjugants containing CTX-M-15 encoding plasmids were extracted from five isolates. Among the five transferable CTX-M-15 producers, the most prevalent Inc type was IncF (n=3) with only one *bla*_{CTX-M-15} being carried on IncII plasmid, and another on IncB/O/K/Z plasmid. IncF plasmids were also positive for other β -lactamase encoding genes such as *bla*_{TEM-1B}, *bla*_{CMY-42}, *bla*_{DHA-1}, and *bla*_{OXA-1}. *bla*_{NDM-5} on the other hand, was detected on a 120 kb plasmid and upon conjugation *E. coli* J53 showed resistance to meropenem, imipenem, and ertapenem.

*bla*_{CTX-M-15} genes were also chromosomal in almost half of the isolates (51.3%) with two (EC30 and EC35) having it on SSU5_Salmonella phage (Supplementary figure 4, Table 4). In addition, the *in silico* analysis revealed that *bla*_{ADC-1} and *bla*_{OXA-132} in *Acinetobacter baumannii*, and *bla*_{SRT-2} and *bla*_{TEM-1B} in *S. marcescens* were all chromosomal (**Error! Reference source not found.**).

We also assessed the mobility of the *bla*_{CTX-M-15} by performing a conjugation assay using EC36 as donor and to EC7 as recipient. The plasmid carrying *bla*_{CTX-M-15} was mobilized through the performed conjugation assay as also shown by the detected cefixime resistance in EC7. PBRT and PCR assays also revealed that EC7 was positive for the *bla*_{CTX-M-15} and had three replicons (IncFIA, IncFIB and IncFII).

Discussion

Antibiotic resistance is a worldwide concern and the spread of drug-resistant bacteria through different environments including water may constitute a route for the dissemination of drug-resistant pathogens (Egervärn *et al.* 2017). The prevalence of antibiotic-resistant

bacteria in surface water in Lebanon is a growing concern with the lack of collection and domestic wastewater management strategies. According to the Central Administration of Statistics, only 37% of the buildings in Lebanon are connected to a sewer networks, the rest either use cesspools or septic tanks or release raw sewage directly into the environment (Ministry of Environment, ECODIT). Community and hospital effluents could contaminate surface water and increase the microbial load (Hourì and El Jeblawì 2007). In this study, we aimed at determining the role of surface water in the mobilization of resistance determinants. Our results showed a common occurrence of drug-resistant organisms in surface water including ESBL- and carbapenemase-producing *Enterobacteriaceae* being exacerbated by population mobility and the absence of wastewater treatment. Sewage contaminated surface water samples from river effluents were collected from across Lebanon. Antibiotic susceptibility testing was performed and MDR isolates (Magiorakos *et al.* 2012), or the ones that showed an interesting resistance profile, were further characterized using whole-genome sequencing. Ninety-one isolates in total were recovered and 25 were whole-genome sequenced and further characterized.

E. coli (39%; n= 36/91) was the most commonly recovered organism. This finding was in accordance with a previous study by Tokajian *et al.* (2018), assessing bacterial loads from sewage contaminated surface water samples from Lebanon. The second most prevalent was *Enterococcus* spp. (13%; n= 12/91) despite the fact that in general they are not frequently recovered from water (Cabral 2010). Their isolation from aquatic environments was linked to livestock, poultry, hospital, and municipal sewage, and wildlife fecal contamination (Byappanahalli *et al.* 2012).

S. marcescens recovered in this study was a non-pigmented biotype and positive for TEM-1B and SRT-2. It's noteworthy that both pigmented and not-pigmented biotypes were previously detected with the non-pigmented being more virulent (Roy, Ahmed and Grover

2014). Hospital outbreaks were also linked to the non-pigmented biotype and which was in contrast to the environmental strains that produced a red pigment, prodigiosin (Kurz *et al.* 2003).

MLST typing, revealed that *E. coli* recovered in this study belonged to 20 unique ST types. *E. coli* typed as ST131, one of the high-risk clones associated with community onset infections (Mathers, Peirano and Pitout 2015), was the most common, and was detected in five out of the six collection sites. ST131 was also previously recovered from dug and spring wells and estuaries from Lebanon as well as different water sources worldwide (Bréchet *et al.* 2014; Chen *et al.* 2016; Egervärn *et al.* 2017). Interestingly, clinical and ecological studies have shown that CTX-M-15 producing ST131 could be rarely recovered from the environmental or veterinary settings (Pitout and DeVinney 2017). Add to it the fact that ST131 was frequently recovered from clinical settings in Lebanon, and which could further suggest contamination linked to human-activities.

Moreover, the five ST131 isolates were of serotype O25b:H4 and of the FimH30 lineage. FimH30 is the most prevalent lineage within ST131; it is named as such because it contains the H30 variant of the type 1 fimbrial adhesin gene (Mathers, Peirano and Pitout 2015). FimH30 also includes around 70% of the recent fluoroquinolone resistant ST131 *E. coli* isolates and which were implicated in the global dissemination of the fluoroquinolone resistant ST131 associated with extra intestinal infections (Johnson *et al.* 2010). In contrast, only one of the recovered isolates in this study (EC7) was of a different serotype namely O16:H5 and which belonged to the FimH41 lineage. A lineage that encompasses only 1-5% of the ST131 *E. coli* (Johnson *et al.* 2014).

Phylogrouping showed that B2 (36%) was the most common followed by phylogroup A (27.8%), and D (17%). Commensal *E. coli* isolates generally belonged to phylogroups A and B1, while the extra intestinal ones which were more pathogenic to B2 and D (Johnson *et al.*

2001). Phylogroups A and B1 however, were more frequently recovered from aquatic environments than phylogroups B2 and D (Figueira, Serra and Manaia 2011).

Amongst the whole-genome sequenced isolates in this study, 80% showed resistance to ticarcillin and 72% to third-generation cephalosporins and aztreonam. *bla*_{CTX-M-15} was the most common detected ESBL in both *E. coli* and *K. pneumoniae*. *bla*_{CTX-M-15} prevalence confirmed previous results (Diab *et al.* 2018; Tokajian *et al.* 2018).

The frequent detection of ESBL in clinical settings and the environment could reveal mobilization through horizontal gene transfer. Such transmission is favored by mobile genetic elements, specifically plasmids. IncF plasmids have a broad host range facilitating mobility and spread of resistance determinants (Villa *et al.* 2010; Zhao and Hu 2013). IncF plasmids were implicated in the spread of TEM, OXA, CMY, and KPC and often conferring multi-drug resistance (Villa *et al.* 2010; Carattoli 2011).

In addition, one NDM-5 positive *E. coli* (EC23, ST361) was recovered from El Qa'a refugee camp. NDM-5 was first described in a clinical *E. coli* isolate in 2011 from the United Kingdom (Hornsey, Phee and Wareham 2011). It was also detected in Montpellier, France from an urban river water sample (Almakki *et al.* 2017) and in India from a hospital-linked sewage water. In Lebanon however, *bla*_{NDM-5} was only linked so far to *K. pneumoniae* with clinical origin (Parvez and Khan 2017; Nawfal Dagher *et al.* 2019). *bla*_{NDM-5} in this study was on a conjugative multi-replicon plasmid (IncFIA, IncFII). *bla*_{NDM-5} commonly was found associated with IncX3 (Almakki *et al.* 2017), but was also linked to IncF plasmids from environmental samples in Myanmar (Sugawara *et al.* 2019) and from wastewater treatment plant effluent in Switzerland (Zurfluh *et al.* 2018).

The detection of MDR *E. coli* in surface water along with mobile element carrying resistance determinants highlights and confirms the role of water and lack of proper sanitation in the propagation and mobilization of resistance determinants. Our results

suggested and confirmed that surface water could act as an important reservoir facilitating the spread of resistance determinants. We have also characterized the genetic basis of antibiotic resistance of ESBLs as well as other antimicrobial agents. Efforts in the future should be focused at building waste treatment plants, minimize pollution from agricultural sources, and control and mitigate human activities linked to introducing antibiotic resistance into the environment, which is an urgent, challenging, and pressing task.

Funding

This work was supported partially by the School of Arts of Sciences Research and Development Council at the Lebanese American University and by the National Council for Scientific Research (Grant #756).

Conflict of Interest: None declared

Acknowledgments

None.

References

1. Almakki A, Almakki A, Maure A *et al.* NDM-5-producing *Escherichia coli* in an urban river in Montpellier, France. *Int J Antimicrob Agents* 2017, DOI: 10.1016/j.ijantimicag.2017.04.003.
2. Andrews S. FastQC - A quality control tool for high throughput sequence data. <http://www.bioinformatics.babraham.ac.uk/projects/fastqc/>. *Babraham Bioinforma* 2010.
3. Arndt D, Grant JR, Marcu A *et al.* PHASTER: a better, faster version of the PHAST phage search tool. *Nucleic Acids Res* 2016, DOI: 10.1093/nar/gkw387.
4. Aziz RK, Bartels D, Best A *et al.* The RAST Server: Rapid annotations using subsystems technology. *BMC Genomics* 2008, DOI: 10.1186/1471-2164-9-75.
5. Baquero F, Martínez JL, Cantón R. Antibiotics and antibiotic resistance in water

418 environments. *Curr Opin Biotechnol* 2008, DOI: 10.1016/j.copbio.2008.05.006.

419 6. Bréchet C, Plantin J, Sauget M *et al.* Wastewater treatment plants release large
420 amounts of extended-spectrum β -lactamase-producing *Escherichia coli* into the
421 environment. *Clin Infect Dis* 2014, DOI: 10.1093/cid/ciu190.

422 7. Brisse S, Passet V, Haugaard AB *et al.* Wzi gene sequencing, a rapid method for
423 determination of capsulartype for *Klebsiella* strains. *J Clin Microbiol* 2013, DOI:
424 10.1128/JCM.01924-13.

425 8. Byappanahalli MN, Nevers MB, Korajkic A *et al.* Enterococci in the Environment.
426 *Microbiol Mol Biol Rev* 2012, DOI: 10.1128/mmbr.00023-12.

427 9. Cabral JPS. Water microbiology. Bacterial pathogens and water. *Int J Environ Res*
428 *Public Health* 2010, DOI: 10.3390/ijerph7103657.

429 10. Carattoli A. Resistance plasmid families in Enterobacteriaceae. *Antimicrob Agents*
430 *Chemother* 2009, DOI: 10.1128/AAC.01707-08.

431 11. Carattoli A. Plasmids in Gram negatives: Molecular typing of resistance plasmids. *Int*
432 *J Med Microbiol* 2011, DOI: 10.1016/j.ijmm.2011.09.003.

433 12. Carattoli A, Zankari E, García-Fernández A *et al.* In Silico detection and typing of
434 plasmids using plasmidfinder and plasmid multilocus sequence typing. *Antimicrob*
435 *Agents Chemother* 2014, DOI: 10.1128/AAC.02412-14.

436 13. Chen PA, Hung CH, Huang PC *et al.* Characteristics of CTX-M extended spectrum β -
437 lactamase-producing *Escherichia coli* strains isolated from multiple rivers in southern
438 Taiwan. *Appl Environ Microbiol* 2016, DOI: 10.1128/AEM.03222-15.

439 14. Clermont O, Christenson JK, Denamur E *et al.* The Clermont *Escherichia coli* phylo-
440 typing method revisited: Improvement of specificity and detection of new phylo-
441 groups. *Environ Microbiol Rep* 2013, DOI: 10.1111/1758-2229.12019.

15. CLSI. (2018). Performance Standards for Antimicrobial Susceptibility Testing. 28th ed. CLSI supplement M100. Wayne, PA: Clinical and Laboratory Standards Institute.
16. Daou C, Salloum M, Legube B *et al.* Characterization of spatial and temporal patterns in surface water quality: a case study of four major Lebanese rivers. *Environ Monit Assess* 2018, DOI: 10.1007/s10661-018-6843-8.
17. Daoud Z, Farah J, Sokhn ES *et al.* Multidrug-Resistant Enterobacteriaceae in Lebanese Hospital Wastewater: Implication in the One Health Concept. *Microb Drug Resist* 2018, DOI: 10.1089/mdr.2017.0090.
18. Egervärn M, Englund S, Ljunge M *et al.* Unexpected common occurrence of transferable extended spectrum cephalosporinase-producing *Escherichia coli* in Swedish surface waters used for drinking water supply. *Sci Total Environ* 2017, DOI: 10.1016/j.scitotenv.2017.02.157.
19. Diab M, Hamze M, Bonnet R *et al.* Extended-spectrum beta-lactamase (ESBL)-and carbapenemase-producing Enterobacteriaceae in water sources in Lebanon. *Vet Microbiol* 2018, DOI: 10.1016/j.vetmic.2018.03.007
20. Faour-Klingbeil D, Kuri V, Fadlallah S *et al.* Prevalence of antimicrobial-resistant *Escherichia coli* from raw vegetables in Lebanon. *J Infect Dev Ctries* 2016, DOI: 10.3855/jidc.7745.
21. Ferreira Da Silva M, Vaz-Moreira I, Gonzalez-Pajuelo M *et al.* Antimicrobial resistance patterns in Enterobacteriaceae isolated from an urban wastewater treatment plant. *FEMS Microbiol Ecol* 2007, DOI: 10.1111/j.1574-6941.2006.00268.x.
22. Figueira V, Serra E, Manaia CM. Differential patterns of antimicrobial resistance in population subsets of *Escherichia coli* isolated from waste- and surface waters. *Sci Total Environ* 2011, DOI: 10.1016/j.scitotenv.2010.12.011.
23. Hornsey M, Phee L, Wareham DW. A novel variant, NDM-5, of the New Delhi

467 metallo- β -lactamase in a multidrug-resistant *Escherichia coli* ST648 isolate recovered
468 from a patient in the United Kingdom. *Antimicrob Agents Chemother* 2011, DOI:
469 10.1128/AAC.05108-11.

470 24. Houri A, El Jeblawi SW. Water quality assessment of Lebanese coastal rivers during
471 dry season and pollution load into the Mediterranean Sea. *J Water Health* 2007, DOI:
472 10.2166/wh.2007.047.

473 25. Inoue M, Itoh J, Mitsuhashi S. pMS76, a plasmid capable of amplification by
474 treatment with chloramphenicol. *Plasmid* 1983, DOI: 10.1016/0147-619X(83)90033-
475 1.

476 26. Jia B, Raphenya AR, Alcock B *et al.* CARD 2017: Expansion and model-centric
477 curation of the comprehensive antibiotic resistance database. *Nucleic Acids Res* 2017,
478 DOI: 10.1093/nar/gkw1004.

479 27. Joensen KG, Scheutz F, Lund O *et al.* Real-time whole-genome sequencing for
480 routine typing, surveillance, and outbreak detection of verotoxigenic *Escherichia coli*.
481 *J Clin Microbiol* 2014, DOI: 10.1128/JCM.03617-13.

482 28. Johnson JR, Clermont O, Johnston B *et al.* Rapid and specific detection, molecular
483 epidemiology, and experimental virulence of the O16 subgroup within *Escherichia*
484 *coli* sequence type 131. *J Clin Microbiol* 2014, DOI: 10.1128/JCM.03502-13.

485 29. Johnson JR, Delavari P, Kuskowski M *et al.* Phylogenetic distribution of
486 extraintestinal virulence-associated traits in *Escherichia coli*. *J Infect Dis* 2001, DOI:
487 10.1086/317656.

488 30. Johnson JR, Johnston B, Clabots C *et al.* *Escherichia coli* sequence type ST131 as the
489 major cause of serious multidrug-resistant *E. coli* infections in the United States. *Clin*
490 *Infect Dis* 2010, DOI: 10.1086/653932.

491 31. Koczura R, Mokracka J, Jabłońska L *et al.* Antimicrobial resistance of integron-

492 harboring *Escherichia coli* isolates from clinical samples, wastewater treatment plant
493 and river water. *Sci Total Environ* 2012, DOI: 10.1016/j.scitotenv.2011.10.036.

494 32. Kurz CL, Chauvet S, Andrès E *et al.* Virulence factors of the human opportunistic
495 pathogen *Serratia marcescens* identified by in vivo screening. *EMBO J* 2003, DOI:
496 10.1093/emboj/cdg159.

497 33. Larsen M V., Cosentino S, Rasmussen S *et al.* Multilocus sequence typing of total-
498 genome-sequenced bacteria. *J Clin Microbiol* 2012, DOI: 10.1128/JCM.06094-11.

499 34. Magiorakos AP, Srinivasan A, Carey RB *et al.* Multidrug-resistant, extensively drug-
500 resistant and pandrug-resistant bacteria: An international expert proposal for interim
501 standard definitions for acquired resistance. *Clin Microbiol Infect* 2012, DOI:
502 10.1111/j.1469-0691.2011.03570.x.

503 35. Mathers AJ, Peirano G, Pitout JDD. The role of epidemic resistance plasmids and
504 international high- risk clones in the spread of multidrug-resistant Enterobacteriaceae.
505 *Clin Microbiol Rev* 2015, DOI: 10.1128/CMR.00116-14.

506 36. Nawfal Dagher T, Azar E, Al-Bayssari C *et al.* First Detection of Colistin-Resistant
507 *Klebsiella pneumoniae* in Association with NDM-5 Carbapenemase Isolated from
508 Clinical Lebanese Patients. *Microb Drug Resist* 2019, DOI: 10.1089/mdr.2018.0383.

509 37. Nurk S, Meleshko D, Korobeynikov A *et al.* MetaSPAdes: A new versatile
510 metagenomic assembler. *Genome Res* 2017, DOI: 10.1101/gr.213959.116.

511 38. Page AJ, Cummins CA, Hunt M *et al.* Roary: Rapid large-scale prokaryote pan
512 genome analysis. *Bioinformatics* 2015, DOI: 10.1093/bioinformatics/btv421.

513 39. Parvez S, Khan, AU. Hospital sewage water: a reservoir for variants of New Delhi
514 metallo- β -lactamase (NDM)-and extended-spectrum β -lactamase (ESBL)-producing
515 Enterobacteriaceae. *Int J Antimicrob Agents* 2018, DOI:
516 10.1016/j.ijantimicag.2017.08.032

- 517 40. Pereira A, Santos A, Tacão M *et al.* Genetic diversity and antimicrobial resistance of
518 *Escherichia coli* from Tagus estuary (Portugal). *Sci Total Environ* 2013, DOI:
519 10.1016/j.scitotenv.2013.04.067.
- 520 41. Pitout JDD, DeVinney R. *Escherichia coli* ST131: A multidrug-resistant clone primed
521 for global domination. *FI000Research* 2017, DOI: 10.12688/f1000research.10609.1.
- 522 42. Roer L, Tchesnokova V, Allesoe R *et al.* Development of a web tool for *Escherichia*
523 *coli* subtyping based on fimh alleles. *J Clin Microbiol* 2017, DOI:
524 10.1128/JCM.00737-17.
- 525 43. Roy P, Ahmed NH, Grover RK. Non-pigmented strain of *Serratia marcescens*: An
526 unusual pathogen causing pulmonary infection in a patient with malignancy. *J Clin*
527 *Diagnostic Res* 2014, DOI: 10.7860/JCDR/2014/8629.4513.
- 528 44. Sanganyado E, Gwenzi W. Antibiotic resistance in drinking water systems:
529 Occurrence, removal, and human health risks. *Sci Total Environ* 2019, DOI:
530 10.1016/j.scitotenv.2019.03.162.
- 531 45. Seemann T. Prokka: Rapid prokaryotic genome annotation. *Bioinformatics* 2014,
532 DOI: 10.1093/bioinformatics/btu153.
- 533 46. Siguier P, Perochon J, Lestrade L *et al.* ISfinder: the reference centre for bacterial
534 insertion sequences. *Nucleic Acids Res* 2006, DOI: 10.1093/nar/gkj014.
- 535 47. Spindler A, Otton LM, Fuentesfria DB *et al.* Beta-lactams resistance and presence of
536 class 1 integron in *Pseudomonas* spp. isolated from untreated hospital effluents in
537 Brazil. *Antonie van Leeuwenhoek, Int J Gen Mol Microbiol* 2012, DOI:
538 10.1007/s10482-012-9714-2.
- 539 48. Sugawara Y, Akeda Y, Hagiya H *et al.* Spreading patterns of NDM-producing
540 Enterobacteriaceae in clinical and environmental settings in Yangon, Myanmar.
541 *Antimicrob Agents Chemother* 2019, DOI: 10.1128/AAC.01924-18.

49. Tokajian S, Moghnieh R, Salloum T *et al.* Extended-spectrum β -lactamase-producing *Escherichia coli* in wastewaters and refugee camp in Lebanon. *Future Microbiol* 2018, DOI: 10.2217/fmb-2017-0093.
50. Villa L, García-Fernández A, Fortini D *et al.* Replicon sequence typing of IncF plasmids carrying virulence and resistance determinants. *J Antimicrob Chemother* 2010, DOI: 10.1093/jac/dkq347.
51. Walsh TR, Weeks J, Livermore DM *et al.* Dissemination of NDM-1 positive bacteria in the New Delhi environment and its implications for human health: An environmental point prevalence study. *Lancet Infect Dis* 2011, DOI: 10.1016/S1473-3099(11)70059-7.
52. Wilbers GJ, Becker M, Nga LT *et al.* Spatial and temporal variability of surface water pollution in the Mekong Delta, Vietnam. *Sci Total Environ* 2014, DOI: 10.1016/j.scitotenv.2014.03.049.
53. Zankari E, Hasman H, Cosentino S *et al.* Identification of acquired antimicrobial resistance genes. *J Antimicrob Chemother* 2012, DOI: 10.1093/jac/dks261.
54. Zhang R, Liu L, Zhou H *et al.* Nationwide Surveillance of Clinical Carbapenem-resistant Enterobacteriaceae (CRE) Strains in China. *EBioMedicine* 2017, DOI: 10.1016/j.ebiom.2017.04.032.
55. Zhao WH, Hu ZQ. Epidemiology and genetics of CTX-M extended-spectrum β -lactamases in Gram-negative bacteria. *Crit Rev Microbiol* 2013, DOI: 10.3109/1040841X.2012.691460.
56. Zurfluh K, Stevens MJA, Stephan R *et al.* Complete and assembled genome sequence of an NDM-5- and CTX-M-15-producing *Escherichia coli* sequence type 617 isolated from wastewater in Switzerland. *J Glob Antimicrob Resist* 2018, DOI: 10.1016/j.jgar.2018.08.015.

Figure 1: Geographical distribution of the chosen surface water sources included in this study; the red asterisk (*) indicates the rivers from which isolates were collected.

Figure 2: Types and distribution of resistance determinants as assigned using ResFinder v1.2 The used antimicrobial categories were labeled as follows: B, β -lactam resistance genes; S, sulfonamide resistance genes; D, trimethoprim resistance genes; F, fosfomycin resistance genes; P, chloramphenicol resistance genes, T, tetracycline resistance genes; H, macrolide resistance genes; O, quinolone resistance genes and A, aminoglycoside resistance genes.

Figure 3: Detailed characterization of all *E. coli* isolates. The dendrogram was generated using Bionumerics software 7.6.2 showing the relatedness of the isolates based on their banding patterns generated by XbaI restriction digestion.

Dark blue= Resistant, white= sensitive and light orange= intermediate resistant.
AMX= Amoxicillin, AMC= Amoxicillin/ Clavulanic Acid, TIC= Ticarcillin, TZP= Piperacillin/Tazobactam, IPM= Imipenem, ETP= Ertapenem, MEM= Meropenem, ATM= Aztreonam, CFT= Cefalotin, CXM= Cefuroxime, FOX= Cefoxitin, CTX= Cefotaxime, CRO= Ceftriaxone, CAZ= Ceftazidime, FEP= Cefepime, CFM= Cefixime, TOB= Tobramycin, GEN= Gentamicin, OFX= Ofloxacin, CIP= Ciprofloxacin, NOR= Norfloxacin, LVX= Levofloxacin, TET= Tetracycline, TGC= Tigecycline, SXT= Trimethoprim/Sulfamethoxazole and CST= Colistin, EC= *E. coli*, ST= sequence type.

Figure 4: Detailed characterization of all *K. pneumoniae* isolates. The dendrogram was generated using Bionumerics software 7.6.2 showing the relatedness of the isolates based on their banding patterns generated by XbaI restriction digestion.

Dark blue= Resistant, white= sensitive and light orange= intermediate resistant.
AMX= Amoxicillin, AMC= Amoxicillin/ Clavulanic Acid, TIC= Ticarcillin, TZP= Piperacillin/Tazobactam, IPM= Imipenem, ETP= Ertapenem, MEM= Meropenem, ATM= Aztreonam, CFT= Cefalotin, CXM= Cefuroxime, FOX= Cefoxitin, CTX= Cefotaxime, CRO= Ceftriaxone, CAZ= Ceftazidime, FEP= Cefepime, CFM= Cefixime, TOB= Tobramycin, GEN= Gentamicin, OFX= Ofloxacin, CIP= Ciprofloxacin, NOR= Norfloxacin, LVX= Levofloxacin, TET= Tetracycline, TGC= Tigecycline, SXT= Trimethoprim/Sulfamethoxazole and CST= Colistin, EC= *E. coli*, ST= sequence type.

601 Supplemental Files

602 **Supplementary figure 1:** Distribution of organisms isolated from river effluents (El Beqaa,
603 Beirut, North and South and El Qa'a) in Lebanon.

604 SM= *Serratia marcescens*, SA= *Salmonella* spp., AB= *Acinetobacter baumannii*, DE= *Delftia*
605 spp., HA= *Hafnia*, KL= *Kluyvera ascorbata*, SH= *Shewanella* spp., PR= *Providencia* spp.,
606 RA= *Raoultella* spp., CI= *Citrobacter* spp., EB= *Enterobacter* spp., KP= *Klebsiella*
607 *pneumoniae*, AE= *Aeromonas* spp., PA= *Pseudomonas* spp. EN= *Enterococcus* spp., EC= *E.*
608 *coli*.

609
610 **Supplementary figure 2:** Pan-genome similarity, isolates data and maximum likelihood tree
611 based on core genome SNPs of all sequenced *E. coli* isolates. Maximum likelihood
612 phylogenetic tree is based on the core genome SNPs and was generated using FastTree 2.
613 Pan-genome is constructed using Roary based on the core and accessory genes showing
614 phylogenetic relatedness between isolates, blue indicates present and white indicates absent
615 fragment.

616 **Supplementary figure 3:** Pan-genome similarity, isolates data, maximum likelihood tree and
617 core genome variation in recombination and SNPs for ST131 *E. coli* isolates. (a) Maximum
618 likelihood phylogenetic tree based on core genome SNPs. (b) Recombination densities were
619 detected between the samples using Gubbins.

620 **Supplementary figure 4:** Phages detected in whole genome sequenced isolates using
621 PHASTER (phaster.ca/). Blue squares indicate the presence of the phage, orange square
622 indicate that the phage carries *bla*_{CTX-M-15} and grey square indicates that the phage carries the
623 increased serum resistance encoding gene *iss*. No phages were detected in isolates EC27,
624 SM1 and PO1.

625 **Supplementary figure 5:** Distribution of the based on the source and time of collection.
626
627
628

629

630

Time	Area	Isolate	Antibiotic Susceptibility																														
			AMX	AMC	TTC	TZP	MEM	ETP	IPM	ATM	CFT	CXM	FOX	CTX	CRO	CAZ	HEP	CFM	TOB	GEM	GEM	ORX	CIP	NOR	LXX	HEI	TGC	SXT	NIT	CST			
2017	North	KP6																															
		KP7																															
		EB2																															
		KP8																															
		EB3																															
		AB1																															
		KP9																															
		AE1																															
		AE7																															
	El Qa'a	AE2																															
		AE3																															
		AE4																															
		EC2																															
		EC3																															
		EC4																															
		EC5																															
		PR2																															
		AE5																															
	AE6																																
	El Beqaa	RA1																															
		CI4																															
		CI1																															
		RA2																															
		RA3																															
		HA1																															
	Beirut	EB4																															
		KL1																															
		CI3																															
		EB1																															
		KP4																															
		DE1																															
	South	SH1																															
		EC9																															
		EC10																															
	El Qa'a	EC11																															
		EC12																															
EB5																																	
North	EC1																																
	CI2																																
	SM1																																
Beirut	PR1																																
	KP1																																
	KP2																																
	EC6																																
	KP3																																
	EC7																																
El Beqaa	EB6																																
	EC8																																
	EB7																																
	EC13																																
	EC14																																
South	SA1																																
	KP10																																
	EC15																																
	KP5																																
	PO1																																
2018	North	EC30																															
		EC31																															
		EC32																															
		EC16																															
	El Qa'a	EC17																															
		EC18																															
		EC19																															
		EC20																															
		EC21																															
		EC22																															
		EC23																															
	Beirut	EC24																															
		EC25																															
	El Beqaa	EC26																															
		EC33																															
EC34																																	
EC35																																	
South	EC36																																
	EC27																																
	EC28																																

Table 1: Antibiotic susceptibility results for all recovered Gram-negative organisms from the river effluents and El Qa'a refugee camp in Lebanon using 29 different antimicrobial agents covering ten different classes.

AMX= Amoxicillin, AMC= Amoxicillin/ Clavulanic Acid, TIC= Ticarcillin, TZP= Piperacillin/Tazobactam, IPM= Imipenem, ETP= Ertapenem, MEM= Meropenem , ATM= Aztreonam, CFT= Cefalotin, CXM= Cefuroxime, FOX= Cefoxitin, CTX= Cefotaxime, CRO= Ceftriaxone, CAZ= Ceftazidime, FEP= Cefepime, CFM= Cefixime, TOB= Tobramycin , GEN= Gentamicin , AMK= Amikacin , OFX= Ofloxacin, CIP= Ciprofloxacin, NOR= Norfloxacin, LVX= Levofloxacin, TET= Tetracycline, TGC= Tigecycline, SXT= Trimethoprim/Sulfamethoxazole, NIT= Nitrofurantoin, FOF= Fosfomycin (*E. coli* urine) and CST= Colistin)

Location	Organism	AMX	KAN High	STR High	GEN	VAN	TEC	LEV	TET	TGC	SXT	ERY
North	<i>Enterococcus</i> spp.											
	<i>Enterococcus</i> spp.											
	<i>Enterococcus</i> spp.											
El Qa'a	<i>Enterococcus</i> spp.											
	<i>Enterococcus</i> spp.											
	<i>Enterococcus</i> spp.											
	<i>Enterococcus</i> spp.											
Beirut	<i>Enterococcus</i> spp.											
El Qa'a	<i>Enterococcus</i> spp.											
North	<i>Enterococcus</i> spp.											
Beirut	<i>Enterococcus</i> spp.											
South	<i>Enterococcus</i> spp.											

Table 2: Antibiotic susceptibility results for all recovered Gram-positive organisms from the river effluents in Lebanon using 12 different antimicrobial agents covering eight different classes.

AMX= Amoxicillin, KAN high= Kanamycin high, STR high= Streptomycin high, GEN high= Gentamicin high, VAN= Vancomycin, TEC= Teicoplanin, LEV= Levofloxacin, TET= Tetracycline, TGC= Tigecycline, SXT= Trimethoprim/Sulfamethoxazole, ERY= Erythromycin.

Date	Location	Isolate	Organism	Accession Number
2017	El Qa'a	EC5	<i>E. coli</i>	PXIF00000000
2017	North	KP8	<i>K. pneumoniae</i>	NZ_CP034540.1
2017	North	AB1	<i>A. baumannii</i>	PXHY00000000
2017	North	KP9	<i>K. pneumoniae</i>	PXHZ00000000
2017	South	EC12	<i>E. coli</i>	PXIA00000000
2017	Beirut	KP4	<i>K. pneumoniae</i>	SDUE00000000
2017	South	EC10	<i>E. coli</i>	SDUD00000000
2017	El Qa'a	EC1	<i>E. coli</i>	SDUF00000000
2017	North	SM1	<i>S. marcescens</i>	PXIB00000000
2017	North	PR1	<i>P. alcalifaciens</i>	PXIC00000000
2017	Beirut	EC7	<i>E. coli</i>	SDUG00000000
2017	Beirut	EC8	<i>E. coli</i>	PXID00000000
2017	El Beqaa	KP10	<i>K. pneumoniae</i>	PXIE00000000
2017	South	PO1	<i>P. otitidis</i>	PXJI00000000
2018	El Qa'a	EC18	<i>E. coli</i>	SDPR00000000
2018	El Qa'a	EC19	<i>E. coli</i>	SDPS00000000
2018	El Qa'a	EC20	<i>E. coli</i>	VAUG01000000
2018	El Qa'a	EC21	<i>E. coli</i>	SDSF00000000
2018	El Qa'a	EC22	<i>E. coli</i>	SDSG00000000
2018	El Qa'a	EC23	<i>E. coli</i>	SDSH00000000
2018	North	EC30	<i>E. coli</i>	SGIV00000000
2018	South	EC27	<i>E. coli</i>	SGIX00000000
2018	Beirut	EC24	<i>E. coli</i>	SGIW00000000
2018	El Beqaa	EC35	<i>E. coli</i>	VDEX00000000
2018	El Beqaa	EC36	<i>E. coli</i>	VDEY00000000

Table 3: Detailed overview on the whole genome sequenced isolates that we chose for further characterization including the: season, year, site of isolation and NCBI accession numbers.

EC= *Escherichia coli*, KP= *Klebsiella pneumoniae*, PR1= *Providencia rettgeri*, PR2= *Providencia alcalifaciens*, SM= *Serratia marcescens*, and PO= *Pseudomonas otitidis*

Isolate	Time	Location	β -lactam	Plasmids	Gene	Genetic location
EC5	2017	El Qa'a	CTX-M-15, TEM-1B	IncI α , IncX3, IncFIB(K), IncFII(pCoo)	CTX-M-15	Chromosome
					TEM-1B	IncI α
KP8	2017	North	SHV-1, CTX-M-15, OXA-1, TEM-1B	No plasmids detected	CTX-M-15, OXA-1, TEM-1B, SHV-1	Chromosome
AB1	2017	North	ADC-25, OXA-132	No plasmids detected	ADC-1, OXA-132	Chromosome
KP9	2017	North	SHV-1	IncFII, IncFIB(K), IncR	SHV-1	Chromosome
EC12	2017	South	CTX-M-15, TEM-1B	IncFIA, IncFIB, IncFII(pRSB107)	CTX-M-15	Chromosome
					TEM-1B	Unknown
KP4	2017	Beirut	TEM-1B, SHV-11	IncFIA, IncFIB(K), IncFII(K), IncR	SHV-11	Chromosome
					TEM-1B	IncFIA, IncR
EC1	2017	El Qa'a	TEM-1B	IncFIB(K)	TEM-1B	IncFIB
SM1	2017	North	SRT-2, TEM-1B	No plasmids detected	SRT-2	Chromosome
					TEM-1B	Chromosome
EC7	2017	Beirut	TEM-1B	IncFIB, IncFII	TEM-1B	IncFIB, IncFII
EC8	2017	Beirut	CTX-M-15, TEM-1B	IncB/O/K/Z, IncFIA, IncFIB, IncFII	CTX-M-15, TEM-1B	IncB/O/K/Z
KP10	2017	El Beqaa	CTX-M-15, SHV-11, TEM-1B	IncB/O/K/Z, IncFIB(K), IncFIB, IncFII(K), IncQ1	SHV-11, CTX-M-15	Chromosome
					TEM-1B	IncFIB(K), IncFII(K)
PO1	2017	South	POM-1	No plasmids detected	POM-1	Chromosome
EC18	2018	El Qa'a	CMY-42	IncI α , IncFIA, IncFIB, IncFII, IncU, IncFII, IncY	CMY-42	IncFIA, IncFIB, IncFII, IncU
EC19	2018	El Qa'a	TEM-1B	IncFIA, IncFIB, IncFII	TEM-1B	IncFIA, IncFIB, IncFII
EC20	2018	El Qa'a	CTX-M-15, TEM-1B	IncFIB KN	CTX-M-15	Chromosome
					TEM-1B	IncFIB KN
EC21	2018	El Qa'a	CTX-M-15	IncFIA, IncFIB, IncFII, IncU, IncFII	CTX-M-15	IncFIA, IncFIB, IncFII
					CTX-M-15	IncI α
EC22	2018	El Qa'a	CTX-M-15, OXA-1, TEM-1B	IncI α , IncFIA, IncFIB, IncFII, IncU, IncFII	TEM-1B	IncFIA, IncFIB, IncFII
					OXA-1	chromosome
					NDM-5	IncFIA, IncFII
EC23	2018	El Qa'a	CMY-42, NDM-5	IncFIA, IncFII, IncFII, IncY	CMY-42	IncI α
					CTX-M-15	Phage
EC30	2018	North	CTXM-15, DHA-1, TEM-1B	IncFIB, IncFII, IncFII(pCoo)	TEM-1B, DHA-1	IncFIB, IncFII
					CTX-M-15, OXA-1	IncFIA, IncFIB, IncFII
EC27	2018	South	CTXM-15, OXA-1	IncFIA, IncFIB, IncFII	CTX-M-15, OXA-1	IncFIA, IncFIB, IncFII
EC24	2018	Beirut	CTXM-15, OXA-1	IncFIA, IncFIB, IncFII	CTX-M-15, OXA-1	Chromosome
EC35	2018	El Beqaa	CTXM-15	IncFIA, IncFII	CTX-M-15	Phage
EC36	2018	El Beqaa	CTXM-15, OXA-1, TEM-1B	IncFIA, IncFII	CTX-M-15, OXA-1, TEM-1B	IncFIA, IncFII

Table 4: Distribution of β -lactamase encoding genes, plasmid typing results and gene locations. *in silico* analysis was done using ResFinder v1.2, while plasmid Inc group determination was through PBRT-based PCR assays combined with *in silico* analysis using PlasmidFinder 1.3. Gene location was determined using conjugation assay and *in silico* analysis.

K-type		47	35	14	18	14	10	31	31	24	New Type
Polypeptides		CST									
others	FOF										
	NIT										
Miscellaneous Agents	SXT										
tetracycline	TET										
fluroquinolone	LVX										
	NOR										
	CIP										
aminoglycosides	OFX										
	GEN										
	TOB										
cephalosporins	CFM										
	FEP										
	CAZ										
	CRO										
	CTX										
	FOX										
	CXM										
	CFT										
monobactam	ATM										
carbapenems	MEM										
	ETP										
	IPM										
Penicillins	TZP										
	TIC										
	AMC										
	AMX										
Location		Beirut	North	North	North	Beirut	El Beqaa	North	North	North	South
Season		2017	2017	2017	2017	2017	2017	2017	2017	2017	2017
Isolate		KP4	KP1	KP2	KP8	KP3	KP10	KP6	KP9	KP7	KP5