Health care seeking behaviour and financial protection of patients with hypertension: a cross-sectional study in rural West Bengal, India

Sandipta CHAKRABORTY, MBBS MPH

MPH Graduate, Institute of Public Health Kalyani, Kalyani, Nadia, West Bengal 741251, India.

MD Candidate, Department of Preventive and Social Medicine, All India Institute of Hygiene and Public Health, West Bengal 700073, India.

Email: dr.sandipta@gmail.com

Rajesh Kumar RAI, MPhil MPH

Senior Research Scientist, Society for Health and Demographic Surveillance, Suri 731101, West Bengal, India.

Lown Scholar, Department of Global Health and Population, Harvard T H Chan School of Public Health, Boston, Massachusetts 02115, United States of America.

Doctoral Candidate, Department of Economics, University of Göttingen, Göttingen 37073, Germany.

Email: rajesh.iips28@gmail.com

Asit Kumar BISWAS, MBBS MAE

Dean, Institute of Public Health Kalyani, Kalyani, Nadia, West Bengal 741251, India.

Email: akbiswasprl@gmail.com

24 **Anamitra BARIK**, MBBS DipPH

25 *Field Director*, Society for Health and Demographic Surveillance, Suri, West Bengal 731101,
26 India.

27 *Senior Medical Officer*, Chest Clinic- Suri District Hospital, and Niramoy TB Sanatorium, Suri,
28 West Bengal 731101, India.

29 Email: anomitro2010@gmail.com

30

31 **Preeti GURUNG**, BSc MPH

32 *Epidemiology Activity Manager*, Medecins Sans Frontieres, Chiengkongpang, Churachandpur,
33 Manipur 795158, India.

34 Email: preetigrg04@gmail.com

35

36 **Devarsetty PRAVEEN**, MBBS MD PhD

37 *Program Head*, Primary Health Care Research, The George Institute for Global Health,
38 Hyderabad, Telangana 500082, India.

39 *Senior Lecturer*, Faculty of Medicine, University of New South Wales, Kensington NSW 2052,
40 Australia.

41 *Conjoint Professor*, Prasanna School of Public Health, Manipal Academy of Higher Education,
42 India

43 Email: dpraveen@georgeinstitute.org.in

44

45

Contributors: SC and AKB conceived and designed the study. SC conceptualized the data collection framework, analysed the data, and prepared the first draft of the manuscript. RKR, AKB, AB, PG, PD critically reviewed the manuscript, and provided feedback. All authors approved content presented in the study.

Abbreviations

aOR: adjusted odds ratio

BIRPOP: Birbhum Population Project

CI: confidence interval

NCDs: non-communicable diseases

OOPs: out of pocket payments

OR: odds ratio

uOR: unadjusted odds ratio

Address for correspondence

Sandipta Chakraborty

Department of Preventive and Social Medicine, All India Institute of Hygiene and Public Health,
West Bengal 700073, India.

Email: dr.sandipta@gmail.com

Abstract

Background: Elevated blood pressure or hypertension is responsible for around 10 million annual deaths globally, and people residing in low and middle-income countries are disproportionately affected by it. India is no exception, where low rate of treatment seeking for hypertension coupled with widespread out-of-pocket payments (OOPs) have been a challenge. This study assessed the pattern of health care seeking and financial protection along with the associated factors among hypertensive individuals in a rural district of West Bengal, India.

Method and findings: A cross-sectional study was conducted in Birbhum district of the state of West Bengal, India during 2017-2018, where 300 individuals with hypertension were recruited randomly from a pre-defined list of individuals with hypertension in the district. Healthcare seeking along with two instance of financial protection –OOPs and relative expense, were analysed. Findings indicated that, of all hypertensives, 47% were not on treatment, 80% preferred private healthcare, and 91% of them had wide-spread OOPs. Cost of medication being a major share of expenses followed by significant transport cost to access public health care facility. Multivariable logistic regression analysis indicated longer duration of disease and private health care seeking were associated with more incident of OOPs. Results from linear regression modelling (generalized linear model) demonstrated association of co-morbidities with higher relative expenditure. Individuals belonging to poor economic group suffered from a high relative expense, compared to the richest.

Conclusion: This study suggested that individuals with hypertension had poor health care seeking, preferred private health care and had suboptimal financial protection. Hypertensives

from economically poorer section had higher burden of health expenditure for treatment of hypertension, which indicated gaps in equitable health care for the control of hypertension.

Introduction

Globally, non-communicable diseases (NCDs) contribute to a major share of the disease burden, where countries with differential level of development and varied phases of epidemiological transition have witnessed a significant rise in overall morbidity and mortality from NCDs [1-3]. Among all NCDs, cardiovascular diseases (ischaemic heart disease and stroke) are listed as the major cause of death worldwide, with hypertension (commonly defined as a systolic blood pressure ≥ 140 or diastolic blood pressure ≥ 90) being the most significant risk factor causing significant amount of premature deaths globally [4, 5]. Despite the high burden of hypertension, health system responses like health service delivery, health information and health financing for hypertension is suboptimal, especially in low and middle-income countries (LMICs) [6-10]. Evidence suggests that people seeking health care for NCDs bear significant and unjustified financial burden characterised by huge out-of-pocket payments (OOPs), often leading to irregular and absence of treatment seeking due to financial difficulties [10, 11]. In addition, studies show that overall health care seeking for blood pressure management is low and shared among public and private facilities [12, 13].

In India, between one-quarter to one-third of adults, aged 18 years or more, have hypertension. This remains a major threat to Indian healthcare system [14-16]. In the year 2010 the federal Indian government introduced the National Programme for Prevention and Control of Cancer, Diabetes, Cardiovascular Diseases and Stroke (NPCDCS) with hypertension and diabetes as the main focus areas. In addition, in 2017, the government launched the National Health Policy targeting 25% reduction in premature mortality occurring from cardiovascular diseases, cancer, diabetes or chronic respiratory diseases by 2025 [17, 18]. The main focus for research on hypertension in India is primarily on the risk factors of hypertension while few actually explored

the health care utilization and service expenses among hypertensive individuals, as evidenced from the PubMed/MEDLINE database search [19-22]. From the perspective of health system strengthening and population health management, understanding the local preferences and health system capacity is essential. In this paper, we present the pattern of health care seeking, financial protection and its predictors among patients with hypertension in rural West Bengal. This study was conducted as a part of a broad study assessing the Capacity of Health Systems to combat the Emergence of Hypertension (COHESION study). In a comprehensive way COHESION study analysed blood pressure control, health care seeking, financial protection and health system responsiveness among adult hypertensive population (Unpublished data) .

Materials and Methods

Study setting, design and sampling

COHESION study is a population-based cross-sectional study, conducted in a population cohort of Birbhum Population Project (BIRPOP), a health and demographic surveillance system (HDSS) functioning under the ambit of Society for Health and Demographic Surveillance (www.shds.co.in), located in the Birbhum district of the state of West Bengal, India, between November 2017 and February 2018. BIRPOP spreads over four administrative blocks (namely Suri I, Sainthia, Mohammad Bazar and Rajnagar) out of a total of 19 blocks in district Birbhum. At its inception in 2008, BIRPOP included a sample of over 12,000 households selected by multistage stratified sampling method and has been periodically collecting information on indicators related to public health and demography. Till date, BIRPOP had completed three rounds of follow-up surveys, in 2008-09, 2012-13, and 2016-17 [23]. COHESION study was based on BIRPOP's 2016-17 survey where blood pressure was measured for 12,255 individuals

aged ≥ 18 years. Those recorded with high blood pressure (systolic blood pressure (SBP) ≥ 140 mm of Hg and/or diastolic blood pressure (DBP) ≥ 90 mm of Hg) or reported taking anti-hypertensive medication of any form were included in the hypertensive cohort [23, 24]. Details about the blood pressure measurement survey at BIRPOP has previously been published elsewhere [25]. Of the hypertensive cohort, 310 individuals were randomly selected for this study. Sample size was calculated using CDC Epi-infoTM version 7.2, assuming 50% prevalence for hypertension control in all hypertensives, 7.5% of error and confidence interval of 99%. With the addition of 5% non-response rate, final sample size was 310 individuals of which 300 interviews were conducted. Terminally ill and mentally challenged individuals, diagnosed by a physician, were not considered for participation in the study. Data were collected by trained surveyors using Computer Assisted Personal Interview (CAPI) technique [26]. A rigorous protocol for survey monitoring was followed to assure the quality of the data being collected.

Outcome measurement

To understand the health care seeking behaviour, patients were asked if they were taking any medication for blood pressure control and have been visiting to any healthcare provider. Patients with a history of intake of daily medication for hypertension in the preceding four weeks were considered to be on ‘regular medication’. Those with a history of visit to any health care provider at least once in the last six months for treatment or follow-up care of hypertension, were noted to have ‘regular medical consultation’. Patients who had both of the above (regular medication and regular visit to physician) were labelled as “having regular treatment for hypertension”. Those reported only regular medication, identified as having “regular medication only”. Patients, currently not on any medication or consultation for last one year or never sought any treatment

for hypertension, were labelled as “not on treatment”. The rest were categorised as ‘patients on irregular treatment’

Two outcomes in relation to cost of treatment, were analysed in this study – i) Out-of-pocket payments (OOPs), and ii) relative cost for hypertension care. Considering the varied practice of health care seeking behaviour, total expected OOPs were calculated considering multiple expenses. Expenses paid for medical consultation, transport and others like food, lodging etc. during the consultation in the last medical visit, and cost of blood pressure lowering medication if taken for a month, all together added to obtain total expected cost. Monthly per capita expenditure (MPCE) was calculated as monthly total consumer expenditure in a household over all items of consumption divided by the household size (total number of persons in the household) and was used as the proxy measure of the economic status [27]. Based on the MPCE, the participants were divided into four quartile classes and categorised into relative economic groups: poorest, lower-middle, upper-middle and richest class. Relative cost of seeking healthcare for hypertension for an individual was defined as percentage of MPCE incurred for OOPs [19].

Covariates

Based on existing literature from developing countries, a range of potential covariates were considered.

Social demographic characteristics: This included age in completed years (<50, 50-63, >63), gender (female and male), educational attainment (secondary and above, upper primary, primary, and illiterate/below primary), social group (other backward classes, scheduled caste/ scheduled tribes and others), religion (Hinduism and Islam), civil status (living with partner, and not living

with partner), employment status (service/business, labourer, homemaker/retired/student, and unemployed), and economic status based on MPCE quartile distribution (high, upper-middle, lower-middle, and poor)

Hypertension related variables: This included duration of hypertension (<5 years, ≥5 years, and not sure/don't know), co-morbidity (no and yes), regularity of treatment of hypertension (as elaborated before), type of health facility accessed (public and non-public), and healthcare provider like, public physician, private physician, AYUSH (*Ayurveda*, Yoga and Naturopathy, *Unani*, *Siddha* and Homoeopathy) doctor, and informal health care practitioner (Quack) [28]. Comorbidity refers to self-report about any of the diseases like diabetes, dyslipidaemia, chronic kidney disease or cardiovascular disease in addition to hypertension.

Statistical analysis

Bivariate and multivariable analyses were performed to attain the study objectives. Means and proportions were presented with 95% confidence intervals. For the purpose of regression modelling, a directed acyclic graph (DAG) was developed, based on causal diagram theory [29] and review of existing literature. Binary logistic regression was deployed to understand the predictors of OOPs, whereas linear regression by generalized linear models (GLM) was used to assess the relative costs. Measures of association were presented as odds ratio (OR) with 95% confidence intervals (CIs) with value “1” as the null point. GLM is preferred because of abundance of zero values in relative cost data and a possible non-parametric distribution [30]. With the linear modelling, the association is expressed with the estimated coefficient (Coeff) and associated 95% CIs, indicating direction of association with value “zero” as the null point. Data analysis were carried out using a statistical package - Stata, version 12.0 and *p* value was

considered to interpret the significance of observed association. Qualitative interpretation based on p value (significant/non-significant based on conventional cut off) was judged cautiously, keeping with the study design and limitations.

Ethics statement

Ethical approval was granted by institutional review board of Society for health and Demographic Surveillance. Written informed consent was obtained from all participants prior to enrolment in the study. Irrespective of their participation status, all, who were approached to participate in the study were provided with a leaflet on healthy lifestyle, health education related to hypertension and other NCDs written in local language.

Results

In total, 310 were approached to participate in this study, and 300 finally participated. **Table 1** outlines the descriptive characteristics of all the participants. The mean age of the participants was 55.99 ± 12.46 years. More than half of the participants were female and were illiterate or had not completed their primary education. Majority of the participants were Hindus and homemaker/retired/students by profession. Over 35% ($n=106$) of participants had hypertension for ≥ 5 years, and 20% ($n= 60$) had a co-morbid condition. Over 47% ($n=141$) of the participants were not on treatment, and among individuals receiving treatment, over 80% ($n=128$) sought healthcare from non-public healthcare provider. Over 90% ($n=144$) of those who sought care for blood pressure treatment incurred some OOPs. Expected cost for seeking complete care for hypertension per month was over ₹ 306 ($> \$4.5$) and relative cost per month was 13.5% of the MPCE (**Table 1**). Further analysis (not shown separately) revealed that the median of relative

cost was higher for those seeking care in non-public healthcare facility (median: 10.7%) compared to the public healthcare provider (2.1%). The median of OOPs was the largest for purchase of medicines (47.7%) in those seeking private healthcare, while it was transport and other costs in those seeking care (51.3%) followed by purchase of medicines (37.5%) from a public healthcare facility. (**Figure 1**).

Fifteen individuals were reported incurring no OOPs for the usual treatment for hypertension. Majority (n=9) were female, aged between 50 to 63 years (n=9), Hindu (n=12), general caste (n=9) with below primary or no formal education (n=10), home maker/ retired (n=10) and belongs to upper-middle class (n=6) of the economic strata of the study population.

Table 2 shows lower odds of having OOPs among participants aged 50-63 years and 63 years and above compared to participants below 50 years. Males when compared to females, and homemaker/retired /student, labourer and unemployed when compared to those in service/business had relatively lower odds of incurring any OOPs. Compared to the richest economic class the poorest had lower odds of having any OOPs, in unadjusted model (uOR_{poorest} 0.22 (CI: 0.04-1.21)). Having hypertension for five years or more (uOR 5.14 (CI: 1.39-19.01) and aOR 5.68 (CI: 1.24-25.99)) and seeking treatment from private establishments (uOR 26.32 (CI: 6.80-101.93) and aOR 34.33 (CI: 4.82-244.68)) were positively associated with OOPs.

Linear regression (**Table 3**) demonstrated lower relative expenses among people with primary or below level of schooling, compared to highest educational group; (Adjusted Coefficient (aCoeff) completed primary -10.65 (CI: -19.78, -1.51) and aCoeff_{no formal education/below primary} -11.60 (CI: -20.88, -2.32)). The unemployed individuals had more relative expenses compared to those engaged in

service/business (Unadjusted Coefficient ($uCoeff$)_{unemployed} 8.71 (CI: 0.04,17.38) and $aCoeff$ _{unemployed} 9.34 (CI: -1.74,20.43)). The poorest, lower-middle and upper-middle class had 11, 8 and 7 units of more relative expenses respectively, compared to the richest economic class ($aCoeff$ _{poorest} 11.27 (CI: 3.82,18.71); $aCoeff$ _{lower-middle} 7.83 (CI: 0.65,15.00) and $aCoeff$ _{upper-middle} 7.25 (CI: 0.80,13.70)) (**Figure 2**). Presence of co-morbidity and seeking treatment from private establishments were both associated with more relative expenses ($aCoeff$ _{one or more co-morbidity} 10.28 (CI: 4.96,15.61); reference group: no co-morbidity and $aCoeff$ _{private establishment} 11.55 (CI: 5.74,17.37); reference group: government institution). Similarly, seeking treatment from private doctors, informal practitioners and AYUSH doctors/others were associated with more relative expenses ($aCoeff$ _{private Doctors} 18.43 (CI: 12.13, 24.73), $aCoeff$ _{informal healthcare provider} 5.96 (CI: -0.36, 12.28), $aCoeff$ _{AYUSH/ Other} 10.28 (CI: 2.56, 17.99)) when compared to those seeking treatment from government doctors.

Discussion

India has witnessed an increasing burden of hypertension, which demands urgent attention from the public health researchers, program and policy makers. To add on to the existing body of literature on prevention of hypertension in India, this study aims to understand characteristics of healthcare seeking and financial protection among hypertensive population in West Bengal, India. The state of West Bengal recorded nearly 25% of total annual deaths and 13% of disability adjusted life years (DALYs) attributed to hypertension [15, 31]. This study revealed poor health care seeking behaviour, preference of private health facilities and high OOPs among patients who sought care for hypertension. Regression analysis adjusted for potential covariates indicate that OOPs are associated with age, sex, occupation, duration of hypertension, and place of

treatment seeking for hypertension, while relative expense is associated with education, occupation, economic class, comorbidity, place of treatment and healthcare provider.

The population under study were relatively older, female predominated, had low education level, and majority were retired/homemaker. This distribution was similar to other studies where hypertension prevalence was more among elderly, females, and in poor socio-economic strata [32, 33]. The findings of poor health care seeking for blood pressure control, attributed to low awareness, affordability and availability of health care services (unpublished data). Among the hypertensives seeking treatment, OOPs were extensively reported. This scenario corroborates with previous findings of sub-optimal health system response for blood pressure control care [10, 11, 13, 14, 20, 34-37]. However better system response was associated with substantial improvement in indicators like awareness, treatment and control of hypertension in a few developed countries [38]. Similar to other studies, private establishments were major places for seeking treatment and Government institutions played a minor role for management of hypertension. Similarly, majority sought consultation from private physicians and informal healthcare providers [13, 20, 22]. The prevalence of OOPs and extent of relative cost varied between service utilization from government to non-government sources as well as with different service providers. The findings related to OOPs in this study are in line with previous reports including a report of the WHO Study on global AGEing and adult health (SAGE) but the significant variation observed in OOPs across government and private institutions in this study is found to be novel [19, 20]. Earlier studies found medicine purchase as the major share for OOPs [19-21] which corroborates with the findings from this study, however transport and other costs are also found to impose a substantial share of OOPs in Government set-up, possibly indicating

better accessibility for the private treatment sources in local level compared to Government sources. This could also justify the increased usage of private facilities for hypertension management. Contrasting with findings from other studies, the present study reported lower incident of OOPs among male and those belonging to 50 years or above age group[10, 20]. Though relative cost (%) and high OOPs was proportional with level of education, the relative cost was found to be inversely related with disadvantageous economic class. These findings point towards potential issues of social justice and inequity which share a complex interrelationship [19, 20]. This might be related to poor treatment seeking behaviour among patients with low education and economic status (jointly the lower socio-economic class) owing to low awareness, financial constraint and limited access to healthcare, which may have led to lower possibility of having OOPs. But despite these barriers, patients who sought treatment experienced inequitable financial burden. Similar explanation may be applied for the unemployed group, having more extent of relative expenses while seeking care but less OOPs. Lower OOPs among homemaker/retired individuals may be due to more utilization of government health facilities, compared to the service holders/businessmen who generally have less opportunity to visit government outpatient services due to its fixed schedule. Longer duration of hypertension and existence of comorbid conditions require more intense therapy resulting in more possibility of OOPs and more relative cost (%) [10].

Limitation of the study should be interpreted in light of the results. Firstly, being a cross-sectional study, temporal ambiguity cannot be ruled out. Secondly, as most of variables under study are information based on recall, some chances for recall errors may be present. Thirdly, measurement of exact expenditure and assessing economic status could be debated. To counter

the variability of health care seeking, health care expenditure related to hypertension management was calculated as expected cost for having complete care. This may have over-represented the relative cost (%) for treatment to some extent. Effects of residual confounding also cannot be ruled out. Within purview of limitations, considering the geographic and demographic uniqueness of the Birbhum population, the findings of this study should be interpreted cautiously for other settings. Despite these limitations, the study contributes tremendously to the existing literature in terms of unique study setting and use of pre-tested and validated study tools. The findings from the study suggest suboptimal financial protection of population for hypertension care. The aspect of awareness generation and evaluation of existing programs on NCDs might be needed for a better financial protection mechanism to people with hypertension.

Data statement

Data of COHESION study and associated codebook can be accessed through [10.5281/zenodo.3911116](https://doi.org/10.5281/zenodo.3911116).

Funding

SC received fellowship from the Department of Health and Family Welfare, Government of West Bengal, India. The fellowship provider had no role in the design/conduct of the study, collection/analysis/interpretation of the data, and preparation/review/approval of the manuscript.

Conflict of interest statement

None of the authors have any competing interest that could influence or bias the study design, settings, conduct, outcome and reporting.

Acknowledgements

Authors are indebted to the members of Society for Health and Demographic Surveillance, West Bengal, India, for helping execute the study.

References

1. Abubakar I, Tillmann T, Banerjee A. Global, regional, and national age-sex specific all-cause and cause-specific mortality for 240 causes of death, 1990-2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet*. 2015;385(9963):117-71.
2. Habib SH, Saha S. Burden of non-communicable disease: global overview. *Diabetes & Metabolic Syndrome: Clinical Research & Reviews*. 2010;4(1):41-7.
3. Murray CJ, Ezzati M, Flaxman AD, Lim S, Lozano R, Michaud C, et al. GBD 2010: design, definitions, and metrics. *Lancet*. 2012;380(9859):2063-6.
4. World Health Organization. The top 10 causes of death 2020 [updated 9 December 2020; cited 2020, 20th December]. [Available from: <https://www.who.int/news-room/fact-sheets/detail/the-top-10-causes-of-death>].

- 384 5. World Health Organization. Hypertension 2019 [updated 13 September 2019; cited 2020,
385 20th December]. [Available from: [https://www.who.int/news-room/fact-](https://www.who.int/news-room/fact-sheets/detail/hypertension)
386 [sheets/detail/hypertension](https://www.who.int/news-room/fact-sheets/detail/hypertension)].
- 387 6. Alshamsan R, Lee JT, Rana S, Areabi H, Millett C. Comparative health system
388 performance in six middle-income countries: cross-sectional analysis using World Health
389 Organization study of global ageing and health. Journal of the Royal Society of Medicine.
390 2017;110(9):365-75.
- 391 7. Feng XL, Pang M, Beard J. Health system strengthening and hypertension awareness,
392 treatment and control: data from the China Health and Retirement Longitudinal Study. Bull
393 World Health Organ. 2014;92(1):29-41.
- 394 8. Ibrahim MM, Damasceno A. Hypertension in developing countries. Lancet.
395 2012;380(9841):611-9.
- 396 9. Peck R, Mghamba J, Vanobberghen F, Kavishe B, Rugarabamu V, Smeeth L, et al.
397 Preparedness of Tanzanian health facilities for outpatient primary care of hypertension and
398 diabetes: a cross-sectional survey. Lancet Glob Health. 2014;2(5):e285-92.
- 399 10. Wang Q, Fu AZ, Brenner S, Kalmus O, Banda HT, De Allegri M. Out-of-pocket
400 expenditure on chronic non-communicable diseases in sub-Saharan Africa: the case of rural
401 Malawi. PLoS One. 2015;10(1):e0116897.

- 402 11. World Health Organization. Impact of out-of-pocket payments for treatment of non-
403 communicable diseases in developing countries: a review of literature. 2011.
- 404 12. Bovet P, Gervasoni JP, Mkamba M, Balampama M, Lengeler C, Paccaud F. Low
405 utilization of health care services following screening for hypertension in Dar es Salaam
406 (Tanzania): a prospective population-based study. BMC Public Health. 2008;8(1):407.
- 407 13. Baliga SS, Gopakumaran PS, Katti SM, Mallapur MD. Treatment seeking behavior and
408 health care expenditure incurred for hypertension among elderly in urban slums of Belgaum
409 City. Community Med. 2013;4(2):227-30.
- 410 14. Anchala R, Kannuri NK, Pant H, Khan H, Franco OH, Di Angelantonio E, et al.
411 Hypertension in India: a systematic review and meta-analysis of prevalence, awareness, and
412 control of hypertension. J Hypertens. 2014;32(6):1170-7.
- 413 15. Indian Council of Medical Research; Public Health Foundation of India and Institute for
414 Health Metrics and Evaluation. India: Health of Nation's States - The India State- level Disease
415 Burden Initiative. New Delhi, India: ICMR, PHFI and IHME; 2017. 2017.
- 416 16. Gupta R, Gaur K, CV SR. Emerging trends in hypertension epidemiology in India. J Hum
417 Hypertens. 2019;33(8):575-87.
- 418 17. Directorate General of Health Services; Ministry of Health & Family Welfare
419 Government of India. National Programme for Prevention and Control of Cancer, Diabetes,

- 420 Cardiovascular Diseases and Stroke (NPCDCS) 2017 [updated 26 August 2020; cited 2021
- 421 February 22]. [Available from:
- 422 http://dghs.gov.in/content/1363_3_NationalProgrammePreventionControl.aspx].
- 423 18. Ministry of Health & Family Welfare GoI. National Health Policy 2017 [cited 2021
- 424 February 22]. [Available from:
- 425 https://www.nhp.gov.in/nhpfiles/national_health_policy_2017.pdf].
- 426 19. Bhojani U, Thriveni B, Devadasan R, Munegowda C, Devadasan N, Kolsteren P, et al.
- 427 Out-of-pocket healthcare payments on chronic conditions impoverish urban poor in Bangalore,
- 428 India. BMC Public Health. 2012;12(1):990.
- 429 20. Brinda EM, Kowal P, Attermann J, Enemark U. Health service use, out-of-pocket
- 430 payments and catastrophic health expenditure among older people in India: The WHO Study on
- 431 global AGEing and adult health (SAGE). J Epidemiol Community Health. 2015;69(5):489-94.
- 432 21. Engelgau MM, Karan A, Mahal A. The Economic impact of Non-communicable
- 433 Diseases on households in India. Global Health. 2012;8(1):9.
- 434 22. Kanungo S, Mahapatra T, Bhowmik K, Saha J, Mahapatra S, Pal D, et al. Patterns and
- 435 predictors of undiagnosed and uncontrolled hypertension: observations from a poor-resource
- 436 setting. J Hum Hypertens. 2017;31(1):56-65.

- 437 23. Ghosh S, Barik A, Majumder S, Gorain A, Mukherjee S, Mazumdar S, et al. Health &
438 Demographic Surveillance System Profile: The Birbhum population project (Birbhum HDSS).
439 Int J Epidemiol. 2015;44(1):98-107.
- 440 24. Chobanian AV, Bakris GL, Black HR, Cushman WC, Green LA, Izzo JL, Jr., et al. The
441 Seventh Report of the Joint National Committee on Prevention, Detection, Evaluation, and
442 Treatment of High Blood Pressure: the JNC 7 report. JAMA. 2003;289(19):2560-72.
- 443 25. Ghosh S, Mukhopadhyay S, Barik A. Sex differences in the risk profile of hypertension:
444 a cross-sectional study. BMJ Open. 2016;6(7):e010085.
- 445 26. Baker RPJSSCR. New technology in survey research: Computer-assisted personal
446 interviewing (CAPI). 1992;10(2):145-57.
- 447 27. Howe LD, Galobardes B, Matijasevich A, Gordon D, Johnston D, Onwujekwe O, et al.
448 Measuring socio-economic position for epidemiological studies in low- and middle-income
449 countries: a methods of measurement in epidemiology paper. Int J Epidemiol. 2012;41(3):871-
450 86.
- 451 28. Das J, Chowdhury A, Hussam R, Banerjee AV. The impact of training informal health
452 care providers in India: A randomized controlled trial. Science (New York, NY).
453 2016;354(6308).

- 454 29. Rothman KJ, Greenland S, Lash TL. Modern epidemiology. 3rd ed: Lippincott Williams
455 & Wilkins; 2008. p. 183-209.
- 456 30. Matsaganis M, Mitrakos T, Tsakloglou P. Modelling health expenditure at the household
457 level in Greece. The European Journal of Health Economics. 2009;10(3):329-36.
- 458 31. Institute for Health Metrics and Evaluation (IHME). GBD India Compare | Viz Hub 2019
459 [cited 2021 February 22]. [Available from: <https://vizhub.healthdata.org/gbd-compare/india>].
- 460 32. Irazola VE, Gutierrez L, Bloomfield GS, Carrillo-Larco RM, Dorairaj P, Gaziano T, et al.
461 Hypertension Prevalence, Awareness, Treatment, and Control in Selected Communities of Nine
462 Low-and Middle Income Countries: Results From the NHLBI/UHG Network of Centers of
463 Excellence for Chronic Diseases. Global heart. 2016;11(1):47.
- 464 33. Sarki AM, Nduka CU, Stranges S, Kandala NB, Uthman OA. Prevalence of
465 Hypertension in Low- and Middle-Income Countries: A Systematic Review and Meta-Analysis.
466 Medicine (Baltimore). 2015;94(50):e1959.
- 467 34. Chow CK, Teo KK, Rangarajan S, Islam S, Gupta R, Avezum A, et al. Prevalence,
468 awareness, treatment, and control of hypertension in rural and urban communities in high-,
469 middle-, and low-income countries. JAMA. 2013;310(9):959-68.

- 470 35. Kaur P, Rao SR, Radhakrishnan E, Rajasekar D, Gupte MD. Prevalence, awareness,
471 treatment, control and risk factors for hypertension in a rural population in South India. Int J
472 Public Health. 2012;57(1):87-94.
- 473 36. Singh AK, Kalaivani M, Krishnan A, Aggarwal P, Gupta SK. Prevalence, awareness,
474 treatment and control of hypertension among elderly persons in an urban slum of Delhi, India.
475 Indian Journal of Medical Specialities. 2014;5(1):7-10.
- 476 37. Tocci G, Ferrucci A, Pontremoli R, Ferri C, Rosei E, Morganti A, et al. Blood pressure
477 levels and control in Italy: comprehensive analysis of clinical data from 2000–2005 and 2005–
478 2011 hypertension surveys. Journal of human hypertension. 2015;29(11):696.
- 479 38. Joffres M, Falaschetti E, Gillespie C, Robitaille C, Loustalot F, Poulter N, et al.
480 Hypertension prevalence, awareness, treatment and control in national surveys from England, the
481 USA and Canada, and correlation with stroke and ischaemic heart disease mortality: a cross-
482 sectional study. BMJ Open. 2013;3(8):e003423.
- 483
- 484
- 485
- 486
- 487
- 488
- 489
- 490

Table 1. Characteristics of sampled hypertensive population.

Background characteristics	n	mean or percentage (95% CI)
Age	300	55.99 (54.58-57.41)
Total expected cost of seeking complete care for hypertension (₹) *	159	306.49 (257.65-355.33)
Relative cost (%) for treatment of hypertension*	159	13.52 (11.13-15.90)
Age group (years)		
< 50	101	33.67(28.29-39.04)
50 - 63	107	35.67(30.22-41.12)
>63	92	30.67(25.42-35.91)
Education		
Completed Secondary or above	48	16.00(11.83-20.17)
Completed Upper-primary	46	15.33(11.23-19.43)
Completed Primary	56	18.67(14.23-23.10)
Illiterate/ Below primary	150	50.00(44.31-55.69)
Sex		
Female	183	61.00(55.45-66.55)
Male	117	39.00(33.45-44.55)
Social group		
Others	140	46.67(40.99-52.34)
OBC	42	14.00(10.05-17.95)
SC/ST	118	39.33(33.77-44.89)
Religion⁺		
Hinduism	225	75.25(70.33-80.17)
Islam	74	24.75(19.83-29.67)
Civil status		
Living with partner	195	65.00(59.57-70.43)
Not living with partner	105	35.00(29.57-40.43)
Occupation		
Service/Business	65	21.67(16.98-26.36)
Labourer	47	15.67(11.53-19.80)
Homemaker/Retired/ Student	160	53.33(47.66-59.01)
Unemployed	28	9.33(6.02-12.64)

Economic Class		
Richest	75	25.00(20.07-29.93)
Upper Middle	79	26.33(21.32-31.35)
Lower-middle	70	23.33(18.52-28.15)
Poorest	76	25.33(20.38-30.28)
Duration of Hypertension (years)		
<5	141	47.00(41.32-52.68)
≥5	106	35.33(29.89-40.77)
Not sure/don't know	53	17.67(13.32-22.01)
Co-morbidity		
No	240	80.00(75.45-84.55)
Yes	60	20.00(15.45-24.55)
Regular treatment for hypertension		
On regular consultation & Medication	71	23.67(18.83-28.50)
On regular medication only	39	13.00(9.17-16.83)
On irregular treatment	49	16.33(12.13-20.54)
Not on treatment	141	47.00(41.32-52.68)
Place of treatment for hypertension*		
Public	31	19.50(13.27-25.72)
Non-public	128	80.50(74.28-86.73)
Health care provider*		
Public physician	30	18.87(12.72-25.02)
Private physician	63	39.62(31.94-47.31)
AYUSH doctor/ Other	19	11.95(6.85-17.05)
Informal healthcare provider	47	29.56(22.39-36.73)
OPP*		
Absent	15	9.43(4.84-14.03)
Present	144	90.57(85.97-95.16)

□: Indian National Rupee; CI: Confidence Interval; OBC: Other backward classes; SC: Scheduled caste; ST: Scheduled tribe; AYUSH: *Ayurveda*, Yoga and Naturopathy, *Unani*, *Siddha* and Homoeopathy; OPP: Out of Pocket Payments

* Sample characteristics is based on 159 participants representing patients seeking treatment for hypertension

⁺ One person did not share information on religion.

498

499

500

501

502

503

504

505

506

Table 2. Odds of out-of-pocket payment.

	Unadjusted		Adjusted	
	OR (95% CI)	p	OR (95% CI)	p
Age group (years)				
< 50	1.00		1.00	
50 - 63	0.14 (0.02-1.13)	0.06	0.14 (0.02-1.38)	0.09
>63	0.29 (0.03-2.60)	0.27	0.47 (0.04-5.88)	0.56
Education				
Completed Secondary or above	1.00		1.00	
Completed Upper-primary	0.89 (0.05-15.00)	0.93	0.39 (0.01-12.94)	0.60
Completed Primary	0.33 (0.03-3.41)	0.35	0.20 (0.01-4.22)	0.30
Illiterate/ Below primary	0.24 (0.03-2.00)	0.19	0.09 (0.00-2.49)	0.15
Sex				
Female	1.00		1.00	
Male	0.70 (0.24-2.10)	0.53	0.08 (0.01-0.71)	0.02
Social group				
Others	1.00		1.00	
OBC	2.19 (0.26-18.37)	0.47	3.13 (0.28-34.91)	0.35
SC/ST	1.09 (0.34-3.43)	0.89	3.36 (0.60-18.97)	0.17
Religion				
Hinduism	1.00		1.00	
Islam	1.40 (0.37-5.22)	0.62	2.30 (0.40-13.39)	0.35
Civil status				
Living with partner	1.00		1.00	
Not living with partner	0.82 (0.28-2.37)	0.71	0.95 (0.25-3.66)	0.94
Occupation				
Service/Business	1.00		1.00	
Labourer	0.16 (0.01-1.91)	0.15	0.08 (0.00-1.65)	0.10
Homemaker/Retired/ Student	0.27 (0.03-2.18)	0.22	0.04 (0.00-0.79)	0.03
Unemployed	0.25 (0.02-2.97)	0.27	0.17 (0.01-3.65)	0.26
Economic Class				
Richest	1.00		1.00	
Upper Middle	0.27 (0.05-1.42)	0.12	0.47 (0.07-3.04)	0.42

Lower-middle	0.57 (0.08-4.28)	0.59	1.14 (0.12-11.18)	0.91
Poorest	0.22 (0.04-1.21)	0.08	0.40 (0.06-2.91)	0.37
Co-morbidity				
No	1.00		1.00	
Yes	0.74 (0.24-2.31)	0.61	0.54 (0.13-2.24)	0.39
Duration of Hypertension (years)				
<5	1.00		1.00	
≥5	5.14 (1.39-19.01)	0.01	5.68 (1.24-25.99)	0.03
Place of treatment seeking for hypertension				
Public	1.00		1.00	
Non-public	26.32 (6.80-101.93)	<0.01	34.33(4.82-244.68)	<0.01

CI: Confidence Interval; OBC: Other backward classes; SC: Scheduled caste; ST: Scheduled tribe; OR: Odds ratio

516

Table 3. Associates of relative expenses.

	Unadjusted		Adjusted	
	β (95% CI)	p	β (95% CI)	p
Age group (years)				
< 50	0.00		0.00	
50 - 63	-2.39 (-8.55,3.77)	0.45	-3.22 (-9.84,3.40)	0.34
>63	0.28 (-5.78,6.33)	0.93	-1.66 (-8.51,5.19)	0.64
Education				
Completed Secondary or above	0.00		0.00	
Completed Upper-primary	-2.58 (-10.82,5.67)	0.54	-5.17 (-14.14,3.80)	0.26
Completed Primary	-3.15 (-11.02,4.73)	0.43	-10.65 (-19.78,-1.51)	0.02
Illiterate/ Below primary	-1.06 (-7.69,5.56)	0.75	-11.60 (-20.88,-2.32)	0.01
Sex				
Female	0.00		0.00	
Male	-0.94 (-5.99,4.10)	0.71	-3.39 (-10.90,4.13)	0.38
Social group				
Others	0.00		0.00	
OBC	0.43 (-6.95,7.82)	0.91	-2.02 (-9.58,5.55)	0.60
SC/ST	3.44 (-1.78,8.66)	0.20	5.36 (-1.27,11.98)	0.11
Religion				
Hinduism	0.00		0.00	
Islam	-0.07 (-5.54,5.40)	0.98	3.23 (-3.37,9.82)	0.34
Civil status				
Living with partner	0.00		0.00	
Not living with partner	1.11(-3.68,5.90)	0.65	1.85 (-3.62,7.32)	0.51
Occupation				
Service/Business	0.00		0.00	
Labourer	5.57 (-4.40,15.54)	0.27	5.88 (-5.04,16.79)	0.29
Homemaker/Retired/ Student	1.59 (-4.38,7.56)	0.60	0.85 (-7.88,9.59)	0.85
Unemployed	8.71 (0.04,17.38)	0.05	9.34 (-1.74,20.43)	0.09
Economic Class				
Richest	0.00		0.00	

Upper Middle	5.54 (-0.38,11.46)	0.07	7.25 (0.80,13.70)	0.03
Lower-middle	5.73 (-0.97,12.43)	0.09	7.83 (0.65,15.00)	0.03
Poorest	10.39 (3.82,16.95)	0.00	11.27 (3.82,18.71)	0.00
Co-morbidity				
No	0.00		0.00	
Yes	7.74 (2.59,12.89)	0.00	10.28 (4.96,15.61)	<0.01
Duration of Hypertension (years)				
<5	0.00		0.00	
≥5	1.64 (-3.10,6.37)	0.50	2.17 (-2.62,6.97)	0.37
Place of treatment seeking for hypertension				
Public	0.00		0.00	
Non-public	9.35(3.56,15.14)	0.00	11.55 (5.74,17.37)	<0.01
Health care provider				
Public physician	0.00		0.00	
Private physician	14.38 (8.24,20.51)	<0.01	18.43 (12.13,24.73)	<0.01
AYUSH doctor/ Other	5.39 (-2.72,13.50)	0.19	10.28 (2.56,17.99)	0.01
Informal healthcare provider	3.40 (-3.07,9.86)	0.30	5.96 (-0.36,12.28)	0.06

CI: Confidence Interval; OBC: Other backward classes; SC: Scheduled caste; ST: Scheduled tribe; AYUSH: *Ayurveda*, Yoga and Naturopathy, *Unani*, *Siddha* and Homoeopathy; OPP: Out of Pocket Payments; β : Coefficient

517
518
519
520
521
522
523
524
525
526
527
528
529
530

Figure 1: Median of OOPs share (%) and relative OOPs (%) across the place of treatment seeking

Figure 2: Relative expenses for care seeking across the economic strata with reference to high income group

543