Title: Community Led Testing among People Who Inject Drugs: A community centered model to find new cases of HIV and Hepatitis C in Nepal

Short title: Community Led HIV and Hepatitis C Testing among People Who Inject Drugs

Authors

Rajesh Didhiya¹, Tara Nath Pokhrel², Sudha Devkota², Purusotam Raj Shedain², Mukunda Sharma⁴, Madan Kumar Shrestha², Deepak Gyenwali¹, Muhammad Imran⁴, Zakir Kadirov⁵, Bhawani Prasad Dahal⁵, Amrit Bikram Rai¹, Prawachan Kumar KC⁶, Sabir Ojha¹, Khem Narayan Pokhrel^{1*}

Author Details

1. National Association of People Living with HIV/AIDS in Nepal (NAP+N)
2. National Centre for AIDS and STD Control (NCASC)
3. National Public Health Laboratory (NPHL)
4. Save the Children, US
5. Save the Children, Nepal Country Office, KPRA Project
6. SPARSHA Nepal

***Corresponding author**

Khem Narayan Pokhrel, PhD,
Senior Research Fellow,
National Association of People Living with HIV/AIDS in Nepal (NAP+N),
House NO.: 174, Anek Marg, Baluwatar, Kathmandu-4, Nepal.
Contact phone: +977-01-4427459, 441783.

Email: pratikjee@gmail.com.

Word Count

Abstract:300

Main text: 3694

31 Abstract

32 Background

33 People Who Inject Drugs (PWIDs) have sub-optimal HIV and HCV testing as the available
34 testing services are inadequate in low and middle-income countries. We examined a model of
35 Community-Led Testing (CLT) in Nepal, exploring the feasibility of HIV and HCV testing
36 by trained lay service providers who had similar backgrounds to those of PWIDs. We also
37 assessed the prevalence of HIV and HCV within this study population and the associated risk
38 factors among PWIDs.

39 Methods

40 A mix-method cross-sectional study was conducted among 1029 PWIDs in five major
41 districts of Nepal from July 2019 to February 2020. Trained PWID peers performed the
42 screening for HIV and HCV using Rapid Diagnostic Test (RDT) kits. Acceptability and
43 feasibility of the testing was assessed. The participants' sociodemographic characteristics and
44 injecting and non-injecting risk characteristics were determined. The association of risk and
45 prevention characteristics with testing results were assessed using multiple logistic
46 regression.

47 Results

48 PWIDs shared that the test providers were friendly and competent in counseling and testing.
49 Of total PWIDs (n=1,029), 20.6% were HCV-positive and 0.2% were HIV-positive. HCV
50 positivity was associated with needle sharing (AOR: 1.83, 95% CI: 1.27, 2.64; p=0.001) and
51 reuse of syringe/needle (AOR: 2.26; 95% CI: 1.34, 3.79; p=0.002). In addition, PWIDs were
52 more likely to be HCV-positive who started opioid substitution therapy (OST) (AOR: 1.88,
53 95% CI: 1.26, 2.80, p=0.002) and attended the rehabilitation center (AOR: 1.66, 95% CI:
54 1.10, 2.53, p=0.017).

56 Conclusions

57 This CLT model was found to be a novel approach of testing of HIV and HCV which was
58 acceptable to PWIDs in Nepal and showed the high prevalence of HCV and its association
59 with injecting-related risk behaviors and being users of OST and rehabilitation. The findings
60 highlight the need of community-led testing in hotspots, OST settings, and rehabilitation
61 centers to screen new HIV and HCV infections.

62 Keywords: Community-led testing, HIV, HCV, Nepal, People who inject drugs

63

64 **Background**

65 People who inject drugs (PWIDs) are a high-risk key population for contracting human
66 immunodeficiency virus (HIV) and Hepatitis C virus (HCV) infections (1-3). This group
67 accounts for the major proportion of HCV infections, globally (3). An estimated 2.3 million
68 people living with HIV are co-infected with HCV globally (3, 4). HCV-induced liver disease
69 has emerged as a major contributor to mortality and morbidity among people living with HIV
70 and are co-infected with HCV (5).

71

72 Current HIV and HCV testing services are inadequate in reaching the hard to reach high risk
73 key populations, particularly PWIDs, which has resulted in the under diagnosis of HIV and
74 HCV in low-and middle-income countries (LMICs) (6). Such gaps in testing may require
75 adopting alternative approaches and strategies to achieve the target for elimination of HIV
76 and hepatitis C as public health threats by 2030 (6, 7).

77

78 Sero-prevalence studies conducted across various sites in Nepal have shown that HCV
79 prevalence among PWIDs is ranging from 21-38% in men and 3% in women in 2017. The
80 HIV/HCV co-infection was 2.5 to 7.4% in male PWIDs and 0.6% in female PWIDs
81 depending upon the study site in the country (8-10). Although Nepal has made progress in
82 improving accessibility of testing and treatment of HIV, an estimated 22% of the HIV-
83 positive people remain undiagnosed (11). In addition, HCV testing services are only available
84 in clinic-based settings, and PWIDs may not feel comfortable to seeking these testing
85 services from the facilities. PWIDs also face social and environmental challenges such as
86 poverty, homelessness, criminalization of drug injecting, lack of confidentiality, stigma, and
87 discrimination (12, 13).

88

89 In order to overcome the prevailing barriers and bridge the gap in testing of HIV and HCV,
90 an innovative model of Community-Led Testing (CLT) was conceived, developed, and tested
91 in Nepal. In this model, trained In-reach Workers (IRWs) who had similar backgrounds to
92 those of PWIDs performed HIV and HCV screening using easy to use rapid diagnostic test
93 kits (RDTs) in the community, bringing testing services closer to those most at risk and who
94 might not otherwise have been tested (14). IRWs are the peer educators and lay service
95 providers to PWIDs who engaged in community service organizations (CSOs) in Nepal.
96 Community led or community driven approaches have increased acceptability and

effectiveness of the interventions employed to improve access to HIV services in some African countries (15-17). Such a community-led model was also found to be effective in overcoming barriers and reaching the hard to reach, vulnerable and stigmatized key populations in HIV testing in Vietnam, and community home-based care support has improved ART adherence in Nepal as well (18, 19). CLT model for HIV screening has been implemented in Nepal since 2018. However, its acceptability and effectiveness is yet to be examined (14). Moreover, there are no PWIDs focused HCV screening programs in community settings.

This study aimed to examine the acceptability and feasibility of a CLT model of testing for HIV and HCV among PWIDs and linking PWIDs to the available treatment, care, and support services. In addition, we assessed the prevalence of HIV, HCV, HIV/HCV co-infection and the associated risk factors with HCV.

Materials and Methods

Study design

A mix-methods cross-sectional CLT study was designed and conducted among PWIDs in five major districts of Nepal. The qualitative approach explored in-depth knowledge on acceptability and feasibility of this CLT model from clients' and providers' perspectives. While the quantitative approach was used to examine the risk characteristics of PWIDs and to measure the prevalence of HIV and HCV, the study spanned from July 2019 to February 2020.

Study setting, sample size and sampling strategy

We purposively selected five districts where an estimated 30% of all PWIDs in Nepal reside (18). Calculation of sample size was based on the current highest prevalence of HCV at 38% according to the integrated biological and behavioural survey (IBBS) conducted in Nepal (9). Sample size was calculated using G*Power 3.1.9.6 for macOS, employing the 95% confidence interval and 80% power, the calculated sample size was 980. In addition to taking the reference prevalence of HCV, injecting related characteristics and their association with HCV positivity might need to be considered. Therefore, we increased the sample size to 1,029. We applied convenience sampling methods to reach the PWIDs peers through IRWs.

Data Collection

Participants were recruited from satellite sites covering hot spots (e.g., the location or secret places where PWIDs generally meet to inject drugs), drop-in centres (DIC), needle/syringe exchange program locations, opioid substitution therapy (OST) sites, and rehabilitation centres. IRWs of the district-based CSOs were trained with five-days intensive training package for conducting CLTs, counselling, and data collection including the practices on KOBO data collection platform using Android tablets (20). Geographical tracking was also done to trace location of the participants from the office to the site where they were interviewed. Trained IRWs and peers were mobilized to identify and contact the participants. For qualitative data collection in-depth interviews (IDIs) were conducted among 70 participants from five study districts who were enrolled in CLT for screening HIV and HCV. Similarly, three Focus Group Discussions (FGDs) were conducted in three project sites among key CLT service providers covering IRWs, lab personnel, and counsellors. Trained professionals were mobilized to conduct IDIs and FGD sessions. Validated IDIs and FGD topic guides were used in the study.

Participant inclusion criteria

PWIDs were considered eligible for the study if they were current or past injecting drug users who, had injected drugs for at least 3 months before they were recruited in the study, and had never tested or tested negative for HIV and /or HCV within the 12 months preceding the survey. Participants aged 16 years and above were enrolled. All the participants meeting inclusion criteria had provided consent to enrol in the study and to provide information. Therefore, no participants from the sample were excluded in the study.

Study variables

Socio-demographic information: Participants' age, ethnicity, educational status, marital status, gender, and occupation, history of HIV and HCV testing were recorded adopting the survey instruments of the Nepal Demographic Health Survey, 2016 (21).

Injection related risk characteristics: We adopted the questionnaires about the injection related risk characteristics covering needle sharing behaviors and other equipment sharing as defined by the IBBS study in Nepal (8, 9).

HIV and HCV risk factors: The risk factors were adopted from the Nepal IBBS study covering tattoos or body piercing, having sexual partner with status of HIV or with chronic hepatitis C/HIV, condom use, and weekly alcohol use (9).

Prevention activities: We interviewed the participants about their exposure to peer-education interventions, opioid substitution therapy, rehabilitation, needle and syringe exchange programs following the harm reduction programs activities of National Center for AIDS and STDs Control (NCASC).

Biological test: HCV and HIV screening used RDTs. Alere Determine™ HIV-1/2 and SD BIOLINE HCV RDTs were used for screening HIV and HCV, respectively, using blood samples from finger pricks in the satellite sites covering various hotspots, OST clinics, drop-in centers, and rehabilitation centers. Trained and certified IRWs performed screening of HIV and HCV of PWIDs and recorded clients' information including behavioural characteristics and test results in client report form (CRF) at satellite sites.

Qualitative information: Qualitative data collection methods such as FGDs and IDIs were applied to test the thematic topics: acceptability, feasibility, challenges/bottlenecks and unplanned effect of CLT.

173 Figure 1 Algorithm : The HIV/HCV CLT model in Nepal (See attached image)

174

175

176

177

178

179

180

181

182

183

184

185

Figure 1: The HIV/HCV CLT model in Nepal

Data analysis

Quantitative data were analyzed using STATA version 14.0. Descriptive analyses were done covering client's characteristics and the prevalence of HIV, HCV and HIV/HCV co-infection stratified by gender. Logistic regression analysis was applied to identify the association of socio-demographic characteristic and risk related characteristics with the prevalence of HCV. Results of HIV testing precluded meaningful analyses, since the number was so low.

The qualitative interviews and focus groups were transcribed and translated into English. The contents of transcripts were analyzed based on the pre-defined and emerging themes and sub-themes. We used MAXQDA 2020 (22) software for data coding, data sorting and extracting quotes according to theme.

Ethical consideration

As PWIDs are vulnerable, socially stigmatized and criminalized for drug use in Nepal, the study was conducted maintaining optimal ethical and human rights standards throughout the study period. Written informed consent was taken from all the participants before enrolling them in the study. Similarly, written assent was taken from senior peers of the participants below 18 years of age. Ethical approval was obtained from ethical review board of the Nepal Health Research Council (NHRC) [Reg No. 552/2019] prior to the commencement of the fieldwork.

Participants, who tested as HIV and HCV negative were counselled on safer behaviors and referred to harm reduction programs. Participants identified as either HIV and/or HCV positive were referred to the demonstration sites for counseling, HIV confirmatory test, linkages for HIV treatment, care and support, and to collect blood sample for HCV RNA/PCR. Participants identified as HCV positive were linked to post-test counselling services, liver fibrosis assessment through APRI scoring and referred to a hepatologist for appropriate follow up.

Results

Sociodemographic characteristics of Participants

Of total participants (n=1,029), 91.6% were men and 8.4% were women. Mean ages were 28.6 (SD=7.0) years for men and 23.4 (SD=3.7) years for women. Proportion of women in the age group of 16-24 years was 66.3% and that of men was 34.3%. The majority of participants were Janajati (Disadvantaged Janajati: 26.3% and Advantaged Janajati: 35.6%). About 63.1% of men and 75.6% of women were unmarried. Nearly two thirds (61.7%) of the participants were unemployed. Those who had ever enrolled in school, their mean year of schooling was 9.8 (SD=2.4) (Table 1).

Risk related characteristics of participants by gender

Of total (n=1029), 56.8% were current drug users. More than half (51.6%) of the participants reported that they had never tested for HIV. Regarding risk behaviors, 75.0% reported that they reused needles, 50.5% shared needles/syringes, and 29.9% shared a cooker/vial/container. About 55% of the participants were recruited from hotspots and 29% were from rehabilitation centers.

About 61.0% of the participants had an injection history of more than five years, and 43.6% reported that they had injected in the last seven days. Regarding needle/syringe sharing, 50% of men and 57.0% of women reported that they shared needles. In addition, 29.9% of participants reported that they had a practice of sharing a cooker/vial/container. Regarding other risk characteristics, 30.4% of men and 5.8% of the women consumed alcohol weekly, and 57.0% of the total participants had done tattooing. Nearly half of the women (47.7%) responded that they did not know their partner's HIV status, and 37.5% of men did not know their partner's status. While having sex with regular partner/sex workers, 57.1% reported that they always used condoms (Table 2).

Past testing, prevention exposure according to gender

Of the total, 17% of the participants reported that they had HCV test at least 12 months prior to the survey. Regarding the reason for not testing for HCV, 43.0 % reported fear of confidentiality breaches and 39.5% were not aware of the risk of HCV and HIV. Other reasons were screening is too expensive (23.5%), carelessness within themselves (18.0%), and no treatment is available (11.5%). About 10.5% reported that they did not know the testing facilities. About 52.9% of men and 62.8% of women participated in the IRW provided

harm reduction intervention, 28.1% started OST, and 63.2% attended a rehabilitation center. About 69.6% reported that they had participated in needle/syringe exchange program. About 61.8% reported that they felt stigma from family; a higher proportion of men compared to women (63.7% vs. 40.7%) felt stigma from family.

The overall prevalence of HCV in this sample was 20.6% (men=21.2%; women=14.0%). All HCV-positive participants were referred to demonstration sites to collect blood samples and the samples were sent to the National Public Health Laboratory [NPHL] for HCV RNA/PCR test. Only two men (0.2%) tested HIV positive and had HCV positive test as well. HIV-positive samples were sent for confirmatory tests using three RDTs as per national HIV testing algorithm. Both participants were confirmed as HIV positive, and both were linked to ART services. Among those who tested for HCV RNA (n=203), 68% had HCV RNA (IU/ml) count ranging from 65 to 1000000 (Table 3).

Descriptive analysis of socio-demographic and risk characteristics and their association with HCV screening result

Significantly higher proportion of participants in the age group of 25-50 years were HCV positive (77.4%) compared with those in the age group of 16-24 years and above 50 years ($p<0.001$). A higher proportion of married participants were HCV positive (57.1%) compared to unmarried and divorced/separated participants ($p<0.001$). HCV negative participants had a higher mean number of years of schooling than HCV positive [negative mean=10.1 (SD=2.2) vs. positive: mean=8.9 (SD=2.7), $p<0.001$].

A higher proportion of participants who reported needle sharing (58.0%) were HCV positive compared to those who did not share needles ($p=0.014$). In addition, a higher proportion participant who shared a cooker/vial/container were HCV positive compared to those who did not share (positive: 37.3% vs. negative: 28.0%, $p=0.009$). In addition, there was a significant difference on HCV positive and HCV negative results between those who reported re-use of needle/syringe compared with who did not reuse needle and syringes. (positive: 86.2% vs. negative: 72.1%, $p<0.001$).

Regarding risky sexual behaviors, those participants who had sexual contact with a person of unknown HIV status had higher proportion of HCV positive results (positive: 30.2% vs. negative: 20.3%, $p=0.002$) compared with participants who had no sexual contact or knew the HIV status of their partner. Those participants who always used condoms had a low proportion of HCV positive results compared to those who did not use condoms and used

them only sometimes (positive: 37.4% vs. negative: 50.6%, $p=0.001$). Also, PWIDs with tattoos had a higher prevalence of HCV positive than those without tattoos (positive: 67.5% vs. negative: 54.2%, $p<0.001$). Similarly, participants who consumed alcohol weekly had a higher proportion of HCV positive results relative to those who did not consume alcohol weekly (positive: 75.9% vs. negative: 67.9%, $p=0.024$). (Table 4).

Descriptive analysis of risk behavior and prevention exposure associated with HCV screening result

A higher proportion of HCV positive PWIDs had started OST compared to HCV negative (positive: 46.2% vs. negative: 23.4%, $p<0.001$). Rehabilitation participants had a higher proportion of HCV positive test results (positive: 76.4% vs. negative: 59.7%, $p<0.001$) relative to non-participants, and a higher proportion of HCV positive results were found among subjects who were involved in needle/syringe exchange programs (positive: 75.9% vs. negative: 67.9%, $p=0.024$). (Table 5).

Multivariate analysis of injecting risk behavior, prevention exposure associated with HCV screening results

Analyses were performed controlling for age, sex, marital, education level, tattooing, and participation in a needle/ syringe exchange program. Higher odds of HCV positive results were found in participants who had practice of needle sharing (AOR: 1.83, 95% CI: 1.27, 2.64, $p=0.001$) and reuse of syringes/needles (AOR: 2.26, 95% CI: 1.34, 3.79, $p=0.002$). In addition, HCV positive results were positively associated with those who started OST (AOR: 1.88, 95% CI: 1.26, 2.80, $p=0.002$) and attended a rehabilitation center (AOR: 1.66, 95% CI: 1.10, 2.53, $p=0.017$). (Table 6).

Qualitative findings

Feasibility of CLT

IRWs shared their experience and perceptions about the feasibility of community led testing. Doing RDT was convenient for them because they found that PWIDs were

“I never thought that I would have this much skills in counseling and testing. The training made me more skilled in testing procedures. I felt myself proud to reach my peer PWIDs who never get tested. It is really the rewarding job testing by the community for the community” ---20’s IRW

ready to get tested and agreed to be tested by their peers. They shared the positive and encouraging aspect of this model was

“testing by the community for the community.” The trained IRWs shared that they developed confidence as they had proper skill for procedure through the training and having received a certificate from a government authority. Another aspect shared by the trained peer was that they felt empowered and employed to test their community for the benefit of the community. The testing was easier for them as they found willingness among the peer PWIDs to get tested and refer their friends for testing for HIV and HCV.

Acceptability of CLT

The community led testing was acceptable among the participants as they reported that they found friendly behavior among test providers and perceived that the trained test providers had competence in counseling and testing. In addition, participants reported that they were happy to have immediate test results and having the testing place of their choice. Regarding privacy and confidentiality, PWIDs felt assured about the privacy maintained by the IRWs for testing. They had also greater trust toward the interviewers and test takers in the field. They had trust on pre-counseling and testing, and felt that their community are the service provider for the benefit of their diagnosis. Their perception was that they received exceptional support from their peers in testing and counseling from trained IRWs.

“I felt praised to meet my friend in the usual injecting site. Got very much impressed with his pre-test counseling and his confidence in testing procedure. Relieved to have result instantly”. - 30’s PWID

Bottlenecks/challenges

A few IRWs perceived that it was difficult for them in finding PWIDs who have not tested. Another problem that the trained peer said that in some cases they had difficulty

“I don’t know how we could overcome the interference of police as our protocol says we can test in the hotspots. I never discouraged from the incident of some sites as we encountered the police during testing. I kept on testing as I felt it is helping to the community” ---20’s IRW

in finding a private testing venue because some were feeling uncomfortable because police might interfere- as drug use is treated as a criminal act in Nepal. However, they never got discouraged and kept on going for testing and finding the community. Some thought that they had problems convincing the peers about the treatment available for HCV infection if they

tested positive. However, they referred positive participants for the consultation with the hepatologist.

Discussion

This study is the novel of its kind in Nepal, which applied CLT model to screen the hidden PWIDs affected by HIV and HCV. The model was proven a feasible testing approach for HIV and HCV in a low-resource setting such as Nepal and was acceptable by PWIDs. The lay test providers and peers/IRWs had confidence in reaching, providing counseling, maintaining the privacy of PWIDs and testing in the satellite sites. The test providers also felt empowered as they were doing something for their own community: “Testing by the community for the community”. Although a few service providers had some difficulties in finding private testing locations and felt fear of interference from the police, they were able to reach the PWIDs in convenient places and provide testing upon request. In addition, PWIDs shared that the test providers were friendly and were competent in counseling and testing. They shared that the best part of the CLT was that they received the result instantly and were referred for further services. This model can be one of the approaches to overcome the access barriers for PWIDs for seeking services from clinical settings (13, 18, 23).

This study found that 0.2% of participants had HIV and 20% had HCV. The lower rate of HIV could be due to better utilization of testing services and their exposure to prevention services. Also, those who had HIV testing prior to 12 months of the survey were not included in this model. The reliability of testing was promising as the result accuracy was 100% while performing confirmatory testing in National Public Health Laboratory (NPHL).

HCV positivity result is in line with the IBBS study findings, which has shown the prevalence ranging from 18.8 to 38% (8-10). The true prevalence of HCV among PWIDs could be even higher than that revealed by the study if the criteria "never tested or tested HCV negative at least for last 12 months" were not used. This statement is supported by the findings of a study conducted in 2015 which had shown almost 50% of the PWIDs with HCV-RNA sought services from organized institutions (24).

PWIDs represent 3% of the total cases of HIV in Nepal and HIV incidence was 0.03% in 2020 and the prevalence among adult population was 0.13% (11). The study results indicated that PWIDs who were not reached by the ongoing program for HIV screening and could be reached through this peer-led approach (16, 18) although detection of HIV is fairly low. This

CLT approach would be complementary to the current HIV/HCV screening services, as the PWIDs feel stigma and fear of confidentiality breaches in the health facilities that have resulted in non-availability or inaccessibility of the screening services (13, 18, 23).

This study also found that needle sharing and needle reuse were major risk factors for HCV infection. Reuse of needles and needle/ syringe sharing may have increased the risk of vulnerability of transmission. In addition, HCV positivity rate was found to be high among PWIDs who started their OST and attended rehabilitation center. Participants seeking services from institutions may have history of years of injecting drug use before enrollment in the services and might have not used the available testing for HCV services with a fear of lack of treatment availability of HCV. Longer duration of exposure is positively associated with HCV infection and the longer duration increases the repeated high risk behaviors such as needle sharing, re-use of needles, and unsafe sexual behaviors (24, 25).

Despite the major significance of this CLT model, the study has two major limitations. First, due to nature of screening hidden PWIDs, we applied non-probability sampling led by peer PWIDs. This may affect the generalizability of the results. However, IRWs were reaching every satellite site with the mapping so that they would be able to reach more hidden and non-diagnosed cases. Second, we were not able to assess the factors associated with HIV-positivity because of the low prevalence in the screened populations. However, the risk characteristics of HCV and HIV could be assumed similar such as needle sharing and sharing of injecting preparation materials.

Conclusion:

This CLT study has shown that peer lay test providers were well accepted by PWIDs for counseling and testing. This model has become promising to test for HCV and HIV among PWIDs in their convenient sites. Also, the study shows that risk behaviour such as needle re-use and needle sharing led to HCV infection. Moreover, higher proportion of PWIDs in rehabilitation center and on OST were also tested positive for HCV. Therefore, this study highlights the expansion of this model to screen for both HCV and HIV together in other parts of the country and similar settings, where PWIDs encounter barriers to access to testing services.

Acknowledgments

We acknowledge the CSOs involved in this research project SPARSHA Nepal-Kathmandu, Community Support Group-Kaski, and Nirmaya-Chitwan. We would like to thank the IRWs, who successfully conducted the testing and field work for quantitative data collection and qualitative researchers who conducted IDIs and FGDs. We owe to all 1029 PWIDs who successfully and willingly participated in this study, provided information, agreed for the test of HIV and HCV in the satellite places. We have also number of consultants to acknowledge who provided the inputs in design. Further, we would like to acknowledge the support received from the administrative, logistic, and financial staffs of Save the children, NCASC, NAP+N, and NPHL.

Author Contributions

Conceptualization: Rajesh Didiya, Tara Nath Pokhrel, Zakir Kadirov, Purusotam Raj Sedai, Mukunda Sharma, Madan Kumar Shrestha, Deepak Gyenwali, Bhawani Prasad Dahal, Amrit Bikram Rai, Khem Narayan Pokhrel

Data curation: Deepak Gyenwali, Khem Narayan Pokhrel

Formal analysis: Khem Narayan Pokhrel, Deepak Gyenwali

Supervision: Rajesh Didiya, Deepak Gyenwali, Tara Nath Pokhrel, Khem Narayan Pokhrel

Validation: Tara Nath Pokhrel, Bhawani Prasad Dahal, Zakir Kadirov, Khem Narayan Pokhrel

Manuscript preparation and finalization: Khem Narayan Pokhrel

Manuscript – review & editing: Rajesh Didiya, Tara Nath Pokhrel, Zakir Kadirov, Muhammad Imran, Purusotam Raj Sedai, Mukunda Sharma, Madan Kumar Shrestha, Deepak Gyenwali, Bhawani Prasad Dahal, Amrit Bikram Rai, Sabir Ojha, Prawachan Kumar KC

References

1. Degenhardt L, Peacock A, Colledge S, Leung J, Grebely J, Vickerman P, et al. Global prevalence of injecting drug use and sociodemographic characteristics and prevalence of HIV, HBV, and HCV in people who inject drugs: a multistage systematic review. *The Lancet Global Health*. 2017;5(12):e1192-e207.
2. Rashti R, Alavian SM, Moradi Y, Sharafi H, Bolbanabad AM, Roshani D, et al. Global prevalence of HCV and/or HBV coinfections among people who inject drugs and female sex workers who live with HIV/AIDS: a systematic review and meta-analysis. *Archives of Virology*. 2020:1-12.
3. WHO. Global hepatitis report 2017: World Health Organization 2017. Report No.: 9241565454.
4. WHO. HIV/AIDS Key facts. 2018:1-7. Available from <https://www.who.int/news-room/fact-sheets/detail/hiv-aids>. Accessed on September 16, 2020.
5. Poudel KC, Palmer PH, Jimba M, Mizoue T, Kobayashi J, Poudel-Tandukar K. Coinfection with hepatitis C virus among HIV-positive people in the Kathmandu Valley, Nepal. *Journal of the International Association of Providers of AIDS Care (JIAPAC)*. 2014;13(3):277-83.
6. WHO. Global health sector strategy on viral hepatitis 2016-2021. Towards ending viral hepatitis: World Health Organization. 2016.
7. WHO. Global health sector strategy on HIV 2016-2021. Towards ending AIDS: World Health Organization. 2016.
8. NCASC. Integrated Biological and Behavioral Surveillance (IBBS) Survey among People Who Inject Drugs (PWID) in Kathmandu Valley. Kathmandu Nepal. Kathmandu, Nepal. 2017.
9. NCASC. Integrated Biological and Behavioural Surveillance (IBBS) Surveys among People Who Inject Drugs (PWID) in Eastern Terai Highway Districts. Kathmandu, Nepal: National Center for AIDS and STDs Control. 2017.
10. NCASC. Integrated Biological and Behavioural Surveillance (IBBS) Surveys among People Who Inject Drugs (PWID-Male) in Pokhara Valley: National Center for AIDS and STDs Control. 2017.
11. HIV Epidemic Update of Nepal [database on the Internet]2020 [cited 12.15.2020]. Available from: <http://www.ncasc.gov.np/WAD2020/Factsheet-2020-S.pdf>.

12. Nelson PK, Mathers BM, Cowie B, Hagan H, Des Jarlais D, Horyniak D, et al. Global epidemiology of hepatitis B and hepatitis C in people who inject drugs: results of systematic reviews. *The Lancet*. 2011;378(9791):571-83.
13. Barocas JA, Brennan MB, Hull SJ, Stokes S, Fangman JJ, Westergaard RP. Barriers and facilitators of hepatitis C screening among people who inject drugs: a multi-city, mixed-methods study. *Harm reduction journal*. 2014;11(1):1-8.
14. NCASC. National Guidelines Community-Led HIV Testing 2017 Nepal. Kathmandu Nepal: National Center for AIDS and STDs Control; 2017.
15. Ndyomugenyi R, Kabali AT. Community-directed interventions for integrated delivery of a health package against major health problems in rural Uganda: perceptions on the strategy and its effectiveness. *International health*. 2010;2(3):197-205.
16. Indravudh PP, Fielding K, Kumwenda MK, Nzawa R, Chilongosi R, Desmond N, et al. Community-led delivery of HIV self-testing to improve HIV testing, ART initiation and broader social outcomes in rural Malawi: study protocol for a cluster-randomised trial. *BMC infectious diseases*. 2019;19(1):814.
17. Sharma M, Barnabas RV, Celum C. Community-based strategies to strengthen men's engagement in the HIV care cascade in sub-Saharan Africa. *PLoS medicine*. 2017;14(4):e1002262.
18. Nguyen VTT, Phan HT, Kato M, Nguyen QT, Le Ai KA, Vo SH, et al. Community-led HIV testing services including HIV self-testing and assisted partner notification services in Vietnam: lessons from a pilot study in a concentrated epidemic setting. *Journal of the International AIDS Society*. 2019;22:e25301.
19. Pokhrel KN, Sharma VD, Pokhrel KG, Neupane SR, Mlunde LB, Poudel KC, et al. Investigating the impact of a community home-based care on mental health and anti-retroviral therapy adherence in people living with HIV in Nepal: a community intervention study. *BMC infectious diseases*. 2018;18(1):263.
20. KOBO Tool Box Cambridge, MA: The Harvard Humanitarian Initiative; 2019 [cited 2021 10.11.2020]; Available from: <https://www.kobotoolbox.org/>.
21. MOHP Nepal. Nepal Demographic and Health Survey 2016. Kathmandu: Ministry of Health 2017.
22. Software V. MAXQDA 2020. 2019.
23. Swan D, Long J, Carr O, Flanagan J, Irish H, Keating S, et al. Barriers to and facilitators of hepatitis C testing, management, and treatment among current and former injecting drug users: a qualitative exploration. *AIDS patient care and STDs*. 2010;24(12):753-62.

- 504 24. Kinkel H-T, Karmacharya D, Shakya J, Manandhar S, Panthi S, Karmacharya P, et al.
505 Prevalence of HIV, hepatitis B and C infections and an assessment of HCV-genotypes
506 and two IL28B SNPs among people who inject drugs in three regions of Nepal. PLoS
507 One. 2015;10(8).
- 508 25. Poudel KC, Poudel-Tandukar K, Yasuoka J, Joshi AB, Jimba M. Correlates of sharing
509 injection equipment among male injecting drug users in Kathmandu, Nepal. International
510 Journal of Drug Policy. 2010;21(6):507-10.
511
512
513

Tables

Table 1: Socio-demographic characteristics of participants by gender

Characteristics	Men (n=943) (n, %)	Women (n=86) (n, %)	Total (n, %)
Age (mean, SD)	28.6 (7.0)	23.4 (3.7)	28.2 (7.0)
Age groups			
16-24 years	323 (34.3)	57 (66.3)	380 (36.9)
25-50 years	610 (64.7)	29 (33.7)	639 (62.1)
Above 50 years	10 (1.0)	0 (0.0)	10 (1.0)
Ethnicity			
Dalit	81 (8.6)	8 (9.4)	89 (8.7)
Disadvantaged Janajati	245 (26.0)	25 (29.0)	270 (26.3)
Disadvantaged non-Dalit Terai	2 (0.2)	1 (1.2)	3 (0.3)
Religious minorities (Muslim)	8 (0.9)	0 (0.0)	8 (0.8)
Advantaged Janajati	340 (36.1)	26 (30.2)	366 (35.6)
Upper caste (Brahmin/Chhetri)	266 (28.2)	26 (30.2)	292 (28.3)
Marital status			
Unmarried	595 (63.1)	65 (75.6)	660 (64.1)
Married	339 (36.0)	21 (24.4)	360 (35.0)
Divorced/separated	9 (1.0)	0 (0.0)	9 (0.9)
Occupation			
Unemployed	562 (59.6)	73 (84.9)	635 (61.7)
Professional/technical/Managerial	7 (0.7)	0 (0.0)	7 (0.7)
Labor	96 (10.2)	2 (2.3)	98 (9.5)
Business	134 (14.2)	6 (7.0)	140 (13.6)
Agriculture	8 (0.9)	0 (0.0)	8 (0.8)
Student	52 (5.5)	3 (3.5)	55 (5.3)
Others	84 (8.9)	2 (2.3)	86 (8.4)
Education level			
Illiterate	0 (0.0)	8 (0.9)	8 (0.8)
Literate but not formal education	12 (1.3)	4 (4.7)	16 (1.6)
Formal education received	923 (97.9)	82 (95.4)	1005 (97.7)
Mean (SD) schooling years (n=1005)	9.4 (2.4)	9.8 (2.4)	9.8 (2.4)

Table 2: Risk characteristics of participants by gender

Characteristics	Men (n=943, %)	Women (n=86, %)	Total (n=1029, %)
Ever drug use			
Current	531 (56.3)	54 (62.8)	585 (56.8)
Past	412 (43.7)	32 (37.2)	444 (43.2)
Last test of HIV Done			
Never	490 (52.0)	41 (47.7)	531 (51.6)
Before 12 months	453 (48.0)	45 (52.3)	498 (48.4)
Risk behavior (multiple response)			

Shared a needle	471 (50.0)	49 (57.0)	520 (50.5)
Shared a cooker/vial/container	288 (30.5)	20 (23.3)	308 (29.9)
Shared a cotton/filter/rinse water	43 (4.6)	1 (1.2)	44 (4.3)
Re-used syringe/needle	717 (76.0)	55 (64.0)	772 (75.0)
Not shared	30 (3.2)	0 (0.0)	30 (2.9)
Type of satellite			
DIC	33 (3.5)	1 (1.2)	34 (3.3)
Hotspots	516 (54.7)	50 (58.1)	566 (55.0)
OST Clinic	71 (7.5)	2 (2.3)	73 (7.1)
Rehabilitation center	272 (28.8)	31 (36.1)	303 (29.5)
ART Clinic	2 (0.2)	0 (0.0)	2 (0.2)
Others	49 (5.2)	2 (2.3)	51 (5.0)
Injecting duration			
<=1 year	25 (2.7)	7 (8.1)	32 (3.1)
Between one year and five years	323 (34.3)	46 (53.5)	369 (35.9)
More than five years	595 (63.1)	33 (38.4)	628 (61.0)
Last injected (days)			
<=7 days	417 (44.2)	32 (37.2)	449 (43.6)
8-30 days	109 (11.6)	22 (22.6)	131 (12.7)
31 days or more	417 (44.2)	32 (37.2)	449 (43.7)
Weekly alcohol use			
Yes	287 (30.4)	5 (5.8)	292 (28.3)
No	656 (69.6)	81 (94.2)	737 (71.6)
Tattoo			
Yes	546 (57.9)	40 (46.5)	586 (57.0)
No	397 (42.1)	46 (53.5)	443 (43.0)
Partner status			
HIV-positive	2 (0.2)	0 (0.0)	2 (0.2)
Hepatitis C	2 (0.2)	1 (1.2)	3 (0.3)
Don't know	354 (37.5)	41 (47.7)	395 (38.4)
None	585 (62.0)	44 (51.2)	629 (61.1)
Condom use with casual partners/SWs			
Never			51 (5.0)
Always			588 (57.1)
Sometimes			326 (31.7)
Never had sex with SWs			64 (6.2)

518 **Table 3: Past testing, prevention, stigma, and current testing results by gender**

Characteristics	Men (n=943,%)	Women (n=86,%)	Total (n=1029)
Past HCV test			
Yes	170 (18.0)	5 (5.8)	175 (17.0)
No	773 (82.0)	81 (94.2)	854 (83.0)
Reason for not testing HCV (multiple response)			
Not aware of the risk	299 (38.6)	41 (50.6)	340 (39.7)
Don't know about the testing	87 (11.2)	3 (3.7)	90 (10.5)

facilities			
Testing facility is too far	18 (2.3)	1 (1.2)	19 (2.2)
Fear of stigma in the facility	60 (7.7)	5 (6.2)	65 (7.6)
Fear of confidentiality	338 (43.6)	30 (37.0)	368 (43.0)
Screening is too expensive	189 (24.4)	12 (14.8)	201 (23.5)
No treatment is available	92 (11.9)	6 (7.4)	98 (11.5)
Carelessness	148 (10.7)	6 (0.7)	154 (18.0)
Participated in IRW intervention			
Yes	499 (52.9)	54 (62.8)	553 (53.7)
No	444 (47.1)	32 (37.2)	476 (46.3)
OST started			
Yes	278 (29.5)	11 (12.8)	289 (28.1)
No	664 (70.5)	75 (87.2)	739 (71.9)
Attended in Rehabilitation center			
Yes	600 (63.6)	50 (58.1)	650 (63.2)
No	343 (36.8)	36 (41.9)	379 (36.8)
Participated in needle/syringe exchange program			
Yes	653 (69.3)	63 (73.2)	716 (69.6)
No	290 (30.8)	23 (26.7)	313 (30.4)
Stigma			
Stigma from family	601 (63.7)	35 (40.7)	636 (61.8)
Stigma in the health facility	64 (6.8)	1 (1.2)	65 (6.3)
Stigma in office/work	42 (4.5)	0 (0.0)	42 (4.1)
Not felt stigma	325 (34.5)	51 (59.3)	376 (36.5)
Stigma felt in other places	38 (4.0)	1 (1.2)	39 (3.8)
HCV screening result			
Positive	200 (21.2)	12 (14.0)	212 (20.6)
Negative	743 (78.8)	74 (86.0)	817 (79.4)
HIV screening result			
Positive	2 (0.2)	0(0.0)	2 (0.2)
Negative	941 (99.8)	86 (100.0)	1027 (99.8)
HIV confirmatory test 3RDT proposed (yes)[as per national HIV testing algorithm]	2 (100.0)	0 (0.0)	2 (100.0)
HCV RNA (IU/ml) (n=203)			
<65			61 (30.0)
65 to 1000000			138 (68.0)
>1000000			4 (2.0)

Table 4: Descriptive analysis of socio-demographic and risk characteristics on HCV screening results

Characteristics	HCV positive (n=212, %)	HCV negative (n=817, %)	p-value
Age groups			<0.001
16-24 years	40 (18.9)	340 (41.6)	
25-50 years	164 (77.4)	475 (58.2)	
Above 50 years	8 (3.8)	2 (0.2)	

Gender			0.111
Men	200 (94.3)	743 (90.9)	
Women	12 (5.7)	74 (9.1)	
Marital status			<0.001
Unmarried	88 (41.5)	572 (70.0)	
Married	121 (57.1)	239 (29.3)	
Divorced/separated	3 (1.4)	6 (0.7)	
Mean (SD) schooling years (n=1005)	8.9 (2.7)	10.1 (2.2)	<0.001
Needle Sharing			0.014
Yes	123 (58.0)	397 (48.6)	
No	89 (42.0)	420 (51.4)	
Shared a cooker/vial/container			0.009
Yes	79 (37.3)	229 (28.0)	
No	133 (62.7)	588 (71.9)	
Reused needle/syringe			<0.001
Yes	183 (86.2)	589 (72.1)	
No	29 (13.7)	228 (27.9)	
Sexual contact with person with HIV unknown status (yes)			0.002
Yes	64 (30.2)	165 (20.3)	
No	148 (69.8)	647 (79.7)	
Condom use			0.001
No	44 (20.9)	162 (19.6)	
Always	79 (37.4)	413 (50.6)	
Sometimes	88 (41.7)	241 (29.5)	
Tattoo done			0.001
No	69 (32.6)	374 (45.8)	
Yes	143 (67.5)	443 (54.2)	
Weekly alcohol consumption			0.024
No	51 (24.0)	262 (32.1)	
Yes	161 (75.9)	555 (67.9)	

Table 5: Descriptive analysis of risk behavior and prevention exposure with HCV screening result

Characteristics	HCV positive (n, %)	HCV Negative (n, %)	p-value
Participated in peer education/IRW intervention			0.869
No	97 (45.8)	379 (46.4)	
Yes	115 (54.3)	438 (53.6)	
Started OST			<0.001
No	114 (53.8)	625 (76.6)	
Yes	98 (46.2)	191 (23.4)	
Attended rehabilitation center			<0.001
No	50 (23.6)	329 (40.3)	
Yes	162 (76.4)	488 (59.7)	
Participated in			0.024

needle/syringe exchange program			
No	51 (24.1)	262 (32.1)	
Yes	161 (75.9)	555 (67.9)	

Table 6: Multivariate analysis of injecting risk behavior, prevention exposure with HCV screening results

Characteristics	Adjusted odds Ratio ¹	95% CI	p-value
Needle sharing	1.83	1.27, 2.64	0.001
Reused syringe/needle	2.26	1.34, 3.79	0.002
OST started	1.88	1.26, 2.80	0.002
Attended Rehabilitation Center	1.66	1.10, 2.53	0.017
Participated needle syringe exchange program	0.67	0.42, 1.08	0.108
Tattooing	1.18	0.76, 1.82	0.455
Age	1.10	1.05, 1.22	<0.001
Sex (male)	0.52	0.25, 1.11	0.093
Marital status (married)	0.84	0.55, 1.22	0.427
Educational level	0.79	0.73, 0.85	<0.001

¹Adjusted for age, sex, marital, education level, tattooing, participated in needle syringe exchange program, needle sharing, need syringe/needle, OST started, attended rehabilitation center