1 **Computerized physical and cognitive training improves the functional architecture**
2 **of the brain in adults with Down Syndrome: a network science EEG study**

3
4 Alexandra Anagnostopoulou^{1*}, Charis Styliadis^{1*}, Panagiotis Kartsidis¹, Evangelia
5 Romanopoulou¹, Vasiliki Zilidou¹, Chrysi Karali², Maria Karagianni¹, Manousos Klados³,
6 Evangelos Paraskevopoulos^{1,4**}, Panagiotis D. Bamidis^{1**}

7
8 ¹ Medical Physics Laboratory, School of Medicine, Faculty of Health Sciences, Aristotle
9 University of Thessaloniki, Greece.

10 ² School of Biology, Faculty of Science, Aristotle University of Thessaloniki, Greece.

11 ³ Department of Psychology, The University of Sheffield International Faculty, City
12 College, Thessaloniki, Greece.

13 ⁴ Department of Psychology, University of Cyprus, Nicosia, Cyprus

14
15 * Joint first authorship

16 ** Joint senior authorship

17
18 **Corresponding Author:** Dr. Charis Styliadis, School of Medicine, Faculty of Health
19 Sciences, Aristotle University of Thessaloniki, Thessaloniki, Greece. Aristotle University
20 of Thessaloniki, University Campus, P.C. 54124, Thessaloniki, Greece. Tel: +30-2310-
21 999310. Fax: +30-2310-999702. Email: cstyliadis@auth.gr

22

23 **Keywords:** Down Syndrome, Physical Training, Cognitive Training,
24 Electroencephalography, Network Science Indices, Adaptive Neuroplasticity

25

26 **Abstract**

27 Understanding the neuroplastic capacity of people with Down Syndrome (PwDS) can
28 potentially reveal the causal relationship between aberrant brain organization and
29 phenotypic characteristics. We used resting-state EEG recordings to identify how a
30 neuroplasticity-triggering training protocol relates to changes in the functional connectivity
31 of the brain's intrinsic cortical networks. Brain activity of 12 PwDS before and after a ten-
32 week protocol of combined physical and cognitive training was statistically compared to
33 quantify changes in directed functional connectivity in conjunction with
34 psychosomatometric assessments. PwDS showed increased connectivity within the left
35 hemisphere and from left to right hemisphere, as well as increased physical and cognitive
36 performance. Our findings reveal a strong adaptive neuroplastic reorganization as a result
37 of the training that leads to a less-random network with a more pronounced hierarchical
38 organization. Our results go beyond previous findings by indicating a transition to a
39 healthier, more efficient, and flexible network architecture, with improved integration and
40 segregation abilities in the brain of PwDS. Resting-state electrophysiological brain activity
41 is used here for the first time to display meaningful relationships to underlying DS
42 processes and outcomes of importance in a translational inquiry. This trial is registered
43 with ClinicalTrials.gov Identifier [NCT04390321](#).

44

Author Summary

The effects of cognitive and physical training on the neuroplasticity attributes of people with and without cognitive impairment have been well documented via neurophysiological evaluations and network science indices. However, there is still insufficient evidence for people with Down Syndrome (PwDS). We investigated the effects of a combinational training protocol on the brain network organization of 12 adult PwDS using EEG and network indices coupled with tests assessing their cognitive and physical capacity. We report evidence of adaptational neuroplastic effects, pointing to a transitional state towards a healthier organization with an increased ability to integrate and segregate information. Our findings underline the ability of the DS brain to respond to the cognitive demands of external stimuli, reflecting the possibility of developing independent-living skills.

Introduction

Neuroplasticity can emerge in both typical and atypical brains and allows for either development, reaction, recovery, or adaptation to internal and external stimuli (Trojan & Pokorny, 1999). In contrast to typically developed (TD) individuals, the Down Syndrome (DS) brain presents atypical levels of inhibition due to gene over-expression, leading to prolonged failure of synaptic plasticity and a reduced capacity for remodeling (Baroncelli et al., 2011). These characteristics, along with morphogenetic modifications, are among the leading causes of brain disability in people with DS (PwDS) (Dierssen, Herault, & Estivill, 2009).

DS individuals, in comparison to age-matched TD individuals, have smaller frontal, amygdala, and cerebellar volumes (but increased parahippocampal volume) (Pinter, Eliez,

Schmitt, Capone, & Reiss, 2001; White, Alkire, & Haier, 2003), a pattern which overpronounces after their 50th year of age (Dierssen, 2012). The current consensus has associated the cognitive capabilities and deficiencies of PwDS with several distinct brain regions emphasizing an abnormal and less efficient DS brain organization (Menghini, Costanzo, & Vicari, 2011).

Electroencephalography (EEG) and magnetoencephalography (MEG) studies have complemented the notion of this atypical organization. PwDS, when compared to TD controls, exhibit slow brain wave, especially in left posterior areas (Babiloni et al., 2010, 2009), with higher delta band and lower alpha and beta band activity, a pattern also evident in patients with Alzheimer's Disease (AD) (Babiloni et al., 2010, 2009; García-Alba et al., 2019; Hemmati, Ahmadlou, Gharib, Vameghi, & Sajedi, 2013). fMRI studies have shown a rather-simplified DS network organization (Anderson et al., 2013), lacking the appropriate efficiency and flexibility (Edgin, Clark, Massand, & Karmiloff-Smith, 2015). Given the limited capacity of the DS brain to consolidate information due to its disorganized architecture of reduced segregation and impaired integration, diffused connectivity (hyper-synchrony) (Anderson et al., 2013), and decreased long-range connectivity, the DS brain network's potential for plasticity is in question (Edgin et al., 2015).

Nevertheless, the DS brain does possess neuroplastic capabilities, at least in the form of compensatory events (Haier, Head, Head, & Lott, 2008; Takashima et al., 2014). The emergence of AD-related characteristics (i.e., altered theta band activity) in PwDS reaffirms the possibility of compensatory mechanisms before and during the expression of AD (Takashima et al., 2014). Characterizing neuroplasticity in PwDS is vital in

understanding causality between aberrant brain circuitry and the cognitive and behavioral phenotype. This step would allow the quantification of the remodeling potential of evidence-based interventions.

Interventional approaches have shifted from a pharmaceutical concept, with, so far, inconclusive results regarding beneficial effects (Costa, 2011; Kálmán et al., 2009; Kishnani et al., 2010; Lott et al., 2011; Mohan, Carpenter, & Bennett, 2009), to non-pharmaceutical interventions of physical and behavioral components, aiming to trigger neuroplasticity and enhance brain health or to protect against neurodegenerative events (Head et al., 2007). Such interventions have shown promise in healthy aging and populations with cognitive impairments and have provided mounting evidence for lifelong brain plasticity (Bamidis et al., 2014; Colcombe et al., 2006, 2004; Cotman, Berchtold, & Christie, 2007; Erickson et al., 2011; Klados, Styliadis, Frantzidis, Paraskevopoulos, & Bamidis, 2016; Kramer et al., 1999; Styliadis, Kartsidis, & Paraskevopoulos, 2015; Styliadis, Kartsidis, Paraskevopoulos, Ioannides, & Bamidis, 2015).

Despite evidence supporting the positive impact of exercise-based (Bertapelli, Pitetti, Agiovlasitis, & Guerra-Junior, 2016; Hardee & Feters, 2017), as well as behavioral and cognitive interventions (Fonseca, Navatta, Bottino, & Miotto, 2015; McGlinchey, McCarron, Holland, & McCallion, 2019) for PwDS, DS literature features a substantial knowledge gap, as there are no reports of neurophysiological investigations of the training-induced effects, and a network-theory based assessment of the training-induced plasticity. Deviations in the brain functionality of PwDS have been previously addressed with the investigation of resting-state networks (RSNs) (Ahmadlou, Gharib, Hemmati, Vameghi, &

Sajedi, 2013; Anderson et al., 2013; Babiloni et al., 2010, 2009; Pujol et al., 2015; Vega, Hohman, Pryweller, Dykens, & Thornton-Wells, 2015).

We hypothesized that combined physical (PT) and cognitive training (CT) would trigger neuroplasticity and reorganize the DS brain network to adapt to the increased cognitive and physical demand. Resting-state EEG data were acquired to characterize plasticity in the DS brain through network science and investigate how the training-induced neuroplasticity affects the organization and characteristics of the DS network, as opposed to the reported random-like architecture (Ahmadlou et al., 2013) with impaired integration and segregation capabilities.

Source analysis was performed using low resolution electromagnetic tomography (LORETA) (Pascual-Marqui, Michel, & Lehmann, 1994), and directed functional connectivity was estimated via phase transfer entropy (PTE) in 0.53-35 Hz. This measure quantifies the direction of the information flow, allowing for a whole-head analysis without requiring an *a priori* head model definition. A non-parametric permutation test was applied to extract the significant, within-network differences (post- vs. pre-intervention) in connectivity. A graph-theoretical analysis was performed to index the training's neuroplastic effects through the statistical comparison of graph measures between the two time-points.

Results

Psychometric and somatometric results

Psychosomatometric score comparisons between the two time-points were performed with the use of non-parametric Wilcoxon tests and paired t-tests. Physical assessments

exhibited a significant improvement in score for the arm curl test (Table 1). Significant differences in the time of completion (decrease in duration) were evident in the Time Up and Go assessment test (Table 1). The comparison of psychometric assessments revealed an increase in the Digits Forward score (Digits Span test), the Mazes test, as well as the Ravens AB and Ravens Total Score (Table 1). The rest of the tests showed no significant post-pre score changes.

Table 1. Somatometric and psychometric tests with significant differences post vs. pre, along with the post-pre mean difference. Results were considered significant for $p < 0.05$.

Somatometric	p-value	mean difference
Arm Curl	0.017	1.75
Time Up and Go	0.007	-1.67
Psychometric		
Digits Span (Digit Forward)	0.032	1.50
Mazes	0.032	0.97
Ravens AB	0.017	2.12
Ravens Total Score	0.015	3.13

PTE results

Statistical comparisons of the PTE matrices between the pre- and post-intervention EEG measurements indicate the reorganization of cortical connections (62 nodes and 40 edges), due to training, in the DS resting-state network ($p < 0.05$, corrected for multiple comparisons via the non-parametric NBS method (Zalesky, Fornito, & Bullmore, 2010),

5000 permutations) (Figure 1). Specifically, the cortical reorganization in the DS brain is characterized by the strengthening of connections within: i) the left hemisphere, from nodes in the occipital (cuneus and lingual gyrus) and the temporal lobe (fusiform gyrus) to the frontal lobe (superior and middle gyrus) and from the frontal lobe to the parietal lobe (superior, precuneus and postcentral gyrus), ii) the right hemisphere, from nodes in the occipital (cuneus, lingual and middle gyrus) and temporal (fusiform and inferior gyrus) lobe to the frontal lobe (superior, cingulate and medial gyrus), and between: i) nodes of the left fusiform gyrus, and inferior temporal lobe and the right frontal lobe (superior, middle and inferior gyrus), ii) the right middle occipital lobe, sub-gyral, parahippocampal and fusiform gyrus to the left frontal lobe (superior, middle, precentral gyrus), iii) the left inferior temporal lobe to the right anterior cingulate, and iv) the right frontal lobe (middle and inferior gyrus) to the left postcentral and middle frontal gyrus (Figure 1). In respect to the core RSNs, as defined by Yeo et al. (Yeo et al., 2011), we have noted increased connectivity within the dorsal attention network (DAN), and between the visual (VIS) and frontoparietal network (FPN), VIS and default-mode-network (DMN), VIS and ventral attention network (VAN), VAN and FPN, FPN and DMN, FPN and somatosensory network (SMN), FPN and DAN, DAN and DMN, as well as DAN and SMN (Table 2).

Figure 1. Cortical connectivity between post- and pre-intervention networks and for each time point (t -value > 3.52). The color scales represent t -values. **A:** Circular graph depicting the cortical reorganization in the DS brain (cyan: left hemisphere, lime: right hemisphere, blue: frontal lobe, green: occipital lobe, purple: parietal lobe, orange: temporal lobe, maroon: limbic lobe). The cortical reorganization is characterized by the strengthening of direct connections within: i) the left hemisphere, from nodes in the occipital and temporal lobe to the frontal lobe, and from the frontal lobe to the parietal lobe, ii) the right hemisphere, from nodes in the occipital and temporal lobe to the frontal lobe, and between: i) nodes of the left fusiform gyrus, and inferior temporal lobe and the right frontal lobe, ii) the right middle occipital lobe, sub-gyral, parahippocampal and fusiform gyrus to the left frontal lobe, iii) the left inferior temporal lobe to the right anterior cingulate, and iv) the right frontal lobe to the left postcentral and middle frontal gyrus. **B:** Depiction of pre- and post-intervention networks in comparison to the null hypothesis. The t -values indicate that the post-

resting-state network has significantly stronger connections than the pre network. **C: Post vs. Pre.** Significant post-pre connectivity differences. Information direction is depicted through line arrows. The visualized networks are significant at a level of $p < 0.05$, corrected for multiple comparisons via the non-parametric NBS method with an internal threshold of 0.8. The difference in nodal size depicts the increase in the node degree centrality; the nodes with the most increased connectivity are located at the left parietal lobe.

Table 2. Significant increase in connectivity of RSNs as defined by Yeo et al. (Yeo et al., 2011). The black color signifies the within-network connectivity, the green color the bilateral connectivity between-networks, and the blue color the unilateral connectivity between-networks from the RSN in the first column to the RSN in the first row.

RSN	VIS	SMN	DAN	VAN	LIM	FPN	DMN
VIS							
SMN							
DAN							
VAN							
LIM							
FPN							
DMN							

Graph measures

The significant changes ($p < 0.05$) in global efficiency (GE), transitivity (TS), characteristic path length (CPL), clustering coefficient (CC), betweenness centrality (BC),

and the small-worldness (SW) measure, sigma (σ), between the pre- and post-networks were investigated using Analysis of Covariance, where graph measures served as the dependent values and density as the covariate (Table 3). Regarding the global measures, we report increases in GE and TS and a decrease in CPL. From the comparison of local measures, CC shows significant differences in 716 nodes, with increased values in 607 of these nodes (Figure 2). From these, 323 nodes belong to the left hemisphere (102 in the frontal lobe (FL), 39 in the central midline (CM), 56 in the parietal lobe (PL), 49 in the temporal lobe (TL), 26 in the limbic lobe (LL), 49 in the occipital lobe (OL)) and 284 to the right hemisphere (76 in FL, 51 in CM, 37 in PL, 49 in TL, 28 in LL, 44 in OL) (Figure 2). From the significant nodes, 109 showcased a significant decrease in value, with 48 nodes localized in the left hemisphere (7 in FL, 6 in CM, 6 in PL, 19 in TL, 2 in LL, 8 in OL) and 61 in the right hemisphere (22 in FL, 2 in CM, 8 in PL, 9 in TL, 3 in LL, 17 in OL). BC increased in 15 nodes with 9 pertaining to the left hemisphere (4 in the medial FL, 2 in PL (precuneus and superior PL), 1 in the inferior TL, 1 in the cingulate gyrus of LL, 1 in the middle OL) and 6 to the right (2 in FL (inferior and medial gyrus), 2 in the precuneus (PL), 1 in the cingulate gyrus (LL), 1 in the lingual gyrus (OL)), while it decreased in 2 nodes of the right angular gyrus (LL). The post vs. pre comparison of the SW measure revealed a significant increase in the σ -value, but the values for both the pre- and post-intervention networks did not exceed the SW threshold of $\sigma > 1$. We performed a multi-scale Louvain community detection, which showed that the networks of the two time-points are organized in different levels of hierarchy. The community analysis on the pre-intervention DS network indicated that its communities are nested within 3 levels of hierarchy with the number of communities across all levels ranging from 70 communities

to 442, while the communities of the post DS network are organized in 4 levels, with their number ranging from 45 to 290 across all levels.

Figure 2. Depiction of the shifts in the CC of nodes. The colormap indicates significance at an alpha-value ($\alpha < 0.05$) coupled with the direction of the shift. The black nodes have no significant shifts in CC values post- and pre-intervention, blue nodes show a significant increase at $\alpha < 0.05$, while magenta nodes show a significant decrease at $\alpha < 0.05$.

Table 3. Significant post-pre changes in global (Global efficiency, Transitivity, Characteristic Path Length) and local (Local Clustering Coefficient, Node Betweenness Centrality) graph measures as well as the small-worldness measure. Results were considered significant for $p < 0.05$. *For the local graph measures, the reported p-value is the mean value of the nodes showing a significant increase and a significant decrease, respectively.

Graph Measure	p-value	Direction of Effect
Global Efficiency	2.91e-12	Increase
Transitivity	3.38e-10	Increase
Characteristic Path Length	3.47e-10	Decrease
Clustering Coefficient	0.0115/0.0119*	Increase/Decrease
Node Betweenness Centrality	0.0386/0.0463*	Increase/Decrease
Small-Worldness	6.28e-08	Increase

Discussion

A ten-week intervention of combined PT and CT in adults with DS can trigger neuroplasticity resulting in cortical reorganization, quantified via EEG indices, and graph measures. Our results reveal that: i) short-term training can modify the physical and cognitive performance of adults with DS, ii) the DS brain can adapt to novel stimulation and challenges by utilizing neuroplasticity and reorganizing itself, and iii) adaptive neuroplasticity may lead to a more flexible and healthier functional network.

Short-term training modifies physical and cognitive performance

Our PT findings indicate the significant improvement of physical capacity for upper body strength and endurance, mobility, static and dynamic balance, and are in line with previous findings on adult PwDS (Hardee & Fetters, 2017; Ruiz-González, Lucena-Antón, Salazar, Martín-Valero, & Moral-Munoz, 2019). However, our protocol targets multiple physical domains, while most studies investigating the effect of PT on the somatic capacity of adult PwDS (Hardee & Fetters, 2017; Ruiz-González et al., 2019) have mainly focused on a specific domain, i.e. resistance training or aerobic training and, less often, on a combination of two or more domains.

With respect to CT, our PwDS exhibited an improved general cognitive capacity, i.e., in the level of general intelligence, planning and organization skills, and in short-term memory, attention, and concentration. Previous cognitive interventions on adult PwDS have mainly focused on memory (Fonseca et al., 2015) and, less often, on executive functions (McGlinchey et al., 2019). In line with the findings of the latter study, adult PwDS can complete a computerized CT program that targets multiple domains of cognition and exhibit significant improvements.

Cortical reorganization in the DS brain and neuroplastic capacity

PwDS present capacity for neuroplasticity. Regarding the type of neuroplasticity that is expressed here, we can rule out the triggering of evolutionary and reactive plasticity since the first is most active during the development of the brain (Trojan & Pokorný, 1999), and the latter is most commonly triggered by a single stimulus (Trojan & Pokorný, 1999) (our training utilizes repetitive stimuli). Adaptational and reparation plasticity can both cause

cortical reorganization through long-term stimulation. Here, combined PT and CT have increased the communication within and between nodes of the two hemispheres (Figure 1A&C). The DS cortical networks are characterized by a simplified organization (Anderson et al., 2013) with hyper-synchrony between brain regions (Anderson et al., 2013; Edgin et al., 2015; Vega et al., 2015). Therefore, the reported shifts in connectivity indicate the emergence of a reorganized network, which can be the result of either adaptational or reparation plasticity. Our finding regarding the increased left intra-hemispheric information flow can be regarded as the result of adaptational neuroplasticity being triggered in the DS brain given the atypical right-hemispheric functional preference (Elliott, Weeks, & Chua, 1994). However, the left-to-right hemisphere directionality is a pattern that has been previously evidenced in young PwDS (Babiloni et al., 2009), and its enhancement in adult PwDS may indicate that the training has triggered reparation neuroplasticity.

The enhanced connectivity within (DAN) and between (VIS and FPN, VIS and DMN, VIS and VAN, VAN and FPN, FPN and DMN, FPN and SMN, FPN and DAN, DAN and DMN, as well as DAN and SMN) the core RSNs (Yeo et al., 2011), further highlights the post-intervention DS functional reorganization. In respect to the aforementioned findings of Anderson and colleagues (Anderson et al., 2013), the strengthening of connections between distinct RSNs may also favor that the induced neuroplastic attributes represent adaptation. Given the evidence pointing towards an inverse relationship between the level of cognitive impairment of PwDS and the within DAN (Pujol et al., 2015; Vega et al., 2015) connectivity (i.e., the lower the DAN connectivity, the higher the cognitive impairment for DS), we suggest that the increased connectivity within DAN may associate

with the increased cognitive capacity due to training in our DS sample. This would rule out the possibility that the type of plasticity represents reparation. In essence, future studies could potentially confirm or rule out this possibility.

The cortical reorganization, in conjunction with the increases in general intelligence, indicates that the DS brain has entered a more flexible state. The DS network has been previously reported to lack the flexibility that characterizes the TD brain (Edgin et al., 2015) and subsequently higher intelligence (Barbey, 2018; Hilger, Fukushima, Sporns, & Fiebach, 2019). Flexibility can reflect the brain's capacity to adapt to novel stimulations (Barbey, 2018). It is plausible that the transition to a healthier organization emerges from the increased DS network flexibility and further points to the triggering of adaptational neuroplastic attributes.

Graph theory characteristics support the neuroplastic transition towards a healthier DS network organization

We suggest that the network's reorganization indexes a transitional state from a random-like network towards a healthier functional architecture, exhibiting less-random characteristics, increased flexibility and robustness, and improved integration and segregation capabilities (please see Figure 3 for an illustration of our theoretical proposal). In close relation, the SW measure, which classifies the pre- and post-intervention networks as non-SW, shows a significant increase in the post DS network and can, in turn, attribute the adult DS functional network (pre-intervention network) with features of a random network (Ahmadlou et al., 2013). Thus, the increased SW can index a transition to a less-random organization. The randomness of the pre-DS network is not surprising, given that

basic cognitive abilities (low level of general intelligence, which also characterizes PwDS) are exemplified by random networks, while broad cognitive abilities (high level of general intelligence) would require a SW organization (Barbey, 2018; Hilger et al., 2019; Watts & Strogatz, 1998). SW networks maintain an optimal balance of integration and segregation (combining random and regular network characteristics) to support general and specific abilities (Barbey, 2018; Hilger et al., 2019; Watts & Strogatz, 1998). In contrast, the absence of SW properties in the DS brain indicates the increased risk of communication loss between connected regions and their causality for cognitive decline (Ahmadlou et al., 2013; Sanz et al., 2010).

Considering the DS brain (lack of functional specialization and impaired integration (Anderson et al., 2013), the simultaneous increment in GE and TS denotes the increase of both integration and segregation and the transition to a healthier network organization, with the rise in TS further supporting the reorganization to a less-random architecture. The decrease of CPL in TD individuals could be interpreted as a contradicting shift towards a more random network that disturbs the SW balance. In our case, the CPL is a step towards the opposite direction since the adult DS network deviates from random networks by displaying a greater CPL, thus characterizing it as “random-like” (Figure 3). Previous fMRI studies report that the brain network of PwDS presents decreased long-range connectivity (Anderson et al., 2013; Pujol et al., 2015) and thus increased CPL (Bassett & Bullmore, 2006). Hence, the decreased CPL supports our hypothesis and, coupled with the previously mentioned shifts, signals the decrease in wiring cost and subsequent increase in cost-efficacy (Achard & Bullmore, 2007; Bullmore & Sporns, 2012).

Our hypothesis of a shift towards a healthier network organization is also evident from a local network perspective. The increased CC in most areas of the brain indicates a rise in robustness, and fault tolerance in the network (Achard & Bullmore, 2007), which can potentially serve in neuroprotection for PwDS (Ahmadlou et al., 2013). Still, the effectiveness of such a mechanism would rely on the trade-off between integration and segregation, mainly maintaining SW network characteristics (Stam, Jones, Nolte, Breakspear, & Scheltens, 2007). The DS network, when compared to a TD network architecture of distributed communities, has been found to exhibit a lobar organization with increased local communication (Anderson et al., 2013) and reduced BC (Hemmati et al., 2013) making the aggregation of information less efficient. The aforementioned evidence, coupled with our increased hierarchical organization of communities in the post DS network, and the increase of BC in nodes, support the transition to a more hierarchical organization, rendering the network more efficient in the coherent distribution of information.

Figure 3. Illustration of our theoretical proposal. Changes in connectivity and graph-theory characteristics, as an outcome of the adaptational neuroplasticity in the DS brain, characterize the post-intervention DS network (**B**→**C**) as a transitional state from the random-like organization of the pre-intervention DS network (Ahmadlou et al., 2013) (**B**) towards a healthier functional structure. Random networks (Erdős–Rényi, random graph) (**A**) are characterized by high global efficiency (over-integration) and low clustering (under-segregation) and exemplify the general abilities of general intelligence (low intelligence level). Small-world networks (**C**) feature the functional organization of TD brain networks and incorporate characteristics of both random and regular networks, achieving an optimal balance between global and local characteristics. SW networks (**C**) are associated with the broad abilities' component of general intelligence (higher intelligence level). The pre-intervention DS network (**B**) showed a random-like, simplified architecture, with impaired segregation and integration, as evidenced by the low CC (random network characteristic) and decreased GE and increased CPL, respectively. This is in line with previous literature (Ahmadlou et al., 2013; Anderson et al., 2013; Pujol et al., 2015; Vega et al., 2015). The DS pre-network's (**B**) integration-related characteristics (lower GE, higher CPL) are not common in random networks. Hence, the DS pre-network (**B**) is classified as random-like and not entirely random, maintaining an equal distance from random (**A**) and SW (**C**) networks. The DS post-network (**B**→**C**) exhibits an increase in integration (random network characteristic), as well as segregation (regular network characteristic), so it is interpreted as a step towards an SW-like architecture that highlights a healthier brain organization.

Future directions and limitations

A limitation of the study was the small sample size of the experimental group. Our results could potentially be interpreted differently in reference to a control group (TD or passive DS). Though this study is the first to provide insights into the benefits of combined

PT and CT in adults with DS through neuropsychological and neurophysiological assessments, a gap remains in our basic understanding of the sole contributions of each training to DS related cognition and brain function. Further studies are necessary to index the influence of each component on the triggering of neuroplasticity and subsequently network organization and cognition. Similarly, the stability of the reported neuroplastic shifts can only be addressed through additional and more in-depth investigations, including follow-up studies, to fill the gaps in the existing literature. Finally, the correlation of the somatometric and/or psychometric scores to the graph measures would increase the strength of our results, as it would indicate that the neuroplastic effect that we report is also reflected in behavior. We explored such a relationship between the changes in graph measures and the shifts in psychosomatic assessments (which showed post-pre significant changes) via a: i) correlation, ii) linear regression, and iii) multiple linear regression. The lack of a significant correlation was probably due to the small sample size, which did not produce enough power to quantify the effect, or due to the greater sensitivity that the neuroimaging methods show with regard to learning-induced neuroplasticity compared to the behavioral ones (Andrés, Parmentier, & Escera, 2006).

Conclusion

The rise of DS prevalence sets a significant challenge in developing innovative health care interventions that provide the best quality of care. Neurocognitive treatments can augment the functional and cognitive abilities of PwDS and thus allow for a more independent, productive, and fulfilling daily life. Our results, based on brain imaging, connectivity, and graph-theory analysis, emphasize the significance of introducing

stimulation and adaptable challenges in the environment of PwDS. We provide evidence that triggering adaptational neuroplasticity in the DS brain provokes the emergence of a less random, more hierarchical, and flexible network, able to integrate and segregate information more efficiently. Designing focalized neurobehavioral interventions can aid in the development of a balanced and stable functional DS phenotype in terms of SW characteristics.

Materials and Methods

Subjects

The study's pool consisted of 12 subjects with DS (age: 29 ± 11 , 6 females). Participants were recruited from a variety of local organizations (please see Acknowledgements). The training procedure took place in the Thessaloniki Active and Healthy Ageing Living Lab (Thess-AHALL) (Konstantinidis, Billis, Bratsas, Siountas, & Bamidis, 2016), and the premises of the Greek Association of Down Syndrome. This study is part of the dsLLM clinical trial, registered with ClinicalTrials.gov, with identifier code NCT04390321. The study protocol was approved by the Bioethics Committee of the Medical School of the Aristotle University of Thessaloniki and was conducted per the Helsinki Declaration of Human Rights. The participants' legal guardians signed written informed consent before their inclusion in the study.

Figure 4. DS-LLM Care design and flow of participants with DS.

LLM Care Intervention

The intervention protocol consisted of physical and cognitive training (LLM Care (Bamidis et al., 2015), <http://www.llmcare.gr/en>). The study protocol was further developed to improve the quality of life and aid in the development of independent living skills in PwDS (Bamidis, Konstantinidis, Billis, & Siountas, 2017; Romanopoulou, Zilidou, Savvidis, Chatzisevastou-Loukidou, & Bamidis, 2018). LLM Care also aims to further improve brain functionality (Bamidis et al., 2015). All training sessions were computerized, center-based, and conducted under supervision. The sequence of training methods was pseudo-randomized and counterbalanced. The details of each training intervention have been previously described in detail (Billis et al., 2010; Smith et al., 2009) and are summarized in Figure 4.

Cognitive training

The CT component of LLM Care uses the BrainHQ software (Posit Science Corporation, San Francisco, CA, USA), an online interactive environment in the Greek language (Romanopoulou, Zilidou, & Bamidis., 2017). It consists of six categories (29 exercises in total), with customizable difficulty levels, utilizing audiovisual stimuli. CT targets memory, attention, cognitive-processing speed, navigation, and people skills. This regime was selected since patients with DS exhibit deficiencies in these processes. CT was conducted for half an hour, with a frequency of 2 days per week for 10 weeks. In every CT session, the participants were required to complete at least one task from each category. During the training, participants were urged to complete as many exercises as they could from each category.

Physical training

The PT component of LLM Care is based on the WebFitForAll protocol (Billis et al., 2010; Konstantinidis, Bamparopoulos, & Bamidis, 2016) adequately adjusted to the needs and capacity of PwDS. It utilizes motion sensor devices (i.e., Kinect). Games and physical exercise are combined, providing a pleasant experience throughout the training. PT sessions lasted for half an hour and were conducted with the same frequency as CT. The training consists of aerobic (cycling, in-place-hiking), flexibility (stretching), strength (resistance, weightlifting), and balance (static, dynamic) exercises. The warm-up and cool-down routines (5-minutes duration) signify the start and completion of every session, respectively. During aerobic exercises, the participants entered a virtual environment, set up in Google maps, and explored cities and landscapes. Upon correct completion of the

flexibility and strength exercises, the trainees were progressively rewarded with an array of pleasing images. The scope of balance exercises was to move their bodies either horizontally or vertically, which was achieved through games.

Psychometric and somatometric assessments

The participants' cognitive and physical capacity was assessed before and after the intervention. The psychometric evaluation consisted of a set of neurocognitive tests that measure memory, attention, concentration (WISC-III: Digits Span) (Woolger, 2001), verbal and non-verbal mental capabilities (Raven) (John & Raven, 2003), processing speed (WISC-III: Digits Span and Picture Arrangement), problem-solving, visuospatial processing, organization skills (WISC-III: Block Design, Picture Arrangement, Mazes), social intelligence (WISC-III: Picture Arrangement), and identification of emotions (Reading the mind in the eyes (Baron-Cohen, Jolliffe, Mortimore, & Robertson, 1997; Baron-Cohen, Wheelwright, Hill, Raste, & Plumb, 2001), and a variation: Reading the mind in the face (emotion recognition from video)).

The somatometric evaluation included the Short Physical Performance Battery (SPPB) (Guralnik et al., 1994), 10 Meter Walk (Bohannon, 1997), Back Scratch (Jones & Rikli, 2002), Sit and Reach (Wells & Dillon, 1952), Arm Curl (Jones & Rikli, 2002), Four Square Step (FSST) (Whitney, Marchetti, Morris, & Sparto, 2007), Stork Balance (for both legs) (Johnson & Jack K. Nelson, 1979), Timed Up and Go (Shumway-Cook, Brauer, & Woollacott, 2000) tests, and Body Mass Index (BMI). These tests appraise functioning mobility, flexibility (in specific areas), dynamic stability, strength, and static and dynamic balance.

EEG recording

Pre- and post-intervention resting-state EEG activity was recorded for 5 minutes, using a high-density Nihon-Kohden EEG device (128 active scalp electrodes) at a sampling rate of 1000Hz. The EEG recordings were performed in an electrically shielded, sound, and light attenuated booth. The electrode impedances were lower than 10 k Ω . The participants were instructed to remain in a resting position while keeping their eyes open. Eyes-closed EEGs were not measured due to the limited capacity of the participants with DS to stay relaxed with their eyes closed.

Figure 5. EEG data analysis schematic. **Pre-processing** (blue): EEG data were interpolated and artifact corrected, visually inspected, high-pass, bandpass, and low-pass filtered. Independent component analysis (ICA), as well as visual inspection, were used to reject artifactual data. 15 segments of 4-s were randomly selected. **Source reconstruction** (orange): The data were processed within 0.53-35 Hz frequency range, source reconstructed (4-D LORETA, for all time points), and a previously used 863-node atlas (Paraskevopoulos et al., 2015) was applied to extract the time-series of every voxel from every segment per subject, which were used to estimate 1000 surrogate time-series per subject/group. **Functional connectivity** (green): Functional connectivity was computed for every segment of every subject, using the phase transfer entropy metric (PTE), and the 15 matrices of every subject were averaged into one. PTE networks were estimated for every surrogate time-series of each subject to create a null-model for the rejection of noise in the real data. A network science approach was taken for the computation of graph measures per subject. Group average statistics were calculated to identify the statistically significant differences between post- and pre-intervention networks.

EEG data analysis

Pre-processing

The raw EEG data were visually inspected, and bad channels were interpolated while any eye-movement related artifacts (blinks and horizontal movement) were corrected through adaptive artifact correction (Ille, Berg, & Scherg, 2002) using the Brain Electrical Source Analysis software (BESA research, version 6, Megis Software, Heidelberg, Germany) (Figure 5, blue section). The artifact-corrected data from each measurement were imported into the Fieldtrip Matlab toolbox (Oostenveld, Fries, Maris, & Schoffelen, 2011) for additional processing (Figure 5, blue section). The signals were denoised by

filtering the data (0.53 Hz high-pass IIR filter, 48-52 Hz notch IIR filter, 97 Hz low-pass IIR filter). The filtered signals were analyzed into independent components (Lee, Girolami, & Sejnowski, 1999), and the artifactual components were rejected. After signal reconstruction, the data was, once again, visually inspected for any remaining artifacts. Fifteen segments (4 seconds each) from each artifact-free EEG recording were randomly selected for additional processing.

Source reconstruction

Each subject's segments (15 segments, 4000 samples, 4 seconds duration each) were imported into BESA (Figure 5, orange section). The current density reconstructions (CDR) were estimated for each sample point, solving the inverse problem using LORETA in the 0.5-35 Hz frequency range. LORETA (Pascual-Marqui et al., 1994) was utilized, as it does not require the *a priori* declaration of the number of sources and is suitable for whole cortex analysis. The CDRs were exported as four-dimensional images (4-D) in the Analyze format (keeping all sampling points), which were, in turn, imported into Matlab. A cortex mask was superimposed on the images. The mask includes only grey matter and excludes the subcortex, the brainstem, and cerebellum, to limit the source space (Paraskevopoulos, Kraneburg, Herholz, Bamidis, & Pantev, 2015). The source space consisted of 863 voxels.

Functional connectivity

After extracting the time-series of each voxel from the 4-D images, they were used to compute the Phase Transfer Entropy (PTE) (Lobier, Siebenhühner, Palva, & Palva, 2014) (Figure 5, green section). The computation resulted in 863×863 adjacency matrices. The

metric is calculated independently for every pair of voxels in each segment. Each voxel represented a node of the brain network, with the node's coordinates corresponding to the center of each voxel. PTE was selected because its results are not based on a specific data model since its computation is reliant on non-linear probability distributions. Therefore, it allows the detection of higher-order relations in the phase information flow and renders the measure resistant to source leakage (Lobier et al., 2014). The algorithm applies the Hilbert transform to estimate the phases of each signal. It determines the number of bins by utilizing the Scott methodology (Scott, 1992), resulting in 37 bins on average (13 samples per bin). Each subject's adjacency matrices were averaged, resulting in two sets of 12 adjacency matrices (Figure 5, green section).

When computing network science indices based on information theory, one should take into account that the derived data contain noise that may mask the “real” result. This potential limitation was dealt with by following a methodology proposed by Timme et al. (Timme et al., 2016). Thus, subject-level null-distributions were computed and then compared with the real PTEs. 2000 surrogate time-series were estimated for each subject (1000 for the pre-intervention measurements and 1000 for the post-intervention ones) via phase randomization, i.e., calculating the Fourier transform of the real times-series, randomizing their phase, inverting the transform, thus resulting in jittered time-series (Timme et al., 2016). PTE networks were estimated from the shifted time-series and then compared edge by edge to the subject-PTE-networks, where any real-edge that had lesser weight than more than 50 (5% significance level) surrogate-edges was considered non-significant and removed from the final PTE network (Timme et al., 2016). To ensure that small differences between the surrogate and real data did not affect the edge-significance

test, the weight difference between them was estimated, and differences less than $e-03$ were ignored. The comparison of the surrogate PTE distributions to the PTE data resulted in networks consisting of 863 nodes and 184720 edges on average for the post-intervention networks and 123980 edges on average for the pre-intervention networks.

Graph measures computation

Using the Brain Connectivity Toolbox (Rubinov & Sporns, 2010), GE, TS (a variation of global CC), CPL, CC, BC, as well as community detection, and the measure of SW (σ) were computed for each participant. In a later step, the centrality Matlab function, which measures node importance, was used to calculate the node degree centrality (DC) of the nodes showcasing significant shifts in the post-intervention network. The density of each graph was estimated by averaging the weights of each graph and multiplying the average by the maximum weight of the network.

These graph measures were chosen to examine the influence of our intervention on the characteristics of the DS brain and its functional organization. To that aim, GE and CPL were utilized to index the effects on integration (Rubinov & Sporns, 2010), TS and CC to measure the impact on segregation (Rubinov & Sporns, 2010), BC and DC to check for shifts in the importance of nodes to the flow of information, and community detection to characterize the hierarchical organization.

GE is the average inverse shortest path length of the network (Latora & Marchiori, 2001) and quantifies the efficacy of information transference and its assimilation in the network (Rubinov & Sporns, 2010). CPL is the average shortest path length of the edges connecting the nodes of the network (Watts & Strogatz, 1998), is influenced by longer paths, and

characterizes the robustness of the network (Rubinov & Sporns, 2010). TS is the ratio of closed triplets to the maximum number of triplets (open and closed) (Newman, 2003), while CC of a node is the ratio of its connected neighbors to the maximum number of possible connections (Watts & Strogatz, 1998). These two measures reflect the clustering organization of the DS brain network on a global and local level, respectively (Rubinov & Sporns, 2010). BC corresponds to the fraction of shortest paths that pass through a node (Freeman, 1978) and is a measure of centrality, measuring the importance of a node in the information flow (Rubinov & Sporns, 2010), and DC estimates the number of links connected to a specific node, i.e., it measures if a node exhibits higher or lower connectivity to other nodes. To characterize the DS network's hierarchical organization, the number of communities that constitute the pre and post DS networks was estimated. Community detection was achieved through the implementation of a multi-iterative generalization of the Louvain community detection algorithm that optimizes modularity (Blondel, Guillaume, Lambiotte, & Lefebvre, 2008). Following the methodology of Akiki & Abdallah (Akiki & Abdallah, 2019), the detection was applied recursively to pinpoint the multi-scale, hierarchical community organization. A single-level detection was performed for 100 iterations for every subject, and from them, a group-representing partition was selected using the partition similarity measure of normalized mutual information. This partition was then used as a base, to extract subgraphs from each community, which were then fed into the algorithm for the next level of community detections. This process was repeated until no new communities were detected. The measure quantifying a network's small-worldness is sigma (σ). It is based on the ratio of the comparison of a network's CC and CPL to that of random networks (Bolaños, Bernat, He, & Aviyente, 2013; Humphries

& Gurney, 2008) with the network characterized as SW when $\sigma > 1$. To estimate the SW measure, we generated 100 random networks from each subject's pruned adjacency matrix while preserving the out-degree, with each edge being reshuffled at least 100 times, and calculated the σ in reference to each random network. Finally, we averaged the σ -values per subject. These random networks were also used to create a personalized null-distribution for each subject to mitigate the effect of density on graph measures when reporting individual results. For a summary of the graph measures used in the analysis, please refer to the Supplementary Material (Supplementary Table 1).

The comparison of our pre- and post-intervention network in relation to the exhibition of random or regular characteristics are discussed on a qualitative level, i.e., whether they show characteristics exemplary for a random organization (small CPL, low CC). These random characteristics are present in many random topologies, and hence we have selected the most simplified random organization, the Erdős–Rényi. The degree distribution of the DS networks is taken into consideration when this comparison is quantified (i.e., estimation of SW), where a generative random model that preserves the degree distribution (Newman, 2002) of the DS networks is used.

Statistical analysis

The pre- and post-intervention somatometric and psychometric assessment scores were compared with the use of non-parametric Wilcoxon tests and paired t-tests. The statistical comparisons were performed using the IBM SPSS 25.0 software. Wilcoxon tests were performed on the psychometric battery tests and somatometric tests with a discrete-values score, while paired t-tests were applied on the remaining somatometric assessment tests.

The Network Based Statistics (NBS) (Zalesky et al., 2010) MATLAB toolbox was employed to estimate the statistically significant differences between the whole-head network of the pre- and post-intervention connectivity networks of our subjects. A paired samples t-test corrected for 5000 random comparisons via the non-parametric NBS method (Zalesky et al., 2010) was performed. The permutation model of the NBS toolbox randomly exchanges the group label of each subject at every permutation level and then recalculates the statistic of interest applying the selected threshold. The maximal component size is derived from each permutation and utilized to create an empirical estimate of the null-distribution of maximal component size (Zalesky et al., 2010). For further information on this process, please refer to the following work (Hayasaka & Nichols, 2004; Nichols & Holmes, 2002). Significant differences between the two time-points were visualized as weighted graphs through the BrainNet Viewer (Xia, Wang, & He, 2013) toolbox (Figure 1). DC was estimated from the outcome of this comparison, and the results were depicted in the same graph.

For the graph measures, Analysis of Covariance (ANCOVA) was used, where each measure serves as the dependent variable with density as the covariate because density seriously affects the values of the other measures. For local measures (i.e., CC and BC), ANCOVA and FDR correction were performed for each of the 863 nodes and p-values for Type I errors, respectively. Results below the 5 percent threshold were considered significant.

Acknowledgments

This study is an extension of the European CIP-ICTPSP. 2008.1.4 Long Lasting Memories (LLM) project (Project no. 238904) (<http://www.longlastingmemories.eu/>). The authors would like to acknowledge the support (i.e. high-performance computing infrastructure and resources) provided by the IT Center of the Aristotle University of Thessaloniki (AUTH) throughout the progress of this research work. The authors would like to thank all the participants, their families, and caregivers, the Down Syndrome Association of Greece, Spring Children Mixed Living Center – Daily Care Activity & Training for Disabled People, The Down Syndrome Association of Serres, Center for the Rehabilitation of Social Support & Creative Employment of People with Disabilities, Child Care Center in Ptolemaida, Centre For Persons With Learning Difficulties - Zoodohos Pigi, Center of Creative Activities - Municipality of Neapoli and Sykies, Special laboratory for vocational education and training in Alexandria, Center for the Rehabilitation of Social Support & Creative Employment of People with disabilities "O Sotir", Center of Creative Activities for People with Disabilities-Municipality of Neapoli and Sykes. Also, the authors would like to thank a list of people assisting in the trial facilitation, as follows: Dimitris Bamidis, Foteini Dolianiti, Ioanna Dratsiou, Sotiria Gilou, Maria Karagianni, Katerina Katsouli, Eleni Manthou, Maria Metaxa, Annita Varella (in alphabetical order), as well as, students from the Dept of Psychology, The University of Sheffield International Faculty, City College (Greece) who were practicing under the supervision of Dr. Anna Vivas and Dr. Elissavet Chrysochoou who were also advising the authoring team on the psychological aspects of the whole endeavor. Last but not least, the authors would like to thank Emeritus Professor of Pediatrics and Genetics Dr. Charikleia Chatzisevastou-

Loukidou for advising the team on all medical aspects of the project and for linking the team with many of the pilot sites.

Conflict of Interest

There are potential conflicts of interest (other, not financial, outside the scope of the submitted work) for the author P. Bamidis in respect of PositScience and the Aristotle University of Thessaloniki. There is a co-marketing agreement between PositScience and the Aristotle University of Thessaloniki to exploit Brain HQ within the LLM Care self-funded initiative that emerged as the non-for-profit business exploitation of the Long Lasting Memories (LLM Project) (www.longlastingmemories.eu) originally funded by the ICT-CIP-PSP Program of the European Commission. Brain HQ now forms part of LLM Care, a technology transfer/self-funded initiative that emerged as the non-for-profit business exploitation of LLM. Additionally, FitForAll (FFA) has been developed in the Aristotle University of Thessaloniki during the Long Lasting Memories (LLM Project) (www.longlastingmemories.eu) originally funded by the ICT-CIP-PSP Program of the European Commission. It now forms part of LLM Care, a technology transfer/self-funded initiative that emerged as the non-for-profit business exploitation of LLM.

References

- Achard, S., & Bullmore, E. (2007). Efficiency and cost of economical brain functional networks. *PLoS Computational Biology*, 3(2), e17.
<https://doi.org/10.1371/journal.pcbi.0030017>
- Ahmadlou, M., Gharib, M., Hemmati, S., Vameghi, R., & Sajedi, F. (2013). Disrupted

small-world brain network in children with Down Syndrome. *Clinical Neurophysiology*, 124(9), 1755–1764.
<https://doi.org/10.1016/J.CLINPH.2013.03.004>

Akiki, T. J., & Abdallah, C. G. (2019). Determining the Hierarchical Architecture of the Human Brain Using Subject-Level Clustering of Functional Networks. *Scientific Reports*, 9(1), 1–15. <https://doi.org/10.1038/s41598-019-55738-y>

Anderson, J. S., Nielsen, J. A., Ferguson, M. A., Burbach, M. C., Cox, E. T., Dai, L., ... Korenberg, J. R. (2013). Abnormal brain synchrony in Down Syndrome. *NeuroImage: Clinical*, 2, 703–715. <https://doi.org/10.1016/J.NICL.2013.05.006>

Andrés, P., Parmentier, F. B. R., & Escera, C. (2006). The effect of age on involuntary capture of attention by irrelevant sounds: A test of the frontal hypothesis of aging. *Neuropsychologia*, 44(12), 2564–2568.
<https://doi.org/10.1016/j.neuropsychologia.2006.05.005>

Babiloni, C., Albertini, G., Onorati, P., Muratori, C., Buffo, P., Condoluci, C., ... Rossini, P. M. (2010). Cortical sources of EEG rhythms are abnormal in down syndrome. *Clinical Neurophysiology*, 121(8), 1205–1212.
<https://doi.org/10.1016/j.clinph.2010.02.155>

Babiloni, C., Albertini, G., Onorati, P., Vecchio, F., Buffo, P., Sarà, M., ... Rossini, P. M. (2009). Inter-hemispheric functional coupling of eyes-closed resting EEG rhythms in adolescents with Down syndrome. *Clinical Neurophysiology*, 120(9), 1619–1627.
<https://doi.org/10.1016/j.clinph.2009.06.017>

Bamidis, P. D., Fissler, P., Papageorgiou, S. G., Zilidou, V. I., Konstantinidis, E. I., Billis, A. S., ... Kolassa, I. T. (2015). Gains in cognition through combined

cognitive and physical training: the role of training dosage and severity of
neurocognitive disorder. *Frontiers in Aging Neuroscience*, 7(152), 152.
<https://doi.org/10.3389/fnagi.2015.00152>

Bamidis, P. D., Konstantinidis, E. I., Billis, A. S., & Siountas, A. (2017). From e-homes
to living labs: founding and organising the Greek active and healthy ageing living
lab (Thess-AHALL) and its networked services. *Hellenic Journal of Nuclear
Medicine*, 20(2), 112–126.

Bamidis, P. D., Vivas, A. B., Styliadis, C., Frantzidis, C. A., Klados, M. A., Schlee, W.,
... Papageorgiou, S. G. (2014). A review of physical and cognitive interventions in
aging. *Neurosci Biobehav Rev.*, 44, 206–220.
<https://doi.org/10.1016/j.neubiorev.2014.03.019>

Barbey, A. K. (2018). Network Neuroscience Theory of Human Intelligence. *Trends in
Cognitive Sciences*, 22(1), 8–20. <https://doi.org/10.1016/j.tics.2017.10.001>

Baron-Cohen, S., Jolliffe, T., Mortimore, C., & Robertson, M. (1997). Another advanced
test of theory of mind: evidence from very high functioning adults with autism or
asperger syndrome. *Journal of Child Psychology and Psychiatry, and Allied
Disciplines*, 38(7), 813–822.

Baron-Cohen, S., Wheelwright, S., Hill, J., Raste, Y., & Plumb, I. (2001). The “Reading
the Mind in the Eyes” Test revised version: a study with normal adults, and adults
with Asperger syndrome or high-functioning autism. *Journal of Child Psychology
and Psychiatry, and Allied Disciplines*, 42(2), 241–251.

Baroncelli, L., Braschi, C., Spolidoro, M., Begenisic, T., Maffei, L., & Sale, A. (2011).
Brain plasticity and disease: A matter of inhibition. *Neural Plasticity*, 2011.

748 <https://doi.org/10.1155/2011/286073>

749 Bassett, D. S., & Bullmore, E. (2006). Small-World Brain Networks. *The Neuroscientist*,
750 12(6), 512–523. <https://doi.org/10.1177/1073858406293182>

751 Bertapelli, F., Pitetti, K., Agiovlasitis, S., & Guerra-Junior, G. (2016). Overweight and
752 obesity in children and adolescents with Down syndrome—prevalence,
753 determinants, consequences, and interventions: A literature review. *Research in*
754 *Developmental Disabilities*, 57, 181–192.
755 <https://doi.org/10.1016/J.RIDD.2016.06.018>

756 Billis, A. S., Konstantinidis, E. I., Mouzakidis, C., Tsolaki, M. N., Pappas, C., &
757 Bamidis, P. D. (2010). A game-like interface for training seniors' dynamic balance
758 and coordination. In *XII Mediterranean Conference on Medical and Biological*
759 *Engineering and Computing* (pp. 691–694). Springer Berlin Heidelberg.

760 Blondel, V. D., Guillaume, J. L., Lambiotte, R., & Lefebvre, E. (2008). Fast unfolding of
761 communities in large networks. *Journal of Statistical Mechanics: Theory and*
762 *Experiment*, 2008(10), P10008. <https://doi.org/10.1088/1742-5468/2008/10/P10008>

763 Bohannon, R. W. (1997). Comfortable and maximum walking speed of adults aged 20-79
764 years: reference values and determinants. *Age and Ageing*, 26(1), 15–19.
765 <https://doi.org/10.1093/ageing/26.1.15>

766 Bolaños, M., Bernat, E. M., He, B., & Aviyente, S. (2013). A weighted small world
767 network measure for assessing functional connectivity. *Journal of Neuroscience*
768 *Methods*, 212(1), 133–142. <https://doi.org/10.1016/j.jneumeth.2012.10.004>

769 Bullmore, E., & Sporns, O. (2012). The economy of brain network organization. *Nature*
770 *Reviews. Neuroscience*, 13(5), 336–349. <https://doi.org/10.1038/nrn3214>

771 Colcombe, S. J., Erickson, K. I., Scalf, P. E., Kim, J. S., Prakash, R., McAuley, E., ...
772 Kramer, A. F. (2006). Aerobic Exercise Training Increases Brain Volume in Aging
773 Humans. *The Journals of Gerontology Series A: Biological Sciences and Medical*
774 *Sciences*, 61(11), 1166–1170. <https://doi.org/10.1093/gerona/61.11.1166>
775 Colcombe, S. J., Kramer, A. F., Erickson, K. I., Scalf, P. E., McAuley, E., Cohen, N. J.,
776 ... Elavsky, S. (2004). Cardiovascular fitness, cortical plasticity, and aging.
777 *Proceedings of the National Academy of Sciences of the United States of America*,
778 101(9), 3316–3321. <https://doi.org/10.1073/pnas.0400266101>
779 Costa, A. C. S. (2011). On the promise of pharmacotherapies targeted at cognitive and
780 neurodegenerative components of Down syndrome. *Developmental Neuroscience*,
781 33(5), 414–427. <https://doi.org/10.1159/000330861>
782 Cotman, C., Berchtold, N. C., & Christie, L. A. (2007). Exercise builds brain health: key
783 roles of growth factor cascades and inflammation. *Trends in Neurosciences*, 30(9),
784 464–472. <https://doi.org/10.1016/j.tins.2007.06.011>
785 Dierssen, M. (2012). Down syndrome: the brain in trisomic mode. *Nature Reviews*
786 *Neuroscience*, 13(12), 844–858. <https://doi.org/10.1038/nrn3314>
787 Dierssen, M., Herault, Y., & Estivill, X. (2009). Aneuploidy: From a Physiological
788 Mechanism of Variance to Down Syndrome. *Physiological Reviews*, 89(3), 887–
789 920. <https://doi.org/10.1152/physrev.00032.2007>
790 Edgin, J. O., Clark, C. A. C., Massand, E., & Karmiloff-Smith, A. (2015). Building an
791 adaptive brain across development: Targets for neurorehabilitation must begin in
792 infancy. *Frontiers in Behavioral Neuroscience*, 9.
793 <https://doi.org/10.3389/fnbeh.2015.00232>

794 Elliott, D., Weeks, D. J., & Chua, R. (1994). Anomalous Cerebral Lateralization and
795 Down Syndrome. *Brain and Cognition*, 26(2), 191–195.
796 <https://doi.org/10.1006/BRCG.1994.1050>

797 Erickson, K. I., Voss, M. W., Prakash, R., Basak, C., Szabo, A., Chaddock, L., ...
798 Kramer, A. F. (2011). Exercise training increases size of hippocampus and improves
799 memory. *Proceedings of the National Academy of Sciences (PNAS)*, 108(7), 3017–
800 3022.

801 Fonseca, L. M., Navatta, A. C. R., Bottino, C. M. C., & Miotto, E. C. (2015). Cognitive
802 Rehabilitation of Dementia in Adults with Down Syndrome: A Review of Non-
803 Pharmacological Interventions. *Dementia and Geriatric Cognitive Disorders Extra*,
804 5(3), 330–340. <https://doi.org/10.1159/000438858>

805 Freeman, L. C. (1978). Centrality in social networks conceptual clarification. *Social*
806 *Networks*, 1(3), 215–239. [https://doi.org/10.1016/0378-8733\(78\)90021-7](https://doi.org/10.1016/0378-8733(78)90021-7)

807 García-Alba, J., Ramírez-Toraño, F., Esteba-Castillo, S., Bruña, R., Moldenhauer, F.,
808 Novell, R., ... Fernández, A. (2019). Neuropsychological and neurophysiological
809 characterization of mild cognitive impairment and Alzheimer’s disease in Down
810 syndrome. *Neurobiology of Aging*, 84, 70–79.
811 <https://doi.org/10.1016/j.neurobiolaging.2019.07.017>

812 Guralnik, J. M., Simonsick, E. M., Ferrucci, L., Glynn, R. J., Berkman, L. F., Blazer, D.
813 G., ... Wallace, R. B. (1994). A short physical performance battery assessing lower
814 extremity function: association with self-reported disability and prediction of
815 mortality and nursing home admission. *Journal of Gerontology*, 49(2), M85-94.

816 Haier, R. J., Head, K., Head, E., & Lott, I. T. (2008). Neuroimaging of individuals with

817 Down's syndrome at-risk for dementia: Evidence for possible compensatory events.
818 *NeuroImage*, 39(3), 1324–1332. <https://doi.org/10.1016/j.neuroimage.2007.09.064>

819 Hardee, J. P., & Fetters, L. (2017). The effect of exercise intervention on daily life
820 activities and social participation in individuals with Down syndrome: A systematic
821 review. *Research in Developmental Disabilities*, 62, 81–103.
822 <https://doi.org/10.1016/j.ridd.2017.01.011>

823 Hayasaka, S., & Nichols, T. E. (2004). Combining voxel intensity and cluster extent with
824 permutation test framework. *NeuroImage*, 23(1), 54–63.
825 <https://doi.org/10.1016/j.neuroimage.2004.04.035>

826 Head, E., Lott, I. T., Patterson, D., Doran, E., & Haier, R. J. (2007). Possible
827 Compensatory Events in Adult Down Syndrome Brain Prior to the Development of
828 Alzheimer Disease Neuropathology: Targets for Nonpharmacological Intervention.
829 *Journal of Alzheimer's Disease*, 11(1), 61–76. [https://doi.org/10.3233/JAD-2007-](https://doi.org/10.3233/JAD-2007-11110)
830 11110

831 Hemmati, S., Ahmadlou, M., Gharib, M., Vameghi, R., & Sajedi, F. (2013). Down
832 syndrome's brain dynamics: analysis of fractality in resting state. *Cognitive*
833 *Neurodynamics*, 7(4), 333–340. <https://doi.org/10.1007/s11571-013-9248-y>

834 Hilger, K., Fukushima, M., Sporns, O., & Fiebach, C. J. (2019). Temporal stability of
835 functional brain modules associated with human intelligence. *Human Brain*
836 *Mapping*, 41(2), 362–372. <https://doi.org/10.1002/hbm.24807>

837 Humphries, M. D., & Gurney, K. (2008). Network “small-world-ness”: A quantitative
838 method for determining canonical network equivalence. *PLoS ONE*, 3(4).
839 <https://doi.org/10.1371/journal.pone.0002051>

840 Ille, N., Berg, P., & Scherg, M. (2002). Artifact correction of the ongoing EEG using
841 spatial filters based on artifact and brain signal topographies. *Journal of Clinical*
842 *Neurophysiology : Official Publication of the American Electroencephalographic*
843 *Society*, 19(2), 113–124. Retrieved from
844 <http://www.ncbi.nlm.nih.gov/pubmed/11997722>

845 John, & Raven, J. (2003). Raven Progressive Matrices. In *Handbook of Nonverbal*
846 *Assessment* (pp. 223–237). Boston, MA: Springer US.

847 Johnson, B. L., & Jack K. Nelson. (1979). *Practical Measurements for evaluation in*
848 *physical education*. (Burgess, Ed.) (4th ed.). Minneapolis.

849 Jones, C. J., & Rikli, R. E. (2002). Measuring Functional Fitness in Older Adults. *The*
850 *Journal of Active Ageing*, 1(1), 24–30.

851 Kálmán, J., Juhász, A., Laird, G., Dickens, P., Járdánházy, T., Rimanóczy, A., ... Janka,
852 Z. (2009). Serum interleukin-6 levels correlate with the severity of dementia in
853 Down syndrome and in Alzheimer's disease. *Acta Neurologica Scandinavica*, 96(4),
854 236–240. <https://doi.org/10.1111/j.1600-0404.1997.tb00275.x>

855 Kishnani, P. S., Heller, J. H., Spiridigliozzi, G. A., Lott, I., Escobar, L., Richardson, S.,
856 ... McRae, T. (2010). Donepezil for treatment of cognitive dysfunction in children
857 with Down syndrome aged 10-17. *American Journal of Medical Genetics Part A*,
858 152(12), 3028–3035. <https://doi.org/10.1002/ajmg.a.33730>

859 Klados, M. A., Styliadis, C., Frantzidis, C. A., Paraskevopoulos, E., & Bamidis, P. D.
860 (2016). Beta-Band Functional Connectivity is Reorganized in Mild Cognitive
861 Impairment after Combined Computerized Physical and Cognitive Training.
862 *Frontiers in Neuroscience*, 10, 55. <https://doi.org/10.3389/fnins.2016.00055>

863 Konstantinidis, E. I., Bamparopoulos, G., & Bamidis, P. D. (2016). Moving Real
864 Exergaming Engines on the Web: The webFitForAll case study in an active and
865 healthy ageing living lab environment. *IEEE J Biomed Health Inform.*, 21(3), 859–
866 866. <https://doi.org/10.1109/JBHI.2016.2559787>

867 Konstantinidis, E. I., Billis, A. S., Bratsas, C., Siountas, A., & Bamidis, P. D. (2016).
868 Thessaloniki Active and Healthy Ageing Living Lab: the roadmap from a specific
869 project to a living lab towards openness. In *9th International Conference Pervasive*
870 *Technologies Related to Assistive Environments (PETRA)* (pp. 1–4).

871 Kramer, A. F., Hahn, S., Cohen, N. J., Banich, M. T., McAuley, E., Harrison, C. R., ...
872 Colcombe, A. (1999). Ageing, fitness and neurocognitive function. *Nature*,
873 400(6743), 418–419. <https://doi.org/10.1038/22682>

874 Latora, V., & Marchiori, M. (2001). Efficient Behavior of Small-World Networks.
875 *Physical Review Letters*, 87(19), 198701.
876 <https://doi.org/10.1103/PhysRevLett.87.198701>

877 Lee, T. W., Girolami, M., & Sejnowski, T. J. (1999). Independent Component Analysis
878 using an Extended Infomax Algorithm for Mixed Sub - Gaussian and Super -
879 Gaussian Sources. *Neural Computation*, 11(2), 409–433.
880 <https://doi.org/10.1162/089976699300016719>

881 Lobier, M., Siebenhühner, F., Palva, S., & Palva, J. M. (2014). Phase transfer entropy: A
882 novel phase-based measure for directed connectivity in networks coupled by
883 oscillatory interactions. *NeuroImage*, 85, 853–872.
884 <https://doi.org/10.1016/j.neuroimage.2013.08.056>

885 Lott, I. T., Doran, E., Nguyen, V. Q., Tournay, A., Head, E., & Gillen, D. L. (2011).

Down syndrome and dementia: A randomized, controlled trial of antioxidant supplementation. *American Journal of Medical Genetics Part A*, 155(8), 1939–1948.
<https://doi.org/10.1002/ajmg.a.34114>

McGlinchey, E., McCarron, M., Holland, A., & McCallion, P. (2019). Examining the effects of computerised cognitive training on levels of executive function in adults with Down syndrome. *Journal of Intellectual Disability Research*, 63(9), 1137–1150.

Menghini, D., Costanzo, F., & Vicari, S. (2011). Relationship Between Brain and Cognitive Processes in Down Syndrome. *Behavior Genetics*, 41(3), 381–393.
<https://doi.org/10.1007/s10519-011-9448-3>

Mohan, M., Carpenter, P. K., & Bennett, C. (2009). Donepezil for dementia in people with Down syndrome. *Cochrane Database of Systematic Reviews*, (1).

Newman, M. E. J. (2002). Random graphs as models of networks. *Handbook of Graphs and Networks*, 35–68.

Newman, M. E. J. (2003). The structure and function of complex networks. *SIAM Review*, 45(2), 167–256. <https://doi.org/10.1137/S003614450342480>

Nichols, T. E., & Holmes, A. (2002). Nonparametric permutation tests for functional neuroimaging: A primer with examples. *Human Brain Mapping*, 15(1), 1–25.
<https://doi.org/10.1002/hbm.1058>

Oostenveld, R., Fries, P., Maris, E., & Schoffelen, J. M. (2011). FieldTrip: Open source software for advanced analysis of MEG, EEG, and invasive electrophysiological data. *Computational Intelligence and Neuroscience*, 2011, 156869.
<https://doi.org/10.1155/2011/156869>

- 909 Paraskevopoulos, E., Kraneburg, A., Herholz, S. C., Bamidis, P. D., & Pantev, C. (2015).
910 Musical expertise is related to altered functional connectivity during audiovisual
911 integration. *Proceedings of the National Academy of Sciences*, 112(40), 201510662.
912 <https://doi.org/10.1073/pnas.1510662112>
- 913 Pascual-Marqui, R. D., Michel, C. M., & Lehmann, D. (1994). Low resolution
914 electromagnetic tomography: a new method for localizing electrical activity in the
915 brain. *International Journal of Psychophysiology*, 18(1), 49–65.
916 [https://doi.org/10.1016/0167-8760\(84\)90014-X](https://doi.org/10.1016/0167-8760(84)90014-X)
- 917 Pinter, J. D., Eliez, S., Schmitt, J. E., Capone, G. T., & Reiss, A. L. (2001).
918 Neuroanatomy of Down’s Syndrome: A High-Resolution MRI Study. *American*
919 *Journal of Psychiatry*, 158(10), 1659–1665.
920 <https://doi.org/10.1176/appi.ajp.158.10.1659>
- 921 Pujol, J., del Hoyo, L., Blanco-Hinojo, L., de Sola, S., Macià, D., Martínez-Vilavella, G.,
922 ... de la Torre, R. (2015). Anomalous brain functional connectivity contributing to
923 poor adaptive behavior in Down syndrome. *Cortex*, 64, 148–156. Retrieved from
924 <http://www.ncbi.nlm.nih.gov/pubmed/25461715>
- 925 Romanopoulou, E. D., Zilidou, V. I., & Bamidis., P. D. (2017). Creating and sustaining a
926 social health care ecosystem: the case of LLM care services in Greece. *Hellenic*
927 *Journal of Nuclear Medicine*, 20(2), 40–48.
- 928 Romanopoulou, E. D., Zilidou, V. I., Savvidis, T., Chatzisevastou-Loukidou, C., &
929 Bamidis, P. D. (2018). Unmet Needs of Persons with Down Syndrome: How
930 Assistive Technology and Game-Based Training May Fill the Gap. *Studies in Health*
931 *Technology and Informatics*, 251, 15–18.

- 932 Rubinov, M., & Sporns, O. (2010). Complex network measures of brain connectivity:
933 uses and interpretations. *NeuroImage*, 52(3), 1059–1069.
934 <https://doi.org/10.1016/j.neuroimage.2009.10.003>
- 935 Ruiz-González, L., Lucena-Antón, D., Salazar, A., Martín-Valero, R., & Moral-Munoz, J.
936 A. (2019). Physical therapy in Down syndrome: systematic review and meta-
937 analysis. *Journal of Intellectual Disability Research*, 63(8), 1041–1067.
938 <https://doi.org/10.1111/jir.12606>
- 939 Sanz, J. M., Schoonheim, M. M., Damoiseaux, J. S., Rombouts, S. A., Maris, E.,
940 Barkhof, F., ... Stam, C. J. (2010). Loss of “Small-World” Networks in Alzheimer’s
941 Disease: Graph Analysis of fMRI Resting-State Functional Connectivity. *PLoS*
942 *ONE*, 5(11). <https://doi.org/10.1371/journal.pone.0013788>
- 943 Scott, D. W. (1992). *Multivariate density estimation : theory, practice, and visualization*.
944 Wiley.
- 945 Shumway-Cook, A., Brauer, S., & Woollacott, M. (2000). Predicting the probability for
946 falls in community-dwelling older adults using the Timed Up & Go Test.
947 *Physical Therapy*, 80(9), 896–903.
- 948 Smith, G. E., Housen, P., Yaffe, K., Ruff, R., Kennison, R. F., Mahncke, H. W., &
949 Zelinski, E. M. (2009). A cognitive training program based on principles of brain
950 plasticity: Results from the improvement in memory with plasticity-based adaptive
951 cognitive training (IMPACT) study. *Journal of the American Geriatrics Society*,
952 57(4), 594–603.
- 953 Sporns, O., & Betzel, R. F. (2016). Modular Brain Networks. *Annual Review of*
954 *Psychology*, 67(1), 613–640. <https://doi.org/10.1146/annurev-psych-122414-033634>

955 Stam, C. J., Jones, B. F., Nolte, G., Breakspear, M., & Scheltens, P. (2007). Small-world
956 networks and functional connectivity in Alzheimer's disease. *Cerebral Cortex*,
957 17(1), 92–99. <https://doi.org/10.1093/cercor/bhj127>

958 Styliadis, C., Kartsidis, P., & Paraskevopoulos, E. (2015). Neuroimaging approaches for
959 elderly studies. In P. D. Bamidis, I. Tarnanas, L. Hadjileontiadis, & M. Tsolaki
960 (Eds.), *Handbook of Research on Innovations in the Diagnosis and Treatment of*
961 *Dementia* (pp. 47–86). IGI Global.

962 Styliadis, C., Kartsidis, P., Paraskevopoulos, E., Ioannides, A. A., & Bamidis, P. D.
963 (2015). Neuroplastic Effects of Combined Computerized Physical and Cognitive
964 Training in Elderly Individuals at Risk for Dementia: An eLORETA Controlled
965 Study on Resting States. *Neural Plasticity*, 2015.
966 <https://doi.org/10.1155/2015/172192>

967 Takashima, M., Takahashi, S., Iwata, O., Iwata, S., Yokoo, M., & Takashima, S. (2014).
968 Neuronal Plasticity in the Developing and Aging Cerebral Cortices of Patients with
969 Down Syndrome. *International Journal of Neurorehabilitation*, 01(02).
970 <https://doi.org/10.4172/2376-0281.1000111>

971 Timme, N. M., Ito, S., Myroshnychenko, M., Nigam, S., Shimono, M., Yeh, F. C., ...
972 Beggs, J. M. (2016). High-Degree Neurons Feed Cortical Computations. *PLoS*
973 *Computational Biology*, 12(5), e1004858.
974 <https://doi.org/10.1371/journal.pcbi.1004858>

975 Trojan, S., & Pokorny, J. (1999). Theoretical aspects of neuroplasticity. *Physiological*
976 *Research*, 48, 87–98.

977 Vega, J. N., Hohman, T. J., Pryweller, J. R., Dykens, E. M., & Thornton-Wells, T. A.

978 (2015). Resting-State Functional Connectivity in Individuals with Down Syndrome
979 and Williams Syndrome Compared with Typically Developing Controls. *Brain*
980 *Connectivity*, 5(8), 461–475. <https://doi.org/10.1089/brain.2014.0266>

981 Watts, D. J., & Strogatz, S. H. (1998). Collective dynamics of ‘small-world’ networks.
982 *Nature*, 393(6684), 440–442. <https://doi.org/10.1038/30918>

983 Wells, K. F., & Dillon, E. K. (1952). The Sit and Reach—A Test of Back and Leg
984 Flexibility. *Research Quarterly. American Association for Health, Physical*
985 *Education and Recreation*, 23(1), 115–118.
986 <https://doi.org/10.1080/10671188.1952.10761965>

987 White, N. S., Alkire, M. T., & Haier, R. J. (2003). A voxel-based morphometric study of
988 nondemented adults with Down Syndrome. *NeuroImage*, 20(1), 393–403.
989 [https://doi.org/10.1016/S1053-8119\(03\)00273-8](https://doi.org/10.1016/S1053-8119(03)00273-8)

990 Whitney, S. L., Marchetti, G. F., Morris, L. O., & Sparto, P. J. (2007). The Reliability
991 and Validity of the Four Square Step Test for People With Balance Deficits
992 Secondary to a Vestibular Disorder. *Archives of Physical Medicine and*
993 *Rehabilitation*, 88(1), 99–104. <https://doi.org/10.1016/j.apmr.2006.10.027>

994 Woolger, C. (2001). Wechsler intelligence scale for children-(WISC-III). In
995 *Understanding Psychological Assessment* (pp. 219–233). Boston, MA: Springer US.
996 https://doi.org/10.1007/978-1-4615-1185-4_11

997 Xia, M., Wang, J., & He, Y. (2013). BrainNet Viewer: a network visualization tool for
998 human brain connectomics. *PloS One*, 8(7), e68910.
999 <https://doi.org/10.1371/journal.pone.0068910>

1000 Yeo, B. T. T., Krienen, F. M., Sepulcre, J., Sabuncu, M. R., Lashkari, D., Hollinshead,

1001 M., ... Buckner, R. L. (2011). The organization of the human cerebral cortex
 1002 estimated by intrinsic functional connectivity. *Journal of Neurophysiology*, 106(3),
 1003 1125–1165. <https://doi.org/10.1152/jn.00338.2011>
 1004 Zalesky, A., Fornito, A., & Bullmore, E. T. (2010). Network-based statistic: identifying
 1005 differences in brain networks. *NeuroImage*, 53(4), 1197–1207.
 1006 <https://doi.org/10.1016/j.neuroimage.2010.06.041>
 1007

A POST vs. PRE CIRCULAR

B

PRE

POST

C

POST vs. PRE

L

R

L

R

A**RANDOM NETWORK**

over-integration
under-segregation
increased wiring cost

small CPL
high GE
low CC
low LE

GENERAL ABILITIES**B****RANDOM-LIKE NETWORK****PRE-INTERVENTION
DS NETWORK**

impaired integration
under-segregation
increased wiring cost

greater CPL
smaller GE
low CC
low LE

B**TRANSITIONAL STATE****POST-INTERVENTION
DS NETWORK**

increased integration
increased segregation
reduced wiring cost

decreased CPL
increased GE
increased CC
increased LE

C**SMALL-WORLD NETWORK**

balanced integration
balanced segregation
optimal wiring cost

small CPL
high GE
high CC
high LE

BROAD ABILITIES

Subjects

12 DS individuals
age: 29 ± 11 , 6 females
10 week protocol

Pre-measurements

Psychometric and somatometric assessment tests
5-minute resting state EEG measurement

Intervention details

2 CT sessions/week, 30 minutes each
audiovisual stimuli, memory, attention,
cognitive processing speed and orientation
average 20 ± 3 sessions

2 PT sessions/week, 30 minutes each
aerobic, flexibility, strength and balance
average 18 ± 4 sessions

Post-measurements

Psychometric and somatometric assessment tests
5-minute resting state EEG measurement

5-min RS EEG recording

128 channels

Interpolations & Artefact corrections

Pre-processing

Filtering

ICA

Visual Inspection

15 random
4s epochs

Source Reconstruction

4-D LORETA calculation for every segment (0.53-35Hz)

Functional Connectivity

Identification of statistically significant network differences

Group average statistics

averaged
PTE per
subject

PTE adjacency matrices for each segment of every subject

Voxel Time-series for each segment/subject

Graph measures computation