Impact of the COVID-19 pandemic on cognitive function in Japanese community-dwelling

older adults in a class for preventing cognitive decline

Short title: Impact of COVID-19 on Japanese community-dwelling older adults

Minoru Kouzuki ^{1*}, Shota Furukawa ¹, Keisuke Mitani ¹, and Katsuya Urakami ¹

¹ Department of Biological Regulation, School of Health Science, Faculty of Medicine,

Tottori University, Tottori, Japan

* Corresponding author

Email: kouzuki@tottori-u.ac.jp

Abstract

We examined the effects of lifestyle and thoughts on cognitive function and change in cognitive function due to restrictions in daily life during the coronavirus disease 2019 (COVID-19) pandemic in community-dwelling older adults with mild cognitive decline. This was a retrospective case-control study. The participants were 88 older adults with mild cognitive decline who participated in a class designed to help prevent cognitive decline. The class was suspended from early-March to end of May 2020 to prevent the spread of COVID-19, and resumed in June 2020. We collected demographic and cognitive function test data (Touch Panel-type Dementia Assessment Scale [TDAS]) before and after class suspension and questionnaire data on their lifestyle and thoughts during the suspension. Change in TDAS scores from before and after the suspension was used to divide the participants into decline (2 or more points worsening) and non-decline (all other participants) groups, with 16 (18.2%) and 72 (81.8%) participants in each group, respectively. A logistic regression model showed that the odds ratio (OR) for cognitive decline was lower in participants whose responses were “engaged in hobbies” (OR = 0.07, $p = 0.015$), “worked on a worksheet about cognitive training provided by the town hall” (OR = 0.19, $p = 0.026$), and “had conversations over the phone” (OR = 0.28, $p = 0.0495$). There was

29 a significant improvement in TDAS scores after class was resumed ($p < 0.01$). A proactive
 30 approach to intellectual activities and social ties may be important for the prevention of
 31 cognitive decline during periods of restrictions due to COVID-19. We found that cognitive
 32 function test scores before class suspension significantly improved after resuming classes.
 33 We speculate that continued participation in the class led to positive behavioral changes in
 34 daily life during periods of restriction due to COVID-19.

Introduction

Recently, local governments throughout Japan have made efforts against dementia. In Houki town (Tottori Prefecture, Japan), classes are held for community-dwelling older adults with suspected mild cognitive decline to provide cognitive decline prevention programs that combine physical exercise, cognitive training, and education on dementia and lifestyle habits [1]. A previous study has shown that combined cognitive and physical training, delivered either simultaneously or sequentially, is effective in promoting both cognitive and physical health in old age [2]. In addition, social isolation is a risk factor for poor cognitive function and dementia [3-7]. Participation in the class leads to the formation of a community. An increase in interpersonal communication as well as physical exercise and cognitive training may contribute synergistically to improving cognitive function. However, due to the COVID-19 pandemic, there were concerns about gathering of people causing infection transmission. The first case of COVID-19 in Japan was reported by the Ministry of Health, Labour and Welfare on January 16, 2020 [8], and COVID-19 has since spread nationwide. A state of emergency was announced by the Japanese government on April 7, 2020, and the government asked people to refrain from going out unless necessary. Houki Town considered measures to prevent the spread of COVID-19 before the

announcement of the state of emergency, and decided to suspend the class from early March 2020. There were concerns around the negative effects of the class suspension such as deterioration of physical and cognitive health due to lack of physical exercise or cognitive training during the class, decreased social contact due to the lack of places to gather, and reduced communication. However, older patients with COVID-19 are more likely to exhibit higher mortality rates and more severe symptoms [9-14]. Furthermore, given the social situation, suspension of the class was an unavoidable decision. Although classes resumed in June 2020, there were concerns around the progression of cognitive impairment that may have occurred during the suspension period. Therefore, when the classes resumed, participants were assessed by the Houki town hall staff using a cognitive function test and a questionnaire on lifestyle and thoughts during the suspension. They grasped the condition of the class participants and provided appropriate support.

There are ongoing concerns about further additional measures that will require restrictions due to COVID-19. It is also possible that the services provided by local governments will be suspended again in the future. Therefore, we thought it was important to determine what can be done to prevent cognitive decline. A previous study has shown that a multidomain lifestyle intervention was able to improve cognition in a group

experiencing cognitive decline [15]. Other studies reported on the relationships among cognitive function and exercise, cognitive training, and social contact [2-7]. In this study, by focusing on lifestyle and awareness of activities, the purpose was to investigate the changes in cognitive function during COVID-19 restrictions. We assessed cognitive function, lifestyle, and thoughts during the class suspension in community-dwelling older adults with mild cognitive decline who participated in a class for preventing cognitive decline. We also compared cognitive function before and after the class suspension to observe changes during the period of self-restraint.

Methods

Study design

This was a retrospective case-control study. Data were collected in September 2020 from information stored at Houki town hall. The ethics committee of Tottori University Faculty of Medicine (Yonago, Japan) approved the study (number: 20A076) and all procedures complied with the Declaration of Helsinki. The need for consent was waived by the ethics committee of Tottori University Faculty of Medicine because this was an

observational study that used pre-collected information. The study was not considered neither invasive nor intervening, and an opt-out approach was employed. All information pertaining to the study was disclosed in advance on the homepage of the Tottori University website and the notice board of Houki town hall to ensure that participants were given the opportunity to decline participation.

Study participants

The participants were community-dwelling older adults in Houki town (Tottori Prefecture, Japan), who were enrolled in classes for prevention of cognitive decline in 2019 and 2020. Participants in the class were evaluated with suspected cognitive decline previously using a cognitive function test on a touch-pad computer (Touch Panel-type Dementia Assessment Scale: TDAS, Nihon Kohden Corporation, Tokyo, Japan [16]). There are three types of classes: a weekly class (held for four months) for new people, and a once or twice a month follow-up class (held throughout the year) for people who have completed the weekly class. Participants are evaluated for cognitive function using the TDAS once a year between January and March. Generally, the classification of the two follow-up classes is based on the degree of cognitive impairment by evaluating the TDAS scores. People with

mild or greater cognitive dysfunction are invited to participate in the twice-a-month class, whereas those with a normal cognition range or extremely mild cognitive dysfunction are invited to participate in the once-a-month class. However, the classes were suspended from early March 2020 to end of May 2020 to prevent the spread of COVID-19. When the classes resumed in June 2020, participants were assessed by the Houki town hall staff using a cognitive function test and a questionnaire on lifestyle and thoughts during the suspension.

In this study, we collected data from participants who completed the TDAS evaluation between January 2020 to March 2020, and those who completed the TDAS evaluation and a questionnaire on lifestyle and thoughts during the class suspension from June 2020 to July 2020. The exclusion criteria were subjects who declined the use of their data.

Data collection

We collected data on the demographic characteristics, results of the cognitive function tests and the questionnaire on lifestyle and thoughts during the class suspension, and the type of class attended (once or twice a month follow-up class or a weekly class).

The TDAS [16] is a modified version of the ADAS-Cog [17], in which participants

enter their answers directly onto a touch-panel computer by following instructions. The nine examination items include “word recognition,” “following simple commands,” “visual-spatial perception,” “accuracy of the order of a process,” “naming fingers,” “orientation,” “money calculation,” “object recognition,” and “clock time recognition.” Scores ranged from 0 (all correct answers) to 101 points (all incorrect answers), so that higher scores indicate greater symptom severity.

The questionnaire was created independently by the Houki town hall staff and included questions about participants’ lifestyle and thoughts during the class suspension (S1 Table), such as anxiety (worrying about physical condition, getting sick, progression of forgetfulness, weakening of lower body, reduced food intake, decreased conversation, difficulty sleeping, and increased stress), awareness (exercising, working on cognitive training, having conversations, eating a well-balanced diet, living a well-regulated lifestyle, engaging in hobbies, and collecting information on dementia), working on cognitive training paper worksheets of the contents covered in the class provided by the town hall (sent by post once each in April and May 2020), opportunities for conversation, and method of conversation (directly, by phone, or using an online communication tool). Although other questions were present, the results of the items that were similar to the above questions

are not shown in this main text. The results are shown in S2 Table. Moreover, descriptive answers were also excluded from the analysis. The questionnaire was self-administered.

Statistical analysis

Statistical analyses were performed using SPSS (version 25, IBM Corporation, Tokyo, Japan) and EZR (version 1.41, Saitama Medical Center, Jichi Medical University, Saitama, Japan), which is a graphical user interface for R (The R Foundation for Statistical Computing, Vienna, Austria) [18]. The Shapiro-Wilk test was used to assess the normality of the data, and Levene's test was used to assess the equality of variance.

In our previous study [1] that evaluated the cognitive function of older adults with suspected cognitive decline using TDAS, we observed an average natural deterioration of 0.74 points (standard error 0.43) after a six month observational period. Based on this result, we set the criteria for cognitive decline as a 2-point or more decrease in TDAS score from the January to March 2020 assessment to the June to July 2020 assessment. The results of the questionnaire were compared by dividing participants into two groups: those whose TDAS score deteriorated (decline group) and those whose TDAS score did not deteriorate (non-decline group). Differences in demographic characteristics between groups

were assessed using Student's t-test or Fisher's exact test, as appropriate. Univariate and multivariate binary logistic regression analyses were used to assess associations between each of the questionnaire's results (the independent variables) and cognitive decline (the dependent variable). Because of fewer participants in the decline group, we needed to consider that the validity of the logistic model becomes problematic when the ratio of the numbers of events per variable analyzed is small [19]. Therefore, with reference to a previous study [20], for multivariable modeling, we used propensity score adjustment for age, sex, TDAS score before the class suspension, and type of class attended. The propensity score was estimated by a multivariate binary logistic regression analysis with the answers to each item in the questionnaire as the dependent variable, and the above adjustment factor as the independent variable. In the logistic regression analysis, unadjusted and adjusted odds ratios (OR) and 95% confidence intervals (CI) were calculated.

Change in the TDAS score between before and after the class suspension was evaluated using the Wilcoxon signed-rank test. We also carried out the above analyses based on the following stratifications: (1) level of cognitive function (based on January to March 2020 scores), which was based on previous studies [21, 22] that showed that the

average or median TDAS score for mild cognitive impairment (MCI) was around 7 points, those with suspected mild or greater cognitive decline had a TDAS score of 7 points or higher, and those with a normal cognition range or extremely mild cognitive decline had a TDAS score of 6 points or lower; (2) type of class attended (once or twice a month); (3) level of anxiety, calculated by the total number of positive responses to items relating to “anxiety during class suspension (question 1 in S1 Table)” in the questionnaire and grouped by the median; (4) level of awareness, calculated by the total number of positive responses to items relating to “awareness during classroom suspension (question 2 in S1 Table)” in the questionnaire and grouped by the median.

All statistical significance tests were two-sided and used an alpha level of 0.05 as statistically significant.

Results

Participant characteristics

We collected data from 88 participants. None of the individuals participated in the weekly class, which is only attended by new participants, and thus our subjects comprised those who participated in the follow-up classes that were held once or twice a month. All

participants attended the follow-up classes in both 2019 and 2020. The participants' characteristics are summarized in Table 1. There were 26 men and 62 women, and the mean age was 81.0 (standard deviation [SD] 6.5) years. The mean TDAS score before the class suspension was 7.1 (SD 6.0), and the average degree of cognitive decline was interpreted as mild. The average time intercurring between the first and second evaluation of TDAS was 139.8 (SD 11.2) days. More than approximately 80% of respondents answered that they worked on the worksheet about cognitive training that was provided by the town hall, and had conversations almost every day, including direct conversations. Less than 5% of respondents answered that they had opportunities for conversation less than once a week or had conversations using an online communication medium.

Table 1. Participants' characteristics

	All participants (n = 88)
Age	81.0 ± 6.5
Sex	

Male	26 (29.5)
Female	62 (70.5)
TDAS score before class suspension	7.1 ± 6.0
Anxiety during class suspension	
Worrying about physical condition	41 (46.6)
Getting sick	10 (11.4)
Progressing of forgetfulness	28 (31.8)
Weakening of lower body	37 (42.0)
Reduced food intake	15 (17.0)
Decreased conversation	44 (50.0)
Difficulty sleeping	13 (14.8)
Increased stress	18 (20.5)
Awareness during class suspension	
Exercising	53 (60.2)
Working on cognitive training	32 (36.4)
Having conversations	37 (42.0)

Eating a well-balanced diet	43 (48.9)
Living a well-regulated lifestyle	47 (53.4)
Engaging in hobbies	36 (40.9)
Collecting information on dementia	22 (25.0)
Worksheet provided by the town hall ^a	
Not working	17 (20.2)
Working	67 (79.8)
Opportunity for conversation ^b	
Rarely	1 (1.2)
Once a week	3 (3.6)
Once every 2–3 days	10 (12.0)
Almost every day	69 (83.1)
Conversation method	
Directly	72 (81.8)
Using a phone	44 (50.0)
Using an online tool	3 (3.4)

200 Data presented as mean $\pm$ standard deviation or number (%).

201 ^a There were 4 non-responders

202 ^b There were 5 non-responders

203 TDAS, Touch Panel-type Dementia Assessment Scale

204

205 **Questionnaire**

206 The results of the questionnaire are shown in Table 2 and S2 Table. In all, 16

207 participants (18.2%) had cognitive decline, and 72 (81.8%) had no cognitive decline. We

208 observed no significant differences in the age and sex ratio ($p = 0.475$, $p = 0.226$,

209 respectively). A logistic regression model showed that the ORs for cognitive decline were

210 lower in participants whose responses were “engaged in hobbies” (OR = 0.07, 95% CI =

211 0.01–0.60, $p = 0.015$), “worked on worksheet about cognitive training provided by the town

212 hall” (OR = 0.19, 95% CI = 0.04–0.82, $p = 0.026$), and “had conversations using a phone”

213 (OR = 0.28, 95% CI = 0.08–0.997, $p = 0.0495$).

214 **Table 2. Comparison of questionnaire results between the cognitive decline and non-decline group.**

	Decline group	Non-decline group	Unadjusted		Adjusted	
	(n = 16)	(n = 72)	OR (95% CI)	P value	OR (95% CI)	P value
Age	82.1 ± 7.1	80.8 ± 6.3				
Sex						
Male	7 (43.8)	19 (26.4)				
Female	9 (56.3)	53 (76.1)				
Anxiety during class suspension						
Worrying about physical condition	8 (50.0)	33 (45.8)	1.18 (0.40–3.49)	0.763	1.40 (0.45–4.36)	0.556
Getting sick	1 (6.3)	9 (12.5)	0.47 (0.05–3.97)	0.485	0.45 (0.05–3.88)	0.465
Progressing of forgetfulness	4 (25.0)	24 (33.3)	0.67 (0.19–2.29)	0.519	0.67 (0.19–2.32)	0.524
Weakening of lower body	9 (56.3)	28 (38.9)	2.02 (0.68–6.04)	0.208	2.12 (0.69–6.49)	0.188

Reduced food intake	4 (25.0)	11 (15.3)	1.85 (0.50–6.79)	0.355	1.78 (0.48–6.62)	0.386
Decreased conversation	9 (56.3)	35 (48.6)	1.36 (0.46–4.04)	0.581	1.51 (0.48–4.74)	0.477
Difficulty sleeping	2 (12.5)	11 (15.3)	0.79 (0.16–3.98)	0.777	0.81 (0.16–4.12)	0.798
Increased stress	5 (31.3)	13 (18.1)	2.06 (0.61–6.96)	0.243	2.10 (0.59–7.47)	0.250
Awareness during class suspension						
Exercising	8 (50.0)	45 (62.5)	0.60 (0.20–1.78)	0.358	0.62 (0.20–1.92)	0.407
Working on cognitive training	6 (37.5)	26 (22.2)	1.06 (0.35–3.26)	0.917	0.86 (0.26–2.84)	0.804
Having conversations	5 (31.3)	32 (44.4)	0.57 (0.18–1.80)	0.337	0.59 (0.18–1.91)	0.378
Eating a well-balanced diet	7 (43.8)	36 (50.0)	0.78 (0.26–2.31)	0.651	0.82 (0.26–2.60)	0.742
Living a well-regulated lifestyle	11 (68.8)	36 (50.0)	2.20 (0.69–6.97)	0.180	2.41 (0.74–7.84)	0.145
Engaging in hobbies	1 (6.3)	35 (48.6)	0.07 (0.01–0.56)	0.012	0.07 (0.01–0.60)	0.015
Collecting information on dementia	4 (25.0)	18 (25.0)	1.00 (0.29–3.49)	1.000	0.85 (0.23–3.09)	0.798

Worksheet provided by the town hall ^a						
Not working	6 (42.9)	11 (15.7)	1 (reference)		1 (reference)	
Working	8 (57.1)	59 (84.3)	0.25 (0.07–0.86)	0.028	0.19 (0.04–0.82)	0.026
Opportunity for conversation ^b						
Rarely	0 (0)	1 (1.4)				
Once a week	0 (0)	3 (4.3)	1 (reference) ^c		1 (reference) ^c	
Once every 2–3 days	1 (7.7)	9 (12.9)				
Almost every day	12 (92.3)	57 (81.4)	2.74 (0.33–23.0)	0.354	3.84 (0.41–35.80)	0.238
Conversation method						
Directly	12 (75.0)	60 (83.3)	0.60 (0.17–2.18)	0.438	0.64 (0.16–2.49)	0.514
Using a phone	4 (25.0)	40 (55.6)	0.27 (0.08–0.91)	0.034	0.28 (0.08–0.997)	0.0495
Using an online tool	0 (0)	3 (4.2)	— ^d		— ^d	

215 Data presented as mean $\pm$ standard deviation or number (%). Propensity score was used as an adjustment covariate.

216 ^a There were 2 non-responders in the decline group, and 2 non-responders in the non-decline group

217 ^b There were 3 non-responders in the decline group, and 2 non-responders in the non-decline group

218 ^c Reference was the total value of “almost none,” “once a week,” and “once every 2–3 days”

219 ^d Analysis was not possible because there were no participants in the decline group that had a conversation using an online

220 communication tool

221 OR, Odds ratio; CI, Confidence interval

Cognitive function test

Fig 1A-1E show the TDAS scores before and after the class suspension. In the analysis of all participants, a significant improvement in TDAS score was observed after the class resumed (Fig 1A, $p < 0.01$). When grouping according to the degree of cognitive function before the class suspension, a significant improvement in cognitive function was observed in the group with mild or greater cognitive decline (TDAS of 7 points or more before the class suspension) (Fig 1B, $p < 0.01$). In addition, when groups were classified according to the type of class attended, a significant improvement in cognitive function was observed in the group that attended classes twice a month (Fig 1C, $p < 0.01$). Furthermore, when grouped according to level of anxiety during class suspension (low anxiety: 0–2 positive responses; high anxiety: 3–8 positive responses), there was significant improvement in cognitive function in both groups (Fig 1D, both $p < 0.05$). When participants were grouped according to level of awareness (high awareness: 3–7 positive responses; low awareness: 0–2 positive responses), a significant improvement in cognitive function was observed in the high awareness group (Fig 1E, $p < 0.01$).

Fig 1. Results of TDAS before class suspension (Pre) and after class resumption (Post).

(A) The results of all participants are shown (n = 88). (B) The results of the mild or greater cognitive decline group (TDAS score of 7 points or more) before class suspension (n = 41 [13 men and 28 women], mean age = 82.3 [SD = 6.2] years), and the normal cognition or extremely mild cognitive decline group (TDAS score of 6 points or less) before class suspension (n = 47 [13 men and 34 women], mean age = 79.9 [SD = 6.5] years). (C) The results of subjects who participated in the class held twice a month (n = 64 [19 men and 45 women], mean age = 81.9 [SD = 6.6] years) and once a month (n = 24 [7 men and 17 women], mean age = 78.6 [SD = 5.5] years). (D) The results of low anxiety participants (n = 53 [16 men and 37 women], mean age = 80.8 [SD = 6.8] years) and high anxiety participants (n = 35 [10 men and 25 women], mean age = 81.3 [SD = 5.9] years). (E) The results of high awareness participants (n = 48 [10 men and 38 women], mean age = 81.2 [SD = 6.5] years) and low awareness participants (n = 40 [16 men and 24 women]), mean age = 80.7 [SD = 6.5] years).

All data indicate mean $\pm$ standard error, and the number on the error bar is the mean value.

*p<0.05, **p<0.01.

TDAS, Touch Panel-type Dementia Assessment Scale; SD, standard deviation

Discussion

In this study, we performed an analysis of lifestyle and awareness during the suspension of cognitive decline prevention classes due to COVID-19. We found that intellectual activities, such as “engaging in hobbies,” “working on the worksheet provided by the town hall,” and “having conversations on the phone” play an important role in preventing cognitive decline during a period of self-restraint. In addition, participants who had high levels of awareness during the class suspension showed significant improvement in TDAS score. Previous studies have reported that certain lifestyle activities may play an important role in the reversal of MCI to normal cognition [23]. Moreover, there is evidence of a significant association between high frequency of stimulating leisure activities in the past and reduced risk of dementia [24, 25]. Therefore, we suggest that the activities of daily life are important in preventing deterioration of cognitive function. In addition, previous studies have reported that having few social ties increased the risk for cognitive decline among elderly people [3, 26]; therefore, it is presumed that the phone was a useful communication medium in situations where the frequency of face-to-face meetings decreased due to the COVID-19 pandemic. Participants who had high awareness and

engaged in many activities showed improvement in cognitive function, which demonstrates that it is beneficial to continue any possible activities while refraining from activities due to COVID-19.

Previous studies have shown that in older adults, the spread of COVID-19 may lead to decreased social [27] and physical activity time [28], and an increasing risk of problematic mental health symptoms such as depression and anxiety due to more days spent staying at home [29]. Physical inactivity, social isolation, and depression in later life (after 65 years) are risk factors for dementia [7]. Moreover, in a study investigating the effects of COVID-19 lockdown in Italy, telephone interviews with caregivers of 31 subjects with MCI revealed a worsening of cognition in 41.9% of people [30]. Although Japan did not impose any lockdown, there was concern that class suspension and staying at home would lead to cognitive decline. However, although 31.8% of the participants answered in the questionnaire that they were worried about the progress of forgetfulness, 18.2% of the participants in our study had cognitive decline. Moreover, cognitive function improved after the classes resumed. Anxiety symptoms have also been reported to increase the risk of developing MCI and dementia [31]; however, cognitive function improved after class resumed, regardless of the levels of anxiety. In addition, subjects who participated in the

290 class held twice a month and subjects with mild or more severe cognitive decline were
 291 concerned about their decline in cognitive function due to class suspension. However, no
 292 adverse effect on cognitive function was observed because of changes in the environment
 293 due to COVID-19. One possible reason can be that the extent of the measures taken to
 294 counteract the COVID-19 pandemic in Japan was on a voluntary basis and presumably
 295 irrelevant in determining a significant detrimental effect on the cognitive functions of
 296 Japanese elderly with mild cognitive decline. Another reason can be that continued
 297 attendance of the classes until the suspension may explain the limited cognitive decline
 298 observed in our study. Although cultural differences and difference in survey years must be
 299 taken into consideration, it has been proposed that both in Japan and other countries, older
 300 adults' knowledge of dementia and cognitive disorders is currently deficient and that public
 301 health approaches such as educational interventions or national campaigns are necessary
 302 [32-35]. Although we did not investigate the degree of knowledge in our participants, we
 303 can infer that the participants in this study had some understanding of the type of lifestyle
 304 that is important to prevent cognitive decline because they had already been participating
 305 in the classes. Therefore, we suspect that many participants carried out beneficial actions
 306 to prevent the deterioration of cognitive function during the period of self-restraint imposed

307 due to COVID-19.

308 We found that few people had conversations using an online communication
 309 medium. According to the results of the 2019 Communications Usage Trend Survey in
 310 Japan released by the Ministry of Internal Affairs and Communications, the usage status of
 311 social networking services was reported to be 40.7% for people aged 70–79 and 42.8% for
 312 people aged 80 and over [36]. In a study examining the compliance rate of computer-based
 313 cognitive training in older adults with increased risk of dementia, 37% did no train at all,
 314 and previous experience with computers, being married or cohabiting, better memory
 315 performance, and positive expectations toward the study predicted greater likelihood for
 316 starting the training [37]. In addition, a study investigating the frequency of exposure to
 317 health information from information and communication technologies reported that older
 318 adult respondents were less frequently exposed to health information from websites and
 319 social media sites [38]. Taken together, we consider it important to provide support for older
 320 adults who are suspected of having cognitive decline by using more traditional methods,
 321 such as telephone and print media, rather than digital mediums, especially during
 322 emergency situations. Furthermore, 79.8% of participants answered that during the class
 323 suspension, they worked on the cognitive training worksheet provided by the town hall.

Although individual awareness was an important factor that influenced whether an individual worked on the worksheet provided by the town hall, the results suggested that it is important to provide information in an appropriate manner such as through paper.

Finally, this study has several limitations. First, the subjects of this study were community-dwelling older adults, who participated in a class for the prevention of cognitive decline. We did not examine older adults who do not participate in such classes. Therefore, we may not be able to generalize these results to those older adults who experienced restrictions due to COVID-19 but do not participate in such programs. The second limitation is that we cannot control for the carryover effect of attending the classes. In a previous study, it was suggested that the effect of the cognitive decline prevention program used in the class may not be sustained for up to 6 months after the intervention is stopped [1]. However, in this study, the class was only suspended for around three months. Therefore, although we observed improvements in cognitive function after the class suspension, it is possible that participation in the class before the suspension may have had some influence. The third limitation is the small sample size. Specifically, the number of participants with cognitive decline was small ($n = 16$). The fourth research limitation is that we did not consider all the possible confounding factors that may affect cognitive

function. For example, the status of infection of COVID-19 in participants may also have an effect on cognitive performance [39, 40] and daily activities. As the data available were limited due to retrospective study, we could only adjust for a few factors. However, since the number of patients with COVID-19 in Tottori Prefecture in Japan was three from March to May 2020 and the classes resumed without any problems, it is estimated that no participants were infected with COVID-19 during the class suspension. The fifth limitation is that the subjects of this study do not include those who did not come to the class when it resumed. The target sample size was set at about 120 participants at the research planning stage, which was the total number of class participants in 2019 and 2020. Participation in the class is voluntary, so there may be various reasons for absence (e.g., having another plan, concerned about COVID-19 infection, etc.).

Conclusions

A proactive approach toward intellectual activities and social ties may be important in preventing cognitive decline during periods of self-restraint due to COVID-19. We observed significant improvement in cognitive function scores following the resumption of the class. We speculate that participants' understanding of the importance of an active

lifestyle (reflected in their continued attendance of the classes) may explain these improvements. We propose that the class is important not only to prevent cognitive decline, but also to prepare for cognitive decline in situations of restrictive living, such as during the COVID-19 pandemic.

Acknowledgments

We thank the staff of the Health measures section, Houki town who cooperated with us during this study. In addition, we would like to thank Editage (www.editage.com) for English language editing.

Author contributions

Conceptualization: Minoru Kouzuki, Katsuya Urakami.

Data curation: Minoru Kouzuki.

Formal analysis: Minoru Kouzuki, Shota Furukawa, Keisuke Mitani.

Supervision: Katsuya Urakami.

Writing – original draft: Minoru Kouzuki.

Writing – review & editing: Minoru Kouzuki, Katsuya Urakami.

375

376 **Data Availability Statement:**

377 The data sets used and/or analyzed during the current study are available from the
378 corresponding author on request. However, access to the data sets will only be granted upon
379 approval by all authors, the data provider (the Health measures section, Houki town), and
380 the ethics committee.

381

382

References

1. Kouzuki M, Kato T, Wada-Isoe K, Takeda S, Tamura A, Takanashi Y, et al. A program of exercise, brain training, and lecture to prevent cognitive decline. *Ann Clin Transl Neurol.* 2020; 7(3): 318-328.
2. Gavelin HM, Dong C, Minkov R, Bahar-Fuchs A, Ellis KA, Lautenschlager NT, et al. Combined physical and cognitive training for older adults with and without cognitive impairment: A systematic review and network meta-analysis of randomized controlled trials. *Ageing Res Rev.* 2021; 66: 101232.
3. Evans IEM, Llewellyn DJ, Matthews FE, Woods RT, Brayne C, Clare L; CFAS-Wales research team. Social isolation, cognitive reserve, and cognition in healthy older people. *PLoS One.* 2018; 13(8): e0201008.
4. Saito T, Murata C, Saito M, Takeda T, Kondo K. Influence of social relationship domains and their combinations on incident dementia: a prospective cohort study. *J Epidemiol Community Health* 2018; 72: 7-12.
5. Kuiper JS, Zuidersma M, Oude Voshaar RC, Zuidema SU, van den Heuvel ER, Stolk RP, et al. Social relationships and risk of dementia: A systematic review and meta-analysis of longitudinal cohort studies. *Ageing Res Rev.* 2015; 22: 39-57.

- 400 6. Evans IEM, Martyr A, Collins R, Brayne C, Clare L. Social Isolation and Cognitive
401 Function in Later Life: A Systematic Review and Meta-Analysis. J Alzheimers Dis.
402 2019; 70(s1): S119-144.
- 403 7. Livingston G, Huntley J, Sommerlad A, Ames D, Ballard C, Banerjee S, et al. Dementia
404 prevention, intervention, and care: 2020 report of the Lancet Commission. Lancet.
405 2020; 396(10248): 413-446.
- 406 8. Ministry of Health, Labour and Welfare. Press releases about a patient diagnosed with
407 pneumonia linked to a new coronavirus(1st case). (in Japanese). 2020 January 16
408 [Cited 2021 February 21]. Available from:
409 https://www.mhlw.go.jp/stf/newpage_08906.html
- 410 9. Chinnadurai R, Ogedengbe O, Agarwal P, Money-Coomes S, Abdurrahman AZ,
411 Mohammed S, et al. Older age and frailty are the chief predictors of mortality in
412 COVID-19 patients admitted to an acute medical unit in a secondary care setting: a
413 cohort study. BMC Geriatr. 2020; 20(1): 409.
- 414 10. Wei C, Liu Y, Liu Y, Zhang K, Su D, Zhong M, et al. Clinical characteristics and
415 manifestations in older patients with COVID-19. BMC Geriatr. 2020; 20(1): 395.

- 416 11. Lian J, Jin X, Hao S, Cai H, Zhang S, Zheng L, et al. Analysis of Epidemiological and
417 Clinical Features in Older Patients With Coronavirus Disease 2019 (COVID-19)
418 Outside Wuhan. Clin Infect Dis. 2020; 71(15): 740-747.
- 419 12. Liu K, Chen Y, Lin R, Han K. Clinical features of COVID-19 in elderly patients: A
420 comparison with young and middle-aged patients. J Infect. 2020; 80(6): e14-18.
- 421 13. Giorgi Rossi P, Marino M, Formisano D, Venturelli F, Vicentini M, Grilli R, et al.
422 Characteristics and outcomes of a cohort of COVID-19 patients in the Province of
423 Reggio Emilia, Italy. PLoS One. 2020; 15(8): e0238281.
- 424 14. Izcovich A, Ragusa MA, Tortosa F, Lavena Marzio MA, Agnoletti C, Bengolea A, et al.
425 Prognostic factors for severity and mortality in patients infected with COVID-19: A
426 systematic review. PLoS One. 2020; 15(11): e0241955.
- 427 15. McMaster M, Kim S, Clare L, Torres SJ, Cherbuin N, D'Este C, et al. Lifestyle Risk
428 Factors and Cognitive Outcomes from the Multidomain Dementia Risk Reduction
429 Randomized Controlled Trial, Body Brain Life for Cognitive Decline (BBL-CD). J Am
430 Geriatr Soc. 2020; 68(11): 2629-2637.

431 16. Inoue M, Jimbo D, Taniguchi M, Urakami K. Touch Panel-type Dementia Assessment
432 Scale: a new computer-based rating scale for Alzheimer's disease. Psychogeriatrics
433 2011; 11: 28-33.

434 17. Rosen WG, Mohs RC, Davis KL. A new rating scale for Alzheimer's disease. Am J
435 Psychiatry 1984; 141: 1356-1364.

436 18. Kanda Y. Investigation of the freely available easy-to-use software 'EZR' for medical
437 statistics. Bone Marrow Transplant 2013; 48(3): 452-458.

438 19. Peduzzi P, Concato J, Kemper E, Holford TR, Feinstein AR. A simulation study of the
439 number of events per variable in logistic regression analysis. J Clin Epidemiol. 1996;
440 49(12): 1373-1379.

441 20. Devasia RA, Blackman A, Gebretsadik T, Griffin M, Shintani A, May C, et al.
442 Fluoroquinolone resistance in Mycobacterium tuberculosis: the effect of duration and
443 timing of fluoroquinolone exposure. Am J Respir Crit Care Med. 2009; 180(4): 365-370.

444 21. Kouzuki M, Ichikawa J, Shirasagi D, Katsube F, Kobashi Y, Matsumoto H, et al.
445 Detection and recognition thresholds for five basic tastes in patients with mild cognitive
446 impairment and Alzheimer's disease dementia. BMC Neurol. 2020; 20(1): 110.

447 22. Kouzuki M, Suzuki T, Nagano M, Nakamura S, Katsumata Y, Takamura A, et al.
448 Comparison of olfactory and gustatory disorders in Alzheimer's disease. *Neurol Sci.*
449 2018; 39(2): 321-328.

450 23. Shimada H, Doi T, Lee S, Makizako H. Reversible predictors of reversion from mild
451 cognitive impairment to normal cognition: a 4-year longitudinal study. *Alzheimers Res*
452 *Ther.* 2019; 11(1): 24.

453 24. Verghese J, Lipton RB, Katz MJ, Hall CB, Derby CA, Kuslansky G, et al. Leisure
454 activities and the risk of dementia in the elderly. *N Engl J Med* 2003; 348: 2508-2516.

455 25. Akbaraly TN, Portet F, Fustini S, Dartigues JF, Artero S, Rouaud O, et al. Leisure
456 activities and the risk of dementia in the elderly: results from the Three-City Study.
457 *Neurology* 2009; 73: 854-861.

458 26. Bassuk SS, Glass TA, Berkman LF. Social disengagement and incident cognitive decline
459 in community-dwelling elderly persons. *Ann Intern Med.* 1999; 131(3): 165-173.

460 27. Yamada Y, Uchida T, Ogino M, Ikenoue T, Shiose T, Fukuma S. Changes in Older
461 People's Activities During the Coronavirus Disease 2019 Pandemic in Japan. *J Am Med*
462 *Dir Assoc.* 2020; 21(10): 1387-1388. e1.

463 28. Yamada M, Kimura Y, Ishiyama D, Ootobe Y, Suzuki M, Koyama S, et al. Effect of the
464 COVID-19 Epidemic on Physical Activity in Community-Dwelling Older Adults in
465 Japan: A Cross-Sectional Online Survey. *J Nutr Health Aging*. 2020; 24(9): 948-950.

466 29. Zhao SZ, Wong JYH, Wu Y, Choi EPH, Wang MP, Lam TH. Social Distancing
467 Compliance under COVID-19 Pandemic and Mental Health Impacts: A
468 Population-Based Study. *Int J Environ Res Public Health*. 2020; 17(18): 6692.

469 30. Baschi R, Luca A, Nicoletti A, Caccamo M, Cicero CE, D'Agate C, et al. Changes in
470 Motor, Cognitive, and Behavioral Symptoms in Parkinson's Disease and Mild Cognitive
471 Impairment During the COVID-19 Lockdown. *Front Psychiatry*. 2020; 11: 590134.

472 31. Liew TM. Subjective cognitive decline, anxiety symptoms, and the risk of mild cognitive
473 impairment and dementia. *Alzheimers Res Ther*. 2020; 12(1): 107.

474 32. Sun F, Gao X, Shen H, Burnette D. Levels and correlates of knowledge about
475 Alzheimer's disease among older Chinese Americans. *J Cross Cult Gerontol*. 2014;
476 29(2): 173-183.

477 33. Glynn RW, Shelley E, Lawlor BA. Public knowledge and understanding of
478 dementia: evidence from a national survey in Ireland. *Age Ageing*. 2017; 46(5): 865-869.

- 479 34. Rosenberg A, Coley N, Soulier A, Kulmala J, Soininen H, Andrieu S, et al. Experiences
480 of dementia and attitude towards prevention: a qualitative study among older adults
481 participating in a prevention trial. BMC Geriatr. 2020; 20(1): 99.
- 482 35. Arai Y, Arai A, Zarit SH. What do we know about dementia?: a survey on knowledge
483 about dementia in the general public of Japan. Int J Geriatr Psychiatry. 2008; 23(4):
484 433-438.
- 485 36. Ministry of Internal Affairs and Communications, JAPAN. Results of FY2019
486 Communication Usage Trend Survey (in Japanese). 2020 May 29 [Cited 2021 February
487 21]. Available from:
488 https://www.soumu.go.jp/johotsusintokei/statistics/data/200529_1.pdf
- 489 37. Turunen M, Hokkanen L, Bäckman L, Stigsdotter-Neely A, Hänninen T, Paaianen T, et
490 al. Computer-based cognitive training for older adults: Determinants of adherence.
491 PLoS One. 2019; 14(7): e0219541.
- 492 38. Shen C, Wang MP, Wan A, Viswanath K, Chan SSC, Lam TH. Health information
493 exposure from information and communication technologies and its associations with
494 health behaviors: Population-based survey. Prev Med. 2018; 113: 140-146.

39. Miners S, Kehoe PG, Love S. Cognitive impact of COVID-19: looking beyond the short term. *Alzheimers Res Ther.* 2020; 12(1): 170.
40. Del Brutto OH, Wu S, Mera RM, Costa AF, Recalde BY, Issa NP. Cognitive decline among individuals with history of mild symptomatic SARS-CoV-2 infection. A longitudinal prospective study nested to a population cohort. *Eur J Neurol.* 2021 Feb 11. doi: 10.1111/ene.14775. Epub ahead of print. PMID: 33576150.

Supporting information

- S1 Table. English translation of the relevant questions used in this study.**
- S2 Table. Comparison of questionnaire results between the cognitive decline and non-decline group (results not shown the main text).**