Trends in homelessness and injection practices among young urban and suburban people who inject drugs: 1997-2017

Anna L. Hotton^{1*}, Mary Ellen Mackesy-Amiti², Basmattee Boodram^{2F}

¹ Department of Medicine, University of Chicago, Chicago, Illinois, USA

² Division of Community Health Sciences, School of Public Health, University of Illinois at Chicago, Chicago, Illinois, USA

*Corresponding author

Email: ahotton@medicine.bsd.uchicago.edu

Preprint, v.1, February 16, 2021

Abstract

Background. Among young people who inject drugs (PWID) homelessness is associated with numerous adverse psychosocial and health consequences, including risk of relapse and overdose, psychological distress and suicidality, limited treatment access, and injection practices that increase the risk of HIV and hepatitis C (HCV) transmission. Homeless PWID may also be less likely to access sterile syringes through pharmacies or syringe service programs.

Methods. This study applied random-effects meta-regression to examine trends over time in injection risk behaviors and homelessness among young PWID in Chicago and surrounding suburban and rural areas using data from 11 studies collected between 1997 and 2017. In addition, subject-level data were pooled to evaluate the effect of homelessness on risk behaviors across all studies using mixed effects logistic and negative binomial regression with random study effects. **Results.** There was a significant increase in homelessness among young PWID over time, consistent with the general population trend of increasing youth homelessness. In mixed-effects regression, homelessness was associated with injection risk behaviors (receptive syringe sharing, syringe mediated sharing, equipment sharing) and exchange sex, though we detected no overall changes in risk behavior over time.

Conclusions. Increases over time in homelessness among young PWID highlight a need for research to understand factors contributing to youth homelessness to inform HIV/STI, HCV, and overdose prevention and intervention services for this population.

Keywords: people who inject drugs; homelessness; meta-regression; injection risk behavior

Background

Homelessness is prevalent among young people who inject drugs (PWID) and has consistently been associated with injection risk behaviors [1-5] and risk for HIV and hepatitis C (HCV) transmission [6-8]. Youth homelessness is also associated with numerous other social adversities and negative health outcomes, including high rates of depression and suicidality, high likelihood of alcohol and substance use problems, poor academic outcomes, sexual risk taking and high rates of sexually transmitted infections (STI), and past and concurrent victimization [9-13]. For young PWID, homelessness is associated with a myriad of adverse effects, including difficulty accessing treatment for substance use [14], risk of relapse [1, 15], blood-borne infections due to syringe and equipment sharing in unsanitary conditions [1, 15], increased risk of overdose [16, 17], vulnerability to violence and sexual exploitation [18, 19], psychological distress and suicidal behavior [20, 21], and general increased morbidity and mortality [22-25]. Homeless PWID may also be less likely to benefit from increased access to sterile syringes at pharmacies in Illinois [26-28] that occurred over the past two decades, either due to lack of resources for purchasing syringes or fear of pharmacy staff or police harassment. PWID from suburban and rural communities who experience homelessness may be at particularly high risk of infectious disease transmission due to less availability of syringe service programs (SSPs) and other harm reduction services compared to urban areas.

Recent data suggest that homelessness has been increasing in the US over the past two decades as a result of economic recessions and rising housing costs [29]. While standard surveys of homelessness do not estimate PWID as a subpopulation, it is possible that increases in homelessness have occurred among PWID as well. Increases in heroin injection among young people as a result of the opioid epidemic may also have resulted in increases in the population of young homeless PWID.

The goal of this study was to examine trends over time in injection risk behaviors and homelessness among young PWID in the Chicago, Illinois and surrounding suburban and rural areas, using data from several federally-funded research studies conducted from 1997 through 2017. Because time was confounded with study, we examined time trends using meta-regression on summary measures. In follow-up analyses, we examined associations between homelessness and risk behaviors across all studies.

Materials and Methods

Data sources. De-identified data were obtained from the principal investigators of eleven research studies that collected detailed data on PWID demographics and injection related behaviors over the past twenty years. These included National HIV Behavioral Surveillance (NHBS) PWID cycles 2-4 conducted by the Chicago Department of Public Health (CDPH), the Collaborative Injection Drug User Studies (CIDUS 2 & 3), and local studies conducted by UIC-COIP investigators shown in Table 1. Detailed descriptions of each of the studies in terms of participants, recruitment strategies, and assessment methods have been previously described in the citations listed in Table 1. Briefly, recruitment approaches included either respondent driven sampling (RDS) or convenience sampling via flyers and advertisements posted at the recruitment sites and/or active recruitment by outreach workers. Our analysis was restricted to participants who reported injection drug use. With the exception of NIHU, which focused exclusively on heroin users, eligibility across studies included injection of any substances, though the most commonly reported drug injected was heroin alone or in combination with other drugs (e.g., cocaine, methamphetamine). Harm reduction and drug treatment resources were made available to all participants as part of the study protocols, and many studies involved recruitment directly from SSP sites. CDPH and the principal investigators disclaim responsibility for any analysis, interpretations, or conclusion of the study. This analysis was restricted to young PWID (18-34) because seven of the studies restricted their samples to this age range.

Table 1. Data sets included in the metaregression

Study	Years	N [ages 18- 35]	Measurement time frame			RDS recruit ment	Percent residence in Chicago
			Homeless	IDU network	Risk behavior		
CIDUS 2 [30]	1997-1999	695	6 months	NA	6 months	No	0.61
CIDUS 3 [31]	2002-2004	775	6 months	3 months	3 months	No	0.32
NATHCV [32]	2002-2005	110	6 months	NA	6 months	No	NA
NIHU [33]	2004-2006	150	NA	30 days	6 months	No	0.56
SATHCAP [34]	2005-2009	453	12 months	6 months	30 days	Yes	0.62
PSYCH [35]	2008-2010	610	6 months	6 months	6 months	Yes ^a	0.28
NHBS IDU2 [36]	2009	143	12 months	30 days	12 months	Yes	0.74
NHBS IDU3 [37]	2012	59	12 months	30 days	12 months	Yes	0.52
SOCNET [38]	2012-2013	163	6 months	6 months	6 months	No	0.80
NHBS IDU4 [39]	2015	263	12 months	30 days	12 months	Yes	0.54
MOOD [4]	2016-2017	180	6 months	6 months	6 months	No	0.61

^a Partial RDS recruitment; 40% of sample was recruited by outreach

RDS: Respondent driven sampling

CIDUS: Collaborative Injection Drug Users Study (CDC)

NATHCV: Early natural history of HCV infection among injection drug users (NIDA)

NIHU: Non-injected heroin use, HIV, and transitions to injection (NIDA)

SATHCAP: Sexual Acquisition and Transmission of HIV Cooperative Agreement Project (NIDA)

PSYCH: Psychiatric disorders and HIV risk behaviors among young injection drug users (NIDA)

SOCNET: Social networks of young injection drug users (Chicago DCFAR)

MOOD: Emotion dysregulation and risky behavior among people who inject drugs (NIDA)

NHBS: National HIV Behavioral Surveillance (CDC)

Measures.

Demographic variables included sex, race/ethnicity (non-Hispanic White, non-Hispanic Black, Hispanic, and Other, including multiracial categories), and age (categorized as 18-24 vs. 25-34). Subjects identifying as transgender were excluded from the analysis due to insufficient analytic sample size.

Harmonization of outcome measures. Data harmonization involved recoding, combining, and collapsing variables as necessary such that constructs were defined and expressed in equivalent scales/units across studies. We did not make any adjustments to account for variation in the time frame for outcome measurement (e.g., past 6 months, past 12 months) across studies (Table 1). Measures from longitudinal studies were summarized across time within individuals such that individuals contributed one observation per study. We created dichotomous indicators for risk behaviors, and converted Likert scale responses on frequency measures to proportions (Never = 0, Rarely = .10, Less than half the time = .25, Half the time = .50, More than half the time = .75, Almost always = .75, Always = 1).

Homelessness was measured as self-perceived homelessness (e.g. have you been homeless, have you considered yourself homeless) within the past 6 or 12 months.

Injection networks. Two measures of injection network size were available: 1) total injection network: the number of PWID that the respondent knows and has seen in past 30 days or 6 months (6 studies); and 2) injection sub-network: the number of people the respondent injected drugs with in the past 30 days, 3 months, or 6 months (6 studies). Three studies included both of these measures.

Injection-related risk behaviors. Any syringe-mediated drug sharing (SMS) in the past 3, 6, or 12 months was defined as having shot up with a needle after someone else squirted drugs into it from another syringe (i.e., “backloading”), or drugs were mixed, measured, or divided using someone else’s syringe. Receptive syringe sharing (RSS) was defined as injecting in the past 30 days to 12 months with a syringe that someone else had previously used to inject.

Frequency of RSS was estimated as the percent of injections that involved RSS based on either Likert scale responses or the number of RSS injections and frequency of injection. For respondents, we also estimated receptive sharing network size (i.e., the number of people respondent used a syringe after) and distributive sharing network size (i.e., the number of people the respondent gave his/her used syringe to). Equipment sharing was defined as any sharing of cookers, cotton, or rinse water for drug injection in the past 30 days to 12 months. *Exchange sex* was defined as having bought or sold sex for drugs, money, shelter, or other goods in the past 3, 6, or 12 months.

Statistical analysis.

Using the harmonized datasets, frequencies of categorical variables and means and standard errors of continuous variables were computed. Standard errors were calculated based on the Poisson distribution for count variables and the binomial distribution for binary variables. Pooled estimates of means and proportions were computed using the *metan* command in Stata 15.1 with random effects based on the DerSimonian & Laird method and heterogeneity estimated via the Mantel-Haenszel method [40].

For the analysis of time trends, it was necessary to employ meta-regression of summary measures rather than analyzing individual level data due to the association between study and time. Meta-regression is an extension of meta-analysis that can be used to examine whether between-study heterogeneity is related to aggregate level study characteristics (e.g., sociodemographic composition, behavior prevalence, network characteristics) [41]. Using the harmonized datasets, we computed aggregate level summary statistics (means and standard errors) by study for sociodemographic variables, injection behaviors, and network variables. We conducted random-effects meta-regression using aggregate means and standard errors for outcomes of interest. Each study contributes one observation to the analysis, with the number of observations ranging from 5-11 depending on the availability of measures across studies. To examine annual time trends, separate models were fit for each outcome with year as the only

independent variable using the *metareg* command in Stata. For ease of interpretation, time was scaled in 10-year intervals so that coefficients represent the average change in the outcome per 10 year period. These analyses were also repeated, stratified by past 6-month homelessness, to determine if trends over time varied according to housing status. Standard errors and confidence intervals for meta-regression coefficients were estimated using the Knapp-Hartung variance estimator [42].

Subject-level data were then pooled to evaluate the effect of homelessness on risk behaviors across all studies. Mixed effects regression models with random study effects were applied, with logistic regression for binary outcomes and negative binomial regression for count outcomes. Analyses were conducted using Stata/SE version 15.1 [43].

Results

Study characteristics are shown in Table 2. Across studies, participants were mostly non-Hispanic White and male. Approximately one-third were less than 25 years of age and over half resided in Chicago. Homelessness was common (41%) as were injection related risk behaviors with average pooled prevalence of receptive syringe sharing, syringe-mediated drug sharing, and equipment sharing of 38%, 26%, and 65% respectively. Average pooled prevalence of exchange sex was 17%.

In the meta-regression analysis of time trends, we found no significant effects on injection sub-network size (number of people injected with) or any of the injection-related risk behaviors (Table 3). Homelessness among young PWID increased significantly over time (Fig 1, Table 3), with time explaining over 60% of between-study variance in homelessness. Total injection network size (number of people you know who inject drugs) also increased over time; however the effect was not statistically significant. In analyses stratified by homelessness, there were no statistically significant time trends among PWID experiencing homelessness. Among PWID not experiencing homelessness there was a non-statistically significant decline over time

in RSS (Coef = -0.10, 95% CI -0.20, 0.004; $p=0.057$), while among PWID who reported homelessness there was no change in RSS over time (Coef = -0.05, 95% CI -0.18, 0.08; $p=0.404$). Among PWID not experiencing homeless, the marginal predicted proportion of PWID engaging in RSS was 0.43 in 1997 (95% CI 0.32-0.54) and 0.23 in 2016 (95% CI 0.14-0.33), while among those experiencing homelessness the marginal predicted proportions were 0.53 in 1997 (95% CI 0.39-0.67) and 0.44 in 2016 (95% CI 0.33-0.54).

Fig 1: Mean proportion of PWID reporting homelessness by study and year

Table 2. Pooled estimates of young PWID characteristics: 1997-2017

	N studies	Pooled mean (95% CI)	Range
Percent white	11	0.64 (0.58, 0.71)	0.41 - 0.78
Percent Black	11	0.07 (0.05, 0.10)	0.02 - 0.30
Percent Hispanic/Latino	11	0.23 (0.19, 0.27)	0.13 - 0.36
Percent young (<25)	10*	0.36 (0.22, 0.50)	0.11 - 0.63
Percent female	11	0.34 (0.32, 0.36)	0.27 - 0.42
Percent urban Chicago	10	0.56 (0.45, 0.67)	0.28 - 0.80
Percent homeless	10	0.41 (0.31, 0.51)	0.15 - 0.68
Percent receptive syringe sharing	11	0.38 (0.33, 0.43)	0.27 - 0.49
Percent syringe mediated sharing	10	0.26 (0.23, 0.30)	0.17 - 0.38
Percent equipment sharing	9	0.65 (0.59, 0.71)	0.51 - 0.77
Percent exchange sex	10	0.17 (0.14, 0.19)	0.10 - 0.31
Mean # partners receptive syringe sharing	7	0.73 (0.59, 0.88)	0.52 - 1.29
Mean # partners distributive syringe sharing	5	1.02 (0.75, 1.30)	0.69 - 1.95
Mean # partners sharing equipment	6	1.89 (1.36, 2.43)	1.04 - 2.87
Mean total injection network size	6	13.75 (9.07, 18.44)	5.08 - 24.48
Mean injection sub-network size	5	4.31 (3.25, 5.36)	2.82 - 5.46

*PSYCH study excluded: all under 25

We conducted sensitivity analyses on homelessness to assess whether increases in homelessness over time could be partly due to expansion of the reference timeframe (i.e. from 6

months to 12 months) by 1) controlling for the timeframe of assessment of homelessness and 2) restricting the analysis to studies that used past 6 months as the timeframe (n=6 studies). Findings were similar in all scenarios. With no adjustment (original): Coef = 0.25, 95% CI 0.10, 0.40; p=0.005; with adjustment for timeframe of 6 vs. 12 months: Coef=0.22, 95% CI 0.07-0.37; p=0.012); with restriction to studies with a timeframe of 6 months: Coef=0.21, 95% CI -0.03-0.46; p=0.073.

Table 3: Effects of time on risk behavior and homelessness in meta-regression analysis

Outcome	N studies	Coefficient ^a	(95% CI)	p-value	Adjusted R-squared ^b
Percent any receptive syringe sharing	11	-0.03	(-0.14, 0.07)	0.471	-3.84%
Percent any syringe mediated sharing	10	0.07	(-0.02, 0.16)	0.095	25.77%
Percent any equipment sharing	9	-0.01	(-0.14, 0.11)	0.825	-15.43%
Percent any exchange sex	10	0.04	(-0.02, 0.10)	0.137	12.60%
Percent homeless	10	0.25	(0.10, 0.40)	0.005	61.60%
Mean # partners receptive syringe sharing	7	0.20	(-0.42, 0.82)	0.453	-6.57%
Mean # partners distributive syringe sharing	5	0.56	(-0.75, 1.87)	0.266	18.34%
Mean # partners sharing equipment	6	1.18	(-0.44, 2.80)	0.113	40.98%
Mean total injection network size	6	12.6	(-13.3, 38.6)	0.248	14.13%
Mean injection sub-network size	5	1.36	(-1.37, 4.09)	0.212	27.81%

^a change in outcome per 10-year interval

^b Adjusted R² is the relative reduction in between-study variance explained by adding independent variables (in this case, time) to the model. It is calculated by comparing tau² from a model with no variables to tau² when variables are added. A negative value indicates that the variables explain less heterogeneity than would be expected by chance.

In subject-level mixed-effects regression of pooled individual-level data with random study intercepts, adjusted for age, race and sex, homelessness was associated with RSS, SMS, equipment sharing, and exchange sex, in addition to total injection network and injection sub-network size and total number of sex partners (Table 4).

Table 4: Associations of homelessness with risk behaviors estimated in multivariable mixed effects regression

	Total obs	n groups	aOR or aRR ^a	(95% CI)	p-value
<i>Binary outcomes</i>					
Receptive syringe sharing	3429	10	1.93	(1.67-2.23)	<0.001
Syringe mediated sharing	2699	9	2.18	(1.81-2.63)	<0.001
Equipment sharing	3326	9	1.71	(1.43-2.05)	<0.001
Exchange sex	3321	9	2.45	(2.17-2.76)	<0.001
<i>Count outcomes</i>					
Total RSS partners	2459	7	2.15	(1.48-3.13)	<0.001
Total DSS partners	2261	5	2.24	(1.60-3.14)	<0.001
Total injection network size	1665	6	1.55	(1.38-1.73)	<0.001
Injection sub-network size	2144	5	1.48	(1.25-1.76)	<0.001
Total sex partners	3288	9	1.41	(1.18-1.69)	<0.001
Total exchange sex partners	1609	5	3.32	(2.57-4.28)	<0.001

a Adjusted for age, race, and sex. Estimates generated from mixed effects logistic or negative binomial regression with random study intercepts.

Discussion

This analysis showed a significant increase in homelessness among young PWID over time, consistent with the general population trend of increasing youth homelessness[29]. Although homelessness was associated with risk behavior across time, we detected no overall change in the level of risk behavior over time. A partial explanation may lie in the expansion of interventions that increased access to sterile syringes and harm reduction education in Chicago and surrounding areas, which may have led to decreased risk behavior, thereby offsetting the increased homelessness among PWID. Stratified samples by homelessness revealed small decreases in RSS over time, but only among young PWID not experiencing homelessness. Although the observed effects were not statistically significant, it suggests that gains in harm reduction may have been countered by worsening social conditions over time. Another potential driving factor may be increasing proportions of suburban PWID over time, who report having less access to SSPs. Increasing homelessness may be explained partly by differences in

sampling, increasing outreach efforts to reach homeless PWID, or increasing representation of socially and economically marginalized populations in study samples over time. Further research is needed to understand causes of the observed time trends in homelessness. Furthermore, explicit homelessness is one outcome along a continuum of housing instability that could include for example couch surfing and doubling up. PWID may experience varying degrees of housing instability, and changes in housing status over time. Future studies could expand definitions of homelessness to gain more nuanced information about how different types of housing instability impact injection risk.

Our findings are consistent with other studies linking homelessness with injection risk behavior, an association that appears to be consistent across age groups, geographic locations, and time periods [1, 8, 44, 45]. Developmentally tailored interventions to increase access to harm reduction services and address the causes of homelessness among young PWID are necessary for prevention of HIV/HCV, overdose, and other negative health and social consequences. Increasing access to mental health services for young PWID is also a priority given the high rates of mental illness among young PWID[4, 35], particularly those experiencing homelessness [46].

Limitations. An important limitation of this study is the small number of samples included in the meta-regression that limited our ability to detect changes in PWID behavior over time. Findings may not be generalizable to PWID in other geographic locations or settings. Participants were also predominantly non-Hispanic white and samples contained small proportions of Black and Hispanic PWID. Thus we did not have sufficient power for comparison of trends over time by race/ethnicity or for subgroup analyses. Understanding risk associated with homelessness and other stressors unique to PWID from Black and Hispanic communities remains a priority for future research. Because many of the included studies recruited participants from an urban SSP, participants likely had more frequent contact with harm reduction services than would be expected in other PWID, which may partly explain why increases in injection risk were not

observed despite increases in homelessness. While measures were harmonized to be as equivalent as possible across studies, differences across studies in time frame of measurement and definitions may have contributed to between study heterogeneity in prevalence of homelessness and injection behaviors and network size. For example, variation in the time frame for assessment of the size of social and injection networks is problematic since networks are often dynamic and may change over short periods due to factors such as incarceration, drug treatment, or mortality. Furthermore, if the timeframe for homelessness increased over time (e.g., from past 6 months to past 12 months), apparent increases in homelessness could have been due to expansion of the reference timeframe. The magnitude of the time trend in homelessness was slightly attenuated with adjustment for the assessment timeframe, but the result remained statistically significant. Restriction to samples with an assessment timeframe of 6 months also yielded similar results. Thus, these findings suggest that the increasing trend in homelessness was not explained solely by differences in the timeframe of measurement.

Conclusions: Increases over time in homelessness among young PWID are concerning and highlight the need to understand the correlates of youth homelessness among general and at-risk populations to inform appropriate targeting of HIV/HCV/STI/overdose prevention and intervention services. Age appropriate interventions for young PWID are critical to prevent them from falling into chronic homelessness.

References

1. Linton SL, Celentano DD, Kirk GD, Mehta SH. The longitudinal association between homelessness, injection drug use, and injection-related risk behavior among persons with a history of injection drug use in Baltimore, MD. *Drug Alcohol Depend.* 2013;132(3):457-65. Epub 2013/04/13. DOI: 10.1016/j.drugalcdep.2013.03.009.
2. Coady M, Latka M, Thiede H, Golub E, Ouellet L, Hudson S, et al. Housing status and associated differences in HIV risk behaviors among young injection drug users (IDUs). *AIDS Behav.* 2007;11(6):854-63. DOI: 10.1007/s10461-007-9248-1.
3. Des Jarlais DC, Braine N, Friedmann P. Unstable housing as a factor for increased injection risk behavior at US syringe exchange programs. *AIDS Behav.* 2007;11(2):78-84. DOI: 10.1007/s10461-007-9227-6.
4. Mackesy-Amiti ME, Boodram B. Feasibility of ecological momentary assessment to study mood and risk behavior among young people who inject drugs. *Drug Alcohol Depend.* 2018;187:227-35. DOI: 10.1016/j.drugalcdep.2018.03.016.
5. Rhodes T, Singer M, Bourgois P, Friedman SR, Strathdee SA. The social structural production of HIV risk among injecting drug users. *Soc Sci Med.* 2005;61(5):1026-44. DOI: 10.1016/j.socscimed.2004.12.024.
6. Robertson MJ, Clark RA, Charlebois ED, Tulsy J, Long HL, Bangsberg DR, et al. HIV seroprevalence among homeless and marginally housed adults in San Francisco. *Am J Public Health.* 2004;94(7):1207-17. Epub 2004/07/01. DOI: 10.2105/ajph.94.7.1207.
7. Cheung RC, Hanson AK, Maganti K, Keffe EB, Matsui SM. Viral hepatitis and other infectious diseases in a homeless population. *J Clin Gastroenterol.* 2002;34(4):476-80.
8. Gelberg L, Robertson MJ, Arangua L, Leake BD, Sumner G, Moe A, et al. Prevalence, distribution, and correlates of hepatitis C virus infection among homeless adults in Los Angeles. *Public Health Rep.* 2012;127(4):407-21. DOI: 10.1177/003335491212700409.

9. Coates J, McKenzie-Mohr S. Out of the frying pan, into the fire: Trauma in the lives of homeless youth prior to and during homelessness. *Journal of Sociology & Social Welfare*. 2010;37(4):65-96.
10. Edidin JP, Ganim Z, Hunter SJ, Karnik NS. The mental and physical health of homeless youth: A literature review. *Child Psychiatry Hum Dev*. 2012;43(3):354-75. DOI: 10.1007/s10578-011-0270-1.
11. Medlow S, Klineberg E, Steinbeck K. The health diagnoses of homeless adolescents: A systematic review of the literature. *J Adolesc*. 2014;37(5):531-42. DOI: <https://doi.org/10.1016/j.adolescence.2014.04.003>.
12. Tyler KA, Melander LA. Child abuse, street victimization, and substance use among homeless young adults. *Youth & Society*. 2015;47(4):502-19. DOI: 10.1177/0044118x12471354.
13. Cutuli JJ, Treglia D, Herbers JE. Adolescent homelessness and associated features: Prevalence and risk across eight states. *Child Psychiatry Hum Dev*. 2020;51(1):48-58. DOI: 10.1007/s10578-019-00909-1.
14. Fortier E, Alavi M, Micallef M, Dunlop AJ, Balcomb AC, Day CA, et al. The effect of social functioning and living arrangement on treatment intent, specialist assessment and treatment uptake for hepatitis C virus infection among people with a history of injecting drug use: The ETHOS study. *Int J Drug Policy*. 2015;26(11):1094-102. DOI: <http://dx.doi.org/10.1016/j.drugpo.2015.06.001>.
15. Corneil TA, Kuyper LM, Shoveller J, Hogg RS, Li K, Spittal PM, et al. Unstable housing, associated risk behaviour, and increased risk for HIV infection among injection drug users. *Health & Place*. 2006;12(1):79-85. DOI: <https://doi.org/10.1016/j.healthplace.2004.10.004>.
16. Richer I, Bertrand K, Vandermeersch J, Roy E. A prospective cohort study of non-fatal accidental overdose among street youth: The link with suicidal ideation. *Drug Alcohol Rev*. 2013;32(4):398-404. Epub 7 NOV 2012. DOI: 10.1111/dar.12003.

17. Sherman SG, Cheng Y, Kral AH. Prevalence and correlates of opiate overdose among young injection drug users in a large U.S. city. *Drug Alcohol Depend.* 2007;88(2–3):182-7. DOI: 10.1016/j.drugalcdep.2006.10.006.
18. Jenness S, Kobrak P, Wendel T, Neaigus A, Murrill C, Hagan H. Patterns of exchange sex and HIV infection in high-risk heterosexual men and women. *J Urban Health.* 2011;88(2):329-41. DOI: 10.1007/s11524-010-9534-5.
19. Walters SM, Rivera AV, Reilly KH, Anderson BJ, Bolden B, Wogayehu A, et al. Exchange sex among persons who inject drugs in the New York metropolitan area: The importance of local context, gender and sexual identity. *AIDS Behav.* 2018;22(9):2773-87. DOI: 10.1007/s10461-018-2039-z.
20. Hodgson KJ, Shelton KH, van den Bree MBM, Los FJ. Psychopathology in young people experiencing homelessness: A systematic review. *Am J Public Health.* 2013;103(6):e24-e37. DOI: 10.2105/ajph.2013.301318.
21. Perlman S, Willard J, Herbers JE, Cutuli JJ, Garg KME. Youth homelessness: Prevalence and mental health correlates. *Journal of the Society for Social Work and Research.* 2014;5(3):361-77. DOI: 10.1086/677757.
22. Binswanger IA, Stern MF, Yamashita TE, Mueller SR, Baggett TP, Blatchford PJ. Clinical risk factors for death after release from prison in Washington State: A nested case–control study. *Addiction.* 2016;111(3):499-510. DOI: 10.1111/add.13200.
23. Spinelli MA, Hessol NA, Schwarcz S, Hsu L, Parisi M-K, Pipkin S, et al. Homelessness at diagnosis is associated with death among people with HIV in a population-based study of a U.S. City. *AIDS.* 2019. DOI: 10.1097/qad.0000000000002287.
24. Zolopa AR, Hahn JA, Gorter R, Miranda J, Wlodarczyk D, Peterson J, et al. HIV and tuberculosis infection in San Francisco's homeless adults: Prevalence and risk factors in a representative sample. *JAMA.* 1994;272(6):455-61. DOI: 10.1001/jama.1994.03520060055032.

25. Zivanovic R, Milloy MJ, Hayashi K, Dong H, Sutherland C, Kerr T, et al. Impact of unstable housing on all-cause mortality among persons who inject drugs. BMC Public Health. 2015;15:106. DOI: 10.1186/s12889-015-1479-x.
26. Fernández-Viña MH, Prood NE, Herpolsheimer A, Waimberg J, Burris S. State Laws Governing Syringe Services Programs and Participant Syringe Possession, 2014-2019. Public Health Rep. 2020;135(1_suppl):128s-37s. Epub 2020/08/01. DOI: 10.1177/0033354920921817.
27. Syringe Distribution Laws. The Policy Surveillance Program: A Law Atlas Project url: <https://lawatlas.org/datasets/syringe-policies-laws-regulating-non-retail-distribution-of-drug-parapherna>. Available from: <https://lawatlas.org/datasets/syringe-policies-laws-regulating-non-retail-distribution-of-drug-parapherna>.
28. State of Illinois General Assembly. An ACT in relation to hypodermic syringes and needles. , (2003).
29. Saulny S. After recession, more young adults are living on street. New York Times. Dec 18, 2012.
30. Thorpe LE, Ouellet LJ, Levy JR, Williams IT, Monterroso ER. Hepatitis C virus infection: Prevalence, risk factors, and prevention opportunities among young injection drug users in Chicago, 1997-1999. J Infect Dis. 2000;182(6):1588-94. DOI: 10.1086/317607.
31. Booth RE, Strathdee SA. Baseline findings from the Third Collaborative Injection Drug Users Study/Drug Users Intervention Trial (CIDUS III/DUIT). Drug Alcohol Depend. 2007;91(Supplement 1):S1-S3. DOI: 10.1016/j.drugalcdep.2007.08.010.
32. Boodram B, Hershov RC, Cotler SJ, Ouellet LJ. Chronic hepatitis C virus infection and increases in viral load in a prospective cohort of young, HIV-uninfected injection drug users. Drug Alcohol Depend. 2011;119(3):166-71. DOI: 10.1016/j.drugalcdep.2011.06.005.
33. Broz D, Ouellet LJ. Prevalence and correlates of former injection drug use among young noninjecting heroin users in Chicago. Substance Use and Misuse. 2010;45(12):2000-25. DOI: 10.3109/10826081003682875.

34. Rothenberg R, Jenkins R, Lambert E. Special Issue: Sexual Acquisition and Transmission of HIV Cooperative Agreement Program (SATHCAP), July 2009. J Urban Health. 2009;86(S1):144-8. Epub Jun 10. DOI: 10.1007/s11524-009-9374-3.
35. Mackesy-Amiti ME, Donenberg GR, Ouellet LJ. Prevalence of psychiatric disorders among young injection drug users. Drug Alcohol Depend. 2012;124(1-2):70-8. Epub 2012/01/10. DOI: 10.1016/j.drugalcdep.2011.12.012.
36. Centers for Disease Control and Prevention. HIV infection and HIV-associated behaviors among injecting drug users -- 20 Cities, United States, 2009. MMWR Morb Mortal Wkly Rep. 2012;61(8):133-8.
37. Centers for Disease Control and Prevention. HIV Infection, Risk, Prevention, and Testing Behaviors among Persons Who Inject Drugs-National HIV Behavioral Surveillance: Injection Drug Use, 20 U.S. Cities, 2012. March. Report No.: HIV Surveillance Special Report 11. Available from: <http://www.cdc.gov/hiv/library/reports/surveillance>.
38. Boodram B, Mackesy-Amiti ME, Latkin C. The role of social networks and geography on risky injection behaviors of young persons who inject drugs. Drug Alcohol Depend. 2015;154:229-35. Epub 2015/07/15. DOI: 10.1016/j.drugalcdep.2015.06.042.
39. Centers for Disease Control and Prevention. HIV Infection, Risk, Prevention, and Testing Behaviors among Persons Who Inject Drugs-National HIV Behavioral Surveillance: Injection Drug Use, 20 U.S. Cities, 2015. May. Report No.: HIV Surveillance Special Report 18. [cited 12/05/2018]. Available from: <https://www.cdc.gov/hiv/library/reports/hiv-surveillance.html>.
40. Harris R, Bradburn M, Deeks J, Harbord R, Altman D, Sterne J. metan: fixed- and random-effects meta-analysis. Stata Journal. 2008;8(1):3-28.
41. Thompson SG, Higgins JPT. How should meta-regression analyses be undertaken and interpreted? Stat Med. 2002;21(11):1559-73. DOI: 10.1002/sim.1187.
42. Knapp G, Hartung J. Improved tests for a random effects meta-regression with a single covariate. Stat Med. 2003;22(17):2693-710. DOI: 10.1002/sim.1482.

43. StataCorp. Stata Statistical Software: Release 15. Version: 15 [software]. Available from: <http://www.stata.com>.
44. Beech BM, Myers L, Beech DJ, Kernick NS. Human immunodeficiency syndrome and hepatitis B and C infections among homeless adolescents. Semin Pediatr Infect Dis. 2003;14(1):12-9. Epub 2003/05/16. DOI: 10.1053/spid.2003.127212.
45. Smereck GA, Hockman EM. Prevalence of HIV infection and HIV risk behaviors associated with living place: on-the-street homeless drug users as a special target population for public health intervention. Am J Drug Alcohol Abuse. 1998;24(2):299-319. Epub 1998/06/27. DOI: 10.3109/00952999809001714.
46. Brown RT, Steinman MA. Characteristics of Emergency Department Visits by Older Versus Younger Homeless Adults in the United States. Am J Public Health. 2013;103(6):1046-51. DOI: 10.2105/ajph.2012.301006.

Figure 1: Mean proportion of PWID reporting homelessness by study and year