

Table S1a. SARS-CoV-2 viral load for each sampling site and swab type

Case number	Viral load (copies/mL)		
	Nasopharyngeal sample	Anterior nasal sample with NP-type swab	Anterior nasal sample with OP-type swab
1	755	3	5
2	94930	357700	145400
3	1804	255800	851300
4	730	2	70
5	490700	962	4018
6	45	42	818
7	2851	35	82
8	340000	41570	133100
9	73090	5532	6415
10	14180	49480	70
11	2139000	18120	683000
12	706800	126100	21140
13	588900	63240	310800
14	136700	78300	216900
15	34030	688	6322
16	8616	114	8763
17	207	Not detected	Not detected
18	119400	91150	69960
19	228	109	19
20	791900	213300	85680
21	886	Not detected	Not detected
22	71	8	Not detected
23	3773000	31	4
24	665500	855800	48210
25	327400	66120	37970
26	22	Not detected	Not detected
27	40	Not detected	Not detected
28	79	Not detected	Not detected
29	48670000	4994000	1386000
30	11	1	8
31	17480000	1047000	4341000
32	1119000	2621	48130

NP-type, nasopharyngeal-type; OP-type, oropharyngeal-type.

Table S2b. SARS-CoV-2 cycle threshold (Ct) value for each sampling site and swab type

Case number	Ct value		
	Nasopharyngeal sample	Anterior nasal sample with NP-type swab	Anterior nasal sample with OP-type swab
1	27.9	35.8	34.8
2	16.6	17.9	19.2
3	24.8	18.4	16.5
4	27.3	35.9	30.8
5	18.3	27.2	25.3
6	31.6	31.5	27.4
7	25.6	31.9	30.7
8	18.9	21.8	20.2
9	21.0	24.7	24.5
10	21.9	19.7	31.2
11	13.1	21.5	15.1
12	15.1	18.1	21.2
13	15.4	19.3	16.5
14	18.0	19.0	17.1
15	20.4	27.4	23.4
16	22.9	30.6	22.8
17	29.5	Not detected	Not detected
18	18.2	18.7	19.2
19	29.3	30.6	33.7
20	14.8	17.2	18.8
21	26.9	Not detected	Not detected
22	34.4	34.3	Not detected
23	14.0	32.1	35.5
24	16.7	16.3	20.8
25	17.8	20.3	21.1
26	32.3	Not detected	Not detected
27	31.5	Not detected	Not detected
28	30.6	Not detected	Not detected
29	12.5	15.6	17.3
30	33.2	36.5	33.6
31	13.9	17.7	15.8
32	17.6	25.9	21.9

NP-type, nasopharyngeal-type; OP-type, oropharyngeal-type.