


**Figure 1:** Flowchart of the study population including response rate for the questionnaire.



<sup>a</sup> Eligibility criteria: patients from Psychiatric Services of Central Denmark Region,  $\geq 18$  years old, having at least one psychiatric contact in the three months (91) days before June 25, 2020. Patients with forensic sanction or diagnosed with an organic mental disorder or mental retardation were not eligible.

<sup>b</sup> A clinical diagnosis of F2x (psychotic disorders), F3x (mood disorders), F4x (anxiety- and stress-related disorders), F5x (eating-, sleeping- and other behavioral syndromes associated with physiological disturbances), F6x (personality disorders), F8x (developmental disorders incl. autism), F90-F98 (child- and adolescent mental disorders), F1x (substance abuse disorders) or Zx (contact with psychiatric health services incl. examination) prior to the nationwide lockdown (March 11, 2020).

<sup>c</sup> Invitation were sent via the electronic mailing system (e-Boks) used by the Danish authorities.

**Table 1:** Sociodemographic and clinical characteristics of respondents versus non-respondents. The p-values are based upon t-test for the continuous values and Fisher's exact test or chi-square test for the categorical variables. Significant p-values are marked in **bold**.

<i>Characteristics</i>	<i>Respondents (n = 1029)</i>	<i>Non-respondents (n = 4258)</i>	<i>P-value</i>
<b><i>Sex, n (%)</i></b>			<b>9.7e-12</b>
Female	708 (68.8)	2439 (57.3)	
<b><i>Age, median (IQR)</i></b>	35 (26;50)	34 (26;48)	0.25
<b><i>Civil status, n (%)</i></b>			<b>9.5e-5</b>
Single	638 (62.0)	2890 (67.9)	
Married or reg. partnership	241 (23.4)	733 (17.2)	
Divorced or repeal of reg. partnership	133 (12.9)	572 (13.4)	
Widow/widower	17 (1.7)	63 (1.5)	
<b><i>Municipality, n (%)</i></b>			<b>6.8e-4</b>
Inhabitants 150,000-350,000	376 (36.5)	1322 (31.0)	
Inhabitants 50,000-150,000	425 (41.3)	2018 (47.4)	
Inhabitants < 50,000 + unknown <sup>a</sup>	228 (22.2)	918 (21.6)	
<b><i>Diagnosis, n (%)</i></b>			<b>9.5e-10</b>
F10-F19	8 (0.8)	50 (1.2)	
F20-F29	204 (19.8)	1226 (28.8)	
F30-F31	172 (16.7)	600 (14.1)	
F32-F33	329 (32.0)	1027 (24.1)	
F40-F48	152 (14.8)	646 (15.2)	
F50-F59	14 (1.4)	41 (1.0)	
F60-F69	64 (6.2)	200 (4.7)	
F90-F98	41 (4.0)	215 (5.0)	
Other F-diagnoses <sup>b</sup>	8 (0.8)	56 (1.3)	
Z-diagnoses	37(3.6)	197 (4.6)	
<b><i>Psychiatric inpatient days, median (IQR)</i></b>			
Pre-corona (01.01.2015-10.03.2020)	0 (0;13)	0 (0;19)	0.39
Corona (11.03.2020-26.06.2020)	0 (0;0)	0 (0;0)	0.12
<b><i>Psychiatric outpatient visits<sup>c</sup>, median (IQR)</i></b>			
Pre-corona (01.01.2015-10.03.2020)	37 (14;77)	33 (13;67)	<b>6.2e-3</b>
Corona (11.03.2020-26.06.2020)	7 (4;12)	6 (4;10)	<b>2.0e-9</b>
<b><i>Adults living in your household besides yourself?</i></b>			
Yes <sup>d</sup>	617 (60.8)		
<b><i>Children living in your household?</i></b>			
Yes <sup>e</sup>	250 (24.7)		
<b><i>Were you born in Denmark?</i></b>			
Yes <sup>f</sup>	919 (91.2)		
<b><i>Where were you born? (condition: previous question = No<sup>g</sup>)</i></b>			
Asia	14 (15.7)		
Europe	42 (47.2)		
Middle East	16 (18.0)		

North America + South America	10 (11.2)
Africa + Oceania + unknown	7 (7.9)
<b>What is your highest attained educational level?<sup>f</sup></b>	
Primary and lower secondary school	214 (21.2)
Upper secondary education	258 (25.6)
Skilled worker/craftsman	64 (6.3)
Short-cycle higher education	157 (15.6)
Medium-cycle higher education including bachelor	206 (20.4)
Long-cycle higher education	109 (10.8)
<b>What describes your current employment status?<sup>f</sup></b>	
Full-time student	92 (9.1)
Part-time student	27 (2.7)
Full-time employed	83 (8.2)
Part-time employed	39 (3.9)
Subsidized employment	73 (7.2)
Absent owing to illness	359 (35.6)
Unemployed	125 (12.4)
Retired	210 (20.8)

<sup>a</sup>: Unknown address.

<sup>b</sup>: F34 + F38 + F39 + F80-F89

<sup>c</sup>: In addition to physical meetings, outpatient contacts also include telephone and video consultations.


<sup>d</sup>: Proportion is taken of 1015


<sup>e</sup>: Proportion is taken of 1012

<sup>f</sup>: Proportion is taken of 1008

<sup>g</sup>: Proportion is taken of 89, since 89 replied no and 919 replied yes (in total 1008) to the question about being born in DK.

**Figure 2:** Visualization of the non-weighted and weighted mean and 95%-confidence interval of the WHO-5 total score, the BSI-18 total score, the three subscales of BSI-18.





**Table 2:** Table displaying the non-weighted (red) and weighted (blue) mean (SD) of the WHO-5 total score, the BSI-18 total score, the three subscales of BSI-18 for all respondents and stratified on the ICD-10 diagnosis: F2, F30-F31, F32-F33, F4 and F6.


<b>Questionnaires</b>	<b>Respondents</b> (n = 992)	<b>F2</b> (n = 204)	<b>F30-F31</b> (n = 172)	<b>F32-F33</b> (n = 329)	<b>F4</b> (n = 152)	<b>F6</b> (n = 64)
<b>WHO-5</b>						
Total score*	37.4 (22.4) 37.9 (22.6)	42.9 (23.3) 42.9 (23.1)	38.6 (23.5) 38.5 (23.6)	34.1 (20.8) 34.2 (20.9)	35.8 (22.1) 35.2 (22.1)	34.9 (19.8) 35.0 (20.0)
<b>BSI-18</b>						
Total score	28.6 (14.9) 28.4 (15.1)	25.9 (14.9) 25.6 (14.8)	25.5 (15.5) 25.9 (15.9)	29.5 (13.4) 29.4 (13.5)	32.7 (16.9) 33.0 (16.7)	31.0 (13.6) 31.0 (13.4)
Scale 1: Somatization	6.5 (4.9) 6.4 (5.0)	5.8 (5.1) 5.6 (5.1)	5.4 (4.7) 5.6 (4.9)	6.5 (4.3) 6.4 (4.3)	8.3 (5.7) 8.3 (5.6)	7.2 (5.3) 7.1 (5.2)
Scale 2: Depression	11.9 (6.3) 11.9 (6.3)	11.2 (6.0) 11.1 (5.9)	11.0 (6.9) 11.1 (6.9)	12.6 (5.8) 12.6 (5.9)	12.3 (6.7) 12.5 (6.7)	12.9 (5.7) 12.9 (5.6)
Scale 3: Anxiety	10.2 (5.7) 10.1 (5.8)	9.0 (5.7) 8.9 (5.7)	9.1 (6.0) 9.2 (6.1)	10.4 (5.2) 10.3 (5.3)	12.1 (6.2) 12.2 (6.2)	10.9 (4.6) 10.9 (4.6)

**Table 3:** Logistic regression with weighted deterioration in mental health as outcome (participants responding either 'Much worse' or 'Slightly worse' to the question: "How do you consider your mental health during the lockdown compared to the period before the corona pandemic came to Denmark?").

	<b>OR</b>	<b>CI<sub>95%</sub>(OR)</b>	<b>z-value</b>	<b>P-value</b>
<b>Clinical and demographic variables</b>				
<b>Sex (reference: male)</b>				
Female	1.17	(0.87;1.59)	1.04	0.298
<b>Age (reference: ≤1<sup>st</sup> quantile of age)</b>				
1 <sup>st</sup> -2 <sup>nd</sup> quantile	1.02	(0.66;1.58)	0.11	0.912
2 <sup>nd</sup> -3 <sup>rd</sup> quantile	0.61	(0.37;1.01)	-1.90	0.057
>3 <sup>rd</sup> quantile	0.66	(0.37;1.17)	-1.43	0.153
<b>Civil status (reference: married &amp; registered partnership)</b>				
Single	0.99	(0.62;1.58)	-0.05	0.963
Divorced, Repeal of registered partnership	1.00	(0.60;1.67)	-0.01	0.992
Widow/widower	0.77	(0.24;2.42)	-0.45	0.656
<b>Municipality (reference: Inhabitants &lt; 50,000)</b>				
Inhabitants 150,000-350,000	1.16	(0.79;1.73)	0.76	0.449
Inhabitants 50,000-150,000	1.03	(0.72;1.48)	0.18	0.855
<b>Diagnosis (reference: F32-F33)</b>				
F10-F19	0.84	(0.21;3.00)	-0.27	0.789
F20-F29	0.57	(0.37;0.87)	-2.58	<b>0.010</b>
F30-F31	1.08	(0.69;1.69)	0.35	0.727
F40-F49	1.46	(0.93;2.30)	1.62	0.105
F50-F59	0.64	(0.16;2.35)	-0.67	0.505
F60-F69	1.41	(0.73;2.78)	1.01	0.311
F90-F98	1.20	(0.58;2.48)	0.49	0.627
Other F diagnoses	0.72	(0.17;2.70)	-0.48	0.629
Z-diagnoses	0.68	(0.31;1.45)	-0.99	0.322
<b>Psychiatric admission pre-corona (reference: no admission)</b>				
Admission	0.93	(0.68;1.29)	-0.41	0.685
<b>Psychiatric outpatient visits pre-corona (reference: 1<sup>st</sup> quantile)</b>				
1 <sup>st</sup> -2 <sup>nd</sup> quantile, n (%)	1.24	(0.83;1.85)	1.04	0.300
2 <sup>nd</sup> -3 <sup>rd</sup> quantile, n (%)	1.56	(1.03;2.37)	2.09	<b>0.037</b>
>3 <sup>rd</sup> quantile, n (%)	1.45	(0.92;2.29)	1.59	0.112
<b>What is your highest attained educational level? (reference: long-cycle higher education)</b>				
Primary and lower secondary school	1.15	(0.64;2.07)	0.48	0.634
Upper secondary education	1.31	(0.75;2.28)	0.95	0.344
Skilled worker/craftsman	1.07	(0.53;2.14)	0.19	0.852
Short-cycle higher education	1.13	(0.64;2.00)	0.41	0.681
Medium-cycle higher education including bachelor	1.20	(0.69;2.08)	0.64	0.521
<b>What describes your current employment status? (reference: full-time employed)</b>				


Full-time student	0.89	(0.44;1.78)	-0.34	0.737
Part-time student	1.73	(0.57;5.49)	0.96	0.337
Part-time employed	0.87	(0.35;2.09)	-0.32	0.748
Subsidized employment	1.95	(0.94;4.08)	1.78	0.076
Absent owing to illness	1.23	(0.71;2.15)	0.73	0.463
Unemployed	0.86	(0.45;1.65)	-0.45	0.654
Retired	1.91	(1.00;3.65)	1.96	0.050
<b>Adults living in your household besides yourself? (reference: yes)</b>				
No	1.39	(1.01;1.91)	2.04	<b>0.042</b>
<b>Children living in your household? (reference: yes)</b>				
No	1.14	(0.77;1.67)	0.65	0.518
<b>Were you born in Denmark? (reference: yes)</b>				
No	1.13	(0.69;1.86)	0.48	0.635

**Figure 3:** Perceived reasons for deterioration during the pandemic lockdown (weighted).



Perceived reasons for deterioration among the 492/960 (51%) of the respondents who reported that their overall mental health had worsened during the nationwide pandemic lockdown.

**Figure 4:** Perceived reasons for deterioration (weighted) among the 492/960 (51%) of the respondents who reported that their overall mental health had worsened during the nationwide pandemic lockdown stratified on the following diagnostic groups: psychotic disorder (ICD-10: F2x), bipolar disorder (ICD-10: F30-31), unipolar depression (ICD-10: F32-33), anxiety disorder (ICD-10: F4x), and personality disorder (ICD-10: F6x). The following categories are removed due to microdata in the subdiagnosis groups: "Home schooling / taking care of children", "Working from home", "Fighting (quarreling) at home", "Violence at home", "Increased consumption of alcohol, addictive medication, or illicit drugs", "Concerns about unemployment", "Other causes for the worsening?" have been removed from the figure due to small numbers and risk of identification of individuals.




**Table 4:** Perceived reasons for deterioration (weighted) among the 492/960 (51%) respondents who reported that their overall mental health had worsened during the nationwide pandemic lockdown and stratified on the following diagnostic groups: psychotic disorder (ICD-10: F2x), bipolar disorder (ICD-10: F30-31), unipolar depression (ICD-10: F32-33), anxiety disorder (ICD-10: F4x), and personality disorder (ICD-10: F6x). “-” indicates microdata.

<i>Deterioration in mental health %</i>	<i>Respondents (n = 492)</i>	<i>F2</i>	<i>F30-F31</i>	<i>F32-F33</i>	<i>F4</i>	<i>F6</i>	<i>Other</i>
Concerns about getting infected with coronavirus	44.9	12.8	5.8	10.2	9.2	2.6	4.4
Concerns about family/friends getting infected with coronavirus	58.3	15.8	7.0	15.7	10.4	4.5	4.8
Concerns about infecting others with coronavirus	40.3	8.8	4.8	11.1	8.0	3.7	3.9
Lack of trust in authorities' management of the corona pandemic	13.8	3.2	2.0	2.2	2.8	2.1	1.4
Home schooling/taking care of children	12.6	-	-	-	-	-	-
Working from home	9.6	-	-	-	-	-	-
No access to support measures	38.7	9.6	5.6	11.6	6.0	2.8	3.0
No access to leisure activities	30.1	9.3	4.1	7.6	5.2	1.5	2.3
No access to fitness center	19.9	4.1	3.0	5.1	3.8	1.4	2.5
Reduced physical activity	43.1	10.7	7.1	10.3	8.5	2.6	3.9
Reduced access to psychiatric treatment	42.2	9.8	6.4	11.9	7.3	3.8	3.0

Disruption of routines	64.8	15.9	10.2	18.2	10.6	5.3	4.5
Cancellation of events	38.2	10.1	5.4	9.7	6.3	2.6	4.2
Media coverage of the corona pandemic	30.1	6.0	4.9	7.6	7.1	1.9	2.6
Social media (e.g. Facebook)	19.2	3.8	3.5	4.2	4.6	1.4	1.7
Other people's management of the corona pandemic	39.3	9.7	5.1	9.0	7.7	4.2	3.7
Fighting (quarreling) at home	13.7	-	-	-	-	-	-
Violence at home	1.1	-	-	-	-	-	-
Increased consumption of alcohol, addictive medication, or illicit drugs	13.0	-	-	-	-	-	-
Concerns about unemployment	14.3	-	-	-	-	-	-
Concerns about financials	33.2	7.4	5.8	8.5	6.6	2.9	1.9
Boredom	43.0	10.5	6.9	10.4	7.8	3.2	4.2
Less contact with friends	58.1	14.3	10.7	14.6	10.4	3.1	5.0
Less contact with family	49.9	11.2	8.6	13.8	9.2	2.7	4.5
Loneliness	66.7	15.7	10.2	17.1	11.7	4.8	7.3
Poor diet	32.8	7.2	5.4	8.1	5.7	3.3	3.1
Other causes for the worsening?	16.3	-	-	-	-	-	-


**Figure 5:** Weighted response to the 14 questions focusing on changes in mental health and compliance to treatment during the COVID-19 lockdown of Denmark in the spring of 2020.




The numbers in the parentheses represent the proportion of respondents endorsing one of the five likert-scale responses (they could also choose: "The symptom was not present (either before or during the lockdown)" or choose not to respond).




**Figure 6: Psychotic disorders**




**Figure 7: Bipolar disorder**




**Figure 8: Unipolar depression**



**Figure 9: Anxiety- and stress-related disorders**



**Figure 10: Personality disorders**



**Table 5: Proportions of response to the 14 mental health questions for all the respondents and stratified on the following diagnostic groups: psychotic disorder (ICD-10: F2x), bipolar disorder (ICD-10: F30-31), unipolar depression (ICD-10: F32-33), anxiety disorder (ICD-10: F4x), and personality disorder (ICD-10: F6x). “-“ indicates microdata.**

<i>All respondents (n=1029)</i> %	<i>Much worse</i>	<i>Slightly worse</i>	<i>Approximately the same</i>	<i>Slightly better</i>	<i>Much better</i>
Mental health (93.2%)	19.5	30.2	34.6	9.2	6.5
Sleep disturbances (86.9%)	14.0	26.0	50.2	7.1	2.7
Appetite disturbances (79.1%)	14.0	28.4	50.8	5.0	1.8
Anxiety symptoms (83.3%)	20.7	30.6	35.8	8.3	4.6
Depression symptoms (87.1%)	23.0	29.6	33.9	9.2	4.2
Obsessions and compulsions (52.5%)	15.9	24.6	53.0	4.0	2.5
Symptoms of somatization (59.8%)	12.3	31.3	50.5	4.2	1.7
Symptoms of mania (49.4%)	9.9	26.7	54.6	5.6	3.2
Psychotic symptoms (40.2%)	13.4	25.7	53.3	3.7	4.0
Negative symptoms (63.9%)	18.8	31.9	40.6	5.7	3.0
Self-harm (33.4%)	14.2	15.8	60.4	5.2	4.3
Symptoms of eating disorders (36.0%)	17.4	26.8	48.4	5.4	2.0
Substance abuse (24.7%)	14.2	25.5	50.8	5.5	4.0
Compliance to medication (78.8%)	2.6	7.1	70.3	3.6	16.5

<b>F2 (n=204)</b> %	<b>Worse</b>	<b>Approximately the same</b>	<b>Approximately the same + better</b>	<b>Better</b>
Mental health (92.2%)	43.8	44.6	-	11.7
Sleep disturbances (81.4%)	38.0	55.7	-	6.3
Appetite disturbances (77.9%)	40.1	53.9	-	5.9
Anxiety symptoms (81.4%)	45.0	46.3	-	8.7
Depression symptoms (81.9%)	46.6	41.6	-	11.9
Obsessions and compulsions (61.8%)	44.9	-	55.1	-
Symptoms of somatization (62.7%)	41.3	-	58.7	-
Symptoms of mania (52.5%)	32.8	-	67.2	-
Psychotic symptoms (70.1%)	39.1	54.0	-	6.9
Negative symptoms (77.9%)	48.3	43.7	-	7.9
Self-harm (40.7%)	17.7	76.2	-	6.1
Symptoms of eating disorders (43.1%)	47.9	45.4	-	6.7
Substance abuse (29.9%)	31.3	59.7	-	8.9
Compliance to medication (83.8%)	11.5	69.2	-	19.3
<b>F30-31 (n=172)</b> %	<b>Worse</b>	<b>Approximately the same</b>	<b>Approximately the same + better</b>	<b>Better</b>
Mental health (92.4%)	53.1	32.4	-	14.5
Sleep disturbances (87.2%)	44.3	47.6	-	8.1
Appetite disturbances (77.9%)	48.3	45.9	-	5.7
Anxiety symptoms (75.6%)	55.2	36.6	-	8.1
Depression symptoms (86.6%)	52.0	33.1	-	14.9
Obsessions and compulsions (39.5%)	38.0	54.9	-	7.1
Symptoms of somatization (52.9%)	46.6	-	53.4	-
Symptoms of mania (68.0%)	35.0	53.3	-	11.7
Psychotic symptoms (32.0%)	36.7	-	63.3	-
Negative symptoms (58.1%)	55.0	-	45.0	-
Self-harm (21.5%)	30.7	-	69.3	-
Symptoms of eating disorders (27.3%)	39.2	51.0	-	9.7
Substance abuse (24.4%)	51.1	-	48.9	-
Compliance to medication (89.5%)	7.8	65.2	-	27.1
<b>F32-33 (n=329)</b> %	<b>Worse</b>	<b>Approximately the same</b>	<b>Better</b>	
Mental health (95.1%)	50.4	32.6	17.1	
Sleep disturbances (91.5%)	37.1	52.8	10.0	
Appetite disturbances (81.5%)	38.4	53.0	8.6	
Anxiety symptoms (87.5%)	50.6	35.7	13.7	
Depression symptoms (94.2%)	53.7	32.4	14.0	
Obsessions and compulsions (49.2%)	33.1	54.3	12.6	
Symptoms of somatization (57.1%)	41.0	50.9	8.1	
Symptoms of mania (41.0%)	32.0	60.1	7.9	
Psychotic symptoms (29.5%)	40.2	50.4	9.4	
Negative symptoms (63.2%)	47.9	40.3	11.8	
Self-harm (34.3%)	33.3	57.8	8.9	
Symptoms of eating disorders (34.0%)	35.6	52.3	12.1	
Substance abuse (21.3%)	42.3	47.2	10.5	
Compliance to medication (86.0%)	6.8	73.4	19.8	
<b>F4 (n=152)</b> %	<b>Worse</b>	<b>Approximately the same</b>	<b>Approximately the same + better</b>	<b>Better</b>
Mental health (91.4%)	58.2	24.1	-	17.7
Sleep disturbances (85.5%)	49.3	34.5	-	16.2
Appetite disturbances (78.9%)	50.7	41.2	-	8.1
Anxiety symptoms (86.8%)	58.0	26.0	-	16.0
Depression symptoms (83.6%)	57.5	27.8	-	14.6

Obsessions and compulsions (60.5%)	50.1	44.5	-	5.4	
Symptoms of somatization (68.4%)	48.0	43.4	-	8.5	
Symptoms of mania (43.4%)	36.0	49.5	-	14.5	
Psychotic symptoms (34.9%)	47.3	-	52.7	-	
Negative symptoms (59.9%)	58.5	30.8	-	10.7	
Self-harm (30.9%)	42.0	44.6	-	13.5	
Symptoms of eating disorders (32.2%)	52.3	-	47.7	-	
Substance abuse (21.7%)	51.5	-	48.5	-	
Compliance to medication (60.5%)	12.3	72.3	-	15.3	
<b><i>F6 (n=64)</i></b>	<b><i>Worse</i></b>	<b><i>Worse +</i></b>	<b><i>Approximately</i></b>	<b><i>Approximately</i></b>	<b><i>Better</i></b>
<b><i>%</i></b>		<b><i>approximately</i></b>	<b><i>the same</i></b>	<b><i>the same +</i></b>	
		<b><i>the same</i></b>		<b><i>better</i></b>	
Mental health (96.9%)	61.3	-	21.6	-	17.1
Sleep disturbances (92.2%)	40.6	-	51.5	-	7.9
Appetite disturbances (82.8%)	58.6	-	-	41.4	-
Anxiety symptoms (92.2%)	57.5	-	24.6	-	17.8
Depression symptoms (95.3%)	64.1	-	24.7	-	11.2
Obsessions and compulsions (65.6%)	45.1	-	-	54.9	-
Symptoms of somatization (73.4%)	50.7	-	37.5	-	11.8
Symptoms of mania (53.1%)	58.7	-	-	41.3	-
Psychotic symptoms (46.9%)	44.6	-	-	55.4	-
Negative symptoms (65.6%)	60.1	-	-	39.9	-
Self-harm (53.1%)	44.3	-	-	55.7	-
Symptoms of eating disorders (56.2%)	57.8	-	-	42.2	-
Substance abuse (25.0%)	48.9	-	-	51.1	-
Compliance to medication (56.2%)	-	83.5	-	-	16.5