1 **Determinants of COVID-19 outcomes: A systematic review**

2 Shirley Crankson*, Subhash Pokhrel, Nana Kwame Anokye
3 Division of Global Public Health, Department of Health Sciences, College of Health,
4 Medicine and Life Sciences, Brunel University London.
5

6 **Abstract**

7 **Background:** The current pandemic, COVID-19, caused by a novel coronavirus SARS-CoV-
8 2, has claimed over a million lives worldwide in a year, warranting the need for more
9 research into the wider determinants of COVID-19 outcomes to support evidence-based
10 policies.

11 **Objective:** This study aimed to investigate what factors determined the mortality and length
12 of hospitalisation in individuals with COVID-19.

13 **Data Source:** This is a systematic review with data from four electronic databases: Scopus,
14 Google Scholar, CINAHL and Web of Science.

15 **Eligibility Criteria:** Studies were included in this review if they explored determinants of
16 COVID-19 mortality or length of hospitalisation, were written in the English Language, and
17 had available full-text.

18 **Study appraisal and data synthesis:** The authors assessed the quality of the included studies
19 with the Newcastle-Ottawa Scale and the Agency for Healthcare Research and Quality
20 checklist, depending on their study design. Risk of bias in the included studies was assessed
21 with risk of bias assessment tool for non-randomised studies. A narrative synthesis of the
22 evidence was carried out. The review methods were informed by the Joanna Briggs Institute
23 guideline for systematic reviews.

24 **Results:** The review included 22 studies from nine countries, with participants totalling
25 239,830. The included studies' quality was moderate to high. The identified determinants
26 were categorised into demographic, biological, socioeconomic and lifestyle risk factors,
27 based on the Dahlgren and Whitehead determinant of health model. Increasing age (ORs
28 1.04-20.6, 95% CIs 1.01-22.68) was the common demographic determinant of COVID-19
29 mortality while living with diabetes (ORs 0.50-3.2, 95% CIs -0.2-0.74) was one of the most
30 common biological determinants of COVID-19 length of hospitalisation.

31 **Review limitation:** Meta-analysis was not conducted because of included studies'
32 heterogeneity.

33 **Conclusion:** COVID-19 outcomes are predicted by multiple determinants, with increasing
34 age and living with diabetes being the most common risk factors. Population-level policies
35 that prioritise interventions for the elderly population and the people living with diabetes may
36 help mitigate the outbreak's impact.
37

38 PROSPERO registration number: CRD42021237063.

39

40 **Strength and limitations of this review**

- 41 • This is the first systematic review synthesising the evidence on determinants of
- 42 COVID-19 LOS outcome.
- 43 • It is also the first review to provide a comprehensive investigation of contextual
- 44 determinants of COVID-19 outcomes, based on the determinants of health model;
- 45 thus, presenting with crucial gaps in the literature on the determinants of COVID-19
- 46 outcomes that require urgent attention.
- 47 • The review was restricted in conducting meta-analysis due to included studies’
- 48 heterogeneity.
- 49 • The review focused on only papers published in the English Language; hence, other
- 50 relevant papers written on other languages could have been omitted.

51

52

53

54

55

56 **Introduction**

57 COVID-19 (SARS-CoV-2, coronavirus) is currently among the leading causes of death

58 globally. As of January 9th, 2021, 87,589,206 cases and 1,906, 606 deaths had been recorded

59 globally (1). While its case fatality ratio (CFR) has been relatively low (CFR=2.2%),

60 compared to CFRs of previous coronavirus outbreaks, notably, MERS (CFR=9.5%) and

61 SARS-COV-1 (CFR=34.4%), its aggressive and alarming transmission rate has posed

62 enormous challenges on global health (2). Even the current reported transmission rate of

63 COVID-19 may be lower than the actual transmission rate because a significant proportion of
64 infected persons may remain undetected because they are asymptomatic (3). And regarding
65 its CFR, current predictions even suggest that mortality ratio may increase since the
66 pandemic is still ongoing (4). The infectivity and fatality rates associated with COVID-19,
67 together with the worldwide panic it generates, make the current coronavirus pandemic a
68 significant threat to public health, and its gargantuan impact unlike anything the world has
69 experienced in the last two decades (5).

70

71 Since its inception, COVID-19 has overburdened the whole global health system, from
72 crippling health resources to causing paradigm shifts in health care delivery (6). The testing
73 process, quarantine and isolation associated with the virus has had dire psychological and
74 financial implications on individuals and institutions (7). Furthermore, lockdowns instituted
75 by affected countries to curb the virus's spread resulted in disrupted formal education,
76 unplanned fiscal costs on emergency reliefs, and decreased productivity, all translating into
77 huge economic costs to governments and organisations (8). The overall COVID-19 burden, in
78 terms of health and fiscal implications, has been consequential in both high-income and low-
79 and middle-income countries, albeit with contextual differences.

80 Regardless of the significant interventions to curb the virus's spread and subsequently reduce
81 its severest outcome, i.e., mortality and morbidity, the outbreak continue to increase. As of
82 January 9th, 2021, the daily global COVID death was 15,522, the highest daily mortality
83 since the pandemic started, and about 3,000 more deaths since the first peak in daily COVID-
84 19 deaths (initial peak April 17th – 12,511 daily deaths) (1). Also, 823,856 cases were
85 confirmed on January 9th, 2021, representing a 0.93% increase from the previous day's case
86 count. The rapid rise in the COVID-19 cases and deaths worldwide necessitates continuous

research on risk factors for COVID-19 outcomes to provide current evidence-based interventions to reduce the outbreak's drastic impact.

Several studies on determinants of COVID-19 outcomes were identified in the literature; however, most of them were primary studies investigating risk factors for COVID-19 mortality (9, 10, 11). The few secondary studies/systematic reviews found in the literature encompassed only papers from high-income countries (12, 13). Also, there is inadequate coverage on other COVID-19 outcomes, specifically, length of hospital stay (LOS). Till date, no review has explored risk factors/determinants of COVID-19 LOS. Findings from such studies will be essential for help health systems to develop contingency plans for bed occupancy and health resources, especially with the swift increase in the COVID-19 cases. Thus, this study aimed to review factors that determine COVID-19 mortality and LOS in individuals diagnosed with COVID-19 to address the literature dearth and contribute to global efforts at curtailing the pandemic. Understanding the risk factors of COVID-19 outcomes based on a comprehensive synthesis of global but rapidly emerging evidence might be useful to implement effective policies to address the disease burden.

Methods

Search strategy

From 21st to 31st December 2020, Scopus, Google Scholar, CINAHL and Web of Science databases were searched for relevant studies using the search terms: 'Determinants' 'Predictors' 'COVID-19' 'SARS-CoV-2' 'Mortality' 'Length of hospital stay' 'Length of hospitalisation'. The search terms were combined with mesh words and Boolean operators to ensure sensitive and targeted search. Full search strategy is shown in supplementary

information 1. First screening of the databases results was conducted independently by two of the reviewers (SC and NKA) to ensure their relevance to this study. The titles and abstract of the identified relevant studies were screened against this review's eligibility using the following predetermined eligibility criteria: population - individuals diagnosed with COVID-19, exposures – demographic, socioeconomic, lifestyle, environmental biological/medical factors, outcome – COVID-19 related mortality and LOS, studies that explored determinants of COVID-19 mortality and LOS in participants with COVID-19, studies whose LOS endpoint was discharge or death, and not hospital transfers, studies written and published in the English Language, and with full-text available. No date restriction was applied in any of the databases since most COVID-19 studies are recent. No database filters were also applied. The references of the identified papers were also tracked for papers eligible for this review. Any disagreements relating to studies' screening was discussed and resolved with the third reviewer (SK) and

Data extraction

Two of the authors (SC and NKA) extracted the relevant data from the included studies using a comprehensive a priori developed set of data extraction questions (Supplementary information 2), informed by the JBI data extraction tools. The questions were categorised under two main themes, i.e., general information (authors name, study settings, study aim, year of publication) and methodology (sample size, sample characteristics, outcomes etc.), to ensure the sensitivity of the questions to the overarching objectives of this review. The data extraction questions were piloted-tested on five selected studies before their final usage. The third author (SP) randomly selected and reviewed 50% of the extracted data from the

included studies to ascertain data extraction quality. Finally, all disagreements were resolved by consensus.

Risk of bias and quality assessment

The Newcastle-Ottawa Scale (NOS) and the Agency for Healthcare Research and Quality (AHRQ) appraisal checklist were used to appraise the quality of the included studies. These checklists were based on a systematic review's recommendation (14). The NOS provides eight items grouped under three main domains: the selection of cohorts, the comparability of cohorts, and outcomes assessment. A star (*) was awarded if a study met an item under the three defined domains. A maximum of one star was given to items within the selection and outcome domain, and a maximum of two stars was given to the item under the comparability domain. Thus, studies with nine stars were rated as high-quality study and those with two stars or less were graded as low quality. Like the NOS, the AHRQ also provides items/checklists (n=11) for assessing the quality of the study's methods and outcomes. A 'yes', 'no' or 'not applicable (NA)' was used to indicate whether a study met the AHRQ requirement. The number of 'yes' from a study represented the study's quality. Consequently, studies with eleven 'yes', suggesting 11 total scores, were ranked as high quality, whereas those with two or less 'yes' were rated as low quality. Also, risk of bias assessment of both the study and outcome level of the included papers was performed with risk of bias assessment tool for non-randomised studies. The quality and risk assessment findings are presented in supplementary information 3.

Data synthesis

Descriptive data synthesis, informed by the JBI manual for evidence synthesis in systematic reviews was conducted to comprehensively describe the methods, findings, and quality of the included studies (15). The studies' methods, its operationalisation and the subsequent findings were compared across the papers to identify common determinants of COVID-19 mortality and LOS. Also, the range of effect sizes (odds ratios/hazard ratios) of the identified determinants in the studies were synthesised to understand the magnitude of the effect on the study outcomes. This was done by reporting the lowest and highest effect sizes from studies that identified common determinants.

Patient and public involvement

This study reviewed already published and available research. Therefore, no patients or the public were directly involved in this review process. The findings of this review will be shared publicly through scientific publication, social media, and conference presentations.

Results

Search result

The database search yielded 1,653 studies. The authors removed 11 duplicates and eliminated a further 1,564 after title screening for relevance to this review. The abstracts of the remaining 78 studies were assessed for eligibility using the predetermined eligibility criteria. Twenty-two studies met the inclusion criteria and were subsequently included in this review, as shown in the Prisma flow diagram below (Fig 1).

Fig 1. Prisma flow diagram showing the literature search.

Overview of included studies

The studies spanned nine major countries, of which eight were high-income countries and one upper-middle-income country. Majority of them were from China (n=6) and USA (n=6), followed by Spain (n=2), and England (n=2), and one each from Kuwait, Mexico, France, Italy, Austria, and one multi-continent study- participants from Africa, Europe, Australia, Asia, and Americas (16). They all used the quantitative research approach, with retrospective cohort design as the predominant study design (n=12). Most of them (n=20) accessed only secondary data, retrieved mainly from the patients' electronic medical records. The remaining two used both secondary and primary data (face to face and telephone interviews) (10, 17).

The studies sample size ranged from 58 to 177,133, totaling 239,830 participants. All the studies included both male and female participants; however, the men dominated, representing 51.4% of the studies' entire population. Only 7 studies were age-specific - limited their inclusion to participants ≥ 18 years old. The majority (n=20) included only persons with confirmed COVID-19 test, either through the reverse transcription-polymerase chain reaction (RT-PCR) or nasopharyngeal swabs. The other two studies included all confirmed, negative, and suspected, and both suspected and confirmed COVID-19 cases (18, 19) (Table 1). The studies' quality on the NOS and AHRQ ranged from 6-9, representing moderate to high quality and the common risk of bias across the studies was the inability to control the influence of unmeasured confounders. This confounding bias inherently influenced the studies' findings and could subsequently affect this review's finding.

Of the COVID-19 outcomes, 19 studies focused on only COVID-19 mortality, 2 solely on COVID-19 LOS and 1 on both COVID-19 mortality and LOS. Mortality was generally described as either in-hospital deaths, i.e., deaths occurring in a hospital or all-cause

230 mortality, i.e., all deaths in COVID-19 patients, regardless of the cause. LOS was also
 231 defined commonly as the number of days in hospital admission due to COVID-19. One study
 232 described it as normal or prolonged, based on their measured average LOS (<17 days –
 233 normal; >17 - prolonged) (20). Consequently, they assessed LOS as a binary outcome. On the
 234 determinants of COVID-19 outcomes, identified risk factors were categorised into
 235 demographic, lifestyle, socioeconomic and biological/medical determinants, based on the
 236 determinants of health model (21). The rationale was to provide contextual analysis and
 237 identify common risk factors for COVID-19 outcomes. The biological/medical determinants
 238 encompassed as comorbidities, laboratory findings, and participants symptoms. The findings
 239 of the studies are synthesised below (Table 2).

240 Table 1: Study characteristics

Studies	Aim	Settings	Sample size	Sample characteristics	Study outcomes (definition)
Alaa et al. (2020) (22)	To investigate the influence of timing of hospital admission on risk of mortality for patients with COVID-19 in England	England	6068	Individuals with COVID-19 in the CHES database during the study period. Average age – 68 years, Men – 61%	Mortality (all cause-mortality)
Almazeedi et al. (2020) (23)	To examine the demographics, clinical manifestations, and outcomes in patients with COVID-19	Kuwait	1096	All COVID-19 patients admitted to Jaber Al Ahmad Al-Sabah hospital in Kuwait, with COVID-19 diagnosis based on WHO guideline and confirmed Polymerase Chain Reaction (PCR) test. Average age – 41 years, Men – 81%.	Mortality (in-hospital mortality)
Bello-Chavolla et al. (2020) (18)	To examine the association between diabetes and SARS-CoV-2 infection and its consequent clinical outcomes	Mexico	177,133	All confirmed, negative, and suspected COVID-19 cases in the Mexican MOH dataset. Average age – 46.7 years, Men – 57.7%	Mortality
Berenguer et al. (2020) (11)	To examine the predictors of death in patients with COVID-19 in Spain	Spain	4035	Males (61%) and females (39%) with COVID-19 confirmed by real-time PCR assay in 127 Spanish centres. Average age – 70 years, Men – 61%	Mortality (all-cause mortality)

Carrasco-Sánchez et al. (2020) (24)	To examine the association between blood glucose levels and in-hospital mortality in non-critically patients with COVID-19.	Spain	11,312	Patients ≥ 18 years with COVID-19 confirmed by Reverse Transcription (RT)-PCR and hospitalized from 1 March 2020 to 31 May 2020. Average age – 67.06 years, Men – 57.1%	Mortality (all-cause mortality during hospitalization)
Halalau et al. (2020) (25)	To provide risk assessment tools for patients with COVID-19.	USA	2025	Patients with positive COVID-19 on nasopharyngeal swabs at any Beaumont Health's eight emergency departments between 1 March 2020 and 1 April 2020. Men – 53.7%	Mortality (in-hospital mortality)
Kaeuffer et al. (2020) (26)	To explore risk factors of severe COVID-19 disease and mortality	France	1,045	Individuals ≥ 18 years with confirmed COVID19, hospitalised in Strasbourg and Mulhouse hospitals - March 2020. Average age – 66 years, Men – 69%	Mortality (in-hospital mortality)
Li et al. (2020) (10)	To investigate severity of COVID-19 outcomes	China	548	Individuals with COVID-19 admitted to Tongji Hospital from 26 January – 5 February 2020. Men – 50.9%.	Mortality (in-hospital mortality)
Okoh et al. (2020) (27)	To examine the clinical features of COVID-19 outcomes in Black/African American and Latino Hispanic	USA	251	Adults ≥ 18 -years admitted between March 10 and April 10, 2020. Men – 51%	Mortality (in-hospital death)
Petrilli et al.	To explore in-hospital COVID-	USA	5279	Laboratory confirmed COVID-19 patients with	Mortality

(2020) (28)	19 outcomes.			average age of 54. Men – 49.5%		
Sourij et al. (2020) (29)	To investigate predictors of in-hospital COVID-19 mortality in patients with prediabetes and diabetes.	Austria	238	People ≥ 18 years with confirmed COVID-19 and a with type 1 diabetes, type 2 diabetes, or prediabetes from 10 hospital sites in Austria. Average age – 71.1 years, Men – 63.6%	Mortality (in-hospital deaths)	
Wang et al. (2020) (30)	To examine the characteristics and prognosis of COVID-19 infections	China	293	Patients with COVID-19 diagnosis based on the NHC- China formulated “Diagnosis and treatment of novel coronavirus pneumonia” Men – 47.1%	Mortality	
Zhang et al. (2020) (31)	To examine the influence of D-dimer levels on COVID-19 mortality	China	343	Adults ≥ 18 years with laboratory-confirmed COVID-19 between 12 January and 15 March 15. Average age – 62 years, Men – 49.3%	Mortality (in-hospital mortality)	
Zhou et al. (2020) (9)	To investigate in-hospital COVID-19 risk factors.	China	191	≥ 18 years old adult inpatients with laboratory confirmed COVID-19 from Jinyintan and Wuhan Pulmonary Hospital. Men – 62%	Mortality (in-hospital death)	
Tartof et al. (2020) (19)	To determine the association between BMI and COVID-19 mortality.	USA	6916	Patients with COVID-19 from 13 February - 2 May 2020 from health care organisations located throughout 9 counties in Southern California. Men – 45%	Mortality (in-hospital death)	
Williamson et al. (2020)	To investigate risk factors of COVID-19 mortality.	England	10,926	≥ 18 years individuals with COVID-19 and currently registered as active patients in GP	Mortality (COVID-19 related death)	

(32)					surgery. Men –49.9%	
Grasselli et al. (2020) (17)	To determine risk factors associated with COVID-19 ICU mortalities.	Italy	3988	Critically ill patients with laboratory-confirmed COVID-19. Average age - 63 years, males- 79.9%	Mortality (COVID-19 death)	
Mikami et al. (2020) (33)	To examine factors associated with COVID-19 mortality.	USA	6493	Patients with laboratory confirmed COVID-19 with from one of the 8 hospitals in New York City metropolitan. Average age 59 years, males - 54.5%	Mortality (in-hospital mortality)	
Albitar et al. (2020) (16)	To explore predictors of COVID-19 mortality among patients from worldwide open access data	Africa Asia America Australia Europe	828	COVID-19 patients with definite outcomes. Average age - 49.4 years, male majority – 59.1%, and majority located in Asia – 69.3%	Mortality)	
Wu et al. (2020) (34)	To examine factors associated with longer length of COVID-19 hospital stay	China	58	Patients with COVID-19 and hospitalised in Qiaokou Fangcang Hospital. Average age – 55.5 years, Men – 37.9%	LOS (number of days spent on admission)	
Guo et al. (2020) (20)	To investigate determinants of COVID-19 prolonged hospital length of stay	China	75	Patients with laboratory-confirmed COVID-19 and discharged from 20 January – 16 March 2020. Average age – 47 years, Men – 57%	LOS (<17 days median Los-normal; >17days median Los prolonged)	

Mendy et al. (2020) (35) To identify factors associated with COVID-19 hospitalization and mortality among ethnically diverse cohort. USA 689 Patients with COVID-19 confirmed with a RT-PCR from the University of Cincinnati health system between 13 March – 31 May 2020. Average age – 49.5 years, Men – 53% Mortality: (COVID-19 death during hospitalisation). LOS: (number of days hospitalised)

Table 2: Synthesised review findings

Variable category	Determinants	Variable description	No. of studies reporting sign. with COVID-19	Range of effect sizes HR/OR/AOR/ β 95% C.I.
-------------------	--------------	----------------------	--	---

			mortality	Low high	high	low
Demographic variables	Age (older/increasing)	Years lived	16(16[+], 0[-])	1.04 22.68	20.6	1.01
	Sex/gender (male)	Male or female	8(8[+], 0[-])	1.15 2.80	1.70	1.00
	Geographic location (America)	Africa/Asia/Australia/America/Europe	7.44	3.55		15.62
Biological/medical variables	CKD (present)	Present or not	5(5[+], 0[-])	1.99 11.08	4.48	1.09
	C-reactive protein (elevated- >5mg/L)	Level of C-reactive protein in blood	4(4[+], 0[-])	1.01 3.20	2.00	1.00
	Diabetes (present)	Present or not	6(6[+], 0[-])	1.18 36.72	12.23	1.00
	Hypertension (present)	Present or not	4(4[+], 0[-])	1.14 7.55	3.58	1.01
	Dyspnoea (present)	Present or not	3(3[+], 0[-])	1.45 3.4	2.1	1.2
	Cancer (present)	Present or not	3(3[+], 0[-])	1.3 2.95	2.46	1.1
	COPD (present)	Present or not	3(3[+], 0[-])	1.27 2.19	1.68	1.08
	Coronary heart disease (present)	Present or not	2(2[+], 0[-])	1.77 17.79	2.14	0.26
	BMI (obesity)	Normal weight/overweight/obese	4(4[+], 0[-])	1.25 8.26	4.18	1.17
	Asthma (present)	Present or not	2(2[+], 0[-])	1.55 23.44	4.29	1.03
	D-dimer ((≥2.0 µg/ml)	Present or not	2(2[+], 0[-])	1.19 175.7	22.4	1.02

252

	Hospitalisation before onset of COVID-19 symptoms	Time of admission relative to symptom onset	1(0[+], 1[-])	0.52 0.61	0.45
Lifestyle variables	Smoking (current smoker)	Smoker or non-smoker	2(2[+], 0[-])	0.89 10.09 83.40	0.28
Socioeconomic variables	Index of multiple deprivation (greatest)	Least to greatest	1(1[+], 0[-])	1.79 1.91	1.68

Variable category	Determinants	Variable description	No. of studies reporting sign. with COVID-19 LOS	*Range of effect sizes HR/OR/AOR 95%C.I.			
Demographic variables	Sex (Male)	Male or female	2(1[+], 1[-])	0.19	0.39	0.05	0.63
Biological variables	Diabetes (present)	Present or not	2(2[+], 0[-])	0.50	3.2	-0.2	0.74
	Fever (present)	Present or not	2(2[+], 0[-])	3.5	8.27	1.39	72.16
	Bilateral pneumonia (present)	Present or not	1(1[+], 0[-])	3.4		0.49	6.25
	CKD	Present or not	1(1[+], 0[-])	3.73		1.95	145.4
	COPD	Present or not	1(1[+], 0[-])	0.45		0.11	0.79
	Asthma	Present or not	1(1[+], 0[-])	0.50		0.20	0.81

Age and sex were the common demographic risk factors. Of the 20 studies on age and COVID-19 mortality, 16 identified increasing age as a significant determinant of COVID-19 mortality, with effect sizes ranging from 1.04 to 20.6 and 95% CI from 1.01 to 22.68 (9, 10, 11, 16, 17, 18, 24, 25, 26, 27, 28, 29, 30, 32, 33, 35). Also, 8 of the 20 studies on gender/sex and COVID-19 found men to have a higher risk of COVID-19 mortality than women (10, 11, 16, 17, 24, 26, 28, 32). On LOS, 2 of the 3 studies assessing LOS found women (AOR=0.19, 95%CI=0.05-0.63) (20) and men (β =0.39, 95% CI=0.16-0.62) (35) as determinants of COVID-19 duration of hospitalisation.

Of biological/medical risk factors, the review identified diabetes (n=6), Chronic kidney/renal disease (CKD) (n=5), hypertension (n=4), C-reactive protein (CRP) (n=4), BMI (n=4), dyspnoea (n=3), COPD (n=3), cancer (n=3), coronary heart disease (n=2), asthma (n=2) and D-dimer (n=2) as determinants of COVID-19 mortality. Of the 10 studies that included CKD in their analysis, 5 found it a significant determinant of COVID-19 mortality (18, 25, 26, 30, 35). Additionally, out of the 5 studies that investigated the influence of CRP on COVID-19 mortality, 4 showed that elevated CRP in the blood (at least >5mg/L) increases the risk of COVID-19 death (11, 24, 26, 29). Also, people with diabetes were found to have a higher risk of COVID-19 mortality in 6 of the 18 studies on diabetes and COVID-19 mortality (16, 17, 18, 26, 30, 32)

Like diabetes, hypertension was also identified as a determinant of COVID-19 mortality in 4 out of the 16 studies (11, 16, 24, 30). Furthermore, 4 of 11 studies on BMI and COVID-19 mortality showed that obesity significantly determines COVID-19 mortality (11, 18, 19, 32). For LOS, fever and diabetes were associated with prolonged LOS (43,35).

On lifestyle factors, smoking was the only assessed determinant of COVID-19 outcomes. It was identified as a risk factor for COVID-19 mortality in 2 of the eleven studies that explored it, with effect size ranging from 0.89-10.09, 95% CIs from 0.28-83.40 (23, 32). None of the studies on LOS reported a significant association between smoking and COVID-19 LOS. Finally, the only study on socioeconomic determinants and COVID-19 reported that greater deprivation determines COVID-19 mortality (HR=1.79, 95% 1.68 – 1.91) (32)

Discussion

This review aimed to investigate the determinants of COVID-19 outcomes. The review found that the specification and subsequent analysis of most of the determinants differed across the studies. For example, Berenguer et al. (2020) described elevated C-reactive protein (CRP) as CRP>5mg/L, while Carrasco-Sánchez et al. (2020) described it as >60mg/L. Also, whilst Almazeedi et al. (2020) and Sourij et al. (2020) assessed CRP as a continuous variable, Kaeuffer et al. (2020) categorised it into two groups: CRP- 100-199mg/L and CRP≥200mg/L. Like CRP, older age was also specified differently across the studies. Bello-Chavolla et al. (2020) and Li et al. (2020) described it as individuals ≥65 years, Zhang et al. (2020) termed it as persons >65 years while Petrilli et al. (2020) defined it as people ≥75 years. Apart from these variations, the study settings also differed across the papers. These contextual differences, which could include disparities in access to healthcare, crowded living, could have inherently influenced the studies' findings (36). Despite these heterogeneities, the result of some of the determinants was similar across the studies. For instance, the significant association between age and COVID-19 mortality was

reported across sixteen out of twenty papers, and four out of five articles examining CRP also showed a significant association between CRP and COVID-19 mortality. Regardless of the differences in methods, these findings could have public health implications for populations worldwide (37). Thus, they can be considered during the planning and implementation of effective policies for COVID-19.

Nonetheless, there were other contrasting findings. For example, Mendy et al. (2020) indicated that men are more likely to stay longer in hospitals due to COVID-19 than women. Conversely, Guo et al. (2020) showed that women are more associated with prolonged COVID-19 hospitalisation than men. Both studies had men dominating their study participants, 53% and 57% respectively; but the proportion of men in Guo et al. (2020) were marginally higher. However, in absolute figures, Mendy et al. (2020) included more male participants (n=365) than Guo et al. (2020) (n=43). Therefore, these sample size variations could account for the differences in their sex and LOS findings due to COVID-19. Moreover, these findings are from only two studies; thus, they may not be enough to conclude the association between sex and COVID-19 LOS. Further discussions on the review findings, based on the determinants of health model, are provided below.

Demographic determinants of COVID-19 outcomes

The underlying mechanism for the association between older age and COVID-19 mortality is unclear; however, several studies indicate that decrease in immune responses coupled with increase comorbid burden with ageing may account for this observation (9, 38, 39). Another study further explained that age-related changes or defects in the immune system, particularly significant defect in cell-mediated immunity, primarily affects immune responses to diseases (40). Also, evolution and ageing theories, like the antagonistic pleiotropy theory, postulate that even beneficial genes at an early age may be less efficient or deleterious with increasing

age, and this may inherently increase susceptibility to previously shielded diseases (41, 42). Moreover, current evidence suggests that increasing age is a common risk factor for several health outcomes, like mortalities and morbidities (43-46).

Even though ageing generally decreases immune responses to diseases and infections, the innate human response is mostly safeguarded (40). Thus, many individual and environmental factors may account for the relationship between ageing and disease outcomes, such as COVID-19 mortality. These factors may include nutritional deficiencies, decreased functionality, exposure to pathogens, vaccinations, an individual's lived experiences and genetic make-up and access to health care (47). Furthermore, there are reports on good COVID-19 prognosis in elderly patients (48). Therefore, it may be imperative to understand how these factors cumulatively affect the immune system, and further mediate ageing and decreased immune system relationship to provide exhaustive literature on the subject. Other studies have also reported severe COVID-19 consequences in children (49, 50). Consequently, there is a need for studies to focus more on children, as much as they have on the adult population to offer a balanced argument to inform COVID-19 and ageing policies.

Like ageing, studies also attribute the sex disparities regarding COVID-19 mortality to sex-based differences in immunological responses to viral infections (51, 52). The X sex chromosome has encoded immune regulatory genes that decrease viral infections' susceptibility (51). Since women have twice X-chromosomes to men, they tend to have higher innate immunity to viral infections, like COVID-19, and by extension a lower risk of severe COVID-19 outcomes than men (51, 53). Similarly, in contrast to oestrogen, testosterone has an immunosuppressive effect; so, it attenuates men's immune responses to viral infections (54). Additionally, it is reported that men are genetically more predisposed to

produce higher levels interleukin (IL)-6, which are unfavourable to longevity, compared to women (55). Apart from these biological reasons accounting for the sex differences in COVID-19 mortality risk, behavioural and lifestyle factors like smoking and alcohol consumption, have been implicated in the gender disparities in COVID-19 outcomes. Data indicate that men are more likely to engage in these lifestyle factors that increase the risk of COVID-19 deaths than women (52). Women are more likely to comply with COVID-19 precautionary measures than men and are more likely to remain confined than men (56). Regarding LOS, the evidence is not enough to indicate whether sex determines longer COVID-19 hospitalisation.

Biological/medical determinants of COVID-19 outcomes

The biological/medical determinants in this review were comorbidities, symptoms, and laboratory findings of the included studies participants. The biological determinants of COVID-19 outcomes included CKD, C-reactive protein, diabetes, hypertension, obesity, cancer, COPD, dyspnea, asthma, and coronary heart disease. Clinical data reveal that chronic conditions, such as the above, decrease innate immune responses in humans (57, 58). For instance, metabolic diseases/disorders, like diabetes, attenuates immunity and increase risk to infections by weakening lymphocyte and macrophage activities (59). Moreover, these chronic conditions are associated with increase pro-inflammatory cytokines resulting from dysregulation of systems, like the hypothalamic-pituitary-adrenal and sympathetic nervous system (58, 60). The accumulation of pro-inflammatory cytokines subsequently impairs systemic and cellular immune functions (57, 61).

Furthermore, studies hypothesize that the use of renin-angiotensin-aldosterone system (RAAS) inhibitors, like angiotensin-converting enzyme-2 (ACE-2), in the management of some of these chronic conditions increase COVID-19 infectivity (62, 63). This is because ACE2 also functions as a receptor for the COVID-19 virus (64). This RAAS and ACE-2 hypothesis recently sparked debate and discourse on gold standard medical management of comorbidities in COVID-19 patients. One study indicated that sudden discontinuation of ACE-2 might have far worse consequences for high-risk-COVID-19 patients, particularly those with cardiovascular conditions, like myocardial infarction (65). Their argument is hinged on the paucity of human studies to corroborate the RAAS and ACE-2 theory. Additionally, experimental studies in mice showed that ACE-2 downregulation facilitates lung injuries and increases viral loads (60, 66). Thus, several human studies are needed to substantiate the harmful effect, or otherwise, of RAAS inhibitors in the management of COVID-19 patients with comorbidities.

Lifestyle determinants of COVID-19 outcomes

The included studies examined only smoking as a lifestyle determinant of COVID-19 mortality. The association between smoking and COVID-19 mortality is biologically plausible because smoking is a risk factor for several conditions, like coronary heart disease and Chronic Obstructive Pulmonary Disease (COPD), that are associated with severe COVID-19 outcomes (67). Also, a cohort study with an average of 9.6 years follow-up by showed that 11% (men) and 13% (women) of pneumonia and COPD deaths were attributable to smoking (68). Additionally, the Centre for Disease Control and Prevention (CDC) also report that smoking is associated with about 113,000 respiratory deaths each year in the United States (69). Since COVID-19 is a respiratory infection and based on the above

evidence on smoking-related respiratory deaths, it may be reasonable to make projections that smoking may be significantly associated with severe COVID-19 outcomes.

Furthermore, data shows that smokers have increased upregulation or expression of ACE-2, the reported enzyme receptor for SARS-CoV-2 (COVID-19) (70), which may increase their risk of severe COVID-19 outcomes compared to non-smokers. A single-cell sequencing experiment further demonstrated that cigarette smoking upregulates ACE-2 in humans' lungs and increases their susceptibility to COVID-19 infections (71). They inferred that smoking cessation could reduce ACE-2 expression and thereby reduce the risk of COVID-19 disease. Thus, their findings advance the above argument on the benefit or otherwise of ACE-2 dysregulation in humans in reducing the burden of COVID-19. Consequently, systematic reviews and meta-analyses of several high-quality studies on ACE-2 and COVID-19 are required to provide empirical evidence to inform policy and clinical practice. All the same, this review is limited in drawing a meaningful conclusion on the association between smoking and COVID-19 mortality because only one of the included studies identified smoking as a risk factor of COVID-19 mortality.

Policy implications

This review findings re-enforce the need for health systems to continue the testing, tracing, and isolation policies to reduce the spread of the virus and subsequently decrease the burden of the pandemic, particularly for high-risk individuals identified in this review. Additionally, countries, such as Nigeria, that are yet to implement crucial public health policies, like immunisation and routine COVID-19 testing, can draw lessons from countries like the UK that have benefited immensely from such policies. Although this may come with increased direct costs, especially the cost of PPEs and testing equipment, the indirect benefits to

populations, such as reduced disease burden and improved productivity, might be enormous for these countries.

However, lockdown policies, which seems to be the go-to policy for most countries, must be evaluated holistically, to ascertain their overall benefits, especially as the pandemic continues to rage. Evidence shows that lockdowns are beneficial when introduced at an earlier stage of an outbreak than later. For example, evaluating the national lockdown response in Norway, USA, Argentina, and the UK shows that the lockdown's timing and not the lockdown itself significantly reduced the burden of the outbreak in Argentina and Norway compared to the UK and the USA. For instance, the UK suffered significant health and economic recession due to the delays in the lockdown response to the viral outbreak. Their delay in implementing the first lockdown resulted in several other lockdowns that have had further economic implications. Therefore, lockdowns must be implemented earlier to prevent dire health and economic consequences.

Even the timing of lockdowns alone may not be enough to radically reduce viral transmission rate and consequently limit the probability of infections in high-risk individuals because data from other jurisdictions that introduced earlier national lockdowns, like Ghana, showed a steady increase in transmissions during the lockdown and a rapid rise few weeks post lockdown. Also, it is still crippling with the lockdown induced economic downfall. This suggests that lockdowns as single policies may be ineffective in plummeting COVID-19 infections, mortalities, and financial hardships. Consequently, global health systems must also place a premium on other equally important policies, like robust testing and tracing. Currently, most countries do not conduct follow-up COVID-19 testing following negative test results at entry borders. A negative test result on arrival at entry borders may not be enough to declare individuals as virus-free since it can take fourteen days maximum for the virus to be detected, per the WHO guidelines. Thus, governments must apply both on arrival

and fourteen days post border arrival COVID-19 testing to ensure comprehensive identification of all positive cases to curtail the virus's potential spread.

Additionally, it will be crucial for countries, like the USA and Brazil, to implement the fourteen-day quarantine policies for all border entries, as it has shown to be effective reducing the COVID-19 impact in places like Argentina. Countries with partial quarantine policies, such as Ghana, will benefit from instituting mandatory quarantine for all border arrivals at government selected facilities since self-isolation education and passenger locator forms may be inadequate in reducing the viral spread. Furthermore, health systems must continuously promote behavioural change interventions to establish significant control over the viral outbreak. Enforcing compliance to behavioural change interventions, like social distancing, nose mask use, social hand washing and cough etiquette, can be the significant catalyst needed to decrease the pandemic's impact. Finally, the continuous increase in the COVID-19 cases and mortality indicate the need for an urgent review of current health policies at both international and national levels to implement suitable context-specific interventions to mitigate the COVID-19 menace effectively.

Concerning this study's strength, this is the first systematic review synthesising the evidence on LOS determinants, to the best of the researchers' knowledge. Therefore, it presents novel findings that could initiate useful interventions to address the burden of prolonged hospitalisation associated with COVID-19. It could also cause a paradigm shift and ensure holistic research coverage on all COVID-19 outcomes. Moreover, this is the first review to provide a comprehensive investigation of contextual determinants of COVID-19 outcomes, based on the determinants of health model. It identified crucial gaps in the literature on the determinants of COVID-19 outcomes that require urgent attention. Additionally, this review

evaluated current public health policies and suggested strategies to augment area-specific efforts at curbing the COVID-19 problem. Regarding limitations, the review was restricted in conducting further analysis, specifically, meta-analysis, to precisely estimate the associations' effect size due to included studies' heterogeneity. Also, most of the included studies (n=12) used retrospective design, thus, there was the possibility of residual confounders that could influence this review's findings. Additionally, all of them used secondary data from medical records of participants. Therefore, any omission or data entry error could affect their results and this review. Hence, caution must be taken when interpreting the findings of this review.

Conclusion and recommendations

This study's overarching aim was to examine the determinants of COVID-19 outcomes. The review findings showed that increasing age and comorbidities are more likely to determine COVID-19 outcomes. Thus, policies, like routine COVID-19 testing and prioritised vaccination to shield these high-risk individuals, must be sustained and extended to other populations yet to implement such important policies. Most importantly, health systems must continually review existing policies to ensure their context-specific relevance, especially with the emergence of a new viral variant and the rapid surge in cases. Based on this review, the authors recommend that future studies also focus on determinants of COVID-19 LOS. Additionally, studies should explore the determinants of COVID-19 outcomes in low-income-countries to ensure holistic and context-specific evidence on risk factors of COVID-19 mortality in the literature.

496

497

498 REFERENCES

- 499 1. WHO. Coronavirus Disease (COVID-19) Dashboard. 2021. Available from:
500 <https://covid19.who.int/>. (Accessed on 10/01/2021).
- 501 2. Azamfirei R. The 2019 novel coronavirus: a crown jewel of pandemics? The Journal
502 of Critical Care Medicine. 2020 Jan 31;6(1):3-4.
- 503 3. Nishiura H, Kobayashi T, Miyama T, Suzuki A, Jung SM, Hayashi K. et al.
504 Estimation of the asymptomatic ratio of novel coronavirus infections (COVID-19).
505 International journal of infectious diseases. 2020 May; 94:154.
- 506 4. Yuan J, Li M, Lv G, Lu ZK. Monitoring transmissibility and mortality of COVID-19
507 in Europe. International Journal of Infectious Diseases. 2020 Jun 1; 95:311-5.
508 <https://doi.org/10.1016/j.ijid.2020.03.050>.
- 509 5. Promislow DE. A geroscience perspective on COVID-19 mortality. The Journals of
510 Gerontology: Series A. 2020 Sep;75(9): e30-3.
511 <https://doi.org/10.1093/gerona/glaa094>.
- 512 6. Maia Chagas A, Molloy JC, Prieto-Godino LL, Baden T. Leveraging open hardware
513 to alleviate the burden of COVID-19 on global health systems. PLoS biology. 2020
514 Apr 24;18(4): e3000730.
- 515 7. Ornell F, Schuch JB, Sordi AO, Kessler FH. “Pandemic fear” and COVID-19: mental
516 health burden and strategies. Brazilian Journal of Psychiatry. 2020 Jun;42(3):232-5.
- 517 8. Amponsah R, Frimpong IA. Ghana in the Face of COVID-19: Economic Impact of
518 Coronavirus (2019-NCOV) Outbreak on Ghana. Open Journal of Business and
519 Management. 2020 Jun 17;8(04): 1404.
520 <http://www.scirp.org/journal/Paperabs.aspx?PaperID=100937>.
- 521 9. Zhou F, Yu T, Du R, Fan G, Liu Y, Liu Z. et al. Clinical course and risk factors for
522 mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort
523 study. The lancet. 2020 Mar 28;395(10229):1054-62.
- 524 10. Li X, Xu S, Yu M, Wang K, Tao Y, Zhou Y. et al. Risk factors for severity and
525 mortality in adult COVID-19 inpatients in Wuhan. Journal of Allergy and Clinical
526 Immunology. 2020 Jul 1;146(1):110-8.

11. Berenguer J, Ryan P, Rodríguez-Baño J, Jarrín I, Carratalà J, Pachón J. et al. Characteristics and predictors of death among 4035 consecutively hospitalized patients with COVID-19 in Spain. *Clinical Microbiology and Infection*. 2020 Nov 1;26(11):1525-36.
12. Sepandi M, Taghdir M, Alimohamadi Y, Afrashteh S, Hosamirudsari H. Factors Associated with Mortality in COVID-19 Patients: A Systematic Review and Meta-Analysis. *Iranian journal of public health*. 2020; 49(7), 1211–1221.
13. Mesas AE, Cavero-Redondo I, Álvarez-Bueno C, Sarriá Cabrera MA, Maffei de Andrade S, Sequí-Dominguez I. et al. Predictors of in-hospital COVID-19 mortality: A comprehensive systematic review and meta-analysis exploring differences by age, sex and health conditions. *PloS one*. 2020 Nov 3;15(11): e0241742
14. Zeng X, Zhang Y, Kwong JS, Zhang C, Li S, Sun F. et al. The methodological quality assessment tools for preclinical and clinical studies, systematic review and meta-analysis, and clinical practice guideline: a systematic review. *Journal of evidence-based medicine*. 2015 Feb;8(1):2-10
15. Aromataris E, Munn Z (Editors). *JBIM Manual for Evidence Synthesis*. JBI, 2020. Available from <https://synthesismanual.jbi.global>. <https://doi.org/10.46658/JBIMES-20-01.#>.
16. Albitar O, Ballouze R, Ooi JP, Ghadzi SM. Risk factors for mortality among COVID-19 patients. *Diabetes research and clinical practice*. 2020 Aug 1; 166:108293.
17. Grasselli G, Greco M, Zanella A, Albano G, Antonelli M, Bellani G. et al. Risk factors associated with mortality among patients with COVID-19 in intensive care units in Lombardy, Italy. *JAMA internal medicine*. 2020 Oct 1;180(10):1345-55.
18. Bello-Chavolla OY, Bahena-López JP, Antonio-Villa NE, Vargas-Vázquez A, González-Díaz A, Márquez-Salinas A. et al. Predicting mortality due to SARS-CoV-2: A mechanistic score relating obesity and diabetes to COVID-19 outcomes in Mexico. *The Journal of Clinical Endocrinology & Metabolism*. 2020 Aug;105(8):2752-61.
19. Tartof SY, Qian L, Hong V, Wei R, Nadjafi RF, Fischer H. et al. Obesity and mortality among patients diagnosed with COVID-19: results from an integrated health care organization. *Annals of internal medicine*. 2020 Nov 17;173(10):773-81.
20. Guo A, Lu J, Tan H, Kuang Z, Lou Y, Yang T. et al. Risk Factors on Admission Associated with Hospital Length of Stay in Patients with COVID-19: A Retrospective Cohort Study. Available at SSRN 3582769. 2020 Apr 20.

21. Dahlgren G., Whitehead M. Tackling social inequalities in health. 1993. Available from: https://www.euro.who.int/_data/assets/pdf_file/0018/103824/E89384.pdf.
22. Alaa A, Qian Z, Rashbass J, Benger J, van der Schaar M. Retrospective cohort study of admission timing and mortality following COVID-19 infection in England. *BMJ open*. 2020 Nov 1;10(11):e042712.
23. Almazeedi S, Al-Youha S, Jamal MH, Al-Haddad M, Al-Muhaini A, Al-Ghimlas F. et al. Characteristics, risk factors and outcomes among the first consecutive 1096 patients diagnosed with COVID-19 in Kuwait. *EClinicalMedicine*. 2020 Jul 1; 24:100448.
24. Carrasco-Sánchez FJ, López-Carmona MD, Martínez-Marcos FJ, Pérez-Belmonte LM, Hidalgo-Jiménez A, Buonaiuto V. et al. Admission hyperglycaemia as a predictor of mortality in patients hospitalized with COVID-19 regardless of diabetes status: data from the Spanish SEMI-COVID-19 Registry. *Annals of medicine*. 2021 Jan 1;53(1):103-16.
25. Halalau A, Imam Z, Karabon P, Mankuzhy N, Shaheen A, Tu J. et al. External validation of a clinical risk score to predict hospital admission and in-hospital mortality in COVID-19 patients. *Annals of medicine*. 2021 Jan 1;53(1):78-86.
26. Kaeuffer C, Le Hyaric C, Fabacher T, Mootien J, Dervieux B, Ruch Y. et al. Clinical characteristics and risk factors associated with severe COVID-19: prospective analysis of 1,045 hospitalised cases in North-Eastern France, March 2020. *Eurosurveillance*. 2020 Dec 3;25(48):2000895.
27. Okoh AK, Sossou C, Dangayach NS, Meledathu S, Phillips O, Raczek C. et al. Coronavirus disease 19 in minority populations of Newark, New Jersey. *International journal for equity in health*. 2020 Dec;19(1):1-8.
28. Petrilli CM, Jones SA, Yang J, Rajagopalan H, O'Donnell L, Chernyak Y. et al. Factors associated with hospital admission and critical illness among 5279 people with coronavirus disease 2019 in New York City: prospective cohort study. *Bmj*. 2020 May 22;369.
29. Sourij H, Aziz F, Bräuer A, Ciardi C, Clodi M, Fasching P. et al. COVID-19 fatality prediction in people with diabetes and prediabetes using a simple score upon hospital admission. *Diabetes, Obesity and Metabolism*. 2020 Nov 16.
30. Wang Z, Ye D, Wang M, Zhao M, Li D, Ye J. et al. Clinical Features of COVID-19 Patients with Different Outcomes in Wuhan: A Retrospective Observational Study. *BioMed research international*. 2020 Oct 5;2020.

31. Zhang L, Yan X, Fan Q, Liu H, Liu X, Liu Z. et al. D-dimer levels on admission to predict in-hospital mortality in patients with Covid-19. *Journal of Thrombosis and Haemostasis*. 2020 Jun;18(6):1324-9.
32. Williamson EJ, Walker AJ, Bhaskaran K, Bacon S, Bates C, Morton CE. et al. OpenSAFELY: factors associated with COVID-19 death in 17 million patients. *Nature*. 2020 Jul 8; 584(7821):430-436. DOI: [10.1038/s41586-020-2521-4](https://doi.org/10.1038/s41586-020-2521-4)
33. Mikami T, Miyashita H, Yamada T, Harrington M, Steinberg D, Dunn A. et al. Risk factors for mortality in patients with COVID-19 in New York City. *Journal of general internal medicine*. 2020 Jun 30:1-0.
34. Wu S, Xue L, Legido-Quigley H, Khan M, Wu H, Peng X. et al. Understanding factors influencing the length of hospital stay among non-severe COVID-19 patients: A retrospective cohort study in a Fangcang shelter hospital. *Plos one*. 2020 Oct 21;15(10): e0240959.
35. Mendy A, Apewokin S, Wells AA, Morrow AL. Factors associated with hospitalization and disease severity in a racially and ethnically diverse population of COVID-19 patients. *MedRxiv*. 2020 Jan 1.
36. van Bavel JJ, Mende-Siedlecki P, Brady WJ, Reinero DA. Contextual sensitivity in scientific reproducibility. *Proceedings of the National Academy of Sciences*. 2016 Jun; 113 (23) 6454-6459; DOI: [10.1073/pnas.1521897113](https://doi.org/10.1073/pnas.1521897113)
37. Ioannidis JP, Axfors C, Contopoulos-Ioannidis DG. Population-level COVID-19 mortality risk for non-elderly individuals overall and for non-elderly individuals without underlying diseases in pandemic epicenters. *Environmental research*. 2020 Sep 1; 188:109890.
38. Mueller AL, McNamara MS, Sinclair DA. Why does COVID-19 disproportionately affect older people? *Aging (Albany NY)*. 2020 May 31;12(10):9959.
39. Chinnadurai R, Ogedengbe O, Agarwal P, et al. Older age and frailty are the chief predictors of mortality in COVID-19 patients admitted to an acute medical unit in a secondary care setting- a cohort study. *BMC Geriatrics*. 2020 Oct;20(1):409. DOI: [10.1186/s12877-020-01803-5](https://doi.org/10.1186/s12877-020-01803-5).
40. Opal SM, Girard TD, Ely EW. The immunopathogenesis of sepsis in elderly patients. *Clinical infectious diseases*. 2005 Nov 15;41(Supplement_7): S504-12.
41. Mitteldorf J. What is antagonistic pleiotropy? *Biochemistry (Moscow)*. 2019 Dec;84(12):1458-68. <https://doi.org/10.1134/S0006297919120058>

42. Williams PD, Day T. Antagonistic pleiotropy, mortality source interactions, and the evolutionary theory of senescence. *Evolution*. 2003 Jul;57(7):1478-88.
43. Australian Institute of Health. Social Distribution of Health Risks and Health Outcomes: Preliminary Analysis of the National Health Survey 2007-08. AIHW; 2012.
44. Eguchi T, Bains S, Lee MC, Tan KS, Hristov B, Buitrago DH. et al. Impact of increasing age on cause-specific mortality and morbidity in patients with stage I non-small-cell lung cancer: a competing risks analysis. *Journal of Clinical Oncology*. 2017 Jan 20;35(3):281.
45. Johansson S, Rosengren A, Young K, Jennings E. Mortality and morbidity trends after the first year in survivors of acute myocardial infarction: a systematic review. *BMC Cardiovascular Disorders*. 2017 Dec;17(1):1-8.
46. Niccoli T, Partridge L. Ageing as a risk factor for disease. *Current biology*. 2012 Sep 11;22(17): R741-52.
47. Hernández DS, Verdecia BG. Immunosenescence: effects of aging process on immune system. *Revista Cubana de Hematología, Inmunología y Hemoterapia*. 2014;30(4):332-45.
48. Ma Y, Hou L, Yang X, Huang Z, Yang X, Zhao N. et al. The association between frailty and severe disease among COVID-19 patients aged over 60 years in China: a prospective cohort study. *BMC medicine*. 2020 Dec;18(1):1-8.
49. Liu J, Zhang S, Wu Z, Shang Y, Dong X, Li G. et al. Clinical outcomes of COVID-19 in Wuhan, China: a large cohort study. *Annals of intensive care*. 2020 Dec;10(1):1-21.
50. Qiu H, Wu J, Hong L, Luo Y, Song Q, Chen D. Clinical and epidemiological features of 36 children with coronavirus disease 2019 (COVID-19) in Zhejiang, China: an observational cohort study. *The Lancet Infectious Diseases*. 2020 Jun 1;20(6):689-96.
51. Conti P, Younes A. Coronavirus COV-19/SARS-CoV-2 affects women less than men: clinical response to viral infection. *J Biol Regul Homeost Agents*. 2020 Apr 7;34(2):339-43.
52. Bwire GM. Coronavirus: why men are more vulnerable to Covid-19 than women?. *SN comprehensive clinical medicine*. 2020 Jul;2(7):874-6. <https://doi.org/10.1007/s42399-020-00341-w>.
53. Elgendy IY, Pepine CJ. Why are women better protected from COVID-19: Clues for men? Sex and COVID-19. *International journal of cardiology*. 2020 Sep 15; 315:105-6.

54. Pradhan A, Olsson PE. Sex differences in severity and mortality from COVID-19: are males more vulnerable?. *Biology of sex Differences*. 2020 Dec;11(1):1-1.
55. Marcon G, Tettamanti M, Capacci G, Fontanel G, Spanò M, Nobili A. et al. COVID-19 mortality in Lombardy: the vulnerability of the oldest old and the resilience of male centenarians. *Aging (Albany NY)*. 2020 Aug 15;12(15):15186.
56. De La Vega R, Ruíz-Barquín R, Boros S, Szabo A. Could attitudes toward COVID-19 in Spain render men more vulnerable than women? *Global public health*. 2020 Sep 1;15(9):1278-91.
57. Odegaard JI, Chawla A. Connecting type 1 and type 2 diabetes through innate immunity. *Cold Spring Harbor perspectives in medicine*. 2012 Mar 1;2(3): a007724.
58. Schulkin J, editor. *Allostasis, homeostasis, and the costs of physiological adaptation*. Cambridge University Press; 2004 Oct 25.
59. Oates JR, McKell MC, Moreno-Fernandez ME, Damen MS, Deepe Jr GS, Qualls JE. et al. Macrophage function in the pathogenesis of non-alcoholic fatty liver disease: the mac attack. *Frontiers in immunology*. 2019 Dec 12; 10:2893.
60. Yang P, Gu H, Zhao Z, Wang W, Cao B, Lai C. et al. Angiotensin-converting enzyme 2 (ACE2) mediates influenza H7N9 virus-induced acute lung injury. *Scientific reports*. 2014 Nov 13;4(1):1-6.
61. McEwen BS. Sex, stress and the hippocampus: allostasis, allostatic load and the aging process. *Neurobiology of aging*. 2002 Sep 1;23(5):921-39.
62. Ferrario CM, Jessup J, Chappell MC, Averill DB, Brosnihan KB, Tallant EA. et al. Effect of angiotensin-converting enzyme inhibition and angiotensin II receptor blockers on cardiac angiotensin-converting enzyme 2. *Circulation*. 2005 May 24;111(20):2605-10.
63. Watkins J. Preventing a covid-19 pandemic. *BMJ*. 2020;368:m810. <https://doi.org/10.1136/bmj.m810>.
64. Hoffmann M, Kleine-Weber H, Schroeder S, Krüger N, Herrler T, Erichsen S. et al. SARS-CoV-2 cell entry depends on ACE2 and TMPRSS2 and is blocked by a clinically proven protease inhibitor. *cell*. 2020 Apr 16;181(2):271-80.
65. Vaduganathan M, Vardeny O, Michel T, McMurray JJ, Pfeffer MA, Solomon SD. Renin–angiotensin–aldosterone system inhibitors in patients with Covid-19. *New England Journal of Medicine*. 2020 Apr 23;382(17):1653-9.

66. Kuba K, Imai Y, Rao S, Gao H, Guo F, Guan B. et al. A crucial role of angiotensin converting enzyme 2 (ACE2) in SARS coronavirus-induced lung injury. *Nature medicine*. 2005 Aug;11(8):875-9.
67. Reddy RK, Charles WN, Sklavounos A, Dutt A, Seed PT, Khajuria A. The effect of smoking on COVID-19 severity: A systematic review and meta-analysis. *Journal of Medical Virology*. 2021 Feb;93(2):1045-56.
68. Katanoda K, Marugame T, Saika K, Satoh H, Tajima K, Suzuki T. et al. Population attributable fraction of mortality associated with tobacco smoking in Japan: a pooled analysis of three large-scale cohort studies. *Journal of epidemiology*. 2008;0812050049-.
69. Centre for Disease Control and Prevention. Tobacco related Mortality. 2020. Available from: https://www.cdc.gov/tobacco/data_statistics/fact_sheets/health_effects/tobacco_related_mortality/index.htm
70. Leung JM, Yang CX, Tam A, Shaipanich T, Hackett TL, Singhera GK. et al. ACE-2 expression in the small airway epithelia of smokers and COPD patients: implications for COVID-19. *European Respiratory Journal*. 2020 May 1;55(5).
71. Smith JC, Sausville EL, Girish V, Yuan ML, Vasudevan A, John KM, Sheltzer JM. Cigarette smoke exposure and inflammatory signalling increase the expression of the SARS-CoV-2 receptor ACE2 in the respiratory tract. *Developmental cell*. 2020 Jun 8;53(5):514-29.
72. Park J, Lee Y, Seo H, Jang B, Son H, Kim S, Shin S, Hahn S. Risk of Bias Assessment tool for Non-randomized Studies (RoBANS): Development and validation of a new instrument. In: *Abstracts of the 19th Cochrane Colloquium*; 2011 19-22 Oct; Madrid, Spain. John Wiley & Sons; 2011.

Contributors: SC and NKA conceptualised the research. SC and NKA conducted the systematic review with inputs from SP. SC wrote the first draft and NKA and SP revised the manuscript.

Funding: The authors did not receive any funding for this study from any funding agency in the public, commercial, or not-for-profit sectors.

727 **Competing interest:** Authors declare no competing interest.

728 **Patient and public involvement:** No direct involvement of patients or the public.

729 **Patient consent for publication:** Not required.

730 **Data availability statement:** All relevant data are either included in this article or added to
731 the supplementary information.

732

733

734

735

736

737

738

739

740

741

742

743

744

745

746

747

748

749

750

751

752

753

754

755

756

757

758

759

760

761

762

763

764

765

766

767