

Reference values and validation of the 1-min sit-to-stand test in healthy 5- to 16-year-old youth

Electronic Supplementary Material Appendix

Sarah R Haile, Thea Fühner, Urs Granacher, Julien Stocker, Thomas Radtke, Susi Kriemler

Table S1 Participation in the various physical fitness test by sex and age group

sex	age group	n	STS	SLJ	CPET
Male	5-7 years	63	73%	38%	16%
Male	8-10 years	72	46%	10%	7%
Male	11-13 years	90	72%	4%	4%
Male	14-16 years	53	92%	8%	8%
Female	5-7 years	59	64%	34%	20%
Female	8-10 years	88	50%	6%	5%
Female	11-13 years	75	75%	8%	5%
Female	14-16 years	47	89%	4%	0%
all	all	547	68%	13%	8%

Table S2 Reference values (median [Q1 to Q3], by sex, age group and definition of STS (first measurement, last measurement, best measurement or mean repetitions)

sex	age group	first	last	best	mean
Male	5-7 years	56 [48 to 62]	60 [52 to 68]	60 [52 to 68]	58 [51 to 65]
Male	8-10 years	59 [50 to 64]	61 [55 to 66]	61 [55 to 66]	60 [53 to 66]
Male	11-13 years	58 [51 to 63]	61 [56 to 66]	61 [57 to 66]	59 [54 to 64]
Male	14-16 years	49 [44 to 55]	54 [49 to 60]	55 [49 to 60]	52 [46 to 56]
Female	5-7 years	53 [48 to 59]	56 [52 to 62]	56 [52 to 63]	54 [51 to 60]
Female	8-10 years	60 [55 to 66]	63 [57 to 67]	64 [58 to 68]	62 [57 to 67]
Female	11-13 years	58 [50 to 64]	61 [53 to 66]	61 [53 to 67]	60 [51 to 64]
Female	14-16 years	47 [38 to 56]	53 [42 to 60]	53 [42 to 60]	50 [40 to 58]




Figure S1 Association of three key measures (A) = sit-to-stand test (STS), (B) = standing long jump test (SLJ / body height), and (C) = cardiopulmonary exercise maximal incremental exercise test on a stationary bike (Wmax / body weight) with age




Figure S2 Scatterplot showing the correlation between the mean of the two sit-to-stand test (STS) results and (A) the maximal incremental exercise test on a stationary bike (Wmax), and (B) the standing long jump test (SLJ), adjusting for age




Figure S3 Association of STS with height (left), leg length (middle), and ratio of leg length to height (right), both unadjusted (top row), and adjusted for age (bottom row)

Working instructions - 1-minute Sit-to-Stand

Equipment

- Height adjustable chair
- Tape measure
- Stopwatch/Smartphone
- Counter (compulsory)
- Heart rate monitor (compulsory)
- Pulse oximeter (compulsory, for chronic disease populations)
- Scales to assess leg fatigue and breathlessness (e.g., Borg Scale)
- Solid footwear/sturdy shoes



Pre-test preparations

- Place the chair against the wall for safety reasons.
- Adjust the seat height to have approximately 90° knee angle flexion.
- Due to the observed learning effect, it is important that the child performs at least one practice test. If two tests are performed on the same day, consider a rest of about 10 minutes [1] between the first and second test to allow for sufficient recovery.

Test instructions

- Children are asked to perform as many repetitions as possible in 1 minute.
- Instruct the child to sit with his/her legs hip-width apart and with the hands placed on the hips.
- Children are not allowed to use their hands or arms to assist movement.
- Start position: Children sitting on a chair.
- Instruct the child to stand completely straight and to touch the chair with his/her bottom when sitting. They don't need to sit fully back on the chair. Correct during the test, if needed. Only correctly performed STS repetitions are counted.
- During the test, children are allowed to have a rest, when needed.
- Do not verbally encourage the child during the test. After 45 s the child is told that "15 s left until the test is over".

Outcomes

- Number of repetitions
- Heart rate
- Oxygen saturation (chronic disease populations)
- Subjective perception of breathlessness and leg fatigue
- STS Power Index (compulsory)

Reference values (N = 547)

Age	Female					Male				
	2.5%	25%	50%	75%	97.5%	2.5%	25%	50%	75%	97.5%
5-7 y	39	52	56	63	72	42	52	60	68	79
8-10 y	41	58	64	68	77	41	55	61	66	77
11-13 y	44	53	61	67	80	43	57	61	66	75
14-16 y	35	42	53	60	76	38	49	55	60	72
all	38	52	60	66	77	41	53	60	66	77

References

- [1] Hebestreit, H.; Mimura, K. & Bar-Or, O. Recovery of muscle power after high-intensity short-term exercise: comparing boys and men. *Journal of Applied Physiology*, 1993, 74(6). <https://doi.org/10.1152/jappl.1993.74.6.2875>