

Quality Control


Genome-wide study of DNA methylation in Amyotrophic Lateral Sclerosis identifies differentially methylated loci and implicates metabolic, inflammatory and cholesterol pathways

March 6, 2021


Contents

1	MinE 450k	2
2	MinE EPIC	9
3	AUS1	16
4	AUS2	23
5	Definition of AUS1 and AUS2 strata	29


1 MinE 450k




QC figure 1: Median unmethylated vs. median methylated signal.




QC figure 2: Median red vs. median green signal, calculated in type I probes.




QC figure 3: Rotated OP (Overall quality) plot based on non-polymorphic control probes as implemented in the *MethylAid* package [1].




QC figure 4: Rotated HC plot based on hybridization control probes as implemented in the *MethylAid* package [1].




QC figure 5: bscon metric from the *wateRmelon* package, which measures bisulfite conversion efficiency [2].




QC figure 6: Percentage of probes with detection P-value $> 1 \times 10^{-16}$




QC figure 7: Percentage of probes with beadNr < 3.




QC figure 8: Median X and Y copy number, as implemented in the *minfi* package [3].




QC figure 9: Difference between median Y-chromosomal and median X-chromosomal signal intensity.




QC figure 10: Identity-by-descent (IBS) comparing DNAm-inferred SNPs and SNPs measured using whole-genome sequencing data.



QC figure 11: Control probe principal components, after first round of QC.




QC figure 12: Array-wide principal components, after first round of QC.




QC figure 13: Chronological age vs. Predicted Age based on the Zhang *et al.* age predictor [4].


2 MinE EPIC




QC figure 14: Median unmethylated vs. median methylated signal.




QC figure 15: Median red vs. median green signal, calculated in type I probes.




QC figure 16: Rotated OP (Overall quality) plot based on non-polymorphic control probes as implemented in the *MethylAid* package [1].




QC figure 17: Rotated HC plot based on hybridization control probes as implemented in the *MethylAid* package [1].




QC figure 18: bscon metric from the *wateRmelon* package, which measures bisulfite conversion efficiency [2].




QC figure 19: Percentage of probes with detection P-value $> 1 \times 10^{-16}$




QC figure 20: Percentage of probes with beadNr < 3.




QC figure 21: Median X and Y copy number, as implemented in the *minfi* package [3].




QC figure 22: Difference between median Y-chromosomal and median X-chromosomal signal intensity.




QC figure 23: Identity-by-descent (IBS) comparing DNAm-inferred SNPs and SNPs measured using whole-genome sequencing data.



QC figure 24: Control probe principal components, after first round of QC.




QC figure 25: Array-wide principal components, after first round of QC.




QC figure 26: Chronological age vs. Predicted Age based on the Zhang *et al.* age predictor [4].


3 AUS1




QC figure 27: Median unmethylated vs. median methylated signal.




QC figure 28: Median red vs. median green signal, calculated in type I probes.




QC figure 29: Rotated OP (Overall quality) plot based on non-polymorphic control probes as implemented in the *MethylAid* package [1].




QC figure 30: Rotated HC plot based on hybridization control probes as implemented in the *MethylAid* package [1].




QC figure 31: bscon metric from the *wateRmelon* package, which measures bisulfite conversion efficiency [2].




QC figure 32: Percentage of probes with detection P-value $> 1 \times 10^{-16}$




QC figure 33: Percentage of probes with beadNr < 3.




QC figure 34: Median X and Y copy number, as implemented in the *minfi* package [3].




QC figure 35: Difference between median Y-chromosomal and median X-chromosomal signal intensity.



QC figure 36: Control probe principal components, after first round of QC.




QC figure 37: Array-wide principal components, after first round of QC.




QC figure 38: Chronological age vs. Predicted Age based on the Zhang *et al.* age predictor [4].


4 AUS2




QC figure 39: Median unmethylated vs. median methylated signal.




QC figure 40: Median red vs. median green signal, calculated in type I probes.




QC figure 41: Rotated OP (Overall quality) plot based on non-polymorphic control probes as implemented in the *MethylAid* package [1].




QC figure 42: Rotated HC plot based on hybridization control probes as implemented in the *MethylAid* package [1].




QC figure 43: bscon metric from the *wateRmelon* package, which measures bisulfite conversion efficiency [2].




QC figure 44: Percentage of probes with detection P-value $> 1 \times 10^{-16}$




QC figure 45: Percentage of probes with beadNr < 3.




QC figure 46: Median X and Y copy number, as implemented in the *minfi* package [3].



QC figure 47: Difference between median Y-chromosomal and median X-chromosomal signal intensity.




QC figure 48: Control probe principal components, after first round of QC.



QC figure 49: Array-wide principal components, after first round of QC.

5 Definition of AUS1 and AUS2 strata



QC figure 50: Median red vs. median green signal in all Australian data, calculated in type I probes. AUS1 samples were defined based on an RG ratio <0.65 , and AUS2 samples were defined based on an RG ratio >0.65 .

References

- [1] van Iterson, M. *et al.* MethylAid: visual and interactive quality control of large Illumina 450k datasets. *Bioinformatics* **30**, 23,435–3437 (2014).
- [2] Pidsley, R. *et al.* A data-driven approach to preprocessing Illumina 450K methylation array data. *BMC genomics* **14**, 293 (2013).
- [3] Aryee, M. J. *et al.* Minfi: a flexible and comprehensive Bioconductor package for the analysis of Infinium DNA methylation microarrays. *Bioinformatics* **30**, 10,1363–1369 (2014).
- [4] Zhang, Q. *et al.* Improved precision of epigenetic clock estimates across tissues and its implication for biological ageing. *Genome Medicine* **11**, 54 (2019).