Community mobilisation approaches to preventing and reducing adolescent multiple risk behaviour: a realist review protocol

Tinner, Laura. laura.tinner@bristol.ac.uk (corresponding author).¹

Caldwell, Deborah. d.m.caldwell@bristol.ac.uk¹

Campbell, Rona. rona.campbell@bristol.ac.uk¹

¹ Population Health Sciences, Bristol Medical School, Canynge Hall, University of Bristol, Bristol, UK, BS8 2PL.

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

Abstract

Background: Adolescent multiple risk behaviour (MRB) continues to be a global health issue, contributing to the burden of non-communicable diseases. Most interventions have focused on the proximal causes of adolescent MRB such as peer or family influence, rather than targeting the wider environmental or structural context. There is increasing recognition that community mobilisation approaches that extend beyond individually-focused educational programmes could be beneficial for adolescent health. Despite this, there are gaps in the current literature, theory and implementation that would benefit from a realist approach due to the suitability of this methodology to analysing complex interventions. In this protocol, we outline our study that aims to understand ‘how, why, for whom and in what circumstances and time periods do community mobilisation interventions work to prevent and/or reduce adolescent multiple risk behaviour?’

Methods: A realist review was chosen as the most suitable review method as it is theory-driven and seeks to understand how, why and for whom interventions work to produce intended and unintended outcomes. A six-stage iterative process is outlined, which includes initial development of a programme theory, systematic searching, study selection and appraisal, data extraction and data synthesis. We will engage with stakeholders at different stages in this process to aid the development of the programme theory.

Discussion: The goal of this realist review is to identify and refine a programme theory for community mobilisation approaches to the prevention and/or reduction of adolescent multiple risk behaviour. Our aim is that the findings surrounding the programme theory

refinement can be used to develop and implement adolescent multiple risk behaviour interventions and maintain collaboration between local policy makers, researchers and community members.

Registration: This realist review is registered on the PROSPERO database (registration number: CRD42020205342).

Keywords: Community mobilisation; adolescence; risk behaviour; realist review; complex interventions; inequalities.

Background

Adolescent health risk behaviours

Health risk behaviours such as tobacco smoking, hazardous alcohol consumption, antisocial behaviour, physical inactivity and unprotected sexual intercourse are global health issues, which are commonly initiated and become habitual in adolescence (1, 2). Adolescents who engage in one risk behaviour are likely to engage in others (3, 4), leading to increased public health interest in multiple risk behaviour (MRB), which refers to the occurrence of two or more risk behaviours directly or indirectly related to health (5, 6). MRB has been found to be associated with a number of adverse health and social outcomes such as poor educational attainment (7), obesity, depression and anxiety in adulthood (8), cancers and premature mortality (7, 9). This has, in turn, led to public health interventions that address multiple as opposed to single behaviours (10).

Most interventions addressing adolescent MRB have focused on the proximal causes such as peer or family influence, rather than targeting the wider environmental, social or structural context (11). For instance, two Cochrane systematic reviews have assessed the impact of individual, family and school-level interventions on adolescent multiple risk behaviour (10, 12). One of those reviews found mixed evidence, concluding that school-based universal interventions are potentially effective in 'preventing engagement in tobacco use, alcohol use, illicit drug use, and antisocial behaviour, and in improving physical activity among young people, but not in preventing other risk behaviours' (12). The authors highlighted that there was no strong evidence of benefit for family-level or individual-level interventions across the risk behaviour outcomes investigated (12). The interventions included in this review were predominantly educational programmes. The effectiveness and equity of these

“downstream” interventions has been questioned (13) because health risk behaviours rarely have a single cause and occur in complex socio-cultural contexts (14). As such, there is increasing recognition that structural changes that extend beyond individually-focused educational programmes could be beneficial for adolescent health (14, 15).

Community mobilisation interventions

Recognition that decisions about health risk behaviours are made within a broad social context has led to the development and implementation of community-engagement interventions (16). There is an extensive range of types of community-engagement public health interventions, varying in the extent to which they emphasise community involvement in determining and delivering the programmes (16). ‘Community mobilisation’ interventions are one such type that work to engage community members to ‘take action towards achieving a common goal’ (17) and have gained traction as a strategy for addressing complex and multifaceted problems (18). Community mobilisation is a collaborative public health effort that is defined by the inclusion of a community coalition made up of diverse stakeholders (such as schools, businesses, residents, youth groups, emergency services and religious leaders) (11). These stakeholders critically analyse the root causes of local problems, identify an array of potential solutions, develop multi-sector partnerships, and implement multi-component strategies for creating local change and more health-promoting environments (14).

Community mobilisation efforts explicitly seek to affect community-level influences through changes of policies, practices, organisations and other features of the social or physical environment that may impact on the health outcome or behaviour (19), signifying a shift away

from individual behaviour change to a focus on the social determinants of health (20). However, these approaches may still include components which address individual behaviours (e.g. health promotion programmes within schools), but they seek to combine these with other structural factors as part of a package of measures that are chosen and monitored from community stakeholders.

There is systematic review evidence suggesting that higher levels of community involvement within an public health intervention is linked to more beneficial effects and positive trends across a range of outcomes (21). There is also some evidence to support the role of community mobilisation efforts in preventing health risk behaviours. For instance, such interventions have resulted in reductions in high risk alcohol consumption and alcohol-related injuries (22); alcohol impaired driving (23); uptake of smoking in young people (16) and youth violence (24). Researchers have highlighted that with adequate resources and training, support from within the community and adoption of evidence-based strategies, community mobilisation approaches have promise as an effective vehicle for addressing adolescent multiple risk behaviour (19). Further, community-mobilisation efforts are also thought to be well suited to achieving health equity (25), due to ‘shared decision making’ (26) and the incorporation of ‘upstream’ or structural elements (27), but this has yet to be explored in relation to adolescent multiple risk behaviour interventions.

There are also significant challenges in implementing and evaluating such approaches, which is unsurprising given the dynamic set of social interactions and relational complexity one might expect in community-centred interventions (28). These implementation challenges include lack of community interest and long-term engagement, design inadequacies, inflated

expectations, and weakness in planning and implementation of the interventions (19, 29, 30). Tensions and different expectations between scientists and community members as well as the practical difficulty in managing multiple components and stakeholder interests have also been cited as issues (11).

Evaluation is equally challenging (31), which is reflected in the lack of empirical evaluations of structural interventions such as community mobilisation compared to those focused at the individual level (14). There is uncertainty around how long it might take to see an impact on behaviours, although it is expected to be a lengthy process. Even if effects are identified, the chain linking any changes in health risk behaviours to the mobilisation efforts is so long and complex that causal attributions become complicated (14). The challenges in evaluating and implementing community mobilisation interventions has meant they are often evaluated through methods such as quasi-experimental studies in addition to randomised controlled trials (RCTs), meaning that they have been missing from systematic reviews such as the aforementioned adolescent MRB review (12). Consequently, the evidence base for community mobilisation efforts is mixed in terms of producing desirable outcomes on a community and individual level.

We aim to address the gap in the literature through investigating community mobilisation interventions aimed at preventing and/or reducing adolescent multiple risk behaviour. There is a strong rationale for an alternative review approach that speaks to the complexities and challenges surrounding the delivery and evaluation of community mobilisation efforts. Further, we are concerned with moving beyond assessing effectiveness of public health interventions to synthesise existing knowledge and articulate *how* community mobilisation

interventions work for adolescent health. Therefore, a realist review was chosen as the most appropriate methodological approach. Realist reviews are theory-driven approaches to evidence synthesis, incorporating diverse data sources to provide insight into the underlying mechanisms and contexts in which the interventions work (32). Realist reviews are ideal for examining social interventions, particularly those in community settings as it is recognised that programmes are rarely delivered in the same way or have exactly the same outcomes, due to contextual factors that can never be fully controlled (33).

Realist inquiry is thus ‘increasingly recognised as an effective process for consolidating evidence and learning from complex social processes and interventions’ (34), with successes in public health and community development (35, 36). Therefore, our realist review aims to contribute to the current adolescent multiple risk behaviour evidence base, which has largely focused on effectiveness of interventions through traditional systematic reviews. Further support for our intended approach comes from a recent PhD thesis which used an adapted realist approach to assessing adolescent multiple risk behaviour programmes, combining realist evaluation with primary data collection (37) (38). The authors did not specifically focus on community mobilisation as an intervention and included fewer health risk behaviours than we do here. This protocol describes our realist methodological approach and intended procedures in the sections that follow.

This realist systematic review is registered on the PROSPERO database (registration number: CRD42020205342). A PRISMA-P check list is included as a supplementary file.

210 Methods

211 Review aim

212 Our aim is to use a theory-driven evidence synthesis to assess how and why community
 213 mobilisation interventions work/do not work to prevent or reduce adolescent multiple risk
 214 behaviour and in what contexts. We are additionally interested in the question of ‘who’ these
 215 interventions work for, in order to understand the impact of these types of interventions
 216 upon existing health inequalities through investigating whether the interventions are
 217 beneficial to disadvantaged communities. Although the focus of the review is adolescent
 218 multiple risk behaviour, we may draw on wider literature to understand the goals of
 219 community mobilisation approaches and the mechanisms by which it is hoped these are
 220 achieved. An additional objective of our review is to develop transferable learning about
 221 community mobilisation approaches in public health research and adolescent health
 222 interventions. The realist review will be guided by the following sub-questions:

- 223 1. What are the outcomes of community mobilisation interventions targeting adolescent
 224 multiple risk behaviour?
- 225 2. What are the mechanisms, acting at the individual, community and societal levels
 226 through which community mobilisation interventions produce outcomes?
- 227 3. What are the key contextual influences that determine whether the mechanisms
 228 produce both intended and unintended outcomes?

229

230 Realist review methodology

231

232 Realist review methodology is a theory-driven, interpretive approach to evidence synthesis
 233 (38) developed by the work of Pawson et al (39-41). It has gained increasing popularity in
 234 addressing the challenge of ‘what works, for whom, under what circumstances and in what

time period' and is considered especially salient when data are complex, multi-layered and there is a need to understand complex relationships, interdependence, and mechanisms (42). A core component of realist reviews is to develop 'middle-range realist programme theory' (43) that explains how an intervention 'works' within what contexts. Realist reviews allow for exploration of complex topics and the inclusion of a wide body of quantitative, qualitative and mixed methods evidence to develop and refine theory (38). Therefore, it is suited to multi-component community-based interventions, for which evaluations may include a range of different data and be published in grey literature.

Realist reviewers view 'causation' as generative, which means that the manifested world is generated (i.e. caused) via underpinning mechanisms (32). They identify where an intervention, under certain contextual conditions (C), triggers a mechanism (M) to achieve a given outcome (O) (38). This CMO configuration is central to analysis and theory development, viewing mechanisms as the integral link between contexts and outcomes. The aim of realist review methodology to move beyond measuring effectiveness of interventions, toward explanation of how and why an intervention works is a key strength that separates it from traditional systematic reviews. Dalkin et al (44) conceptualise a mechanism as a resource or reasoning, which triggers an outcome, but may only be active in certain contexts. Under certain contextual conditions, the mechanisms are triggered, while in others they 'fire' to a lesser degree or not at all (44). Realist review methodology is highly applicable to complex public health interventions and is an approach that can build 'common ground' between researchers and policy makers through providing accessible recommendations on how interventions might be delivered in different contexts (41).

We have conceptualised multi-component community mobilisation approaches as complex interventions in which outcomes and mechanisms will be context sensitive (39). Therefore, the realist review approach will allow us to investigate in what contexts community mobilisation interventions are effective. This realist review will follow the practice guidelines outlined by the Realist and Meta-narrative Evidence Synthesis: Evolving Standards (RAMESES) framework (40, 41).

Study design

This review is structured around the five review stages outlined by Pawson et al (39) and has been informed by other realist review protocols in the field (38, 45). Figure 1 is a diagram of the review process adapted from Power et al (38).

Figure 1 to go here.

Figure 1: Summary of stages of realist review adapted from Power et al (38). This depicts the steps for developing the initial programme theory, searching for evidence and synthesising the data with the input of key stakeholders. Retrodution refers to inferences made through interpreting the data about the underlying causal mechanisms.

Stages of realist review

(1) Locating existing theories

The first step in a realist review is to conduct scoping searches that begin to identify theories that might explain how community mobilisation interventions may work to address adolescent multiple risk behaviour. The search will include academic databases (MEDLINE, PubMed, Web of Science), UK health websites and grey literature databases (OpenGrey, the King's Fund, The Health Foundation) as well as Google Scholar. Broad search terms will be

used at this initial stage (e.g. “community mobilisation”, “community coalition”, “youth”, “adolescence”, “health risk behaviour”, “substance use”, “antisocial behaviour”) and back and forth citation tracking will be utilised until we develop a core set of empirical studies to help build the initial programme theory framework (40). This initial search is not designed to be exhaustive: this stage in the theory development is expected to be a ‘rough starting point’ that will be refined throughout the realist review process (43).

It is advisable to include the expertise of those delivering or evaluating the interventions. At this stage we will engage with key stakeholders identified through the literature to provide guidance on the development of the programme theory. For example, they may provide insight into the different contexts and mechanisms that impact on adolescent risk behaviour outcomes from their experience in the field. These stakeholders may also highlight other relevant studies or individuals we should engage with to further develop the programme theory. When we have developed an initial programme theory, we will move onto stage 2 and the more structured and systematic searching.

(2) Search strategy

In Stage 2 we will conduct more formal searches, which will be informed by the initial programme theory development in stage 1. The objective in this stage will be to identify literature and evidence capable of informing the refinement of a more detailed programme theory (45). We will develop search terms from the initial background search in Stage 1 and discussions with a subject librarian, leading to systematic searches being undertaken to collect evidence to refine the programme theory. We will include the following databases:

308 PubMed; MEDLINE; PsycINFO; Web of Science; CINAHL; Sociological Abstracts. Grey literature
309 will also be searched on OpenGrey and on external expert organisations and charity websites.
310 ProQuest will be searched for unpublished theses and dissertations. Google Scholar will be
311 used for citation searching as well as reference lists of relevant papers.

312

313 Search terminology and syntax will be informed by the initial programme theory
314 identification, known literature and collaboration between the research team and a subject
315 librarian. We will draw on the search terms used in previous systematic reviews on individual-
316 level adolescent multiple risk behaviour interventions. Search terms will include MeSH terms
317 and free text related to “community mobilisation”, “adolescence” and a range of multiple
318 health risk behaviours. No date restrictions will be used and only studies in the English
319 Language will be assessed for eligibility.

320

321 While formalised and systematic, the sampling approach in realist reviews remains purposive
322 to answer specific questions and develop theories (39). Therefore, the process will likely be
323 iterative and need to be repeated (39), with back and forth citation tracking remaining a key
324 part of the iterative search strategy (46). Corresponding authors of selected articles may also
325 be contacted for further examples that may be relevant to the question. The search terms
326 and strategies will be documented in a log-book as the review progresses.

327

328 (3) Study selection

329 We will use the following inclusion criteria to determine if a document is likely to contribute
330 to the programme theory development:

- Type of intervention: Community mobilisation must form a core part of the intervention, most commonly identified by the development of a community coalition group involving a diverse range of community stakeholders as identified above. In many cases, the community coalition will select intervention components from a ‘menu’ of strategies and adapt them to fit local needs. As mentioned, although these types of interventions derive from the desire to move away from individual behaviour change, they are also likely to include a range of “upstream” and “downstream” components, which may include education delivery. The key criterion remains that the intervention should include community mobilisation at its core, and should include at least two components (e.g. an educational programme and local policy enforcement).
- A range of document types, study designs and data types may be relevant to the development of the initial programme theory. All intervention evaluation study designs and data types, from all time periods, may be included in the review to test and refine the programme theory (e.g. randomised controlled trials (RCTs), quasi-experimental studies, case studies).
- Participants: the community mobilisation efforts should be targeted (at least predominately) at young people age 10-19 years, although this age range remains flexible¹. Adult stakeholders (such as parents, community members, school staff) will likely be included in the community coalition but should not be the focus of the intervention. All sampling decisions will be transparently reported.

¹ For instance, a ‘pre-teen’ intervention targeting children age 9-12 would be included as the population are only one year out of our pre-specified age range of 10-19 years. However, an older adolescent/young adult intervention targeting individuals aged 18-25 would not be included as only two years (18 and 19) of the eight year range addressed by the intervention falls within our age banding.

- Aim of intervention: the intervention should have a primary focus on prevention and reduction of adolescent multiple risk behaviour and include at least two health risk behaviours from a wide range including: regular tobacco smoking; regular alcohol drinking; binge drinking (alcohol); cannabis use; recent or regular illicit drug use; risky sexual behaviours; anti-social behaviour and offending; vehicle-related risk behaviours (e.g. cycling without a helmet; not using a car seatbelt, joy riding); self-harm; gambling; unhealthy diet; and physical inactivity.
 - Outcome measures: Primary outcomes of interest include reduction and/or prevention of the wide range of multiple risk behaviours mentioned above. A range of additional medium- and long-term outcomes within health and social domains are expected given the number of health risk behaviours the interventions can cover as well as the multi-component nature of these types of interventions. Secondary outcomes of interest in this review include: Education and employment: educational qualifications; truancy and school exclusion; employment; not being in education, employment or training (NEET); Crime: Criminal record/offending; long-term addictive behaviours; Gambling; Teenage pregnancy or parenthood; Sexually transmitted infections; Injuries; Morbidity (e.g. Hepatitis C, HIV, anxiety and depression, obesity, type II diabetes, fatty liver disease, liver cirrhosis); Suicide/self-harm; and Premature mortality. Realist reviews are interested in intended and unintended outcomes related to the context, mechanism, and outcomes of the intervention therefore other unknown outcomes may become relevant to the programme theory development.
- Studies will be excluded if they relate to any of the following:

- A single component intervention (such as an educational programme) that is delivered in the community but does not incorporate community mobilisation as we have defined it.
- Interventions targeted at participants outside of the age range. Some interventions may include other populations, but youth should be the primary focus.
- Interventions aimed at preventing and reducing a single adolescent health risk behaviour (e.g. alcohol misuse).
- Clinical and pharmaceutical interventions including ‘community outreach’ services such as the provision of mobile clinics.
- Studies not described in the English language.

We will use Raayan (QCRI) software for screening and management of the studies at this stage in the review. The RAMESES guidelines will be used to appraise the studies (41). The selection of evidence will be made based on judgements around their *relevance* (contribution to the programme theory development and refinement) and *rigour* (credibility and trustworthiness of methods) (45, 47). Any exclusions based on these appraisals will be documented.

(4) Data extraction

Study characteristics will be extracted into a table to provide a descriptive overview of the types of community mobilisation interventions included, based on a ‘bespoke’ set of data extraction forms informed by the relevant literature (39). Realist reviews are structured through Context-Mechanisms-Outcome (CMO), comparable to PICO for traditional systematic reviews (42). Context, mechanisms and outcomes are extracted during the realist

review and can be conceptualised as the ‘data’ that support evidence to support, reject or refine the programme theory (33). However, this process is not as rigid as with traditional systematic reviews and different sources may provide different information that contributes to the programme theory development. We will extract the following information:

- Study details: authors, year of publication, country of intervention delivery, study aims, study design, participant characteristics, quality appraisal.
- Context: background of the intervention, aims of the intervention, type of intervention, setting (e.g. type and size of community), age range, number of components, policy context for the area, historical context.
- Mechanism: descriptions of the processes through which the intervention influenced outcomes, who the intervention worked for and who it did not, author-identified mechanisms.
- Outcomes: adolescent multiple health risk behaviour outcomes but also a range of health and social outcomes mentioned previously.
- Additional study information and researcher comments.

The above is not an exhaustive list and information on the mechanisms and context will be expected to change through experience with the studies and input from expert stakeholders. Any disagreements on extracted data will be resolved through discussion with the research team. Realist reviews assimilate information more through note taking and documents are scoured for ideas about how the intervention might work (39). To approach the more complex and iterative process of examining study sources, we will follow the guidance from Pawson et al (39) and will also upload the documents to NVivo for organising and coding to aid development of the programme theory and to keep a record of our procedures (45).

422

423 (5) Data synthesis

424 The goal of data synthesis in a realist review is to consolidate the data from the previous steps
 425 to refine the initial programme theory (45). Analysis will involve interpretation of the
 426 researchers and judgement of the data. Coding the data will involve deductive (informed by
 427 the initial programme theory), inductive (emerging from the data within the identified
 428 interventions) and retroductive approaches (inferences made through interpreting the data
 429 about the underlying causal mechanisms) (45). These approaches will guide the review to
 430 interpret and explain the findings and outline the contextual conditions and mechanisms that
 431 may need to be present for outcomes to occur. Data to inform our interpretation of the
 432 relationships between the contexts, mechanisms and outcomes will be analysed within and
 433 across the documents. For example, mechanisms inferred from one document will be applied
 434 to other documents to test if they can explain the way contexts influence outcomes in another
 435 intervention (45). This interpretive process will lead to programme theory development of
 436 community mobilisation approaches to preventing and reducing adolescent multiple risk
 437 behaviour.

438

439 (5) Refine programme theories

440 The final product of a realist theory is not a statement of effectiveness, but a refinement of
 441 middle-range theory that answers the questions of what works, for whom, under what
 442 circumstances, in what time period, why and how? (48). Evidence may include primary
 443 outcome data, but also rich description that conveys the contextual elements and
 444 interpretation of the interactions between the context, mechanisms and outcomes by the
 445 researchers (48). Stakeholders involved in stage one will be contacted again, with the
 446 potential for the inclusion of newly identified individuals, for input into the final programme

theory(ies). The aim of this process is to confirm that the programme theory makes sense to those involved, in order to enhance our ability to make practice recommendations from our findings (39). If needed, we will re-scrutinise elements of the review based on the stakeholder involvement. We intend to develop a final logic model that will visually present the programme theory/theories and the relationships between the CMO.

Discussion

The realist review approach will allow us to explore the relationships between contexts, mechanisms and outcomes and synthesise evidence surrounding adolescent multiple risk behaviour interventions that incorporate community mobilisation. We seek to gain a greater understanding of ‘what works, for whom, in what circumstances, in what time period and why’, in terms of preventive interventions to improve adolescent health and reduce inequalities. Our aim is that the findings surrounding the programme theory refinement can be used to develop and implement adolescent multiple risk behaviour interventions and maintain collaboration between local policy makers, researchers and community members. We will recognise any limitations to our study and the realist review approach in the final synthesis. The dissemination of the findings of this review will follow the RAMESES reporting guidelines (40, 41).

472

473 [Abbreviations](#)

474 MRB: Multiple risk behaviour

475

476 NEET: not being in education, employment or training

477

478 HIV: human immunodeficiency virus

479

480 RCTs: Randomised Controlled Trials

481

482 CMO: Context, Mechanism, Outcome

483

484 RAMESES: Realist and meta-narrative evidence syntheses: evolving standards

485

486 [Declarations](#)

487 [Ethics approval and consent to participate](#)

488 Not applicable.

489

490 [Consent for publication](#)

491 Not applicable.

492

493 [Availability of data and materials](#)

494 Not application.

495

496 [Competing interests](#)

497 The authors declare no competing interests.

498

499 [Funding](#)

500 This work is funded through LT's post-doctoral post, which is funded by Bristol, North

501 Somerset and South Gloucestershire Clinical Commission Group (CCG) Research Capacity

502 Funding via RC's Senior Investigator Award. The funder had no involvement with the

503 development of the protocol.

504

505 [Authors' contributions](#)

506 LT, DC and RC developed the project idea, chose the realist review approach and developed

507 the inclusion criteria. LT wrote the first draft of the protocol. All authors contributed to writing

508 and refining the protocol and will be involved in the review. LT is the corresponding author.

Acknowledgements

Not applicable.

References

1. Jackson CA, Henderson M, Frank JW, Haw SJ. An overview of prevention of multiple risk behaviour in adolescence and young adulthood. *Journal of Public Health*. 2012;34(suppl_1):i31-i40.
2. Patton GC, Sawyer SM, Santelli JS, Ross DA, Afifi R, Allen NB, et al. Our future: a Lancet commission on adolescent health and wellbeing. *The Lancet*. 2016;387(10036):2423-78.
3. Kipping RR, Campbell RM, MacArthur GJ, Gunnell DJ, Hickman M. Multiple risk behaviour in adolescence. *Journal of Public Health*. 2012;34(suppl_1):i1-i2.
4. Brooks FM, Magnusson J, Spencer N, Morgan A. Adolescent multiple risk behaviour: an asset approach to the role of family, school and community. *Journal of Public Health*. 2012;34(suppl_1):i48-i56.
5. Hurrelmann K, Richter M. Risk behaviour in adolescence: the relationship between developmental and health problems. *Journal of Public Health*. 2006;14(1):20-8.
6. Tinner L, Caldwell D, Hickman M, MacArthur GJ, Gottfredson D, Perez AL, et al. Examining subgroup effects by socioeconomic status of public health interventions targeting multiple risk behaviour in adolescence. *BMC Public Health*. 2018;18(1):1180.
7. Wright C, Kipping R, Hickman M, Campbell R, Heron J. Effect of multiple risk behaviours in adolescence on educational attainment at age 16 years: a UK birth cohort study. *BMJ Open*. 2018;8(7):e020182.
8. Campbell R, Wright C, Hickman M, Kipping RR, Smith M, Pouliou T, et al. Multiple risk behaviour in adolescence is associated with substantial adverse health and social outcomes in early adulthood: Findings from a prospective birth cohort study. *Preventive Medicine*. 2020;106:157.
9. Hausdorf K, Eakin E, Whiteman D, Rogers C, Aitken J, Newman B. Prevalence and correlates of multiple cancer risk behaviors in an Australian population-based survey: results from the Queensland Cancer Risk Study. *Cancer Causes & Control*. 2008;19(10):1339-47.
10. Hickman M, Caldwell DM, Busse H, MacArthur G, Faggiano F, Foxcroft DR, et al. Individual-, family-, and school-level interventions for preventing multiple risk behaviours relating to alcohol, tobacco and drug use in individuals aged 8 to 25 years. *Cochrane Database of Systematic Reviews*. 2014(11).
11. Fagan AA, Hawkins JD, Catalano RF. Interventions For Addiction: Chapter 90. *Mobilizing Communities for Alcohol, Drug, and Tobacco Prevention*: Elsevier Inc. Chapters; 2013.
12. MacArthur G, Caldwell DM, Redmore J, Watkins SH, Kipping R, White J, et al. Individual-, family-, and school-level interventions targeting multiple risk behaviours in young people. *Cochrane Database of Systematic Reviews*. 2018(10).
13. Adams J, Mytton O, White M, Monsivais P. Why are some population interventions for diet and obesity more equitable and effective than others? The role of individual agency. *PLoS Medicine*. 2016;13(4).

14. Miller RL, Janulis PF, Reed SJ, Harper GW, Ellen J, Boyer CB, et al. Creating youth-supportive communities: outcomes from the Connect-to-Protect® (C2P) structural change approach to youth HIV prevention. *Journal of Youth and Adolescence*. 2016;45(2):301-15.
15. Aguirre-Molina M, Gorman DM. Community-based approaches for the prevention of alcohol, tobacco, and other drug use. *Annual review of public health*. 1996;17(1):337-58.
16. Carson KV, Brinn MP, Labiszewski NA, Esterman AJ, Chang AB, Smith BJ. Community interventions for preventing smoking in young people. *Cochrane Database of Systematic Reviews*. 2011(7).
17. Lippman SA, Neilands TB, Leslie HH, Maman S, MacPhail C, Twine R, et al. Development, validation, and performance of a scale to measure community mobilization. *Social Science & Medicine*. 2016;157:127-37.
18. Watson-Thompson J, Fawcett SB, Schultz JA. A framework for community mobilization to promote healthy youth development. *American Journal of Preventive Medicine*. 2008;34(3):S72-S81.
19. Flewelling RL, Austin D, Hale K, LaPlante M, Liebig M, Piasecki L, et al. Implementing research-based substance abuse prevention in communities: Effects of a coalition-based prevention initiative in Vermont. *Journal of Community Psychology*. 2005;33(3):333-53.
20. Kim-Ju G, Mark GY, Cohen R, Garcia-Santiago O, Nguyen P. Community mobilization and its application to youth violence prevention. *American Journal of Preventive Medicine*. 2008;34(3):S5-S12.
21. Brunton G, Caird J, Stokes G, Stansfield C, Kneale D, Richardson M, et al. Review 1: Community engagement for health via coalitions, collaborations and partnerships: a systematic review. 2015.
22. Holder HD, Gruenewald PJ, Ponicki WR, Treno AJ, Grube JW, Saltz RF, et al. Effect of community-based interventions on high-risk drinking and alcohol-related injuries. *Jama*. 2000;284(18):2341-7.
23. Shults RA, Elder RW, Nichols JL, Sleet DA, Compton R, Chattopadhyay SK, et al. Effectiveness of multicomponent programs with community mobilization for reducing alcohol-impaired driving. *American journal of preventive medicine*. 2009;37(4):360-71.
24. Massetti GM. Preventing violence among high-risk youth and communities with economic, policy, and structural strategies. *MMWR supplements*. 2016;65.
25. Trickett EJ, Beehler S. The ecology of multilevel interventions to reduce social inequalities in health. *American Behavioral Scientist*. 2013;57(8):1227-46.
26. Durand M-A, Carpenter L, Dolan H, Bravo P, Mann M, Bunn F, et al. Do interventions designed to support shared decision-making reduce health inequalities? A systematic review and meta-analysis. *PloS one*. 2014;9(4):e94670.
27. Nickel S, von dem Knesebeck O. Do multiple community-based interventions on health promotion tackle health inequalities? *International Journal for Equity in Health*. 2020;19(1):1-13.
28. Adhikari B, Vincent R, Wong G, Duddy C, Richardson E, Lavery JV, et al. A realist review of community engagement with health research. *Wellcome open research*. 2019;4.
29. Wandersman A, Florin P. Community interventions and effective prevention. *American Psychologist*. 2003;58(6-7):441.
30. Gabriel RM. Methodological challenges in evaluating community partnerships & coalitions: Still crazy after all these years. *Journal of Community Psychology*. 2000;28(3):339-52.

31. Bonell C, Hargreaves J, Strange V, Pronyk P, Porter J. Should structural interventions be evaluated using RCTs? The case of HIV prevention. *Social Science & Medicine*. 2006;63(5):1135-42.
32. Jagosh J. Realist synthesis for public health: building an ontologically deep understanding of how programs work, for whom, and in which contexts. *Annual Review of Public Health*. 2019.
33. Usher AM, McShane KE, Dwyer C. A realist review of family-based interventions for children of substance abusing parents. *Systematic Reviews*. 2015;4(1):1-12.
34. Richardson EZL, Bandewar SVS, Boulanger RF, Mehta R, Lin T, Vincent R, et al. Addressing diversity and complexity in the community engagement literature: The rationale for a realist review. *Wellcome Open Research*. 2020;5(1):1.
35. Wong G, Pawson R, Owen L. Policy guidance on threats to legislative interventions in public health: a realist synthesis. *BMC Public Health*. 2011;11(1):222.
36. Jagosh J, Macaulay AC, Pluye P, Salsberg J, Bush PL, Henderson JIM, et al. Uncovering the benefits of participatory research: implications of a realist review for health research and practice. *The Milbank Quarterly*. 2012;90(2):311-46.
37. Cooper C, Lhussier M, Shucksmith J, Carr SM. Protocol for a realist review of complex interventions to prevent adolescents from engaging in multiple risk behaviours. *BMJ open*. 2017;7(9):e015477.
38. Power J, Gilmore B, Vallières F, Toomey E, Mannan H, McAuliffe E. Adapting health interventions for local fit when scaling-up: a realist review protocol. *BMJ Open*. 2019;9(1).
39. Pawson R, Greenhalgh T, Harvey G, Walshe K. Realist review-a new method of systematic review designed for complex policy interventions. *Journal of Health Services Research & Policy*. 2005;10(1_suppl):21-34.
40. Greenhalgh T, Wong G, Westhorp G, Pawson R. Protocol-realist and meta-narrative evidence synthesis: evolving standards (RAMESES). *BMC Medical Research Methodology*. 2011;11(1):115.
41. Wong G, Greenhalgh T, Westhorp G, Pawson R. Development of methodological guidance, publication standards and training materials for realist and meta-narrative reviews: the RAMESES (Realist And Meta-narrative Evidence Syntheses—Evolving Standards) project. 2014.
42. Booth A, Briscoe S, Wright JM. The “realist search”: A systematic scoping review of current practice and reporting. *Research Synthesis Methods*. 2020;11(1):14-35.
43. Weetman K, Wong G, Scott E, Schnurr S, Dale J. Improving best practice for patients receiving hospital discharge letters: a realist review protocol. *BMJ Open*. 2017;7(11):e018353.
44. Dalkin SM, Greenhalgh J, Jones D, Cunningham B, Lhussier M. What’s in a mechanism? Development of a key concept in realist evaluation. *Implementation Science*. 2015;10(1):1-7.
45. Abrams R, Wong G, Mahtani KR, Tierney S, Boylan A-M, Roberts N, et al. Understanding the impact of delegated home visiting services accessed via general practice by community-dwelling patients: a realist review protocol. *BMJ Open*. 2018;8(11).
46. GRADE Working Group. Grading of Recommendations, Assessment, Development, and Evaluation. Available from: <https://www.gradeworkinggroup.org/>.
47. Pawson R. Digging for nuggets: how ‘bad’ research can yield ‘good’ evidence. *International Journal of Social Research Methodology*. 2006;9(2):127-42.

- 646 48. Jagosh J, Bush PL, Salsberg J, Macaulay AC, Greenhalgh T, Wong G, et al. A realist
647 evaluation of community-based participatory research: partnership synergy, trust building
648 and related ripple effects. BMC Public Health. 2015;15(1):725.
649