Title page

Title: Defining Vascular Cognitive Impairment in India—A Descriptive Study of MRI Volumetry vs. Cognition in ‘Normal’ Adults

Authors:

Uma Sundar;^a Amita Mukhopadhyay;^b Sheelakumari Raghavan;^c Ramshekhar N. Menon;^d Chandrasekharan Kesavadas;^e Nilesh Shah;^f Balkrishna B. Adsul;^g Anagha R. Joshi.^h

Affiliations:

^a Department of Medicine, Lokmanya Tilak Municipal Medical College and General Hospital (LTMMC&GH), Sion, Mumbai 400022, Maharashtra, India. Email address: umasundar2@gmail.com

^b Department of Community Medicine, Dr Chandramma Dayananda Sagar Institute of Medical Education and Research, Kanakapura 562112, Ramanagara District, Karnataka, India. Email address: dr.amukho@gmail.com

^c Department of Radiology, Sree Chitra Tirunal Institute for Medical Sciences and Technology (SCTIMST), Thiruvananthapuram 695011, Kerala, India. Email address: sheela82nair@gmail.com

^d Department of Neurology, R Madhavan Nayar Centre for Comprehensive Epilepsy Care, Cognition & Behavioral Neurology, SCTIMST, Thiruvananthapuram. Email address: rsnmenon@sctimst.ac.in

^e Department of Radiology, SCTIMST, Thiruvananthapuram. Email address: chandkesav@yahoo.com

^f Department of Psychiatry, LTMMC&GH, Mumbai. Email address: drmilshah@hotmail.com

^g Department of Community Medicine, Hinduhradaysamrat Balasaheb Thackeray Medical College and Dr. RN Cooper Municipal General Hospital, Juhu, Mumbai 400056, Maharashtra, India. Email address: adsulbb@gmail.com

^hDepartment of Radiodiagnosis, LTMMC&GH, Mumbai. Email address:

anaghajoshi2405@gmail.com

Corresponding author: Uma Sundar, umasundar2@gmail.com

Word count (abstract): 250; Word count (text): 3507; References: 50; Tables: 5; Figures: 3.

Study funding: Fully funded by the Cognitive Science Research Initiative, Department of Science & Technology, Government of India (Grant ID: CSRI 129/2012).

Ethical approval: Approved by the Institutional Ethical Committee of Lokmanya Tilak Municipal Medical College and General Hospital, Sion, Mumbai (approval no. 15/2015).

Declarations of interest: None.

Abstract

Background: It is important to establish criteria to define Vascular Cognitive Impairment (VCI) in India as VCI is an image-based diagnosis and MRI changes resulting from age with prevalent vascular risk factors may confound MRI interpretation.

Objective: To establish normative community data for MRI volumetry including white matter hyperintensity volume (WMHV), correlated with age-stratified cognitive scores and vascular risk factors in adults ≥ 40 years old.

Materials and Methods: We screened 2651 individuals without known neurological morbidity, living in Mumbai and nearby rural areas, using validated Marathi translations of Kolkata Cognitive Battery (KCB) and Geriatric depression score (GDS). We stratified 1961 persons with GDS ≤ 9 by age and cognitive score, and randomly selected ten percent from each subgroup for MRI brain volumetry. Crude volumes were standardized to reflect percentage of intracranial volume.

Results: MRI volumetry studies were done in 199 individuals (F/M = 90/109; 73 with BMI ≥ 25 ; 44 hypertensives; 29 diabetics; mean cognitive score 76.5). Both grey and white matter volumes decreased with increasing age. WMHV increased with age and hypertension. Grey matter volume decreased with increasing WMHV. Positive predictors of cognition included standardized hippocampal volume (HCV), urban living, education, and BMI, while WMHV and age were negative predictors. Urban dwellers had higher cognitive scores than rural, and (paradoxically) smaller HCV.

Conclusions: Further study is warranted into sociodemographic and biological factors that mutually influence cognition. Our findings contribute to baseline diagnostic criteria for VCI and could help in early diagnosis and control of cognitive decline and its key risk factors.

Keywords: Vascular Cognitive Impairment; Cognitive aging; Community-based Study; Cognitive Testing; Volumetric MRI; White Matter Hyperintensities

Manuscript

1. Introduction:

India is a highly populated, rapidly ageing nation. The proportion of the Indian population aged over 60 is projected to increase from 8 percent in 2015 to 19 percent in 2050.¹ This transition can significantly change the Indian morbidity profile, with chronic conditions such as age-related cognitive decline and comorbidities exacting a formidable toll on the healthcare apparatus. Hypertension and diabetes mellitus, and other cardiovascular risk factors such as smoking, are known risk factors for cognitive impairment.²⁻⁴ While definitional criteria exist for Alzheimer's Disease and Vascular Dementia, the same is not true for Vascular Cognitive Impairment (VCI). There is a lack of clarity on what constitutes VCI. To diagnose VCI clinically, one must have some degree of clinical cognitive impairment (short of dementia), corroborated by imaging findings of vascular compromise. The latter would comprise subclinical lacunes, microhemorrhages, periventricular white matter hyperintensities (WMH), and at times even non-lacunar small embolic infarcts or bleeds which may not be clinically overt. However, it is well known that some of these imaging features, especially WMH, can coexist with vascular risk factors (VRFs) such as advancing age, hypertension, diabetes mellitus, obesity and hyperlipidemia, without overt clinical motor-sensory deficits. In regions with a high prevalence of undetected or untreated VRF, such as India, these factors can confound the definition of VCI.

Thus, it is necessary to know the baseline values of cognition in different age groups in populations without prior stroke or dementia in particular regions, in correlation with prevalent VRFs and brain volumetry, to lay down the 'normative data' specific to that population. Based on this information, the definition of VCI can be formulated for that population.

Hence, our objectives in this study were to determine cognitive scores in an age-stratified 'normal' population (with no known history of prior clinical strokes or cognitive/behavioral disorders) from the western part of Maharashtra state in India (both rural and urban areas), to obtain data on VRFs in

this population (hypertension, diabetes mellitus, alcohol and tobacco use, body mass index), and after stratification for cognition in each age bin, to obtain volumetry data on MRI in each stratified bin as a representative sample of that bin. These baseline parameters would serve as ‘normative data’ of cognitive scores in relation to age, VRFs and brain volumetry (including volume of WMHs, if noted), and would hence enable a definition of VCI specific to this population, in further studies.

2. Materials and Methods:

2.1. Study design:

Initially, we translated the Kolkata Cognitive Battery (KCB), a partially modified Hindi cognitive screening battery used by Ganguli et al.⁵ and the Geriatric Depression Score (GDS),⁶ into Marathi and validated both for field use. The KCB is a screening battery consisting of eight tests designed to test different cognitive domains. Test 1 is for Verbal Fluency, 2 is for Object Naming, 3 is the Mini Mental State Examination (MMSE), 4 is a Calculation task, 5 tests Immediate Recall, 6 tests Visuo-Constructional Ability, 7 tests Delayed Recall, and 8 tests Delayed Word Recognition. We used these instruments to conduct a cross-sectional interview-based survey in several urban and rural communities in Maharashtra state. We selected the urban study population from Dharavi and other neighborhoods in Mumbai city. The rural population was chosen from villages that had ongoing primary health care projects under the community health department of the parent hospital, which made it feasible to obtain information on the number of households, their family size, and the number of people over 40 years of age in these areas. We conducted awareness programs about the study in these communities in the month preceding the survey, through pamphlets and doorstep campaigning with the help of community health workers.

2.2. Data collection:

We invited all persons aged 40 and over, and living in the selected localities, to take part in the survey. We screened 2752 persons who consented to participate in our study, from March 2015 to July 2016, and excluded 19 with known intellectual developmental disorder, 31 with possible previous stroke, 36 with previous psychiatric illness, and 15 persons reported by family to have

previously diagnosed dementia or significant memory impairment impeding daily activities. We interviewed the remaining 2651 participants and collected their sociodemographic data, noted current daily use of tobacco and alcohol, and measured their height, weight, blood pressure (BP) and fasting blood glucose. Participants were classified as having hypertension, diabetes mellitus, or other cardiovascular conditions based on self-report corroborated by medical records and prescriptions. Participants who had high blood glucose or high BP detected for the first time during examination, had repeat BP measurements, and fasting and post prandial blood sugar study on a later date, to confirm the diagnoses in accordance with standard criteria.^{7,8}

Finally, trained personnel screened the participants using the validated Marathi KCB and GDS tools and identified 1961 individuals with GDS scores in the normal range ($GDS \leq 9$). We stratified this group into four subgroups using ten-year age bins (40–49, 50–59, 60–69 and ≥ 70). Each age group was further sub-stratified into three cognitive classes based on their KCB score (< 80 , 80–100 and > 100). Our final sample for MRI brain imaging ($n=199$) comprised ten percent of the individuals from each cognitive category within each age bin (12 subgroups in total), selected using computer generated random numbers.

2.3. Volumetry:

We used a Siemens 1.5T MRI system (1.5 T Avanto, Siemens Healthcare, Germany), equipped with a 32-channel head coil, for MRI scanning. For each participant, we acquired 3D Fluid sensitive T2 space Inversion Recovery (FLAIR) images with repetition time = 6000 ms, echo time = 369 ms, flip angle = 90° , field of view = 260 mm $\times$ 208 mm $\times$ 174 mm, slice thickness = 1.14 mm, 128 slices per slab, matrix = 256 $\times$ 174, and voxel size = 1.01 mm $\times$ 1.19 mm $\times$ 1.14 mm.

2.4. Image processing:

Anatomical images were segmented into grey matter, white matter and cerebrospinal fluid, followed by spatial normalization of images into stereotactic space using the voxel-based morphometry tool box in SPM12.⁹ The normalized images were smoothed with an 8 mm isotropic Gaussian kernel. The hippocampal regional volume was extracted by atlas based parcellation using automated anatomical

labelling, AAL atlas.^{10,11} Automated white matter hyperintensity (WMH) detection was done using FLAIR images with the lesion segmentation tool box (LST) implemented for SPM12 package.^{12,13} We determined WMH volume using lesion prediction algorithm (LPA SPM12). The processing methodology is described in detail elsewhere.¹⁴ Post processing was done by an independent team with prior experience in volumetric studies of Indian populations,^{14,15} who were blinded to the neuropsychological and demographic data.

2.5. Statistical methods:

In order to adjust for differences in cranial size, we standardized crude volumetric measurements and expressed them as a percentage of the total intracranial volume. We analyzed the data using IBM SPSS Statistics for Windows, Version 20.0. (Armonk, NY: IBM Corp) and JASP.¹⁶ We used the Mann Whitney U test, the Chi square test, and Welch's unequal variances t-test, to study differences in sociodemographic and vascular risk factors. We used the Wilcoxon signed rank test to compare left and right hippocampal volumes, Spearman's rank correlation coefficient for non-parametric correlations after testing for normality and equality of variances, and Pearson product-moment correlation to study correlation between variables after adjusting for confounders. Our analysis is primarily explorative, without a definite outcome variable. Therefore, we did not correct our obtained P values for multiple comparisons. While reporting results, we have replaced P values with estimates of effects or association and corresponding 95% confidence intervals. We used rank biserial correlation coefficients as indicators of effect size for Wilcoxon T and Mann Whitney U tests, Cramer's V for the Chi square test, and Cohen's d for Welch's t test. We used linear regression to identify key predictors of volumetric variables and cognition scores. We entered area of residence, sex, age, education, body mass index (BMI), vascular risk factors, blood glucose and pressure readings, and brain regional volumes found to be associated in bivariate analysis, as independent factors for volumetric dependent variables. For cognitive dependent variables, we entered area of residence, sex, 10-year age groups, educational level, BMI, vascular risk factors, blood glucose and pressure readings, and standardized brain region volumes as independent factors. Each regression

model was checked for multicollinearity, and independent variables with variance inflation factor (VIF) more than five, or tolerance less than 0.4 were removed from the final models. We have reported P values as an additional data point for predictor variables in the regression models. However, we have selected predictors to report based on 95% confidence intervals, in place of P value.

2.6. Glossary of abbreviations:

Cognitive scores: KCB = Kolkata Cognitive Battery; KCB Memory = Memory component: Test scores 5, 7, 8 combined; KCB Executive = Executive component: Test scores 1, 2, 4, 6 combined.

Crude regional brain volumes (measured in milliliters): GMV = Grey matter volume; WMV = White matter volume; CSFV = Cerebrospinal fluid volume; LHCV = Left hippocampal volume; RHCV = Right hippocampal volume; HCV = Total hippocampal volume; WMHV = White matter hyperintensity volume; TIV = Total intracranial volume.

Standardized volumes: All expressed with a % sign suffixed to the corresponding crude volume abbreviation (such as GMV for grey matter volume and GMV% for standardized grey matter volume).

Vascular risk factors: HTN = Hypertension; DM = Diabetes mellitus; IHD = Ischemic heart disease; BMI = Body mass index; SBP = Systolic BP; DBP = Diastolic BP; PPR = Pulse pressure; Tobacco = Current daily use of tobacco in any form; Alcohol = Current daily use of alcohol in any form.

Statistical tests: MWU = Mann Whitney U test; Wilcoxon T = Wilcoxon signed rank test; ρ = Spearman's rank correlation coefficient; r = Pearson product-moment correlation; X^2 = Chi square; Welch t = Welch's unequal variances t-test; RC = Rank biserial correlation coefficient; OR = Odds ratio; 95% CI = 95% confidence intervals.

2.7. Standard Protocol Approvals, Registrations, and Patient Consents:

The study design was approved by the Cognitive Science Research Initiative of the Department of Science & Technology under the Government of India, and ethical approval was granted by the

institutional ethical committee of Lokmanya Tilak Municipal Medical College and General Hospital, Sion, Mumbai. Written informed consent was obtained from all participants in our study.

2.8.Data availability:

The master dataset has not been made publicly available as it contains confidential sociodemographic and clinical data of the participants. Anonymized data will be shared by reasonable request from any qualified investigator.

3. Results:

3.1.Demography: (Table 1) Our study sample comprised 199 individuals (109 men), three-fourths of whom (150/199) were between 40 and 59 years of age. A similar proportion (146/199) were poorly educated, having completed primary school or less. More than one third were overweight or obese (73/199; 36.9%).

3.2.Cognitive scores: The mean cognitive score (KCB) of our sample was 76.5 (range 21–125). The KCB score decreased with increasing age and increased with education. Average KCB score was notably lower in rural as compared to urban participants (Tables 2, 3).

3.3.Volumetry and its associations:

On average, both crude and standardized volumes were smaller for the left hippocampus than the right (Table 2).

3.3.1. Age: GMV% and WMV% decreased with increasing age. CSFV and WMHV, both crude and standardized, increased with age (Table 3, Fig 1).

3.3.2. Sex: Crude volumes were lower in women than men, but after standardization, this difference disappeared for all variables save WMV% (higher in women) and CSFV% (higher in men); WMHV did not show a definite sex difference (Tables 2, 3).

3.3.3. Locality: Crude and standardized CSFV and HCV were on average larger in rural participants, while average WMV% was larger in city dwellers (Table 3).

3.3.4. Education: We did not note a uniform increase or decrease in regional brain volumes with years of education. However, compared to literate participants, illiterate participants (n = 29) had smaller mean GMV% (38.5 vs 40.9), higher mean CSFV% (16.3 vs 13.6) and higher mean WMHV% (0.3 vs 0.1). Illiterate participants also had higher average CSFV (198.9 vs 166.6 ml) and WMHV (3.8 vs 1.4 ml). Mean crude and standardized volumes of the right hippocampus were higher in illiterate participants than literate participants (3.5 vs 3.2 ml; and 0.29 vs 0.26 percent). Among our literate participants, crude and standardized hippocampal volumes increased with years of education after controlling for age (Fig 2).

3.3.5. VRFs: Diastolic BP (DBP) correlated positively with crude and standardized hippocampal volumes. (Table 4, Fig 3). Correlation controlling for age showed both Systolic BP (SBP) and pulse pressure (PPR) to correlate positively with WMH volumes (Table 4).

3.3.6. Cognitive tests: High scores in Test 8 of the KCB (Delayed Word Recognition) were correlated with higher hippocampal volumes (Table 4).

3.4. Inter-relationships of regional volumes, controlled for age:

GMV% decreased with increasing WMHV% ($r = -0.60$, 95% CI = -0.77 – -0.26; see Fig 1). CSFV% correlated negatively with both GMV% ($r = -0.41$, 95% CI = -0.55 – -0.29) and WMV% ($r = -0.49$, 95% CI = -0.61 – -0.39).

3.5. Multivariate analysis:

(Table 5) Urban living, higher education and BMI predicted higher KCB scores, while age and WMHV% predicted reduced scores. We noted that HCV% and education predicted an increase in the executive component of the KCB, while BMI predicted increase in the memory component.

Female sex and WMHV were the key predictors of decreased GMV. Post standardization, however, female sex and hypertension predicted increase in GMV%, while WMHV%, HCV% and age predicted decrease. For both crude and standardized HCV, education predicted increase in volumes while urban living predicted decrease. Age, hypertension and SBP predicted increased WMHV.

4. Discussion:

Image-based definition of VCI poses difficulties in India, with its high prevalence of vascular risk factors, often undetected and untreated, resulting in subclinical vascular findings on the MRI. It is difficult to determine whether these MRI findings indicate or predict a true VCI, or whether they are normal for age and associated vascular risk factors. Our study sample consisted of participants who had no prior history of stroke or clinical dementia, and who, after screening, did not suffer from depression. Our data would thus enable an evaluation of MRI volumetry in correlation with vascular risk factors on the one hand, and with age-stratified cognitive scores on the other.

Among our study participants, 22.11% were hypertensive and 14.57% were diabetic. Compared to rural participants, urban dwellers had a higher percentage of hypertensives (26.2% vs 15.6%) as well as diabetics (19.7% vs 6.5%; $X^2 = 6.59$, Cramer's $V = 0.18$, $OR = 0.28$, 95% $CI_{OR} = 0.10 - 0.78$). This finding is consistent with the results of a recent nationally representative study, although our urban-rural differences are wider.¹⁷

The men in our sample had larger mean volumes than women for all brain regions. This can be explained by difference in cranial size, as standardized volumetric values were similar for both.

The average right hippocampal volume (both crude and standardized) was larger than the left among our participants; similar findings have been reported by researchers from India and abroad.¹⁸⁻²⁰

Our study shows an increase in hippocampal volume with diastolic BP. Lower DBP has been shown to be associated with smaller right hippocampal volume in healthy elderly individuals.²¹ Lower SBP, DBP, and mean arterial pressure (MAP) have also been associated with smaller volumes of the thalamus and putamen, with lower MAP associated with reduced HCV in elderly hypertensives, and

281 it has been suggested that in order to ensure adequate perfusion, the target BP should not be too low
 282 in elderly hypertensives.²²

283 Our observation of literate participants having a higher mean GMV% than illiterate, is consistent
 284 with reports of volume and metabolism increases in GMV with education in the healthy elderly,²³
 285 and similar volumetric findings by other researchers.²⁴

286 Literature indicates a positive effect of education on hippocampal volume.²⁵ It has been
 287 demonstrated that education may exert a “dose-dependent” effect on HCV, with age-related decrease
 288 in volumes being steepest among those least educated, and less pronounced in individuals with
 289 higher educational attainment.²⁶ The results of our bivariate and multivariate analyses (a positive
 290 correlation between HCV and years of education in literate participants, and education predicting
 291 increased HCV in linear regression) provide further evidence of this association. It is possible that, in
 292 our sample of healthy individuals, higher levels of education were able to counteract typical age-
 293 related changes in hippocampal structure. The differential rates of decline may be functionally
 294 significant, resulting in greater retention of the cognitive functions, including declarative memory
 295 performance, which relies on this structure.²⁷

296 Hippocampal volume loss has been reported to be specifically associated with decline in delayed
 297 word recall.²⁸ This is indirectly supported by our observation of the association of hippocampal
 298 volumes with the delayed word recognition score from the KCB.

299 Our illiterate participants were older than literate (mean age 60.7 vs 51.0 years, Welch $t = 4.47$,
 300 Cohen’s $d = 0.96$, 95% $CI_d = 0.50 - 1.41$), with higher percentages of alcohol (24.1% vs 0.6%) and
 301 tobacco (27.6% vs 2.9%) users. We also noted a greater proportion of hypertensives (27.6% vs
 302 21.2%) and diabetics (24.1% vs 12.9%) among them. These factors may explain their higher
 303 observed WMHV.

Our correlations of WMHV with age, blood pressure, and illiteracy are consistent with literature establishing age, hypertension, diabetes mellitus, smoking and low educational level as important risk factors for WMHV.^{29–34}

Urban versus rural living deserves close examination as a possible determinant of vascular and neurological morbidity. Our findings are concordant with several Indian and international studies which have reported association of urban residence with higher cognitive scores.^{35–41} Our finding of lower mean cognitive scores and HCVs in women could not have influenced this association, as women were equally represented in both subsets (45.5% rural vs 45.1% urban).

It is noteworthy that our urban participants had smaller volumes than rural for all volumetric variables including HCV, despite their higher cognitive scores, and in contradiction to our regression analysis showing that HCV% predicts high executive function. Urban living alters brain structures, and has been linked to mental disorders such as schizophrenia.^{42,43} Animal models^{44,45} and observational studies of post-traumatic stress disorder patients,^{46–48} demonstrate that hippocampal volume is inversely correlated with exposure to stress. Our urban participants, most of whom were of a low socio-economic status, would possibly have faced the challenges of underprivileged backgrounds in their daily lives, which could explain their comparatively smaller HCV. The discrepancy between volume and cognition in urban dwellers could also be explained by the fact that age, education, WMHV and BMI, along with urban residence itself, also independently predicted cognition in this study. Our urban participants were younger than their rural counterparts (mean age 51.41 vs 54.04 years), and had longer years in school (5.32 vs 3.03 Welch $t = -4.94$, Cohen's $d = -0.711$, 95% $CI_d = -1.005 - -0.415$), smaller mean WMHV (1.52 vs 2.00 ml, 0.13 vs 0.16 %), and higher mean BMI (24.22 vs. 23.62), all of which could override the effect of smaller HCV. Results from a study with a participant profile similar to ours (healthy, non-demented, relatively young), do not indicate any association of smaller HCV with cognitive decline.⁴⁹ In another study, it was found that baseline hippocampal volume, albeit strongly related to baseline cognition, did not predict

subsequent cognitive change.⁵⁰ Researchers have noted that not all subjects with hippocampal decline will develop dementia, either due to death preceding symptoms, or because hippocampal volume decline can be part of healthy ageing.²⁸ The apparent paradox of smaller HCV with higher cognition in city dwellers may be explained by the (possibly synergistic) effect of sociodemographic, behavioral, and environmental factors that compensate for possible stress-induced hippocampal volume loss.

4.1.Limitations:

Persons aged over 70 years and rural dwellers were inadequately represented in our sample due to the logistical difficulties of transporting the elderly, and of bringing people over long distances from rural areas to our urban MRI center. Our study is cross-sectional. While we have established associations between cognition, brain regional volumes and vascular risk factors, we cannot comment on causality, or the longitudinal ramifications of observed volumetric differences on cognitive decline, without follow-up investigation.

5. Conclusions:

Our population-based study of adults without neuropsychiatric morbidity attempts to place cognitive scores stratified by age and education, within the context of MRI brain volumetry findings. Our study shows that both GMV and WMV decrease, and WMHV increases, with increasing age. Our findings also suggest a shrinkage of GMV with increasing WMHV. Urban living, education, and standardized hippocampal volume predict higher cognitive scores, while WMHV and age predict lower scores. We noted higher cognitive scores in city dwellers with, paradoxically, smaller hippocampal volumes. Further study is warranted into sociodemographic and biological factors that mutually influence cognition. Our findings, particularly the correlation of KCB scores with WMHV and age, could help guide the definition of VCI in India.

Funding information: This study was fully funded by the Cognitive Science Research Initiative of the Department of Science & Technology, Government of India.

Acknowledgment: The authors thank Dr Kuldeep Singh, Radiologist at Gurmeet Imaging, Mumbai, for his role in obtaining MRI volumetry sequences.

Declarations of interest: None.

6. References:

1. India Ageing Report - 2017 (Final Version).pdf [online]. Accessed at: <https://india.unfpa.org/sites/default/files/pub-pdf/India%20Ageing%20Report%20-%202017%20%28Final%20Version%29.pdf>. Accessed December 14, 2019.
2. Dregan A, Stewart R, Gulliford MC. Cardiovascular risk factors and cognitive decline in adults aged 50 and over: a population-based cohort study. *Age Ageing*. 2012;42:338–345.
3. Baumgart M, Snyder HM, Carrillo MC, Fazio S, Kim H, Johns H. Summary of the evidence on modifiable risk factors for cognitive decline and dementia: A population-based perspective. *Alzheimers Dement*. 2015;11:718–726.
4. Kaffashian S, Dugravot A, Elbaz A, et al. Predicting cognitive decline. *Neurology*. 2013;80:1300.
5. Ganguli M, Chandra V, Gilby JE, et al. Cognitive test performance in a community-based nondemented elderly sample in rural India: the Indo-U.S. Cross-National Dementia Epidemiology Study. *Int Psychogeriatr*. 1996;8:507–524.
6. Yesavage JA, Brink TL, Rose TL, et al. Development and validation of a geriatric depression screening scale: a preliminary report. *J Psychiatr Res*. 1982;17:37–49.

- 375 7. James PA, Oparil S, Carter BL, et al. 2014 evidence-based guideline for the management of
376 high blood pressure in adults: report from the panel members appointed to the Eighth Joint
377 National Committee (JNC 8). JAMA. 2014;311:507–520.
- 378 8. Diagnosis and Classification of Diabetes Mellitus. Diabetes Care. 2010;33:S62–S69.
- 379 9. SPM - Statistical Parametric Mapping [online]. Wellcome Cent. Hum. Neuroimaging UCL
380 Queen Sq. Inst. Neurol. Lond. UK. Accessed at: <https://www.fil.ion.ucl.ac.uk/spm/>. Accessed
381 July 18, 2020.
- 382 10. Tzourio-Mazoyer N, Landeau B, Papathanassiou D, et al. Automated anatomical labeling of
383 activations in SPM using a macroscopic anatomical parcellation of the MNI MRI single-subject
384 brain. NeuroImage. 2002;15:273–289.
- 385 11. CyCeron - Biomedical Imaging Platform. 2015. Accessed at:
386 <http://www.cyceron.fr/index.php/en/plateforme-en/freeware>. Accessed July 18, 2020.
- 387 12. Schmidt P, Gaser C, Arsic M, et al. An automated tool for detection of FLAIR-hyperintense
388 white-matter lesions in Multiple Sclerosis. NeuroImage. 2012;59:3774–3783.
- 389 13. Paul Schmidt. LST – Lesion segmentation for SPM [online]. Accessed at: [https://www.applied-](https://www.applied-statistics.de/lst.html)
390 [statistics.de/lst.html](https://www.applied-statistics.de/lst.html). Accessed July 18, 2020.
- 391 14. Sheelakumari R, Kesavadas C, Lekha VS, Justus S, Sarma PS, Menon R. Structural correlates
392 of mild cognitive impairment: A clinicovolumetric study. Neurol India. 2018;66:370.
- 393 15. Sheelakumari R, Sarma SP, Kesavadas C, et al. Multimodality Neuroimaging in Mild Cognitive
394 Impairment: A Cross-sectional Comparison Study. Ann Indian Acad Neurol. 2018;21:133–139.

- 395 16. JASP Team. JASP (Version 0.13)[Computer software] [online]. 2020. Accessed at: [https://jasp-](https://jasp-stats.org/)
396 stats.org/. Accessed July 15, 2020.
- 397 17. Geldsetzer P, Manne-Goehler J, Theilmann M, et al. Diabetes and Hypertension in India: A
398 Nationally Representative Study of 1.3 Million Adults. JAMA Intern Med. 2018;178:363–372.
- 399 18. Mohandas A, Bharath R, Prathyusha P, Gupta A. Hippocampal volumetry: Normative data in
400 the Indian population. Ann Indian Acad Neurol. 2014;17:267–271.
- 401 19. Embong MF, Yaacob R, Abdullah MS, Abdulkarim AH, Ghazali AK, Jalaluddin W. MR
402 Volumetry of Hippocampus in Normal Adult Malay of Age 50 Years Old and Above. Malays J
403 Med Sci MJMS. 2013;20:25–31.
- 404 20. Pruessner JC, Li LM, Serles W, et al. Volumetry of Hippocampus and Amygdala with High-
405 resolution MRI and Three-dimensional Analysis Software: Minimizing the Discrepancies
406 between Laboratories. Cereb Cortex. 2000;10:433–442.
- 407 21. Elcombe EL, Lagopoulos J, Duffy SL, et al. Hippocampal volume in older adults at risk of
408 cognitive decline: the role of sleep, vascular risk, and depression. J Alzheimers Dis JAD.
409 2015;44:1279–1290.
- 410 22. Foster-Dingley JC, van der Grond J, Moonen JEF, et al. Lower Blood Pressure Is Associated
411 With Smaller Subcortical Brain Volumes in Older Persons. Am J Hypertens. Oxford Academic;
412 2015;28:1127–1133.
- 413 23. Arenaza-Urquijo EM, Landeau B, La Joie R, et al. Relationships between years of education
414 and gray matter volume, metabolism and functional connectivity in healthy elders. NeuroImage.
415 2013;83:450–457.

- 416 24. Liu Y, Julkunen V, Paajanen T, et al. Education increases reserve against Alzheimer's
417 disease—evidence from structural MRI analysis. *Neuroradiology*. 2012;54:929–938.
- 418 25. Nobis L, Manohar SG, Smith SM, et al. Hippocampal volume across age: Nomograms derived
419 from over 19,700 people in UK Biobank. *NeuroImage Clin*. 2019;23:101904.
- 420 26. Noble KG, Grieve SM, Korgaonkar MS, et al. Hippocampal volume varies with educational
421 attainment across the life-span. *Front Hum Neurosci* [online serial]. 2012;6. Accessed at:
422 <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3494123/>. Accessed July 25, 2020.
- 423 27. Grieve SM, Korgaonkar MS, Clark CR, Williams LM. Regional heterogeneity in limbic
424 maturational changes: evidence from integrating cortical thickness, volumetric and diffusion
425 tensor imaging measures. *NeuroImage*. 2011;55:868–879.
- 426 28. den Heijer T, van der Lijn F, Koudstaal PJ, et al. A 10-year follow-up of hippocampal volume
427 on magnetic resonance imaging in early dementia and cognitive decline. *Brain J Neurol*.
428 2010;133:1163–1172.
- 429 29. Valdés Hernández M del C, Booth T, Murray C, et al. Brain white matter damage in aging and
430 cognitive ability in youth and older age. *Neurobiol Aging*. 2013;34:2740–2747.
- 431 30. Breteler MM, van Swieten JC, Bots ML, et al. Cerebral white matter lesions, vascular risk
432 factors, and cognitive function in a population-based study: the Rotterdam Study. *Neurology*.
433 United States; 1994;44:1246–1252.
- 434 31. Longstreth WTJ, Manolio TA, Arnold A, et al. Clinical correlates of white matter findings on
435 cranial magnetic resonance imaging of 3301 elderly people. The Cardiovascular Health Study.
436 *Stroke*. United States; 1996;27:1274–1282.

- 437 32. Ylikoski A, Erkinjuntti T, Raininko R, Sarna S, Sulkava R, Tilvis R. White matter
438 hyperintensities on MRI in the neurologically nondiseased elderly. Analysis of cohorts of
439 consecutive subjects aged 55 to 85 years living at home. *Stroke*. United States; 1995;26:1171–
440 1177.
- 441 33. Modir R, Gardener H, Wright CB. Blood Pressure and White Matter Hyperintensity Volume -
442 A Review of the Relationship and Implications for Stroke Prediction and Prevention. *Eur*
443 *Neurol Rev*. 2012;7:174.
- 444 34. Habes M, Erus G, Toledo JB, et al. White matter hyperintensities and imaging patterns of brain
445 ageing in the general population. *Brain J Neurol*. England; 2016;139:1164–1179.
- 446 35. Xu H, Ostbye T, Vorderstrasse AA, Dupre M, Wu B. Place of Residence and Cognitive
447 Function among the Adult Population in India. *Neuroepidemiology*. 2018;50:119–127.
- 448 36. Patel RM, Singh US. Prevalence study of cognitive impairment and its associated
449 sociodemographic variables using mini-mental status examination among elderly population
450 residing in field practice areas of a medical college. *Indian J Community Med*. 2018;43:113.
- 451 37. Poddar K, Kant S, Singh A, Singh TB. An epidemiological study of dementia among the
452 habitants of Eastern Uttar Pradesh, India. *Ann Indian Acad Neurol*. 2011;14:164.
- 453 38. Nadel JL, Ulate D. Incidence and risk factors for cognitive impairment in rural elderly
454 populations in Costa Rica. *Rev Biol Trop*. 2014;62:869–876.
- 455 39. Nunes B, Silva RD, Cruz VT, Roriz JM, Pais J, Silva MC. Prevalence and pattern of cognitive
456 impairment in rural and urban populations from Northern Portugal. *BMC Neurol*. 2010;10:42.
- 457 40. Saenz JL, Downer B, Garcia MA, Wong R. Cognition and Context: Rural–Urban Differences in
458 Cognitive Aging Among Older Mexican Adults. *J Aging Health*. 2018;30:965–986.

- 459 41. Xiang Y, Zare H, Guan C, Gaskin D. The impact of rural-urban community settings on
460 cognitive decline: results from a nationally-representative sample of seniors in China. BMC
461 Geriatr. 2018;18:323.
- 462 42. Besteher B, Gaser C, Spalthoff R, Nenadić I. Associations between urban upbringing and
463 cortical thickness and gyrification. J Psychiatr Res. 2017;95.
- 464 43. Haddad L, Schäfer A, Streit F, et al. Brain Structure Correlates of Urban Upbringing, an
465 Environmental Risk Factor for Schizophrenia. Schizophr Bull. 2014;41.
- 466 44. Ohl F, Michaelis T, Vollmann-Honsdorf GK, Kirschbaum C, Fuchs E. Effect of chronic
467 psychosocial stress and long-term cortisol treatment on hippocampus-mediated memory and
468 hippocampal volume: a pilot-study in tree shrews. Psychoneuroendocrinology. 2000;25:357–
469 363.
- 470 45. Lee T, Jarome T, Li S-J, Kim JJ, Helmstetter FJ. Chronic stress selectively reduces
471 hippocampal volume in rats: a longitudinal magnetic resonance imaging study. Neuroreport.
472 2009;20:1554–1558.
- 473 46. Smith ME. Bilateral hippocampal volume reduction in adults with post-traumatic stress
474 disorder: a meta-analysis of structural MRI studies. Hippocampus. 2005;15:798–807.
- 475 47. Woon FL, Hedges DW. Hippocampal and amygdala volumes in children and adults with
476 childhood maltreatment-related posttraumatic stress disorder: a meta-analysis. Hippocampus.
477 2008;18:729–736.
- 478 48. Villarreal G, Hamilton DA, Petropoulos H, et al. Reduced hippocampal volume and total white
479 matter volume in posttraumatic stress disorder. Biol Psychiatry. 2002;52:119–125.

49. Vibha D, Tiemeier H, Mirza SS, et al. Brain Volumes and Longitudinal Cognitive Change: A Population-based Study. *Alzheimer Dis Assoc Disord.* 2018;32:43–49.
50. Carmichael O, Mungas D, Beckett L, et al. MRI predictors of cognitive change in a diverse and carefully characterized elderly population. *Neurobiol Aging.* 2012;33:83–95.

Tables

Table 1: Socio-demography and Vascular Risk Factors

Demographic variables		Men		Women		Total	
		Count	%	Count	%	Count	%
Sample Total ^a		109	54.8	90	45.2	199	100
10-year age groups	40–49	45	41.3	42	46.7	87	43.7
	50–59	35	32.1	28	31.1	63	31.7
	60–69	20	18.3	15	16.7	35	17.6
	70+	9	8.3	5	5.6	14	7.0
Educational level (years of schooling)	Illiterate	14	12.8	15	16.7	29	14.6
	Primary (1–5)	64	58.7	53	58.9	117	58.8
	Secondary (6–12)	30	27.5	20	22.2	50	25.1
	Graduate (≥13)	1	0.9	2	2.2	3	1.5
Area of residence	Rural	42	38.5	35	38.9	77	38.7
	Urban	67	61.5	55	61.1	122	61.3
Vascular Risk Factors	HTN	26	23.9	18	20.0	44	22.1
	DM	18	16.5	11	12.2	29	14.6
	IHD ^b	2	1.8	2	2.2	4	2.0
	Tobacco	12	11.0	1	1.1	13	6.5
	Alcohol	7	6.4	1	1.1	8	4.0
BMI ^c	Underweight (<18.5)	7	6.4	7	7.9	14	7.0
	Overweight (25–30)	31	28.4	25	28.1	56	28.3
	Obese (>30)	9	8.3	8	9.0	17	8.6

^a Row percentage for sample total; column percentages for all other variables.

^b IHD = Ischemic heart disease; ^c BMI measurements were available for 89 of 90 women.

508 **Table 2: Descriptive Statistics for MRI Volumetry and Cognitive Scores**

	Mean	Median	SD	Range	Minimum	Maximum	Mean (M)	Mean (F)	Mean (R)	Mean (U)
Crude Volumes (ml)										
GMV	487.8	489.9	73.5	582.4	161.6	744.0	515.2	454.6	493.4	484.2
WMV	545.2	545.9	73.0	605.8	214.5	820.3	568.5	517.1	546.8	544.2
CSFV	171.3	162.2	64.2	307.7	55.4	363.2	193.5	144.3	186.5	161.6
LHCV ^a	3.1	2.8	0.9	4.4	0.9	5.3	3.3	2.9	3.5	2.9
RHCV	3.2	2.9	0.9	4.8	0.8	5.6	3.4	2.9	3.6	3.0
HCV	6.3	5.7	1.8	9.0	1.7	10.7	6.7	5.8	7.1	5.8
WMHV	1.7	0.3	4.4	39.6	0	39.6	2.4	0.9	2.0	1.5
TIV	1204.2	1218.7	141.7	1161.9	458.1	1620.0	1277.2	1116.0	1226.7	1190.1
Standardized Volumes (percentage of TIV)										
GMV%	40.6	40.8	4.3	46.7	13.1	59.8	40.4	40.8	40.3	40.8
WMV%	45.4	45.5	4.4	44.8	21.5	66.3	44.6	46.4	44.7	45.9
CSFV%	14.0	13.4	4.5	22.2	6.4	28.6	15.1	12.8	15.1	13.4
LHCV% ^b	0.260	0.246	0.069	0.397	0.080	0.477	0.259	0.260	0.28	0.24
RHCV%	0.266	0.249	0.071	0.393	0.071	0.464	0.270	0.261	0.29	0.25
HCV%	0.525	0.490	0.138	0.790	0.151	0.941	0.529	0.521	0.58	0.49
WMHV%	0.140	0.026	0.357	3.204	0	3.204	0.187	0.083	0.16	0.13
Cognitive Scores										
KCB Score	76.5	79	21.8	104	21	125	78.3	74.2	65.3	83.5
KCB Memory	23.6	24	9.1	40	4	44	24.4	22.6	20.9	25.3

KCB	30.1	31	9.8	48	5	53	31.0	29.1	25.3	33.2
Executive										

SD = Standard Deviation; M = Male; F = Female; R = Rural; U = Urban.

^a LHCV<RHCV: Wilcoxon T = 6232.5, z = -3.93, RC = -0.33, 95% CI_{RC} = -0.46 – -0.17

^b LHCV%<RHCV%: Wilcoxon T = 6315, z = -3.83, RC = -0.32, 95% CI_{RC} = -0.46 – -0.17

KCB consists of eight tests of different cognitive domains. Test1: Verbal Fluency, Test2: Object

Naming, Test3: MMSE, Test4: Calculation, Test5: Immediate Recall, Test6: Visuo-Constructional

Ability, Test7: Delayed Recall, Test8: Delayed Word Recognition.

KCB(Memory) = Memory component (Test scores 5, 7, 8 combined)

KCB(Executive) = Executive component (Test scores 1, 2, 4, 6 combined).

532 **Table 3: Sociodemographic Associations of Volumetry and Cognition**

	Age		Years of Education		Sex			Residence		
	ρ	95% CIs for ρ	ρ	95% CIs for ρ	MWU	RC	95% CIs for RC	MWU	RC	95% CIs for RC
Crude Volumes										
GMV	-0.14	-0.30 – 0.03	0.02	-0.13 – 0.15	7754	0.58	0.46 – 0.68	4997	0.06	-0.10 – 0.23
WMV	-0.13	-0.27 – 0.01	0.01	-0.14 – 0.15	7384.5	0.51	0.38 – 0.62	4687.5	-0.002	-0.17 – 0.16
CSFV	0.46	0.36 – 0.57	-0.04	-0.17 – 0.10	7149	0.46	0.32 – 0.58	5897.5	0.26	0.10 – 0.40
LHCV	0.11	-0.03 – 0.28	0.01	-0.12 – 0.13	6277	0.28	0.13 – 0.42	6373	0.36	0.21 – 0.49
RHCV	0.07	-0.07 – 0.22	0.04	-0.09 – 0.16	6477	0.32	0.17 – 0.46	6421.5	0.37	0.22 – 0.50
HCV	0.09	-0.05 – 0.24	0.02	-0.11 – 0.14	6402.5	0.31	0.15 – 0.44	6447	0.37	0.22 – 0.51
WMHV	0.48	0.37 – 0.59	-0.12	-0.26 – 0.03	5532	0.13	-0.03 – 0.28	4942.5	0.05	-0.11 – 0.21
TIV	0.08	-0.07 – 0.21	0.03	-0.10 – 0.15	8556	0.74	0.66 – 0.81	5240	0.12	-0.05 – 0.27
Standardized Volumes										
GMV%	-0.29	-0.42 – -0.16	0.03	-0.12 – 0.18	4726	-0.04	-0.20 – 0.12	4359	-0.07	-0.23 – 0.09
WMV%	-0.33	-0.45 – -0.20	0.01	-0.15 – 0.17	3368	-0.31	-0.45 – -0.16	3748	-0.20	-0.35 – -0.04
CSFV%	0.52	0.41 – 0.62	-0.05	-0.20 – 0.09	6380	0.30	0.15 – 0.44	5926	0.26	0.10 – 0.41

LHCV%	0.11	-0.03 – 0.24	0.04	-0.10 – 0.17	4917.5	0.003	-0.16 – 0.16	6271.5	0.34	0.18 – 0.47
RHCV%	0.08	-0.07 – 0.23	0.06	-0.08 – 0.18	5229.5	0.07	-0.10 – 0.22	6318.5	0.35	0.19 – 0.48
HCV%	0.09	-0.06 – 0.24	0.05	-0.09 – 0.18	5030.5	0.03	-0.14 – 0.19	6346.5	0.35	0.20 – 0.49
WMHV%	0.48	0.37 – 0.59	-0.11	-0.26 – 0.02	5358	0.09	-0.07 – 0.25	4904	0.04	-0.12 – 0.21
Cognitive Scores										
KCB Score	-0.27	-0.39 – -0.13	0.33	0.20 – 0.45	5642	0.15	-0.01 – 0.30	2363	-0.50	-0.61 – -0.36
Memory	-0.26	-0.39 – -0.12	0.10	-0.05 – 0.22	5512.5	0.12	-0.04 – 0.28	3512.5	-0.25	-0.40 – -0.09
Executive	-0.19	-0.33 – -0.04	0.45	0.32 – 0.56	5604.5	0.14	-0.02 – 0.30	2425	-0.48	-0.6 – -0.35

ρ = Spearman's rank correlation coefficient (rho); RC = Rank Biserial Correlation Coefficient.

Sex and residence were respectively coded as: Group 1 = Male, Group 2 = Female, and Group 1 = Rural, Group 2 = Urban. A negative RC thus indicates that on average, values for Group 2 were greater than those for Group 1.

537 **Table 4: Notable Associations of Volumetry with Cognition and VRFs**

Volumetry	Factor	Test	Test statistic	Effect size (RC)	95% CI
GMV%	Illiteracy	MWU ^a	1909	-0.23	-0.43 —0.002
WMHV%			3448	0.40	0.19 – 0.57
CSFV%			3416	0.39	0.18 –0.56
RHCv%			3040.5	0.23	0.01 – 0.44
CSFV			3339	0.36	0.14 – 0.54
WMHV			3458	0.40	0.20 – 0.58
RHCv			3027.5	0.23	0.004 – 0.43
LHCv%	DBP	Spearman rho	0.25		0.11– 0.38
RHCv%			0.24		0.10– 0.39
HCV%			0.25		0.11– 0.39
LHCv			0.20		0.06– 0.33
RHCv			0.20		0.06– 0.34
HCV			0.19		0.06– 0.33
LHCv%	SBP		0.16		0.03– 0.29
WMHV%	SBP	Pearson r, controlled for age	0.22		0.06– 0.38
WMHV			0.22		0.05– 0.38
WMHV%	PPR		0.29		0.10– 0.46
WMHV			0.30		0.11– 0.46
LHCv%	KCB test 8	Pearson r, controlled for age and education	0.17		0.04– 0.30
RHCv%			0.21		0.09– 0.34
HCV%			0.20		0.07– 0.33
LHCv			0.18		0.04– 0.32

RHCV			0.21		0.08– 0.35
HCV			0.20		0.06– 0.34

^a 95% CIs are for effect size (RC)

561 **Table 5: Regression analyses**

Dependent Variable	Predictors	B	95% CI _B		Beta	P value
			Lower	Upper		
Predictors of Brain Region Volumes						
GMV	HCV	9.64	4.31	14.97	0.24	<0.001
	Female sex	-58.46	-77.90	-39.01	-0.40	<0.001
	WMHV	-4.85	-7.08	-2.62	-0.29	<0.001
GMV%	Female sex	1.19	0.30	2.07	0.14	0.009
	HTN	1.58	0.50	2.66	0.15	0.004
	Age	-0.18	-0.23	-0.13	-0.41	<0.001
	WMHV%	-2.57	-3.92	-1.22	-0.21	<0.001
	HCV%	-8.03	-11.62	-4.44	-0.25	<0.001
	WMV%	-0.71	-0.83	-0.60	-0.71	<0.001
HCV	Education	0.15	0.07	0.23	0.28	<0.001
	GMV	0.007	0.003	0.01	0.31	<0.001
	Urban residence	-1.26	-1.83	-0.68	-0.34	<0.001
HCV%	Education	0.01	0.005	0.02	0.28	<0.001
	Urban residence	-0.10	-0.15	-0.05	-0.35	<0.001
WMHV	Age	0.13	0.05	0.21	0.28	0.002
	HTN	2.06	0.33	3.79	0.18	0.02
	SBP	0.05	0.005	0.09	0.21	0.03
	Tobacco	-3.17	-6.14	-0.20	-0.16	0.04
	GMV	-0.02	-0.03	-0.006	-0.26	0.002
WMHV%	Age	0.01	0.006	0.02	0.31	<0.001

	Hypertension	0.16	0.02	0.29	0.18	0.02
	SBP	0.004	0.001	0.007	0.22	0.02
	GMV%	-0.02	-0.04	-0.01	-0.31	<0.001
	Tobacco	-0.27	-0.51	-0.03	-0.17	0.03
Predictors of Cognitive Score						
KCB score: Memory (Tests 5, 7, 8)	BMI	0.41	0.06	0.75	0.17	0.02
	WMHV%	-5.04	-9.70	-0.39	-0.19	0.03
	10-year age increment	-3.07	-5.05	-1.09	-0.25	0.003
KCB score: Executive (Tests 1, 2, 4, 6)	HCV%	12.26	1.61	22.92	0.16	0.02
	Educational level	5.01	2.69	7.33	0.32	<0.001
	Urban residence	4.22	0.94	7.50	0.20	0.01
	WMHV%	-5.31	-9.78	-0.85	-0.19	0.02
Total KCB Score	Urban residence	11.53	3.99	19.07	0.24	0.003
	Educational level	7.07	1.74	12.41	0.20	0.01
	BMI	0.86	0.10	1.62	0.16	0.03
	10-year age increment	-5.90	-10.27	-1.53	-0.20	0.008
	WMHV%	-11.55	-21.81	-1.29	-0.18	0.03

562

563

564

565

566

567

568

569

Figures

Figure 1: Scatterplot Matrix of Grey Matter and White Matter Hyperintensity Volumes (Standardized), and Age

Diagonal cells show histogram distributions of their respective variables.

GMV% = Standardized Grey Matter Volume; WMHV% = Standardized White Matter Hyperintensity Volume

584 **Fig 2: Scatterplots of Hippocampal Volumes vs. Years of Education in Literate Participants (n = 170)**

585

586 Results of correlation analyses controlling for age: LHCV: $r = 0.17$, 95% CI = 0.007 –0.32; RHCV: $r = 0.22$, 95% CI =0.06 –0.36; HCV: $r =$
 587 0.19, 95% CI =0.04 –0.34; LHCV%: $r = 0.18$, 95% CI =0.05 –0.30; RHCV%: $r = 0.23$, 95% CI =0.10 –0.36; HCV%: $r = 0.21$, 95% CI = 0.08 –
 588 0.34.

Figure 3: Scatterplot Matrix of Standardized Hippocampal Volumes vs. Systolic and Diastolic Blood Pressures

Blood pressure readings (both systolic and diastolic values) were available for 164 participants.

Diagonal cells show histogram distributions of their respective variables.

LHCV% = Standardized Left Hippocampal Volume; RHCV% = Standardized Right Hippocampal Volume; SBP = Systolic BP; DBP = Diastolic BP.