1 Occupation as a risk factor for progression of chronic kidney disease: retrospective
2 cohort study
3

4 Daisuke Takada¹, Susumu Kunisawa¹, Akira Kikuno², Tomoko Iritani², and Yuichi
5 Imanaka, PhD*¹.
6

7 ¹Department of Healthcare Economics and Quality Management, Graduate School of
8 Medicine, Kyoto University, Yoshida Konoe-cho, Sakyo-ku, Kyoto City, Kyoto, Japan

9 ² Japan Health Insurance Association, Kyoto
10

11 *Corresponding author

12 EMAIL: imanaka-y@umin.net (YI)
13
14
15
16
17
18

19 **Abstract**

20 **Background**

21 Several social determinants of health are associated with the progression of chronic
22 kidney disease (CKD). We compared the risk of CKD progression among 18
23 occupational classifications using an annual health-checkup database.

24 **Methods**

25 We used the annual health checkup data and health insurance claims data of the Japan
26 Health Insurance Association in Kyoto prefecture between April 2012 and March 2016.
27 The primary outcome for survival analysis was defined as a more than 30% change in
28 the estimated glomerular filtration rate (eGFR) from the first health checkup. We used
29 the Cox proportional-hazards model for time-to-event analyses to estimate the hazard
30 ratio, and 95% CIs for the primary outcome, adjusting for age, sex, eGFR, body mass
31 index, blood pressure, blood sugar, dyslipidemia, uric acid, urinary protein, and
32 existence of kidney diseases at first health checkup.

33 **Results**

34 We analyzed 239,506 employees, and 1,736 (0.7%) individuals whose eGFR had
35 decreased by 30% or more; the mean follow-up period was 2.8 years (standard
36 deviation: 1.2 years). When we compared the risk for “manufacturing,” five categories

37 of industries (“information and communications”; “transport and postal services”;
38 “accommodations, eating and drinking services”; “living-related and personal services
39 and amusement service”; “medical, health care and welfare”) were associated with a
40 decline in the risk of eGFR after adjusting for the confounding factors and/or
41 mediators.

42 **Conclusions**

43 We provided evidence that the risk of CKD progression depends on occupation type.
44 Further research is needed to confirm the mechanism and causal relationships
45 involved.

46

47

48

49

50 **Keyterms:** survival analysis, occupational category, decline of estimated glomerular

51 filtration ratio, cohort study, annual health checkup data, health insurance claims data

52

53 **Introduction**

54 Chronic kidney disease (CKD) is recognized as one of the most serious problems in
55 public health. The global prevalence of CKD reached approximately 10% worldwide,
56 affecting 500 million people. [1] Currently, the estimated glomerular filtration ratio
57 (eGFR) is a good marker for kidney function, as a gradual reduction in eGFR leads to
58 kidney failure in the general population. [2] Traditional lifestyle diseases such as
59 diabetes and hypertension influence the development of CKD, increasing the risk of
60 cardiovascular and cerebrovascular events, and some cancers. [1, 3]

61 Recently, the etiology of non-traditional disorders has gained attention,
62 including novel risk factors for CKD. Systematic reviews have shown that
63 occupational status, including agricultural and livestock-related work, is associated
64 with an increased risk of CKD progression. [4-8] The causal role of occupation in renal
65 dysfunction might be accounted for by several mechanisms, including a greater
66 prevalence of hyperphosphatemia [9], work-related heat stress [10], and lower
67 health-related quality of life both mentally and physically. [11, 12] However, this field
68 of study is only just emerging, and more studies are needed to clarify such associations.
69 Here, we report the association between occupation and the progression of CKD
70 through an extensive health-checkup database in Japan.

71

72 **Methods**

73 **Database, medical records and target populations**

74 We conducted a retrospective database analysis using annual health checkup data
 75 between April 2012 and March 2016 and health insurance claims data, together with
 76 information on the subject's employer, from a database of the Japan Health Insurance
 77 Association in Kyoto prefecture, Japan[13]. Annual specific medical checkups for
 78 employees aged ≥ 40 years were established in the following law: "Act on Assurance
 79 of Medical Care for Elderly People". Accordingly, the Japan Health Insurance
 80 Association get employees aged ≥ 35 years to have annual health checkups until they
 81 lose their eligibility, such as their changing jobs, being fired, or passed away. We
 82 enrolled employees who were aged more than 35 years of age and had undergone two
 83 or more health checkups during the observation period. In our analysis, we excluded
 84 participants: 1) Whose eGFR was less than 30 mL/min/1.73m²; 2) who had several
 85 kidney diseases; and 3) those who were missing data from their first health checkup.
 86 Kidney diseases were defined using ICD-10 codes recorded in the claims data, such as
 87 glomerular disease, renal artery stenosis, and polycystic kidney disease (N00-08, I70,
 88 and Q61).

89

90 **Covariates**

91 Covariates include patient sex, age, body mass index (BMI), eGFR, urinary protein,
92 abdominal circumference, blood pressure (BP), dyslipidemia, diabetes, uric acid and
93 kidney disease. All the covariates were collected at the first health checkup, and eGFR
94 was calculated by the formula for Japanese patients devised by Matsuo et al. [14]

95 The patients were classified as follows: four groups based on age (35–44,
96 45–54, 55–64, and ≥65 years); four groups based on BMI (<18.5 kg/m², thin; ≥18.5
97 and <25 kg/m², normal range; ≥25 and <30 kg/m², pre-obesity; and ≥30 kg/m²,
98 obesity) according to the World Health Organization [15]; five groups based on eGFR
99 (>60, 60–45, 45–30 mL/min/1.73m²); three groups based on urinary protein (using
100 dipsticks ≥ 1+ (positive), ± (trace), and negative), abdominal circumference (if male ≥
101 85 cm, female ≥ 90 cm); five groups based on BP (systolic blood pressure (sbp) ≥180
102 mmHg or diastolic blood pressure (dbp) ≥110 mmHg without antihypertensives, sbp
103 ≥160 mmHg or dbp ≥100 mmHg without drugs, sbp ≥140 mmHg or dbp ≥90 mmHg
104 without drugs, normal without drugs, and with drugs); and four groups based on
105 dyslipidemia (triglyceride ≥150 mg/dL or high-density lipoprotein cholesterol <40
106 mg/dL defined as abnormality with hypolipidemic drugs, abnormality without

107 hypolipidemic drugs, normal with hypolipidemic drugs, normal without hypolipidemic
108 drugs), diabetes (fasting blood sugar ≥ 110 mg/dL or hemoglobin A1c $\geq 5.6\%$ defined as
109 abnormality with antidiabetic drugs, abnormality without antidiabetic drugs, normal
110 with antidiabetic drugs, normal without antidiabetic drugs), and hyperuricemia
111 (defined uric acid ≥ 8 mg/dl without anti-hyperuricemias or using anti-hyperuricemias).

112 Occupational information was acquired from notifications when companies
113 were established and submitted information to the Japan Pension Service, so that their
114 employees were covered by health insurance. The occupational information was
115 classified into 18 categories according to the Japan Standard Industrial Classification,
116 as modified from the International Standard Industrial Classifications of All Economic
117 Activities. The classifications are as follows: 1) agriculture and forestry and fishers, 2)
118 mining and quarrying of stone, 3) construction, 4) manufacturing, 5) electricity, gas,
119 heat supply and water production, transmission and distribution of electricity, 6)
120 information and communications, 7) transport and postal services, 8) wholesale and
121 retail trade, 9) finance and insurance, 10) real estate and goods rental and leasing, 11)
122 scientific research, professional and technical service, 12) accommodations, eating and
123 drinking services, 13) living-related and personal services and amusement service, 14)
124 education, learning support school education, 15) medical, health care and welfare, 16)

125 compound services, such as establishments engaged in administrative or ancillary
126 economic activities, 17) other services, such as waste disposal business, 18)
127 government, except where elsewhere classified as national government service. The
128 primary industry is classified as category “1”, the secondary industry is classified as
129 category “2” to “4”, and the tertiary industry is classified as category “5” to “18”.

130

131 **Statistical analysis**

132 The primary outcome for the survival analysis per person-year was defined as a
133 decrease of 30% or more in eGFR from the baseline eGFR at the time of their first
134 medical check-up. [2] The eGFR was calculated by the equation used by the Japanese
135 Society of Nephrology. [14] Patients were followed until there was a decline in their
136 eGFR of at least 30% (event), or censored due to death, a change in their insurance
137 contract, or the date of their last health check-up, whichever came first.

138 The Cox proportional-hazards model was used for time-to-event analyses to
139 estimate the hazard ratio; 95% CIs were calculated for the primary outcome. Follow-up
140 period data for patients were censored on the date of the last health checkup. The
141 analysis used two types of models: Model 1, adjusted for age, sex, and eGFR at
142 baseline (without medication factor); and Model 2, adjusted for age, sex, eGFR, body

143 mass index, blood pressure, blood sugar, dyslipidemia, uric acid, and urinary protein at
144 baseline (with medication factor).

145 Subgroup analysis was performed in Model 2, where the analysis population
146 was limited according to eGFR from G1 to G2 (eGFR >60 mL/min/1.73m²).

147

148 **Ethical considerations**

149 The research protocol was approved by the ethics committee of the Kyoto University
150 Graduate School and Faculty of Medicine (approval number: R1631) and by the ethics
151 committee of the Japan Health Insurance Association. The participants were informed
152 to use their data in epidemiological studies that dealt with personal information in a
153 manner that was not identifiable, for example, we cannot show less than 10 people in
154 the tables. Due to the law: “Act on the Protection of Personal Information”, the data
155 cannot be released to the public. The data will be shared on reasonable request to the
156 corresponding author.

157

158 **Results**

159 A total of 253,673 employees were enrolled and fulfilled the inclusion criteria; 14,218
160 (5.6%) were excluded due to missing data. We analyzed the remaining 237,987

161 employees (Fig 1). By the end of follow-up period, there were 1,736 (0.7%)
162 individuals whose eGFR had decreased by 30% or more; the mean follow-up period
163 was 2.8 years (standard deviation: 1.2 years).

164

165 **Fig 1. Flow chart of study participants' selection**

166

167 The baseline characteristics of the patients are shown in Table 1. As
168 manufacturing was the most popular category, we used manufacturing as the reference
169 category.

170

171 **Table.1 selected baseline characteristics according to the occupational**

172 **information categories**

173

174 Cox regression analysis was performed to identify the predictors of renal survival in
175 the two models (Table 2). After adjusting for all the confounding factors, five
176 categories of industries were associated with the risk of eGFR decrease. The hazard
177 ratios and 95% confidence estimated in Model 2 were 1.69 (1.18-2.42) for
178 “information and communications”; 1.37 (1.15-1.64) for “transport and postal

179 services”; 1.55 (1.18-2.04) for “accommodations, eating and drinking services”; 1.44
180 (1.06-1.97) for “Living-related and personal services and amusement service”; 1.56
181 (1.34-1.82) for “medical, health care and welfare”.

182

183 **Table.2 The incident rate per 1000 person-years and hazard ratios for a decrease**
184 **of 30% or more in estimated glomerular filtration ratio in crude and adjusted**
185 **models**

186

187 The other hazard ratios for a decrease of 30% or more in eGFR in Model 2 were shown
188 in Figure 2.

189

190 **Fig.2 The other hazard ratios for a decrease of 30% or more in estimated**
191 **glomerular filtration ratio, adjusted for sex, age, body mass index, estimated**
192 **glomerular filtration ratio, urinary protein, abdominal circumference, blood**
193 **pressure, dyslipidemia, diabetes, uric acid and kidney disease at first health**
194 **checkup**

195

196 The results for the subgroup analysis were similar to those for the main analysis.

197

198 **Discussion**

199 We showed that type of occupation substantially influences CKD progression. In
200 particular, when we compared risks to the risk of “manufacturing,” we found that five
201 categories of industries (“Information and communications”; “Transport and postal
202 services”; “Accommodations, eating and drinking services”; “Living-related and
203 personal services and amusement service”; “Medical, health care and welfare”) were
204 significantly associated with the risk of eGFR decline after adjusting for confounding
205 factors and/or other mediators such as medication factors. Then we can interpret that
206 these five categories include other risk factors than such medication factors.

207 Interestingly, we found that these categories share two working conditions:
208 nonstandard work schedule and sedentary time. In the case of nonstandard work
209 schedule, night shift work and rotating shift work may influence renal function by
210 disrupting the endogenous circadian system with several hormonal and metabolic
211 disturbances and negatively impacting physical and mental health. [16, 17] In fact,
212 nonstandard work schedule is likely a risk factor for several vascular diseases [18, 19],
213 and might result in hospitalization for cardiovascular disease, metabolic syndrome or
214 various kinds of cancer. [16] Similarly, shift works has been shown to increase the risk

215 of CKD onset in Japanese workers. [20] Thus, the four identified types of industry may
216 decrease an employee's renal function based on the employee's nonstandard work
217 schedule.

218 The other condition, sedentary time, is a feature of occupation categories such
219 as "transport and postal services" and "information and communications," both of
220 which require people to sit long hours each day. Recently, several researchers have
221 reported that prolonged sedentary time is associated with diseases such as diabetes and
222 cardiovascular events and leads to an increase in total mortality [21-24], although the
223 mechanism remains unknown. Likewise, prolonged sedentary time has also been
224 associated with higher mortality due to kidney diseases. [25] Remarkably, we could
225 change these conditions by changing the way to work, future researches need to
226 elucidate the causal relationships.

227 According to a previous meta-analysis, agriculture is also related to kidney
228 damage. [4] The authors of the study raised the possibility that the cause could be
229 agrochemicals or sugar cane. The present study showed that agriculture, including
230 forestry and fishers, significantly increased the risk of CKD progression in Model 1 but
231 not in Model 2, suggesting that lifestyle disease, rather than occupation, could have an
232 impact on the renal function in Japanese agricultural workers.

233 There were some limitations to our study. First, competing events, including
234 office relocation and employees' events, were regarded as censored in our analysis, and
235 some of these might result in a healthy-worker bias. For example, the decline in an
236 employee's eGFR might cause other complications that result in them result in their
237 quitting jobs, changing their jobs, or dying because of their poor health. Second, our
238 study did not consider several potential factors, including socio-economic status and
239 dietary and exercise habits. Lifestyle habits may be a mediator variable, so further
240 studies are needed to confirm the impact of such factors, even though it is difficult to
241 accurately measure them.

242

243 **Conclusion**

244 In summary, we found that certain types of occupations substantially influence
245 CKD progression. Further research in a clinical setting is needed to clarify the
246 mechanisms and causal relationships involved.

247

248 **Acknowledgements**

249 We thank all the staff members in Japan Health Insurance Association.

250

251 **Funding source**

252 The publication charges for this article were funded from a Grant-in-Aid for Scientific
253 Research from the Japan Society for the Promotion of Science (16H02634, 19H01075).
254 The funders had no role in study design, data collection and analysis, decision to
255 publish, or preparation of the manuscript.

256

257 **Reference**

- 258 1. Levin A, Tonelli M, Bonventre J, Coresh J, Donner JA, Fogo AB, et al. Global
259 kidney health 2017 and beyond: a roadmap for closing gaps in care, research, and policy.
260 Lancet. 2017;390(10105):1888-917. Epub 2017/04/25. doi: 10.1016/s0140-6736(17)30788-2.
261 PubMed PMID: 28434650.
- 262 2. Kanda E, Usui T, Kashihara N, Iseki C, Iseki K, Nangaku M. Importance of
263 glomerular filtration rate change as surrogate endpoint for the future incidence of
264 end-stage renal disease in general Japanese population: community-based cohort study.
265 Clin Exp Nephrol. 2018;22(2):318-27. Epub 2017/09/09. doi: 10.1007/s10157-017-1463-0.
266 PubMed PMID: 28884361; PubMed Central PMCID: PMC5838149.
- 267 3. Webster AC, Nagler EV, Morton RL, Masson P. Chronic Kidney Disease. Lancet.
268 2017;389(10075):1238-52. Epub 2016/11/27. doi: 10.1016/s0140-6736(16)32064-5. PubMed
269 PMID: 27887750.
- 270 4. Chapman E, Haby MM, Illanes E, Sanchez-Viamonte J, Elias V, Reveiz L. Risk factors
271 for chronic kidney disease of non-traditional causes: a systematic review. Revista
272 panamericana de salud publica = Pan American journal of public health. 2019;43:e35. Epub

2019/05/17. doi: 10.26633/rpsp.2019.35. PubMed PMID: 31093259; PubMed Central PMCID: PMCPMC6461065.

5. Lunyera J, Mohottige D, Von Isenburg M, Jeuland M, Patel UD, Stanifer JW. CKD of Uncertain Etiology: A Systematic Review. *Clinical journal of the American Society of Nephrology : CJASN*. 2016;11(3):379-85. Epub 2015/12/30. doi: 10.2215/cjn.07500715. PubMed PMID: 26712810; PubMed Central PMCID: PMCPMC4791820.

6. Plantinga LC. Socio-economic impact in CKD. *Nephrologie & therapeutique*. 2013;9(1):1-7. Epub 2013/01/16. doi: 10.1016/j.nephro.2012.07.361. PubMed PMID: 23318113.

7. Landrigan PJ, Goyer RA, Clarkson TW, Sandler DP, Smith JH, Thun MJ, et al. The work-relatedness of renal disease. *Archives of environmental health*. 1984;39(3):225-30. Epub 1984/05/01. doi: 10.1080/00039896.1984.9939529. PubMed PMID: 6380428.

8. Sponholtz TR, Sandler DP, Parks CG, Applebaum KM. Occupational exposures and chronic kidney disease: Possible associations with endotoxin and ultrafine particles. *American journal of industrial medicine*. 2016;59(1):1-11. Epub 2015/11/18. doi: 10.1002/ajim.22541. PubMed PMID: 26572099; PubMed Central PMCID: PMCPMC4715760.

9. Gutierrez OM, Anderson C, Isakova T, Scialla J, Negrea L, Anderson AH, et al. Low socioeconomic status associates with higher serum phosphate irrespective of race. *Journal of the American Society of Nephrology : JASN*. 2010;21(11):1953-60. Epub 2010/09/18. doi: 10.1681/asn.2010020221. PubMed PMID: 20847142; PubMed Central PMCID: PMCPMC3014009.

10. Kupferman J, Ramirez-Rubio O, Amador JJ, Lopez-Pilarte D, Wilker EH, Laws RL, et al. Acute Kidney Injury in Sugarcane Workers at Risk for Mesoamerican Nephropathy. *American journal of kidney diseases : the official journal of the National Kidney Foundation*. 2018;72(4):475-82. Epub 2018/07/26. doi: 10.1053/j.ajkd.2018.04.014. PubMed

- 298 PMID: 30042041.
- 299 11. Kusek JW, Greene P, Wang SR, Beck G, West D, Jamerson K, et al. Cross-sectional
300 study of health-related quality of life in African Americans with chronic renal
301 insufficiency: the African American Study of Kidney Disease and Hypertension Trial.
302 American journal of kidney diseases : the official journal of the National Kidney
303 Foundation. 2002;39(3):513-24. Epub 2002/03/06. doi: 10.1053/ajkd.2002.31401. PubMed
304 PMID: 11877570.
- 305 12. Correa-Rotter R, Wesseling C, Johnson RJ. CKD of unknown origin in Central
306 America: the case for a Mesoamerican nephropathy. American journal of kidney diseases :
307 the official journal of the National Kidney Foundation. 2014;63(3):506-20. Epub
308 2014/01/15. doi: 10.1053/j.ajkd.2013.10.062. PubMed PMID: 24412050.
- 309 13. Takada D, Kunisawa S, Kikuno A, Iritani T, Imanaka Y. Stages of a transtheoretical
310 model as predictors for the decline of estimated glomerular filtration rate: a
311 retrospective cohort study. Journal of epidemiology. 2021. Epub 2021/01/26. doi:
312 10.2188/jea.JE20200422. PubMed PMID: 33487611.
- 313 14. Matsuo S, Imai E, Horio M, Yasuda Y, Tomita K, Nitta K, et al. Revised equations
314 for estimated GFR from serum creatinine in Japan. American journal of kidney diseases :
315 the official journal of the National Kidney Foundation. 2009;53(6):982-92. Epub
316 2009/04/03. doi: 10.1053/j.ajkd.2008.12.034. PubMed PMID: 19339088.
- 317 15. [cited 2019 0309]. Available from:
318 <http://www.assessmentpsychology.com/icbmi.htm>.
- 319 16. Boivin DB, Boudreau P. Impacts of shift work on sleep and circadian rhythms.
320 Pathologie-biologie. 2014;62(5):292-301. Epub 2014/09/24. doi:
321 10.1016/j.patbio.2014.08.001. PubMed PMID: 25246026.
- 322 17. Costa G. The impact of shift and night work on health. Applied ergonomics.

1996;27(1):9-16. Epub 1996/02/01. doi: 10.1016/0003-6870(95)00047-x. PubMed PMID: 15676307.

18. Knutsson A. Health disorders of shift workers. Occupational medicine (Oxford, England). 2003;53(2):103-8. Epub 2003/03/15. doi: 10.1093/occmed/kqg048. PubMed PMID: 12637594.

19. Biggi N, Consonni D, Galluzzo V, Sogliani M, Costa G. Metabolic syndrome in permanent night workers. Chronobiology international. 2008;25(2):443-54. Epub 2008/05/20. doi: 10.1080/07420520802114193. PubMed PMID: 18484373.

20. Sasaki S, Yoshioka E, Saijo Y, Bannai A, Kita T, Tamakoshi A, et al. A prospective cohort study of insomnia and chronic kidney disease in Japanese workers. Sleep & breathing = Schlaf & Atmung. 2018;22(1):257-65. Epub 2017/05/24. doi: 10.1007/s11325-017-1508-3. PubMed PMID: 28534280.

21. Wilmut EG, Edwardson CL, Achana FA, Davies MJ, Gorely T, Gray LJ, et al. Sedentary time in adults and the association with diabetes, cardiovascular disease and death: systematic review and meta-analysis. Diabetologia. 2012;55(11):2895-905. Epub 2012/08/15. doi: 10.1007/s00125-012-2677-z. PubMed PMID: 22890825.

22. Ford ES, Caspersen CJ. Sedentary behaviour and cardiovascular disease: a review of prospective studies. International journal of epidemiology. 2012;41(5):1338-53. Epub 2012/05/29. doi: 10.1093/ije/dys078. PubMed PMID: 22634869; PubMed Central PMCID: PMC4582407.

23. Schmid D, Leitzmann MF. Television viewing and time spent sedentary in relation to cancer risk: a meta-analysis. Journal of the National Cancer Institute. 2014;106(7). Epub 2014/06/18. doi: 10.1093/jnci/dju098. PubMed PMID: 24935969.

24. Ekelund U, Steene-Johannessen J, Brown WJ, Fagerland MW, Owen N, Powell KE, et al. Does physical activity attenuate, or even eliminate, the detrimental association of

348 sitting time with mortality? A harmonised meta-analysis of data from more than 1 million
 349 men and women. Lancet. 2016;388(10051):1302-10. Epub 2016/08/01. doi:
 350 10.1016/s0140-6736(16)30370-1. PubMed PMID: 27475271.

351 25. Patel AV, Maliniak ML, Rees-Punia E, Matthews CE, Gapstur SM. Prolonged Leisure
 352 Time Spent Sitting in Relation to Cause-Specific Mortality in a Large US Cohort. American
 353 journal of epidemiology. 2018;187(10):2151-8. Epub 2018/06/28. doi: 10.1093/aje/kwy125.
 354 PubMed PMID: 29947736; PubMed Central PMCID: PMC6454412.

355

356

Table.1 selected baseline characteristics according to the occupational information categories

	Total number	Sex :female (%)	Age :mean(s d)	Serum creatinine :mean(s d) mg/dl	Estimated glomerular filtration rate:mean (sd) mL/min/1.73m ²	Urinary protein(%) trace positive		Body mass index :mean(s d)	Systolic blood pressure :mean(s d) mmHg	Diastolic blood pressure :mean (sd) mmHg	Abdominal circumference (%)(male ≥85, female ≥90)	Diabetes ^a (%) high, no drug normal or high with drug		Dyslipidemia ^b (%) high, no drug normal or high with drug		Hyperuricemia ^c (%)
Agriculture and forestry and fishers	697	130 (18.7%)	50 (9.9)	0.77 (0.15)	83 (15)	64 (9.2%)	50 (7.2%)	23.1 (3.3)	124 (19)	76 (13)	210 (30.1%)	69 (9.9%)	25 (3.6%)	152 (21.8%)	41 (5.9%)	29 (4.1%)
Mining and quarrying of stone	293	46 (15.7%)	49 (9.0)	0.81 (0.15)	79 (14)	29 (9.9)	21 (7.1)	24.0 (3.6)	127 (19)	79 (12)	129 (44.0)	38 (12.9)	13 (4.4)	81 (27.6)	11 (3.8)	13 (4.1)
Construction	14,191	2,064 (14.5%)	48 (9.7)	0.80 (0.23)	81 (14)	1,971 (13.8)	1,056 (7.4)	23.6 (3.5)	124 (18)	77 (13)	5,696 (40.1)	1,412 (9.9)	495 (3.5)	3,927 (27.7)	481 (3.4)	936 (6.6)
Manufacturing	65,421	16,496 (25.2%)	47 (9.3)	0.78 (0.21)	81 (14)	9,362 (14.2)	4,316 (6.6)	22.9 (3.6)	122 (17)	76 (12)	18,610 (28.4)	5,380 (8.2)	1,693 (2.6)	12,379 (18.9)	3,619 (5.5)	2,788 (4.2)

Electricity, gas, heat supply and water production, transmission and distribution of electricity	1,290	220 (17.1%)	48 (9.8)	0.82 (0.48)	80 (14)	197 (15.1)	99 (7.6)	23.3 (3.5)	123 (17)	77 (13)	464 (36.0)	114 (8.8)	50 (3.9)	291 (22.6)	82 (6.3)	56 (4.3)
Information and communications	3,967	852 (21.5%)	43 (7.8)	0.81 (0.19)	82 (14)	473 (11.9)	359 (9.0)	23.2 (3.8)	118 (16)	73 (12)	1,334 (33.6)	275 (6.9)	89 (2.3)	891 (22.5)	162 (4.0)	241 (6.0)
Transport and postal services	20,985	2,115 (10.1%)	52 (10.1)	0.82 (0.29)	79 (15)	2,912 (13.8)	2,344 (11.1)	23.6 (3.7)	127 (18)	79 (13)	8,973 (42.8)	3,046 (14.5)	1,143 (5.4)	5,980 (28.5)	1,526 (7.2)	1,138 (5.4)
Wholesale and retail trade	41,725	15,058 (36.1%)	48 (9.4)	0.75 (0.19)	82 (14)	4,809 (11.5)	3,362 (8.0)	22.8 (3.7)	121 (18)	75 (13)	11,178 (26.8)	3,420 (8.1)	1,157 (2.8)	7,507 (18.0)	2,284 (5.5)	1,638 (3.9)
Finance and insurance	1,851	653 (35.3%)	46 (8.5)	0.77 (0.36)	81 (15)	165 (8.9)	170 (9.1)	23.1 (3.6)	118 (17)	74 (13)	609 (32.9)	148 (7.9)	39 (2.1)	401 (21.7)	106 (5.7)	101 (5.4)
Real estate and goods rental and leasing	5,751	1,852 (32.2%)	48 (10.3)	0.77 (0.24)	81 (15)	709 (12.3)	488 (8.4)	22.9 (3.6)	120 (17)	74 (13)	1,776 (30.9)	511 (8.8)	161 (2.8)	1,177 (20.5)	409 (7.1)	307 (5.3)
Scientific research, professional and technical service	7,267	2,242 (30.9%)	47 (9.5)	0.77 (0.26)	80 (14)	900 (12.3)	522 (7.1)	22.9 (3.5)	118 (17)	73 (12)	2,172 (29.9)	588 (8.0)	180 (2.4)	1,443 (19.9)	414 (5.7)	309 (4.2)
Accommodations, eating and drinking services	7,077	2,253 (31.8%)	48 (9.8)	0.74 (0.22)	85 (16)	851 (11.9)	571 (8.0)	22.6 (3.7)	121 (18)	74 (13)	1,922 (27.2)	668 (9.4)	173 (2.4)	1,197 (16.9)	363 (5.1)	315 (4.4)
Living-related and personal services	5,540	2,238 (40.4%)	46 (9.7)	0.75 (0.17)	83 (15)	637 (11.4)	486 (8.7)	22.5 (3.5)	119 (18)	73 (13)	1,338 (24.2)	429 (7.7)	133 (2.3)	1,004 (18.1)	270 (4.9)	199 (3.6)

and amusement service																
Education, learning support school education	3,261	1,559 (47.8%)	47 (9.3)	0.76 (0.30)	80 (15)	418 (12.7)	235 (7.2)	22.5 (3.6)	116 (17)	72 (13)	725 (22.2)	205 (6.2)	93 (2.9)	536 (16.4)	161 (4.9)	110 (3.3)
Medical, health care and welfare	33,789	23,711 (70.2%)	47 (9.0)	0.69 (0.18)	82 (15)	3,125 (9.2)	2,844 (8.4)	22.4 (3.7)	117 (16)	72 (12)	4,111 (12.2)	2,681 (7.9)	693 (2.1)	4,158 (12.3)	2,025 (6.0)	641 (1.9)
Compound services, such as establishments engaged in administrative or ancillary economic activities	774	283 (36.6%)	49 (9.7)	0.76 (0.16)	80 (14)	94 (12.0)	51 (6.5)	22.9 (3.5)	121 (18)	75 (13)	211 (27.3)	74 (9.5)	27 (3.5)	147 (19.0)	45 (5.7)	30 (3.9)
Other services, such as waste disposal business	20,637	6,581 (31.9%)	50 (10.3)	0.77 (0.32)	81 (15)	2,328 (11.2)	1,727 (8.3)	23.0 (3.7)	121 (18)	75 (13)	6,036 (29.2)	2,111 (10.2)	794 (3.8)	3,973 (19.3)	1,446 (6.9)	874 (4.2)
Government, other national government service	3,471	2,283 (65.8%)	48 (9.6)	0.73 (0.23)	76 (14)	499 (14.2)	318 (9.1)	22.4 (3.5)	119 (17)	73 (12)	582 (16.8)	377 (10.7)	129 (3.7)	443 (12.8)	379 (10.8)	68 (1.9)

^a diabetes “high”: fasting blood sugar ≥ 110 mg/dL or hemoglobin A1c $\geq 5.6\%$.
^b dyslipidemia “high”: triglyceride ≥ 150 mg/dL or high-density lipoprotein cholesterol < 40 mg/dL.
^c hyperuricemia: uric acid ≥ 8 mg/dL

1 Table 2 The incident rate per 1000 person-years and hazard ratios for a decrease of 30% or more in estimated glomerular filtration ratio in crude and adjusted
2 models

Sector of industry	Japan Standard Industrial Classification	Number	Incident rate per 1000 person-years (95% CI)	Model 1 ^a	Model 2 ^b
Primary industry	1) Agriculture and forestry and fishers	697	4.1 (1.9–7.5)	1.93 (0.96–3.88)	1.90 (0.94–3.83)
Secondary industry	2) Mining and quarrying of stone	293	2.1 (0.4–6.6)	1.03 (0.26–4.14)	0.88 (0.22–3.53)
	3) Construction	14,191	2.5 (2.1–3.1)	1.29* (1.04–1.60)	1.23 (0.99–1.52)
	4) Manufacturing	65,421	2.0 (1.8–2.2)	1.0 (ref)	1.0 (ref)
Tertiary industry	5) Electricity, gas, heat supply and water production, transmission and distribution of electricity	1,290	2.1 (1.5–5.0)	1.51 (0.83–2.75)	1.51 (0.83–2.75)
	6) Information and communications	3,967	3.0 (2.0–4.1)	1.74* (1.21–2.50)	1.69* (1.18–2.42)
	7) Transport and postal services	20,985	3.4 (3.0–3.9)	1.60* (1.34–1.90)	1.37* (1.15–1.64)
	8) Wholesale and retail trade	41,725	2.3 (2.0–2.6)	1.12 (0.96–1.31)	1.10 (0.94–1.28)
	9) Finance and insurance	1,851	2.0 (0.9–3.8)	1.09 (0.54–2.19)	1.06 (0.53–2.14)
	10) Real estate and goods rental and leasing	5,751	2.6 (1.9–3.2)	1.25 (0.91–1.73)	1.25 (0.91–1.73)
	11) Scientific research, professional and technical service	7,267	2.5 (1.9–3.2)	1.24 (0.92–1.66)	1.25 (0.93–1.67)
	12) Accommodations, eating and drinking services	7,077	3.3 (2.5–4.2)	1.59* (1.21–2.09)	1.55* (1.18–2.04)
	13) Living-related and personal services and amusement service	5,540	3.0 (2.2–4.0)	1.48* (1.08–2.01)	1.44* (1.06–1.97)
	14) Education, learning support school education	3,261	1.6 (0.7–2.6)	0.78 (0.47–1.31)	0.77 (0.46–1.29)
	15) Medical, health care and welfare	33,789	3.6 (3.2–4.0)	1.59* (1.37–1.86)	1.56* (1.34–1.82)
	16) Compound services, such as establishments engaged in administrative or ancillary economic activities	774	2.7 (1.1–5.5)	1.28 (0.57–2.86)	1.25 (0.56–2.81)
	17) Other services, such as waste disposal business	20,637	2.7 (2.3–3.1)	1.21* (1.0–1.47)	1.18 (0.98–1.43)
	18) Government, other national government service	3,471	2.5 (1.6–3.7)	0.95 (0.63–1.53)	0.97 (0.62–1.50)

Hazard ratio (95% confidence interval)* in bold showed p-value <0.05

Abbreviation eGFR, estimated glomerular filtration rate; CI, confidence interval

^a Adjusted for sex, age and estimated glomerular filtration ratio at first health checkup

^b Adjusted for sex, age, body mass index, estimated glomerular filtration ratio, urinary protein, abdominal circumference, blood pressure, dyslipidemia, diabetes, uric acid and kidney disease at first health checkup

369

370 Figure legends

371

372 Figure 1: Flow chart of patient selection

373 N: number of patients

374

375 Figure 2: The other hazard ratios for a decrease of 30% or more in estimated

376 glomerular filtration ratio, adjusted for sex, age, body mass index, estimated

377 glomerular filtration ratio, urinary protein, abdominal circumference, blood pressure,

378 dyslipidemia, diabetes, uric acid and kidney disease at first health checkup

379

380

Employees who aged 35 years or over and
had two or more annual health checkups
between April 2012 and March 2016

N=253,673

Exclusion criteria

13,620: estimated glomerular filtration
ratio was less than 30 mL/min/1.73m²

1,346: had several kidney diseases

Missing data (duplicated)

Sex and Age: 833

Body mass index: 12

Blood pressure: 9,518

Serum creatinine: 972

Urinary protein: 1,411

Abdominal circumference: 437

Triglyceride or high-density lipoprotein: 163

Blood sugar or hemoglobin A1c: 968

Serum uric acid: 121

Occupations: 1,738

N=237,987

Sex: male

Age: 35-45 years (ref)

Age: 46-55 years

Age: 56-65 years

Age: ≥66 years

Body mass index: 18.5≥ kg/m²

Body mass index: 18.5-25 kg/m² (ref)

Body mass index: 25-30 kg/m²

Body mass index: >30 kg/m²

Estimated glomerular filtration rate: ≥60 mL/min/1.73m² (ref)

Estimated glomerular filtration rate: 45-60 mL/min/1.73m²

Estimated glomerular filtration rate: 30-45 mL/min/1.73m²

Blood pressure: less than 140/90 mmHg (ref)

Blood pressure: 140-159/90-99 mmHg

Blood pressure: 160-179/100-109 mmHg

Blood pressure: 180/110 or more mmHg

Blood pressure: Prescribed anti-hypertention drug

Dyslipidemia: normal (ref)

Dyslipidemia: high triglyceride/low high-density lipoprotein without drug

Dyslipidemia: normal with drug

Dyslipidemia: high triglyceride/low high-density lipoprotein with drug

Diabetes: normal (ref)

Diabetes: high HbA_{1c}/blood sugar without drug

Diabetes: normal with drug

Diabetes: high HbA_{1c}/blood sugar with drug

Abdominal circumference (male ≥85 cm, female ≥90 cm)

Uric acid abnormal or prescribed antihyperuricemic drug

Urinary protein: negative (ref)

Urinary protein: trace

Urinary protein: positive