Has the age distribution of hospitalized Covid-19 patients changed in Brazil?

**Carla Lourenço Tavares de Andrade, Sheyla Maria Lemos Lima, Monica Martins,
Claudia Cristina de Aguiar Pereira, Margareth Crisóstomo Portela**

**Department of Health Administration and Planning, Sergio Arouca National School of
Public Health, Oswaldo Cruz Foundation**

**Corresponding author:
Margareth Crisóstomo Portela
mportela@ensp.fiocruz.br**

Abstract

The aim of this study was to compare the age profile of hospitalized Covid-19 patients during the first year of the pandemic, as well as hospital mortality and use of ICUs, by age group, in large geographic regions of Brazil. We used data from the Influenza Epidemiological Surveillance Information System for patients who presented the first symptoms of the disease between the epidemiological weeks 8 of 2020 and 7 of 2021, which were divided into three periods.

779,257 records of patients hospitalized by Covid-19 were obtained. Of this total, 720,363 (92.4%) referred to discharged hospitalizations, considered in the analysis of ICU use and death. Among 244,611 hospitalizations (34.0%) with indication for use of ICU, 190,833 allowed the calculation of the time in ICU. There was variation in the age profile of hospitalized patients between the three periods, but there was no evidence in favor of the hypothesis of an increase, in the last period, in the participation of adults between 18 and 50 years old in hospitalizations by Covid-19. A differentiated increase in the mortality of young adults in the North suggests the possibility of greater severity of the P1 variant in this population. The results also show that the participation of young adults in hospitalizations and hospital deaths was never negligible and is related to hospital mortality rates close to or above 10%.

The Covid-19 “youthening” phenomenon in Brazil is based on the country's own sociodemographic and economic characteristics and may have been strengthened by the increasing circulation of viral variants. It is important to continue monitoring its progression and effects.

Keywords: Covid-19; hospitalizations; healthcare utilization; hospital mortality

Introduction

Brazil ended the first year of the Covid-19 pandemic in a scenario of increasing number of cases, with almost simultaneous growth in the rates of incidence and mortality due to the disease and high levels of occupancy of intensive care beds (ICU) in the 26 states and the Federal District, often greater than 90%¹. During the first year, regions of the country were most affected at different times, with maintenance of the daily moving average at more than a thousand deaths between June and August 2020. There was, then, a drop in the average of deaths until the beginning of November, but, in January 2021, the moving average again exceeded one thousand deaths, reaching, in March, more than two thousand daily deaths, a figure never seen before². Currently, Brazil is the second country with the highest absolute number of cases (11,998,233) and deaths (294,042)³, and accounts for more than 1/4 of the registered deaths globally, with only 2.7% of the world population⁴.

After a year of pandemic, the most critical moment of the health and care crisis is experienced^{1,5}. The low adherence of the population to the pandemic control and prevention measures, aggravated during the last months of 2020, and the appearance of new variants of the SARS-CoV-2 virus, which are more transmissible than the original version⁶, are plausible explanations for the high growth of cases and deaths.

In this context, there is frequent conjecture about a possible change in the age profile of patients hospitalized by Covid-19 in the country, with younger people presenting more severe conditions of the disease, higher mortality and more prolonged use of ICU beds⁷. The aim of this study was to compare, during the first year of the pandemic, the age profile of patients hospitalized by Covid-19, as well as hospital mortality and use of ICUs, by age group, in large geographic regions of Brazil.

Methods

This study used data from the Influenza Epidemiological Surveillance Information System (SIVEP-Gripe) from DATASUS, Ministry of Health, which is public and unrestricted. The system is nationwide and includes compulsory notifications about severe acute respiratory syndrome, including demographic and clinical variables, related to the use of outpatient and hospital services and the outcome of the case. Hospital data include the use and dates of entry and exit from the ICU. The data were accessed on the website <https://opendatasus.saude.gov.br> on March 17, 2021. Of the records from 2020 and 2021, those with a final classification of 'SRAG by COVID-19' that involved hospitalization were considered.

For each region of the country, the age profiles of hospitalized patients were compared, and in each age group, the occurrence of hospital mortality and ICU use, as well as the time (days) in the ICU, considering three periods during the first year of the pandemic. Cases were included for

those with first symptoms occurring between the eighth epidemiological week of 2020 and the seventh epidemiological week of 2021. The first period included 18 epidemiological weeks, between 16 February and 20 June 2020; the second included 18 epidemiological weeks, between 21 June and 24 October 2020; and the third included 17 epidemiological weeks, between October 25, 2020 and February 20, 2021. The time frame until the seventh epidemiological week of 2021 allowed us to examine the first year of the pandemic, mitigating the problem of dealing with the underreporting of cases in other available weeks and excessive loss of unfinished records.

For comparisons of the age profiles of hospitalized patients, between periods, all records were counted in the calculation of absolute and relative frequencies of hospitalizations, using the chi-square test. The comparisons of ICU use and the occurrence of death by age group considered only the records with discharge information. For comparing the ICU length of stay, the closed records that informed the date of admission and discharge from ICU. In the absence of the date of exit and occurrence of death, it was assumed that the date of exit was the date of death. In all three cases, the Kruskal-Wallis non-parametric test was applied.

Results

779,257 records of patients hospitalized by Covid-19 were obtained. Of this total, 720,363 (92.4%) had data (type and / or date) of the patient's evolution or had been closed, making up the set of observations used in the analysis of ICU use and death. Among these, 244,611 (34.0%) contained an indication of ICU use, and 190,833 had data that allowed the calculation of ICU length of stay.

Table 1 shows that there was variation in the age profile of hospitalized patients between the three periods considered in the regions of the country, however it does not provide evidence in favor of the hypothesis of an increase, in the last period, of hospitalizations of people between 18 and 50 years, compared to previous periods of the pandemic. In the North and Northeast regions, the proportion of individuals aged between 18 and 50 years old, in the last period, approached those observed in the first period, when they were also substantially affected by the pandemic. In the Southeast and South, there was a consistent reduction in these age groups over the periods. In the Central West region, which was mostly affected only after the second period, the participation of adults between 18 and 50 years old in hospitalizations was higher in the first period, decreasing in the second and remaining relatively close to the second in the third.

Table 2 compares the occurrence of hospital deaths and ICU use by region and age group between the three defined periods. The North region, which experienced collapse in the provision of healthcare in the first and last periods, emerged in the last, with high percentages of mortality in all age groups, compatible with those observed in the first, in the groups above 60 years, but even greater than those among adults up to 60 years old. It also showed greater use of ICU among

patients aged 18-60 and > 80 years. In the Northeast, except for individuals aged 18-30 years, whose mortality remained at a relatively stable level (close to 14%), hospital mortality decreased among adults in the periods, while the percentages of ICU use increased significantly. In the Southeast, there was a decrease in hospital mortality throughout the year for patients aged 60 years and over, and an increase, after falling in the second period, among younger adults, with the highest mortality among those aged 18-30 years. The percentages of ICU use did not vary significantly in most age groups, with an increase in use among patients aged 31-40 years and a reduction between patients aged 71-80 and over 80 years being significant. In the South, the percentages of ICU use dropped significantly, over the periods, among adults over 40 years old, maintaining statistical stability among adults aged 18-40 years. The highest percentage of mortality in the third period among adults aged 18-70 years is highlighted reflecting the severity of the pandemic in the region. In Central West, there were statistically non-significant variations in hospital mortality for adults between 18 and 50 years old, and a drop among adults over 50 years old, with the pattern of ICU use being higher in the third period than in the second, but lower than that in the first period.

Table 3 should be viewed with caution because of the likelihood of underestimating the length of stay in the ICU, especially for patients who fell ill in the third period. Nevertheless, it presents results that suggest that the length of stay in the ICU by Covid-19 has not been prolonged in the last period.

Final considerations

Considering the analyses undertaken, based on data referring to February 20, 2021, this study does not provide evidence to support a greater participation of young adults in hospitalizations in the third period. However, the differentiated rise in Covid-19 hospital deaths among adults aged 18-50 and even 50-60 years in the North, compared especially to the first period, also marked by a collapse in the health system in some states, is a warning which meets the hypothesis of greater severity of the disease caused by the P1 variant in this population. In any case, the results showed that the participation of young adults in hospitalizations for Covid-19 was never negligible and was always related to hospital mortality rates close to or above 10%. In this study, the extension of the ICU use time was not observed.

The risk of an increase in severe cases among adults up to 50 years of age is open to multifactorial explanations and confronts a certain common sense that the disease would offer a low risk for younger people. The Covid-19 “youthening” phenomenon in Brazil is based on the country's own socio-demographic and economic characteristics and may have been strengthened by the increasing circulation of viral variants. It is important to continue monitoring its progression and effects. Analyses from SIVEP itself and from the Civil Registry are beginning to point out the

rapid growth of Covid-19 cases and deaths in individuals between 20 and 59 years old since the beginning of the year^{8,9}. The focus on hospitalizations is more affected by data delays. The results, however, provide a baseline. Explorations at less aggregate levels, in capitals and large cities, which suffer markedly from the effects of large agglomerations, may be able to add more specific and circumscribed findings about the rejuvenation of the pandemic reported by health and media professionals^{8,9}.

References

1. Fiocruz. Observatório Covid-19. Boletim Extraordinário, 02 de março de 2021. Available at <https://portal.fiocruz.br/documento/boletim-extraordinario-covid-19> [accessed on 26 March 2021].
2. Mortes e casos de coronavírus nos estados. Portal G1. Available at <https://especiais.g1.globo.com/bemestar/coronavirus/estados-brasil-mortes-casos-media-movel/> [accessed on 03 March 2021].
3. Johns Hopkins University & Medicine. Coronavirus Resource Center. Disponível em <https://coronavirus.jhu.edu/map.htm> [acessado em 22/03/2021].
4. WHO. [WHO Coronavirus \(COVID-19\) Dashboard](https://covid19.who.int). Available at <https://covid19.who.int> [accessed on 22 March 2021].
5. Conselho Nacional de Secretários de Saúde (CONASS). Carta dos Secretários Estaduais de Saúde à Nação Brasileira. Publicada em 1 de março de 2021. Available at <https://www.conass.org.br/carta-dos-secretarios-estaduais-de-saude-a-nacao-brasileira/> [accessed em 22 March 2021].
6. Faria NR, Mellan TA, Whittaker C, Claro IM, Candido DS, Mishra S et al. Genomics and epidemiology of a novel SARS-CoV-2 lineage in Manaus, Brazil. Available at jornalggn.com.br-final-p1-manuscript-25-02-2021-combined.pdf [accessed on 22 March 2021].
7. Ferrari M. Especialistas apontam diminuição na idade dos internados por Covid-19. CNN, São Paulo, 06/03/2021. Available at <https://www.cnnbrasil.com.br/saude/2021/03/06/especialistas-apontam-diminuicao-na-idade-dos-internados-por-covid-19> [accessed on 06 March 2021].
8. Freire VT. Proporção de jovens mortos por Covid-19 cresce em SP. Folha de São Paulo, 20/03/2020. Available at <https://www1.folha.uol.com.br/equilibrioesaude/2021/03/proporcao-de-jovens-mortos-por-covid-19-cresce-em-sp.shtml> [accessed on 22 March 2021].
9. Fiocruz. Observatório Covid-19. Boletim das semanas epidemiológicas 10 e 11, de 7 a 20 de março de 2021. Available at https://agencia.fiocruz.br/sites/agencia.fiocruz.br/files/u34/boletim_covid_2021-semanas_10-11-red.pdf [accessed on 26 March 2021].

Table 1. Comparison of the distribution of Covid-19 hospitalizations (N = 779,257) by age groups among three periods of the pandemic's first year in the regions of Brazil

Region/ Age group (years)	Epidemiologic weeks						χ^2 (p)
	08/2020 – 25/2020		26/2020 – 43/2020		44/2020 – 07/2021		
	(16 Feb 2020 – 20 Jun 2020)		(21 Jun 2020 – 24 Oct 2020)		(25 Oct 2020 – 20 Feb 2021)		
	N	%	N	%	N	%	
North							<0.0001
0-5	351	1.5	793	4.8	744	3.0	
6-17	341	1.4	499	3.1	275	1.1	
18-30	1374	5.8	1310	8.0	1245	5.0	
31-40	2646	11.2	1765	10.8	2749	11.1	
41-50	3476	14.8	2178	13.3	3735	15.0	
51-60	3990	17.0	2614	16.0	4491	18.1	
61-70	4751	20.2	2968	18.1	5262	21.2	
71-80	3990	17.0	2594	15.8	3916	15.8	
>80	2622	11.1	1662	10.1	2416	9.7	
Total	23541	100.0	16383	100.0	24833	100.0	
Northeast							<0.0001
0-5	804	1.4	980	2.3	782	1.9	
6-17	530	0.9	713	1.7	378	0.9	
18-30	2806	4.9	1981	4.6	1683	4.0	
31-40	5539	9.7	3659	8.6	3993	9.6	
41-50	7527	13.2	4887	11.4	5434	13.1	
51-60	9744	17.1	6711	15.7	7342	17.6	
61-70	11238	19.8	8206	19.2	8468	20.3	
71-80	10646	18.7	8437	19.8	7609	18.3	
>80	8049	14.2	7128	16.7	5935	14.3	
Total	56883	100.0	42702	100.0	41624	100.0	
Southeast							<0.0001
0-5	948	0.9	1099	0.9	1185	0.8	
6-17	573	0.5	881	0.7	653	0.4	
18-30	4541	4.1	4648	3.7	4611	3.1	
31-40	11962	10.8	11294	9.0	12509	8.5	
41-50	17410	15.8	16986	13.5	19636	13.3	
51-60	21627	19.6	23316	18.5	27908	18.9	
61-70	22341	20.2	27419	21.8	33711	22.8	
71-80	17400	15.8	22707	18.0	27429	18.5	
>80	13559	12.3	17551	13.9	20326	13.7	
Total	110361	100.0	125901	100.0	147968	100.0	
South							<0.0001
0-5	64	0.7	301	0.7	317	0.5	
6-17	41	0.4	232	0.5	225	0.3	
18-30	570	5.8	1725	4.0	2160	3.3	
31-40	1166	11.8	4083	9.4	5678	8.7	
41-50	1722	17.5	6127	14.0	9075	13.8	
51-60	2048	20.8	8648	19.8	12952	19.8	
61-70	1937	19.7	9740	22.3	15279	23.3	
71-80	1408	14.3	7528	17.2	12096	18.5	
>80	886	9.0	5290	12.1	7719	11.8	
Total	9842	100.0	43674	100.0	65501	100.0	
Central West							<0.0001
0-5	65	0.7	368	1.0	146	0.6	
6-17	49	0.5	232	0.6	119	0.5	
18-30	506	5.5	1562	4.2	1032	4.3	
31-40	1272	13.9	3762	10.2	2528	10.6	
41-50	1684	18.4	5580	15.1	3727	15.6	
51-60	1883	20.5	7119	19.2	5014	21.0	
61-70	1656	18.1	7842	21.2	5020	21.0	
71-80	1245	13.6	6322	17.1	3808	15.9	
>80	808	8.8	4214	11.4	2481	10.4	
Total	9168	100.0	37001	100.0	23875	100.0	

Source: <https://opendatasus.saude.gov.br>. The periods were defined based on the epidemiologic week of onset of the first symptoms.

Table 2. Covid-19 hospitalization discharges (N=720,363), and comparison of hospital mortality and ICU use by age groups among three periods of the pandemic's first year in the regions of Brazil

Reg./ Age group (Ys)	Epidemiologic weeks 8/2020 – 25/2020 (16/02/2020 – 20/06/2020)					Epidemiologic weeks 26/2020 – 43/2020 (21/06/2020 – 24/10/2020)					Epidemiologic weeks 44/2020 – 7/2021 (25/10/2020 – 20/02/2021)					χ^2 (p) (1)	χ^2 (p) (2)
	N	Inpatient deaths		ICU use		N	Inpatient deaths		ICU use		N	Inpatient deaths		ICU use			
		N	%	N	%		N	%	N	%		N	%				
North																	
0-5	335	49	14,6	70	20,9	736	40	5,4	87	11,8	645	59	9,1	97	15,0	0,00	0,00
6-17	324	41	12,7	46	14,2	450	28	6,2	52	11,6	236	24	10,2	32	13,6	0,01	0,52
18-30	1274	159	12,5	140	11,0	1200	105	8,8	152	12,7	1096	201	18,3	202	18,4	0,00	0,00
31-40	2493	423	17,0	408	16,4	1623	175	10,8	277	17,1	2451	613	25,0	533	21,7	0,00	0,00
41-50	3323	883	26,6	627	18,9	2023	365	18,0	404	20,0	3356	1094	32,6	801	23,9	0,00	0,00
51-60	3815	1491	39,1	958	25,1	2459	653	26,6	597	24,3	4073	1761	43,2	1120	27,5	0,00	0,01
61-70	4570	2549	55,8	1346	29,5	2767	1080	39,0	795	28,7	4804	2687	55,9	1490	31,0	0,00	0,08
71-80	3879	2514	64,8	1168	30,1	2420	1172	48,4	786	32,5	3622	2352	64,9	1171	32,3	0,00	0,06
>80	2570	1881	73,2	686	26,7	1587	928	58,5	465	29,3	2276	1664	73,1	703	30,9	0,00	0,01
Northeast																	
0-5	765	118	15,4	183	23,9	911	82	9,0	254	27,9	720	61	8,5	225	31,3	0,00	0,01
6-17	507	66	13,0	135	26,6	655	44	6,7	159	24,3	334	42	12,6	89	26,6	0,00	0,58
18-30	2647	385	14,5	486	18,4	1776	243	13,7	413	23,3	1419	192	13,5	383	27,0	0,59	0,00
31-40	5290	919	17,4	1116	21,1	3295	548	16,6	852	25,9	3335	492	14,8	959	28,8	0,01	0,00
41-50	7215	1763	24,4	1694	23,5	4461	972	21,8	1300	29,1	4532	932	20,6	1507	33,3	0,00	0,00
51-60	9295	3222	34,7	2681	28,8	6178	1816	29,4	2033	32,9	6186	1745	28,2	2242	36,2	0,00	0,00
61-70	10807	5159	47,7	3499	32,4	7622	3028	39,7	3009	39,5	7251	2721	37,5	3072	42,4	0,00	0,00
71-80	10273	6008	58,5	3535	34,4	7897	3893	49,3	3388	42,9	6636	3199	48,2	3061	46,1	0,00	0,00
>80	7814	5310	68,0	2537	32,5	6733	4008	59,5	2885	42,8	5312	3104	58,4	2405	45,3	0,00	0,00
Southeast																	
0-5	909	65	7,2	308	33,9	1026	45	4,4	283	27,6	1006	44	4,4	299	29,7	0,01	0,01
6-17	545	52	9,5	158	29,0	812	55	6,8	257	31,7	563	23	4,1	167	29,7	0,00	0,53
18-30	4396	427	9,7	1071	24,4	4368	366	8,4	1103	25,3	3948	424	10,7	1002	25,4	0,00	0,50
31-40	11624	1327	11,4	3113	26,8	10643	995	9,3	2853	26,8	10691	1177	11,0	3006	28,1	0,00	0,04
41-50	16838	2815	16,7	4833	28,7	16069	2236	13,9	4724	29,4	16848	2421	14,4	5006	29,7	0,00	0,11
51-60	20977	5548	26,4	6925	33,0	22077	4733	21,4	7287	33,0	24043	5387	22,4	7974	33,2	0,00	0,92
61-70	21710	8834	40,7	8053	37,1	26056	8898	34,1	9831	37,7	29455	10090	34,3	10973	37,3	0,00	0,31
71-80	16949	8994	53,1	7013	41,4	21713	10050	46,3	8923	41,1	24464	11052	45,2	9838	40,2	0,00	0,04
>80	13238	8639	65,3	5602	42,3	16837	9669	57,4	6885	40,9	18534	10655	57,5	7388	39,9	0,00	0,00
South																	
0-5	64	2	3,1	17	26,6	292	11	3,8	66	22,6	292	19	6,5	64	21,9	0,24	0,72
6-17	41	4	9,8	17	41,5	223	14	6,3	53	23,8	197	20	10,2	44	22,3	0,33	0,03
18-30	555	33	5,9	115	20,7	1683	142	8,4	388	23,1	1812	165	9,1	387	21,4	0,06	0,36
31-40	1145	93	8,1	284	24,8	3962	299	7,5	977	24,7	4706	510	10,8	1142	24,3	0,00	0,88
41-50	1695	192	11,3	521	30,7	5958	768	12,9	1719	28,9	7593	1117	14,7	2002	26,4	0,00	0,00
51-60	2014	351	17,4	718	35,7	8418	1624	19,3	2843	33,8	10957	2382	21,7	3350	30,6	0,00	0,00
61-70	1904	576	30,3	816	42,9	9507	3225	33,9	3929	41,3	13174	4620	35,1	4951	37,6	0,00	0,00
71-80	1395	620	44,4	657	47,1	7371	3424	46,5	3212	43,6	10770	5069	47,1	4412	41,0	0,16	0,00
>80	883	541	61,3	394	44,6	5229	3041	58,2	2003	38,3	7164	4193	58,5	2448	34,2	0,22	0,00
Central West																	
0-5	64	7	10,9	20	31,3	344	16	4,7	98	28,5	126	6	4,8	31	24,6	0,12	0,58
6-17	47	4	8,5	15	31,9	222	12	5,4	63	28,4	101	5	5,0	33	32,7	0,66	0,70
18-30	489	37	7,6	124	25,4	1486	143	9,6	309	20,8	888	90	10,1	213	24,0	0,28	0,05
31-40	1226	135	11,0	386	31,5	3573	383	10,7	855	23,9	2142	201	9,4	546	25,5	0,20	0,00
41-50	1621	243	15,0	542	33,4	5310	833	15,7	1386	26,1	3149	464	14,7	882	28,0	0,47	0,00
51-60	1810	422	23,3	707	39,1	6777	1609	23,7	2135	31,5	4220	907	21,5	1415	33,5	0,02	0,00
61-70	1605	665	41,4	704	43,9	7518	2666	35,5	2763	36,8	4278	1450	33,9	1635	38,2	0,00	0,00
71-80	1215	631	51,9	605	49,8	6077	2814	46,3	2557	42,1	3334	1560	46,8	1538	46,1	0,00	0,00
>80	799	522	65,3	410	51,3	4083	2451	60,0	1862	45,6	2243	1301	58,0	1058	47,2	0,00	0,01

Source: <https://opendatasus.saude.gov.br>. The periods were defined based on the epidemiologic week of onset of the first symptoms. (1) Inpatient deaths; (2) ICU use

Table 3. Comparison of ICU length of stay among Covid-19 patients (N=190,833) by region and age group, among three periods of the pandemic's first year In Brazil

Reg./ Age group (Ys)	Epidemiologic weeks 8/2020-25/2020					Epidemiologic weeks 26/2020-43/2020					Epidemiologic weeks 44/2020-7/2021					KW Test (p)
	(16/02/2020 – 20/06/2020)					(21/06/2020 – 24/10/2020)					(25/10/2020 – 20/02/2021)					
	N	Mean	Std	Md	Max	N	Mean	Std	Md	Max	N	Mean	Std	Md	Max	
North																
0-5	54	13.3	26.4	6	157	42	10.1	14.9	5	70	52	11.1	15.8	5.5	92	0.90
6-17	36	11.2	15.2	7	84	32	5.7	5.5	5	22	19	8.9	7.3	6	25	0.11
18-30	113	11.2	10.8	8	57	113	11.6	12.9	7	86	147	9.8	8.6	7	58	0.93
31-40	324	9.7	9.2	7	50	194	10.2	9.5	7	63	426	9.6	8.6	7	46	0.88
41-50	529	10.5	12.0	7	108	318	10.9	10.2	8	57	636	10.8	10.3	7.5	68	0.23
51-60	847	10.3	12.7	7	130	495	11.7	10.6	9	71	939	10.7	10.2	8	107	0.00
61-70	1220	10.2	13.5	7	223	672	12.8	14.6	8	146	1283	10.8	9.4	8	67	0.00
71-80	1082	9.2	10.2	6	98	677	11.9	18.0	7	357	1042	9.5	8.4	7	63	0.00
>80	625	7.7	8.7	5	67	418	10.2	12.1	7	77	641	8.5	8.1	6	60	0.00
Northeast																
0-5	109	11.4	15.8	5	96	110	12.3	16.9	6	109	85	11.8	15.0	5	89	0.67
6-17	88	10.6	13.5	6	70	82	10.1	13.3	5.5	81	47	10.9	14.0	5	59	0.98
18-30	342	10.3	16.5	5	175	264	10.4	16.1	5	187	256	8.8	10.2	6	78	0.91
31-40	768	10.2	11.9	7	105	536	9.9	11.0	6.5	89	588	8.6	9.4	6	83	0.15
41-50	1265	11.0	14.1	7	147	876	10.3	11.8	7	118	998	9.1	9.0	7	98	0.81
51-60	2094	12.0	14.0	8	149	1388	11.4	12.4	8	98	1528	11.0	9.9	8	68	0.42
61-70	2879	11.3	13.1	8	176	2247	11.9	12.9	8	146	2221	11.0	10.2	8	102	0.07
71-80	3062	11.0	12.7	7	155	2679	11.9	13.2	8	161	2319	11.0	10.2	9	93	0.00
>80	2217	9.8	13.3	6	276	2352	9.6	10.7	7	155	1943	9.4	8.8	7	75	0.02
Southeast																
0-5	157	13.4	24.1	6	187	141	10.3	15.9	5	103	123	7.9	9.8	4	49	0.10
6-17	100	7.7	9.7	4	62	144	10.4	20.3	5	190	71	8.4	9.3	6	55	0.78
18-30	690	9.5	12.7	6	183	665	9.4	11.7	5	108	611	8.6	9.5	5	55	0.10
31-40	1923	10.1	12.3	6	151	1765	10.1	12.5	6	164	1792	9.2	9.3	6	83	0.78
41-50	3110	10.8	12.3	7	204	3005	11.3	13.0	7	182	3128	9.8	11.3	7	374	0.64
51-60	4942	12.6	14.3	9	225	5062	13.1	14.5	9	212	5393	11.4	10.6	8	108	0.76
61-70	6330	13.4	15.9	9	289	7511	13.4	13.7	9	164	8059	12.1	10.9	9	97	0.06
71-80	5807	12.6	15.1	8	240	7291	12.7	13.1	9	174	7615	11.8	10.4	9	97	0.00
>80	4783	11.3	14.4	7	209	5806	10.3	11.5	7	191	6003	10.0	9.2	8	102	0.02
South																
0-5	16	11.7	14.3	5.5	44	58	15.5	23.1	7.5	129	49	10.3	9.5	7	43	0.73
6-17	15	18.5	13.7	19	53	47	12.9	14.5	6	61	33	10.5	12.4	7	58	0.09
18-30	103	11.0	12.8	6	83	352	10.0	10.6	7	71	321	9.8	10.8	7	65	0.97
31-40	255	12.3	12.2	8	85	859	12.0	14.0	7	110	946	10.4	10.0	7	68	0.27
41-50	477	12.2	13.2	8	108	1575	12.9	13.8	9	177	1727	11.7	11.2	8	105	0.17
51-60	661	14.4	14.7	9	105	2600	14.3	14.9	10	169	2955	12.7	11.0	10	99	0.55
61-70	761	15.5	14.7	11	120	3693	14.9	14.9	11	138	4577	13.2	11.5	10	97	0.00
71-80	631	14.8	16.9	10	222	3086	13.9	13.9	10	156	4170	12.3	11.1	9	105	0.00
>80	384	11.6	12.0	8	89	1933	10.2	11.7	7	185	2335	9.7	9.1	7	95	0.01
Central West																
0-5	13	7.4	7.4	3	20	63	7.9	10.6	4	63	27	8.7	10.0	5	40	0.64
6-17	6	6.8	7.3	4.5	20	40	9.7	9.9	6	46	26	8.0	8.7	4	33	0.44
18-30	85	8.8	10.4	4	47	198	10.2	12.9	6	110	156	7.7	6.8	5	42	0.38
31-40	252	11.4	16.7	6.5	177	538	9.3	9.7	6	53	368	9.1	8.5	7	72	0.64
41-50	357	11.2	12.5	7	112	972	11.2	12.9	7.5	125	649	10.4	10.5	7	96	0.85
51-60	499	12.5	16.1	8	202	1572	12.5	14.3	8	163	1063	11.3	9.9	9	83	0.46
61-70	580	13.5	14.1	9	112	2192	12.7	13.7	9	159	1308	11.9	10.0	9	89	0.62
71-80	510	13.4	14.0	9	99	2137	11.8	12.7	8	156	1317	12.0	10.6	9	85	0.03
>80	366	11.2	12.4	8	112	1637	10.3	11.8	7	212	937	9.8	8.8	7	66	0.66

Source: <https://opendatasus.saude.gov.br>. The periods were defined based on the epidemiologic week of onset of the first symptoms. Std=Standard deviation; Md=Median; Max=Maximum. KW=Kruskal-Wallis Test.