Full title: Exploring the causal effect of maternal pregnancy adiposity on offspring adiposity: Mendelian randomization using polygenic risk scores

Short title: Exploring the causal effect of maternal pregnancy adiposity on offspring adiposity

Tom A Bond^{1,2,3,4,5,*}, Rebecca C Richmond^{4,5}, Ville Karhunen^{1,6,7}, Gabriel Cuellar-Partida^{3,8}, Maria Carolina Borges^{4,5}, Verena Zuber^{1,9}, Alexessander Couto Alves^{1,10}, Dan Mason¹¹, Tiffany C Yang¹¹, Marc J Gunter¹², Abbas Dehghan^{1,2}, Ioanna Tzoulaki^{1,2,13}, Sylvain Sebert⁶, David M Evans^{3,4}, Alex M Lewin^{1,14}, Paul F O'Reilly¹⁵, Deborah A Lawlor^{4,5,§}, Marjo-Riitta Järvelin^{1,2,6,16,17,§}

¹Department of Epidemiology and Biostatistics, Imperial College London, London, UK.

²MRC-PHE Centre for Environment and Health, School of Public Health, Imperial College London, London, UK.

³The University of Queensland Diamantina Institute, The University of Queensland, Brisbane, Australia.

⁴MRC Integrative Epidemiology Unit at the University of Bristol, Bristol, UK.

⁵Population Health Sciences, Bristol Medical School, University of Bristol, Bristol, UK.

⁶Center for Life-course Health Research, Faculty of Medicine, University of Oulu, Oulu, Finland.

⁷Research Unit of Mathematical Sciences, University of Oulu, Oulu, Finland.

⁸23andMe, Inc., Sunnyvale, CA, USA.

⁹MRC Biostatistics Unit, School of Clinical Medicine, University of Cambridge, Cambridge, UK.

¹⁰School of Biosciences and Medicine, Faculty of Health and Medical Sciences, University of Surrey, Guildford, UK.

¹¹Born in Bradford, Bradford Institute for Health Research, Bradford Teaching Hospitals NHS Foundation Trust, Bradford, UK.

¹²Section of Nutrition and Metabolism, IARC, Lyon, France.

¹³Department of Hygiene and Epidemiology, University of Ioannina Medical School, Ioannina, Greece

¹⁴Department of Medical Statistics, London School of Hygiene and Tropical Medicine, London, UK.

¹⁵Genetics and Genomic Sciences, Icahn School of Medicine at Mount Sinai, New York, USA.

¹⁶Unit of Primary Care, Oulu University Hospital, Oulu, Finland.

¹⁷Department of Life Sciences, College of Health and Life Sciences, Brunel University London, London, UK.

*Corresponding author

§These authors contributed equally to this work

Abstract

Background

It has been hypothesised that greater maternal adiposity before or during pregnancy causes greater offspring adiposity in childhood and adulthood, via causal intrauterine or periconceptional mechanisms. Previous Mendelian randomization (MR) estimates were imprecise, with wide confidence intervals that included potentially important protective or adverse effects, and may have been biased by collider effects or imperfect adjustment for genetic inheritance. Here we use an improved MR approach to investigate whether associations between maternal pre-/early pregnancy body mass index (BMI) and offspring adiposity from birth to adolescence are causal, or are instead due to confounding.

Methods and findings

We undertook confounder adjusted multivariable (MV) regression and Mendelian randomization (MR) using mother-offspring pairs from two UK cohorts: Avon Longitudinal Study of Parents and Children (ALSPAC) and Born in Bradford (BiB). In ALSPAC and BiB the outcomes were birthweight (BW; $N = 9339$) and BMI at age 1 ($N = 8659$) and 4 years ($N = 7575$), and in ALSPAC only we investigated BMI at 10 ($N = 4476$) and 15 years ($N = 4112$) and dual-energy X-ray absorptiometry (DXA) determined fat mass index (FMI) from age 10–18 years ($N = 2659$ to 3855). We compared MR results from several polygenic risk scores (PRS), calculated from maternal non-transmitted alleles at between 29 and 80,939 single nucleotide polymorphisms (SNPs). MV and MR showed a consistent positive association of maternal BMI with BW, but for adiposity at most older ages MR estimates were weaker than MV estimates. In MV regression a one standard deviation (SD) higher maternal BMI was associated with a 0.13 (95% confidence interval [CI]: 0.10, 0.16) SD increase in offspring BW. The corresponding MR estimate from the strongest PRS (including up to 80,939 SNPs) was 0.14 (95% CI: 0.05, 0.23), with no difference between the two estimates ($P_{\text{difference}} = 0.84$). For 15 year BMI the MV and MR estimates (80,939 SNPs) were 0.32 (95% CI: 0.29, 0.36) and 0.13 (95% CI: 0.01, 0.24) respectively ($P_{\text{difference}} = 1.0 \times 10^{-3}$). Results for FMI were similar to those for adolescent BMI. As the number of SNPs included in the PRS increased, the MR confidence intervals narrowed and the effect estimates for adolescent adiposity became closer to the MV estimates. Sensitivity analyses suggested the stronger effects with more SNPs were explained by horizontal pleiotropic bias away from zero. Consequently, the unbiased difference between the MV and MR estimates is probably greater than shown in our main analyses. Furthermore, MR estimates from IVs with fewer SNPs provided no strong evidence for a causal effect on adolescent adiposity.

Conclusions

Our results suggest that higher maternal pre-/early-pregnancy BMI is not a key driver of higher adiposity in the next generation. Thus, they support interventions that target the whole population for reducing overweight and obesity, rather than a specific focus on women of reproductive age.

Introduction

It has been hypothesised that prenatal exposure to greater maternal adiposity during or prior to pregnancy causes greater adiposity in the offspring throughout life, via intrauterine effects or periconceptional mechanisms (for example effects on the oocyte) (1-4). There are well replicated observational associations between maternal body mass index (BMI) before or during pregnancy and offspring adiposity and cardiometabolic outcomes in childhood, adolescence and adulthood (5-8). Furthermore, evidence from animal experiments suggests that such associations are plausibly due to causal biological effects in the intrauterine period (9, 10). If true, this could have important implications for obesity prevention policy, because interventions to reduce maternal obesity before pregnancy might reduce offspring obesity risk in later life (1, 2, 6).

Triangulated epidemiological evidence from different study designs (11) suggests that associations between maternal BMI and offspring childhood/adolescent adiposity may *not* reflect a causal effect. For example, negative paternal exposure control studies (12-18) and studies examining associations within sibling groups (19, 20) suggest that confounding by genetic and/or environmental factors shared within families may be an important explanation for the associations. In addition, two Mendelian randomization (MR) (21, 22) studies, which used genetic variants as instrumental variables (IVs) for maternal BMI, provided no strong evidence for a causal effect (14, 23). However, the primary analysis in the most recent MR study (23) was adjusted for an offspring weighted allele score, and simulations suggest that the use of a *weighted* allele score may not be the optimal approach to avoid bias due to genetic inheritance (24). Also, previous studies (21, 22) were only able to adjust for offspring genetic variants, and it may be necessary to control for paternal genotype as well to avoid collider bias (25) (see **Supplementary information S1**). Furthermore, the causal estimates from previous MR studies were imprecise (14, 23, 25). For example, in the largest study ($N = 6057$) a one SD higher maternal BMI was associated with a 0.05 SD increase in mean offspring BMI at age 7, but the 95% confidence interval was consistent with a 0.11 SD reduction or a 0.21 SD increase (23). If a positive causal effect is present this could have important public health implications, because it could lead to an accelerating intergenerational cycle of obesity that is difficult to break (1, 26). It is therefore important to conduct further MR investigations with improved methods, in order to obtain more precise estimates that are not subject to the aforementioned biases.

An alternative approach for avoiding bias due to genetic inheritance is to use maternal alleles that were not inherited by the offspring (maternal non-transmitted alleles) (27, 28), which avoids the need to adjust for offspring genotype. The precision of estimates could be improved by increasing the study sample size, but the feasibility of this is limited by the small number of studies worldwide with the necessary data for maternal non-transmitted allele analyses: maternal and offspring genotype, maternal pregnancy exposures and offspring outcomes. An alternative way to increase precision would be to increase the strength of the genetic IV (29). Previous MR studies followed the widely used practice of using only genome wide significant (GWS; $P < 5e-8$) genetic variants (hereafter referred to as single nucleotide polymorphisms [SNPs]) to calculate the genetic risk score to be used as an IV (14, 23). However, for polygenic traits such as BMI it is known that substantially improved phenotypic prediction can be achieved by including more SNPs in the genetic risk score (i.e. more weakly associated SNPs that individually are not GWS) (30-33). Such polygenic risk scores (PRS) have been shown to associate strongly with BMI throughout the life course (31).

We aimed to use maternal non-transmitted allele PRS as IVs in a one-sample MR design, to explore the causal effect of maternal BMI on offspring adiposity from birth to adolescence,

and to compare those results with confounder adjusted multivariable (MV) regression estimates. Based on previous MR studies (23, 28, 34) we hypothesised that greater maternal BMI would cause increased offspring BW, but that the causal effect would attenuate over childhood and adolescence. **Supplementary information S1** shows our MR analyses diagrammatically.

Methods

Study design

We conducted one-sample MR and compared these results with confounder adjusted multivariable (MV) regression analyses. We analysed data from two population based prospective birth cohorts: the Avon Longitudinal Study of Parents and Children (ALSPAC) and Born in Bradford (BiB). ALSPAC enrolled pregnant women who resided in and around the city of Bristol in the South West of England and had an expected delivery date between April 1, 1991 and December 31, 1992. The enrolled cohort included 15,247 pregnancies resulting in 14,775 live born babies. Ethical approval was obtained from the ALSPAC Ethics and Law Committee and the Local Research Ethics Committees. The study website contains details of all the data that are available through a fully searchable data dictionary and variable search tool and details of the study methodology have been reported previously (35, 36). BiB enrolled women who resided in the city of Bradford in the North of England who attended an antenatal booking clinic between March 2007 and December 2010. The recruited cohort included 13,776 pregnancies resulting in 13,740 live born babies. Ethical approval was obtained from Bradford National Health Service Ethics Committee (ref 06/Q1202/48) and details of the study methodology have been reported previously (37).

Selection of participants

Details of sample selection for each cohort are given in the flow charts in **Supplementary information S2**. We included live-born singletons with maternal and offspring genotype data, maternal BMI data and at least one offspring adiposity measure available, and selected one offspring from any sibling groups for inclusion (chosen at random in ALSPAC or to maximise the sample size with data available in BiB). As the effects we were exploring may differ by ethnicity (38) we limited analyses to two ethnic groups: White European and South Asian. There were very few participants from other ethnic groups in either cohort, therefore these participants were excluded. ALSPAC (93% White European) contributed only to the analyses in White Europeans and we meta-analysed these results with those from models fitted separately for BiB South Asians and BiB White Europeans. In ALSPAC, ethnicity was assessed by genetic multidimensional scaling (MDS) analysis during the quality control procedure for maternal, paternal and offspring genotype data. In BiB, mothers reported their ethnicity and the place of birth of their parents at the baseline interview and this was augmented by information from General Practitioner records (**Supplementary information S3**). For the offspring phenotypes the overall sample size for MR analyses ranged from 2659–5085 for ALSPAC, 1566–2262 for BiB South Asians and 1339–1992 for BiB White Europeans. The sample sizes for confounder adjusted MV estimates were somewhat smaller due to missing confounder data (1884–3265 for ALSPAC, 325–449 for BiB South Asians and 442–604 for BiB White Europeans). To enable comparison between the confounder adjusted MV estimates from models that adjusted for different covariates, we fitted all the models for each outcome using an identical sample with non-missing data for all relevant variables.

Parental anthropometric variables

In ALSPAC, maternal pre-pregnancy weight and height were retrospectively reported by the women during pregnancy (at a mean gestational age of 24.7 weeks [SD 6.3]) or postnatally for 11.2% of mothers (at a mean of 22.0 weeks after birth [SD 6.7]); the reported weights correlated highly with weight recorded at the first antenatal clinic (Pearson correlation coefficient = 0.96). Paternal height and weight were reported by the fathers during their partner's pregnancy (or postnatally for a minority of fathers). In BiB, early pregnancy BMI was calculated from height reported by the mothers at recruitment (26–28 weeks gestation) and weight extracted from the first antenatal clinic records (median 12 weeks gestation). Paternal height and weight were reported by the fathers at recruitment, which for the majority of fathers was at the time of their partner's pregnancy.

Offspring anthropometric variables

Offspring outcomes included BW and BMI at age 1 and 4 years (in ALSPAC and BiB), BMI at age 10 and 15 years (ALSPAC only) and FMI at age 10, 12, 14, 16 and 18 years (ALSPAC only). The assessment of these outcomes is described in **Supplementary information S4**. Offspring BW was extracted from the birth record in BiB, and extracted from the birth record/notification or measured by research staff in ALSPAC. Birth length in ALSPAC was obtained similarly, and is not available for BiB participants. In ALSPAC, child and adolescent height and weight were obtained from clinical examination by study staff using a Harpenden Stadiometer (Holtain Limited, Dyfed, UK) and Tanita Body Fat Analyser (Model TBF 305; Tanita UK Limited, Viewsley, UK) respectively, or from child health records or maternal/offspring questionnaire responses (**Supplementary information S5**). In BiB, childhood height and weight were obtained from a variety of sources including clinical measurement by study staff or measurement at around age 4 years as part of the UK Government National Child Measurement Programme (NCMP) (in both cases using a Leicester Height Measure [Seca] and Seca digital scales), child health records, primary care records and school nurse records (**Supplementary information S5**). In ALSPAC, we calculated FMI as fat mass (kg) / height (m)² using fat mass measured by whole body DXA carried out with a Lunar Prodigy DXA scanner (GE Medical Systems Lunar, Madison, WI, USA). The scans were visually inspected and realigned where necessary. Once complete, the tester examined the scan to ensure its quality, and if necessary repeated the scan.

Anthropometric variable standardisation

In each of the three samples (ALSPAC, BiB White Europeans and BiB South Asians) we internally standardised exposure and outcome variables to give measures in standard deviation (SD) units. We standardised maternal BMI by maternal age (at delivery), in one year age categories. We standardised offspring BW by sex, and offspring BMI and FMI by sex and age (in one month categories).

Confounder adjusted multivariable regression

We considered the following variables to be potential confounders: maternal age (which was adjusted for in the standardised exposure by calculating z-scores within maternal age strata), parity, maternal smoking during pregnancy, parental occupation, maternal educational attainment, paternal educational attainment and paternal BMI. Standard protocols for assessing these variables were used in each cohort, and full details are provided in **Supplementary information S6**. We fitted three MV regression models: in model one we adjusted for maternal age, offspring age and offspring sex, in model two we additionally adjusted for the potential confounders listed above except for paternal BMI, and in model three (which was the main multivariable model of interest and is presented in **Results**) we additionally adjusted for paternal BMI. We took a complete case approach and

excluded individuals with any missing data, therefore models one to three were fitted using identical samples. In sensitivity analyses we adjusted all models for gestational age at delivery, and for 20 genetic principal components (PCs) which we calculated from genome-wide SNPs separately for each of the three samples (**Supplementary information S7**), in order to adjust for ancestry. In BiB we had to exclude a large number of individuals from the main MV models due to missing paternal BMI data. We therefore refitted models one and two without first excluding individuals with missing paternal BMI data (i.e. on larger samples), in order to explore potential selection bias.

Genotyping, quality control and imputation

Mothers, offspring and (in ALSPAC only) fathers were genotyped using genome-wide arrays, followed by standard quality control (QC) measures (**Supplementary information S8**). Array genotypes were then imputed to the Haplotype Reference Consortium (HRC), 1000 Genomes or UK10K reference panels (39-41) (**Supplementary information S8**). In order to maximize the sample size we did not exclude cryptically related individuals for the primary analyses. As a sensitivity analysis we removed cryptic relatedness at a level corresponding to first cousins by applying a KING (48) kinship coefficient threshold of 0.044 to the offspring using the PLINK software package version 2.00 (49, 50).

Inference of maternal non-transmitted alleles

After conversion of imputed genotypes to hard calls (integer valued allele dosages) and application of QC filters (**Supplementary information S8**), we phased offspring imputed SNPs (for the sample of genotyped mother-offspring duos) using the duoHMM method implemented in the SHAPEIT v2 (r904) software package, with a window size of 5 Mb as per the authors recommendations for parent-offspring duos (42). This yielded maternal transmitted alleles (i.e. maternal alleles that were inherited by the offspring), which we used (along with the maternal genotypes) to infer the maternal non-transmitted alleles. For MR analyses we used PRS calculated from the maternal non-transmitted alleles (27), having first estimated SNP weights using maternal genotypes (see below).

Polygenic risk score calculation

In order to maximise statistical power we used genome-wide SNPs to calculate a BMI PRS, as a weighted sum of BMI-increasing maternal non-transmitted alleles at SNPs across the genome. We tested four PRS methods (clumping and thresholding (43), LDpred (44), lassosum (45, 46) and the BOLT-LMM linear predictor (47)) (**Supplementary information S9**). Of these four methods, lassosum explained the highest proportion of variance (R^2) for maternal BMI in both ALSPAC and BiB (which we refer to as the target datasets), therefore we used the lassosum PRS for subsequent MR analyses. Lassosum requires summary statistics from a genome wide association study (GWAS), which we refer to as the base dataset. We conducted a GWAS in the UK Biobank (UKB), a prospective cohort of 502,628 volunteers (with 5% response rate of those invited), recruited from across the UK at age 40–69 years between 2006 and 2010 (48, 49) (**Supplementary information S9**). In order to avoid overfitting due to overlap between the base and target samples we excluded attendees of the Bristol or Leeds UKB assessment centres. We meta-analysed the summary statistics from the UKB GWAS with a published BMI GWAS from the GIANT consortium (50), giving a total base sample size of up to 756,048. We applied the lassosum algorithm to the meta-analysed base dataset; lassosum uses penalised regression to carry out shrinkage and selection on the base GWAS SNP effects and accounts for LD information from a reference panel. We used the ALSPAC or BiB datasets as the reference panels as per the authors' recommendations) (45). PLINK was used to calculate the PRS for ALSPAC and BiB

individuals using the lassosum SNP weights for around 80,000 SNPs (see **Table 2** for the exact number of SNPs for each cohort). We also calculated three PRS from fewer SNPs, to be used in sensitivity analyses to explore potential pleiotropic effects (see below). These PRS used (i) around 30 GWS SNPs identified in a 2010 BMI GWAS (51), (ii) around 90 GWS SNPs identified in a 2015 BMI GWAS (50), and (iii) around 500 GWS SNPs identified as primary signals in a 2018 BMI GWAS (30). Full details of these analyses, including the exact number of SNPs used to calculate each PRS (which varied between samples) are given in **Supplementary information S10**.

Mendelian randomization

For the primary MR analyses we used the lassosum non-transmitted allele BMI PRS as an IV for maternal BMI and fitted models using the two-stage least squares (TSLS) method (22) (i.e. one sample MR). We included 20 genetic PCs as covariates in order to adjust for population stratification. We tested for a difference between the most extensively confounder adjusted MV estimates (model three) and MR estimates using a z-test (**Supplementary information S11**), and used a bootstrapping procedure to estimate the covariance between MV and MR estimates in order to calculate the z-statistic. Evidence for a difference between the two could reflect residual confounding in the MV analyses or violation of one or more of the MR assumptions.

Meta-analysis

We examined the point estimates, I^2 statistics and Cochran's Q test P-values for the MV and MR associations and found little evidence for heterogeneity between ALSPAC, BiB South Asians and BiB White Europeans (**Supplementary information S12**). We therefore meta-analysed estimates from the three samples using a fixed effects model. Results were similar when we instead used a random effects model. For the meta-analyses we used the ratio estimator (calculated as the meta-analysed PRS-outcome regression coefficient $\hat{\beta}_{ZY}$ divided by the meta-analysed PRS-exposure regression coefficient $\hat{\beta}_{ZX}$; in the present study both coefficients were estimated in the same sample) which gives equivalent results to TSLS (52). We calculated the standard errors for the pooled MR estimates using a Taylor series approximation (53).

Checking MR assumptions

We checked the assumptions made by MR analyses (**Supplementary information S1**); if these assumptions are met then our MR estimates can be interpreted as causal effect estimates (22). We first assessed whether the PRS were associated with maternal BMI using the R^2 and F-statistics. Next, we explored whether the PRS-outcome associations were confounded by ancestry (population stratification) using a linear mixed model (LMM). LMMs have been widely used in GWAS to adjust for population stratification and cryptic relatedness (54). We fitted models for the numerator and denominator of the ratio estimator separately, using the `--reml-est-fix` command in the GCTA software package (version 1.91.7beta) (55). Further details of the LMM approach are given in **Supplementary information S13**. Finally, we conducted several analyses to explore whether the maternal PRS influences offspring adiposity via mechanisms other than intrauterine or periconceptional exposure to increased maternal BMI (horizontal pleiotropy). We first tested for associations of the PRS with other potential risk factors for the offspring outcomes (56). We would expect that the risk of pleiotropic bias might decrease as fewer SNPs are included in the IV. We therefore repeated MR analyses with IVs calculated from a single BMI-associated SNP (rs9939609 at the *FTO* locus, the locus at which there is currently the strongest evidence for association with BMI (30)), as well as the three PRS calculated from

only strongly BMI-associated (GWS) SNPs, as described above. Furthermore, most of the SNPs included in the lassosum BMI PRS had small effect sizes, and the consequences of this for the extent of horizontal pleiotropic effects are unclear (57), so we explored how MR estimates varied with varying SNP effect size distributions. We also tested for evidence of between-SNP MR estimate heterogeneity (Cochran's Q test) and used MR Egger regression (58) to investigate horizontal pleiotropy, for the analyses based on GWS SNPs. Finally, to investigate collider bias and bias due to assortative mating we examined the association between the maternal and paternal lassosum BMI PRS, in the subset of ALSPAC participants with paternal genotype data available ($N = 1325$).

Other sensitivity analyses

We explored departure from linearity of the MV and MR associations by examining augmented partial residual plots with overlaid linear regression lines and nonparametric loess smoothers (59). The residuals from several models involving adolescent BMI and FMI variables were somewhat positively skewed so we repeated MV and MR analyses using the natural log of the relevant variables. We examined whether results differed for BW, BMI and ponderal index (weight [kg] / length [m]³) at birth (in ALSPAC only as birth length was not available in BiB). Finally, we tested for interaction by offspring sex for the MV and MR models. We carried out statistical analyses in R version 3.5.1 (60), and Stata version 13.1 (StataCorp, College Station, TX, USA).

Results

Participant characteristics

Table 1 shows the participant characteristics. The prevalence of maternal obesity (maternal BMI ≥ 30) was 5.5% (95% confidence interval [CI]: 4.9%, 6.1%) in ALSPAC and markedly higher in BiB South Asians (20.5% [95% CI: 18.9%, 22.2%]) and BiB White Europeans (26.0% [95% CI: 24.1%, 28.0%]). There were similar differences in mean maternal BMI between samples. Mean maternal age was highest in ALSPAC, lower in BiB South Asian participants and the lowest in BiB white participants, whilst mean offspring BW was greatest in ALSPAC, lower in BiB white participants and markedly lower in BiB South Asian participants (P -values for all comparisons described here $< 6.9 \times 10^{-7}$). The samples for our analyses were smaller than those for the full cohorts at birth due to missing data, particularly for the MV associations. Despite this there were not large differences in the distributions of BW, maternal BMI or offspring sex between the baseline samples and those from which we calculated the MV estimates (**Supplementary information S14**). Furthermore, when we fitted MV models one and two on a larger sample (retaining individuals with missing paternal BMI) there were not large differences in the primary MV results (**Supplementary information S21**).

Table 1: Characteristics of the mothers and offspring in ALSPAC and BiB

Cohort	N	Exposure/outcome			Age at exposure/outcome assessment			Female offspring (%)
			Mean	SD		Mean	SD	
ALSPAC	5085	Maternal BMI (kg/m ²)	23.0	3.8	Maternal age at offspring birth (years)	29.3	4.5	
BiB (SA)	2262		25.8	5.5		28.8	5.0	
BiB (WE)	1992		26.9	6.0		27.4	6.0	
ALSPAC	5085	Birth weight (kg)	3.45	0.52	Gestational age at birth (weeks)	39.6	1.7	50.5
BiB (SA)	2262		3.12	0.49		39.1	1.5	47.9
BiB (WE)	1992		3.36	0.53		39.4	1.6	47.9
ALSPAC	4838	1yr BMI (kg/m ²)	17.5	1.5	Age at BMI measurement (years)	0.9	0.2	50.6
BiB (SA)	2023		16.7	1.6		0.8	0.3	48.1
BiB (WE)	1798		17.4	1.6		0.8	0.3	47.8
ALSPAC	4670	4yr BMI (kg/m ²)	16.1	1.5		4.1	0.7	50.2
BiB (SA)	1566		15.9	1.8		4.5	0.7	48.5
BiB (WE)	1339		16.4	1.5		4.5	0.7	48.6
ALSPAC	4476	10yr BMI (kg/m ²)	17.7	2.8		9.9	0.5	51.3
ALSPAC	4112	15yr BMI (kg/m ²)	21.0	3.5		15.0	0.8	51.7

SA: South Asians, **WE:** White Europeans, **SD:** standard deviations

353

354 Associations of genetic IVs with maternal BMI and offspring genotype

355 As we included more SNPs in the IV the R^2 for maternal BMI increased markedly, from <1%
 356 for the *FTO* IV to ~3–7% for the lassosum IV (**Table 2**). First-stage *F*-statistics were >75 for
 357 all lassosum MR models (**Supplementary information S15**). The lassosum maternal non-
 358 transmitted allele BMI PRS was not correlated with the offspring's PRS (results available
 359 from the authors on request).

360 Associations of maternal BMI and BMI PRS with confounders/outcome risk factors

361 There was strong evidence in all three samples for associations between maternal BMI and
 362 several other potential risk factors for the offspring outcomes, including parental occupation,
 363 educational attainment, maternal parity and paternal BMI (results are summarised in **Table**
 364 **2**, and full regression results including the direction of associations are given in
 365 **Supplementary information S16–S19**). Genetic IVs based on fewer SNPs (i.e. <100
 366 SNPs) were generally not associated with the outcome risk factors. In ALSPAC however
 367 there was strong evidence for association of the lassosum IV (based on 80,939 SNPs) with
 368 parental occupation, parental educational attainment, parental age, maternal smoking and
 369 paternal BMI. These associations were mostly present for BiB White Europeans but absent
 370 for BiB South Asians.

Table 2: Associations of maternal BMI with outcome risk factors, and of the genetic IVs (maternal non-transmitted alleles) with maternal BMI and outcome risk factors

			Correlation with outcome risk factors ^a							
			Parental occupation	Maternal education	Paternal education	Maternal smoking	Parity	Paternal BMI	Maternal age	Paternal age
ALSPAC										
<i>N</i>		5157	4807	4826	4572	4891	5042	3766	5157	3593
Maternal BMI ^b			0.11***	-0.12***	-0.11***	-0.01	0.06**	0.16***	0.00 ^c	-0.01 ^c
Genetic IV										
<i>FTO</i>	1	0.36%	0.03*	-0.01	-0.01	0.00	0.02	-0.01	-0.01	0.00
Speliotes	31	0.89%	0.02	0.00	-0.03*	-0.01	0.02	0.01	0.00	0.01
Locke	87	1.02%	0.03	-0.01	-0.04*	-0.01	0.03*	0.01	0.00	0.00
Yengo	497	2.37%	0.04*	-0.04*	-0.05**	0.02	0.03	0.01	-0.02	-0.01
Lassosum	80939	6.61%	0.08***	-0.07***	-0.08***	0.07***	0.02	0.03*	-0.08***	-0.06**
BiB (SA)										
<i>N</i>		2267	1689	2259	2258	2262	2215	475	2267	583
Maternal BMI ^b			-0.10**	-0.08**	-0.03	0.01	0.15***	0.12*	-0.02 ^c	0.02 ^c
Genetic IV										
<i>FTO</i>	1	0.77%	-0.01	-0.01	0.00	-0.03	0.00	0.00	0.02	0.06
Speliotes	29	1.71%	-0.01	0.01	0.01	-0.04	0.00	-0.01	0.00	-0.05
Locke	82	1.33%	-0.04	0.02	0.03	-0.05*	-0.01	-0.06	0.00	0.02
Yengo	446	1.64%	-0.04	-0.04*	0.01	0.01	0.01	-0.03	-0.02	-0.03
Lassosum	79101	3.46%	-0.01	-0.02	-0.02	0.01	-0.02	-0.04	-0.04	-0.06
BiB (WE)										
<i>N</i>		2000	1587	2000	1999	1999	1951	639	2000	788
Maternal BMI ^b			-0.09**	-0.07*	0.01	0.01	0.11***	0.22***	0.00 ^c	0.04 ^c
Genetic IV										
<i>FTO</i>	1	0.56%	0.03	0.02	-0.01	0.00	0.00	0.03	0.00	-0.02
Speliotes	31	0.92%	0.05	0.01	0.03	-0.03	0.00	0.02	0.00	-0.01
Locke	86	1.16%	0.03	0.00	0.04	-0.01	-0.01	0.04	-0.02	-0.02
Yengo	453	1.78%	-0.01	0.00	0.05*	0.01	0.00	0.00	-0.03	0.00
Lassosum	79101	5.21%	-0.08*	-0.05*	0.04	0.07*	0.04	0.03	-0.07*	0.01

373

374 **a:** Pearson correlation coefficients are presented here to give an indication of the direction and
375 magnitude of associations; full regression results are presented in **Supplementary information S16–**
376 **S19**, * $P < 0.05$, ** $P < 0.001$, *** $P < 1e-5$, **b:** age-standardised z-scores for maternal BMI, as per the
377 primary analyses, **c:** maternal BMI is not correlated with maternal or paternal age because it was age-
378 standardised, **SA:** South Asians, **WE:** White Europeans, ***R*²:** proportion of maternal BMI variance
379 explained by the IV (maternal non-transmitted allele scores), ***FTO*:** rs9939609 at the *FTO* locus,
380 **Speliotes, Locke, Yengo:** PRS calculated from SNPs that reached genome wide significance in the
381 BMI GWAS with the indicated first author, **Lassosum:** PRS calculated by the lassosum method

Confounder adjusted MV regression and MR results

In confounder adjusted MV regression models maternal BMI was positively associated with all offspring outcomes (**Figure 1, Supplementary information S20**; meta-analysis heterogeneity statistics are given in **Supplementary information S12**). When we meta-analysed MV estimates from all three samples, a 1 SD higher age-adjusted maternal BMI was associated with a 0.13 (95% CI: 0.10, 0.16) SD higher sex adjusted BW; the corresponding estimates for 1 and 4 year age- and sex-adjusted BMI were 0.07 (0.04, 0.10) and 0.18 (0.15, 0.21) respectively. For ALSPAC, the MV estimate for 15 year BMI was 0.32 (0.29, 0.36), and estimates were similar for 10 year BMI and FMI from 10–18 years. Adjustment for potential confounders had a negligible impact on the estimates, aside from a small attenuation on adjustment for paternal BMI for outcomes after birth. Results were similar when we refitted MV models one and two on larger samples without excluding individuals with missing paternal BMI data (**Supplementary information S21**). Additional adjustment for gestational age at birth or 20 genetic PCs had a negligible effect (**Supplementary information S22, S23**), and there was not a large difference when BMI or ponderal index at birth was substituted for BW in ALSPAC (**Supplementary information S24**).

For BW the MR estimate for the lassosum PRS for all three samples meta-analysed was 0.14 (0.05, 0.23), which was similar to the MV estimate; $P_{\text{difference (MV vs. MR)}} = 0.84$) (**Figure 1**). The corresponding lassosum MR estimates for 1 year BMI and 4 year BMI were -0.02 (-0.11, 0.07) and 0.01 (-0.08, 0.10) respectively, and there was moderate to strong evidence for an MR-MV difference ($P_{\text{difference}} = 0.10$ and $1.3\text{e-}3$ respectively). The MR estimates for 10 and 15 year BMI in ALSPAC (0.10 [-0.01, 0.21] and 0.13 [0.01, 0.24] respectively) were also smaller than the MV estimates ($P_{\text{difference}} = 1.4\text{e-}4$ and $1.0\text{e-}3$ respectively). Results for FMI were similar to those for BMI (**Supplementary information S25**). We did not observe strong evidence for non-linearity or interaction by sex for either the MV or MR models (results available from the authors), and results were similar when we (i) substituted BMI or ponderal index at birth for BW (**Supplementary information S25**), (ii) natural log transformed skewed variables (results available from the authors), (iii) removed cryptic relatedness from the sample (results available from the authors), and (iv) used linear mixed models to adjust for population structure (**Supplementary information S26**). In linear regression models (as opposed to two-stage least squares regression) there was strong to moderate evidence that the lassosum maternal non-transmitted allele BMI PRS was associated with offspring BW and adolescent adiposity (**Supplementary information S27, S28**).

Figure 1: Mean difference in offspring outcomes (SD) per 1SD increase in maternal BMI, from MR (lassosum) and confounder adjusted multivariable regression (MV) models

Confounder adjusted multivariable regression (MV) estimates are from model three (**Methods**). P : P -value for the null hypothesis that the effect equals zero, P_{dif} : P -value for the null hypothesis that MR effect equals the MV effect.

MR estimates for IVs with fewer SNPs

When we replaced the lassosum PRS with alternative IVs calculated from fewer SNPs, our MR estimates varied in a manner that was specific to the offspring outcome (**Figure 2**). For BW, including fewer SNPs in the IV did not result in large differences in the MR estimates, although the precision reduced markedly as we used fewer SNPs. For 1 and 4 year BMI, MR estimates increased as we used fewer SNPs, whereas for 10 year BMI they largely remained stable and for 15 year BMI they decreased. The patterns for FMI were similar to those for BMI (**Supplementary information S25**). For all outcomes apart from BW, including more SNPs in the IV resulted in stronger evidence that MR estimates differed from MV estimates (i.e. smaller P_{dif}).

MR estimates for SNPs with differing effect size distributions, between-SNP heterogeneity and MR Egger results

For the majority of outcomes (particularly in adolescence) there was moderate to strong statistical evidence that SNPs with smaller effect sizes gave larger (more positive) MR estimates, and this was not driven by weak instrument bias (**Supplementary information S29**). When using only large-effect (GWS) SNPs, in general there was not strong statistical evidence for between-SNP MR estimate heterogeneity (Cochran's Q), nor was there strong evidence that the MR-Egger intercept differed from zero (**Supplementary information S30**).

Figure 2: Mean difference in offspring outcomes (SD) per 1SD increase in maternal BMI, from MR models using different SNP sets and confounder adjusted multivariable regression (MV) models

Confounder adjusted multivariable regression (MV) estimates are from model three (**Methods**). **P**: **P**-value for the null hypothesis that the effect equals zero, **P_{diff}**: **P**-value for the null hypothesis that MR effect equals the MV effect, **FTO**: rs9939609 at the *FTO* locus, **Speliotes**, **Locke**, **Yengo**: GWS SNPs from the GWAS with the indicated first author, **Lassosum**: PRS calculated by the lassosum method. Colours denote outcomes

Parental phenotypic and PRS correlations

In ALSPAC there was strong evidence for correlation between maternal and paternal BMI (Pearson's r : 0.22, 95% CI: 0.16, 0.28, $P = 7.9\text{e-}14$), but no evidence for correlation between maternal non-transmitted allele and paternal lassosum BMI PRS (r : 0.02, 95% CI: -0.04, 0.07, $P = 0.55$). For comparison, a maternal lassosum BMI PRS that was calculated from both transmitted and non-transmitted alleles was slightly more strongly correlated with the paternal PRS (r : 0.04, 95% CI: -0.01, 0.10, $P = 0.14$).

Discussion

We applied a Mendelian randomization (MR) approach using PRS calculated from maternal non-transmitted alleles, to explore the causality of associations between maternal pre-/early-pregnancy BMI and offspring birth weight (BW) and child/adolescent adiposity. For the association between maternal BMI and offspring BW, our MR and confounder adjusted multivariable regression (MV) estimates were similar. In contrast, for offspring adiposity outcomes beyond 1 year of age the MR estimates were weaker than the MV estimates. These results markedly strengthen evidence that confounder adjusted observational associations between maternal BMI and offspring adolescent adiposity are subject to residual confounding. We found no strong evidence for a causal effect of maternal BMI on offspring adiposity beyond birth, although based on the present results we cannot rule out a small to moderate causal effect.

Our data build on two previous MR studies which investigated associations between maternal BMI and offspring child/adolescent adiposity (14, 23), and a methodological paper which presented a limited investigation of adiposity outcomes as an empirical example (25). Although the previous studies provided no strong evidence for a causal effect, they were limited by wide confidence intervals and/or potential biases (see **Strengths and Limitations** below). The present study overcame these limitations by using more powerful PRS and a maternal non-transmitted allele score approach. For the association between maternal BMI and offspring BW, our MR and MV estimates were highly concordant, in agreement with previous MR studies that supported a causal effect of greater maternal BMI on greater offspring BW (28, 34). We have previously shown that genetic confounding (i.e. confounding due to direct effects of maternal alleles inherited by the offspring) is unlikely to explain the association of maternal BMI with BW, but may potentially be important for the association with adolescent BMI (61). The present results are consistent with this, as well as with the balance of evidence from negative paternal exposure control studies (12-18) and within sibship analyses (19, 20), which suggests that familial confounding is an important explanation of the maternal BMI-offspring child/adolescent adiposity association. Studies that examined the effect of extreme maternal obesity using a pre- and post-bariatric surgery design (62-65) have small sample sizes and have not been entirely consistent, therefore do not provide strong evidence against this conclusion.

Although we found no strong evidence for a causal effect in late childhood/adolescence, we cannot rule out a small to moderate causal effect, due to the imprecision of our MR estimates. Indeed, the primary lassosum MR estimates were greater than zero for 15 year BMI and 14–18 year FMI. We do not interpret these as unbiased estimates for the causal effect of maternal BMI, because of the possibility of pleiotropic bias (see below). These results do suggest however that some maternal exposure(s) that are correlated with the maternal BMI PRS have a causal effect on offspring child/adolescent adiposity, although our analyses are unable to distinguish whether this is a pre or postnatal effect. Plausible mechanisms include intrauterine effects such as fetal overnutrition (1) and postnatal effects such as maternal influence on offspring eating behaviour (66), but other mechanisms have

been hypothesised, including periconceptional effects (such as altered oocyte structure or function (3)). In linear regression analyses we found moderate to strong evidence for associations between the maternal non-transmitted allele BMI PRS and offspring adolescent adiposity. Such associations are due to indirect effects of maternal genotype on offspring adiposity, mediated via the offspring's pre or postnatal environment (67) (also known as genetic nurture (68), dynastic effects (69) and social genetic effects (70)). These observed maternal genetic effects merit further investigation in other datasets, particularly as previous studies have not found evidence for parental genetic effects on BMI in childhood (71) or adulthood (68).

In ALSPAC and BiB White Europeans we observed associations between the maternal BMI PRS and variables that are often considered to be potential confounders in epidemiological studies, including parental occupation, education, age and maternal smoking. Such associations were largely absent for the IVs calculated from fewer SNPs, and in BiB South Asians (although it is notable that in BiB South Asians the genetic IVs explained less maternal BMI variance). These results invite careful consideration of which of the ever-increasing number of GWS associated BMI SNPs are likely (in combinations) to be the most valid instruments for MR studies, having taken account of all IV assumptions. Such considerations may also extend to other exposures, in line with concerns raised by other authors (32, 57, 72, 73).

Strengths and limitations

Our study has several key strengths. We studied two prospective birth cohorts with maternal and offspring genome-wide genotype data, maternal BMI measurements and offspring adiposity outcomes available, allowing us to conduct mother-offspring MR analyses. We used state-of-the-art methods to calculate a powerful PRS from around 80,000 SNPs. This yielded a substantial increase in statistical power over previous MR studies, which analysed similar ALSPAC datasets to ours, but employed either a single SNP in the *FTO* gene (14) or allele scores calculated from up to 97 SNPs (23, 25) (similar to the "Speliotes" and "Locke" IVs in the present analysis). Our primary lassosum PRS explained 3–7% of maternal BMI variance, compared to ~1.5% for the strongest IVs used previously (power calculations are given in **Supplementary information S31**).

Another strength over previous work is our use of maternal non-transmitted allele PRS, thereby avoiding the need to control for genetic inheritance by adjusting for offspring genotype. A previous methodological paper made limited use of this approach (25), but investigated a far smaller subset of adiposity outcomes than that which we have explored here. Controlling for offspring genotype may be suboptimal for two distinct reasons: (i) it may introduce collider bias if paternal genotype influences the offspring outcome independently of offspring genotype (i.e. if paternal genetic effects exist) (25, 74), and (ii) if the investigator adjusts for a *weighted* allele score, this may introduce bias by inadequately blocking the genetic inheritance path (24). Because these two biases may be in opposite directions, the net direction of any bias affecting the largest previous study (23) is uncertain. We acknowledge that our primary MR estimates may be affected by pleiotropic bias due to the large number of SNPs, many of which had small effect sizes, that we used to calculate the PRS. This possibility is also suggested by the associations that we observed between the lassosum BMI PRS and several potential confounders of the maternal BMI-offspring adiposity association. However, sensitivity analyses suggested that for most outcomes such bias is likely to be away from zero, which would weaken the apparent evidence for an MR-MV difference (**Supplementary information S29, S30**). Thus, our primary MR results are conservative, in that they may overstate the size of the causal effect (which we hypothesised

to be zero). The fact that for 10 and 15 year BMI, using more SNPs yielded increased precision and stronger evidence for an MR-MV difference (**Figure 2**), despite the potential pleiotropic bias away from zero, illustrates the benefit of our approach.

We also conducted extensive sensitivity analyses to explore other potential biases in our results. When we used a linear mixed model (LMM) to adjust for population structure the results were similar to our primary estimates. We did not remove cryptic relatedness for our primary analyses, in order to maximise the sample size and because the LMM controls for bias due to cryptic relatedness (54). However, results were similar when we removed cryptic relatedness at a level corresponding to first cousins. Finally, we found no strong evidence that maternal and paternal lassosum BMI PRS were correlated, suggesting that our results are not importantly biased due to assortative mating.

We acknowledge several other limitations of our study. First, although the results in BiB and ALSPAC were similar, replication in other cohorts with suitable data, and in particular with adolescent adiposity measures (which we could only examine in ALSPAC) would be valuable. A previous study meta-analysed data from ALSPAC and the Generation R cohort using 32 maternal SNPs (23), but we were unable to extend our approach to Generation R due to the unavailability of maternal genome-wide SNP data. We have only studied UK participants. However, the similarity of findings between White European and South Asian BiB participants, and between BiB (a cohort with high levels of deprivation born during the obesity epidemic) and ALSPAC (more affluent than the UK average) suggest that our findings may be generalisable to other populations. Second, BMI (especially self-reported BMI) is an imperfect proxy measure for adiposity. However, it has been shown previously in ALSPAC that self-reported pre-pregnancy BMI is strongly correlated with BMI measured in early pregnancy (23), and that any misreporting does not markedly differ by mean weight (75). There is also evidence that the correlation with directly measured adiposity is strong for child and adult BMI (76, 77) and moderate for neonatal weight (78); furthermore, our results were similar for DXA derived FMI. Third, we assumed that causal relationships between exposures and outcomes were linear. Although our data provided no evidence for non-linearity, a slight plateauing of the observational association between maternal BMI and offspring child/adolescent BMI at higher maternal BMI levels was previously observed in a large meta-analysis (6). MR estimates such as ours, which assume linearity, nevertheless approximate the population-averaged causal effect (which is the average effect resulting from a unit increase in the exposure for all individuals in the population, regardless of their initial exposure level) (79). However, given the shape of the observational association (6) it is plausible that our MR estimates overstate the true causal effect for mothers with overweight/obesity. Finally, the samples used for some of our analyses were smaller than the full samples at baseline due to missing data and loss to follow up. This was particularly the case for BiB MV models, because in BiB BMI data were only available for a minority of fathers. This raises the possibility that our results are affected by selection bias, which can occur even for studies such as ours which estimate genetic effects (80). However, the distributions of maternal BMI, BW and offspring sex were similar for the samples used for our analyses and the samples at baseline, and MV results were similar when we refitted models on larger samples without excluding individuals with missing paternal BMI data. It therefore seems unlikely that selection bias would be of sufficient magnitude to alter our conclusions.

Conclusion

We explored the causality of associations between maternal pre-/early-pregnancy BMI and offspring BW and child/adolescent adiposity, using an MR approach with PRS calculated

from maternal non-transmitted alleles. This approach yielded narrower confidence intervals compared with previous studies, and avoided sources of bias that may have affected previous work. We found no strong evidence for a causal effect of maternal BMI on offspring adiposity beyond birth, but strong evidence that confounder adjusted observational associations between maternal BMI and adolescent adiposity are affected by residual confounding. Although we cannot rule out a small or moderate causal effect on child/adolescent adiposity, the present study suggests that higher maternal pre-/early-pregnancy BMI is not a key driver of greater adiposity in the next generation. Thus, our results support interventions that target the whole population for reducing overweight and obesity, rather than a specific focus on women of reproductive age.

Author contributions

Conceptualization: TAB, MRJ, DAL

Methodology: DAL, VZ, PFO, MG, AL, SS, DME, MRJ

Software: TAB, RCR, VK, GCP

Formal analysis: TAB

Data curation: TAB, IT, AD, ACA, DM, TY, MCB

Writing- original draft preparation: TAB, MRJ, DAL, RCR, PFO

Writing- review and editing: TAB, MRJ, DAL, RCR, PFO, VZ, MG, AL, SS, IT, AD, DM, TY, VK, ACA, GCP, DME, MCB

This publication is the work of the authors and TAB, MRJ and DAL will serve as guarantors for the contents of this paper.

Conflict of interest

DAL has received support from numerous national and international government and charity funders and from Medtronic LTD and Roche Diagnostics for research unconnected with that presented in this study. GCP is an employee of 23andMe Inc and may hold stock or stock options. All other authors report no conflict of interest.

Acknowledgements

We thank Tom Palmer, Eleanor Sanderson and Stephen Burgess for helpful discussions, Mark Iles for provision of BiB ethnicity variables and Amanda Hill and David Hughes for support in delivery and management of the ALSPAC data. We gratefully acknowledge the GIANT (Genetic Investigation of ANthropometric Traits) Consortium for making GWAS summary statistics available. This publication is the work of the authors and may not reflect the views of those acknowledged here.

We are extremely grateful to all the families who took part in ALSPAC, the midwives for their help in recruiting them, and the whole ALSPAC team, which includes interviewers, computer and laboratory technicians, clerical workers, research scientists, volunteers, managers, receptionists and nurses. Born in Bradford is only possible because of the enthusiasm and commitment of the Children and Parents in BiB. We are grateful to all the participants, health professionals and researchers who have made Born in Bradford happen. We gratefully acknowledge the contribution of TPP and the TPP ResearchOne team in completing study participant matching to GP primary care records and in providing ongoing informatics support. This research has been conducted using the UK Biobank Resource under applications 10035, 13436 and 236 granting access to the corresponding UK Biobank genetic and phenotype data. We are extremely grateful to all the UK Biobank participants, investigators and team members.

Data availability

The ALSPAC study website (<http://www.bristol.ac.uk/alspac/researchers/our-data/>) contains details of all the data that are available through a fully searchable data dictionary and variable search tool. Scientists are encouraged and able to use BiB data. Data requests are made to the BiB executive using the form available from the study website <http://www.borninbradford.nhs.uk> (please click on 'Science and Research' to access the form). Guidance for researchers and collaborators, the study protocol and the data collection

schedule are all available via the website. All requests are carefully considered and accepted where possible. UK Biobank data are available from the UK Biobank (<http://biobank.ndph.ox.ac.uk/showcase/>).

Funding

The UK Medical Research Council and Wellcome (102215/2/13/2) and the University of Bristol provide core support for ALSPAC. Genotyping of the ALSPAC maternal samples was funded by the Wellcome Trust (WT088806) and the offspring samples were genotyped by Sample Logistics and Genotyping Facilities at the Wellcome Trust Sanger Institute and LabCorp (Laboratory Corporation of America) using support from 23andMe. A comprehensive list of grants funding is available on the ALSPAC website (<http://www.bristol.ac.uk/alspac/external/documents/grant-acknowledgements.pdf>). BiB receives core infrastructure funding from the Wellcome Trust (WT101597MA), a joint grant from the UK Medical Research Council (MRC) and UK Economic and Social Science Research Council (ESRC) (MR/N024397/1), the British Heart Foundation (CS/16/4/32482) and the National Institute for Health Research (NIHR) under its Collaboration for Applied Health Research and Care (CLAHRC) for Yorkshire and Humber. Further support for genome-wide data is from the UK Medical Research Council (G0600705) and the National Institute of Health Research (NF-SI-0611-10196). The work presented here was also supported by the US National Institute of Health (R01 DK10324), the European Research Council under the European Union's Seventh Framework Programme (FP7/2007-2013)/ERC grant agreement (669545) and the British Heart Foundation (AA/18/7/34219). TAB is supported by the Medical Research Council (MRC) (UK) (MR/K501281/1), TAB and DME are supported by the NHMRC (Australia) (GNT1183074 and GNT1157714), DAL, TAB and RCR work in/are affiliated with a unit that is supported by the UK Medical Research Council (MC_UU_00011/1 & MC_UU_00011/6) and DAL is a NIHR Senior Investigator (NF-0616-10102). DAL and TAB are supported by the British Heart Foundation Accelerator Award at the University of Bristol (AA/18/7/34219). VK is funded by the European Union's Horizon 2020 research and innovation program under the Marie Skłodowska-Curie grant (721567). MRJ is funded by EU-H2020 LifeCycle Action (733206), EU-H2020 EDCMET (825762), EU-H2020 EUCAN Connect (824989), EU H2020-MSCA-ITN-2016 CAPICE Marie Skłodowska-Curie grant (721567) and the MRC (UK) (MRC/BBSRC and MR/S03658X/1 [JPI HDHL]). RCR is a de Pass Vice Chancellor's Research Fellow at the University of Bristol. AL and MRJ are supported by the MRC (UK) (MR/M013138/1) and the European Union Horizon 2020 programme (633595). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

References

1. Lawlor DA. The Society for Social Medicine John Pemberton Lecture 2011. Developmental overnutrition—an old hypothesis with new importance? *Int J Epidemiol.* 2013;42(1):7-29.
2. Godfrey KM, Reynolds RM, Prescott SL, *et al.* Influence of maternal obesity on the long-term health of offspring. *Lancet Diabetes Endo.* 2017;5(1):53-64.
3. Fleming TP, Watkins AJ, Velazquez MA, *et al.* Origins of lifetime health around the time of conception: causes and consequences. *The Lancet.* 2018;391(10132):1842-52.
4. Larqué E, Labayen I, Flodmark C-E, *et al.* From conception to infancy—early risk factors for childhood obesity. *Nature Reviews Endocrinology.* 2019:1.
5. Voerman E, Santos S, Golab BP, *et al.* Maternal body mass index, gestational weight gain, and the risk of overweight and obesity across childhood: An individual participant data meta-analysis. *PLoS Med.* 2019;16(2):e1002744.

6. Heslehurst N, Vieira R, Akhter Z, *et al.* The association between maternal body mass index and child obesity: A systematic review and meta-analysis. *PLoS Med.* 2019;16(6):e1002817.
7. Reynolds RM, Allan KM, Raja EA, *et al.* Maternal obesity during pregnancy and premature mortality from cardiovascular event in adult offspring: follow-up of 1 323 275 person years. *BMJ.* 2013;347:f4539.
8. Lahti-Pulkkinen M, Bhattacharya S, Wild SH, *et al.* Consequences of being overweight or obese during pregnancy on diabetes in the offspring: a record linkage study in Aberdeen, Scotland. *Diabetologia.* 2019:1-8.
9. Friedman JE. Developmental programming of obesity and diabetes in mouse, monkey, and man in 2018: where are we headed? *Diabetes.* 2018;67(11):2137-51.
10. Poston L. Maternal obesity, gestational weight gain and diet as determinants of offspring long term health. *Best Practice & Research Clinical Endocrinology & Metabolism.* 2012;26(5):627-39.
11. Lawlor DA, Tilling K, Davey Smith G. Triangulation in aetiological epidemiology. *Int J Epidemiol.* 2016;45(6):1866-86.
12. Fleten C, Nystad W, Stigum H, *et al.* Parent-offspring body mass index associations in the Norwegian Mother and Child Cohort Study: a family-based approach to studying the role of the intrauterine environment in childhood adiposity. *Am J Epidemiol.* 2012;176(2):83-92.
13. Patro B, Liber A, Zalewski B, Poston L, Szajewska H, Koletzko B. Maternal and paternal body mass index and offspring obesity: a systematic review. *Ann Nutr Metab.* 2013;63(1-2):32-41.
14. Lawlor DA, Timpson NJ, Harbord RM, *et al.* Exploring the developmental overnutrition hypothesis using parental-offspring associations and FTO as an instrumental variable. *PLoS Med.* 2008;5(3):e33.
15. Gaillard R, Steegers EA, Duijts L, *et al.* Childhood cardiometabolic outcomes of maternal obesity during pregnancy: The Generation R Study. *Hypertension.* 2014;63(4):683-91.
16. Santos Ferreira DL, Williams DM, Kangas AJ, *et al.* Association of pre-pregnancy body mass index with offspring metabolic profile: Analyses of 3 European prospective birth cohorts. *PLoS Med.* 2017;14(8):e1002376.
17. Zalbazar N, Najman J, McIntyre HD, Mamun A. Parental pre-pregnancy BMI influences on offspring BMI and waist circumference at 21 years. *Aust N Z J Public Health.* 2016;40(6):572-8.
18. Sørensen TI, Ajslev TA, Ängquist L, Morgen CS, Ciuchi IG, Smith GD. Comparison of associations of maternal peri-pregnancy and paternal anthropometrics with child anthropometrics from birth through age 7 y assessed in the Danish National Birth Cohort. *Am J Clin Nutr.* 2016;104(2):389-96.
19. Branum AM, Parker JD, Keim SA, Schempf AH. Prepregnancy body mass index and gestational weight gain in relation to child body mass index among siblings. *Am J Epidemiol.* 2011:kwr250.
20. Lawlor DA, Lichtenstein P, Långström N. Association of Maternal Diabetes Mellitus in Pregnancy With Offspring Adiposity Into Early Adulthood: Clinical Perspective. *Circulation.* 2011;123(3):258-65.
21. Davey Smith G, Ebrahim S. 'Mendelian randomization': can genetic epidemiology contribute to understanding environmental determinants of disease? *Int J Epidemiol.* 2003;32(1):1-22.
22. Lawlor DA, Harbord RM, Sterne JA, Timpson N, Davey Smith G. Mendelian randomization: using genes as instruments for making causal inferences in epidemiology. *Stat Med.* 2008;27(8):1133-63.
23. Richmond RC, Timpson NJ, Felix JF, *et al.* Using genetic variation to explore the causal effect of maternal pregnancy adiposity on future offspring adiposity: a Mendelian randomisation study. *PLoS Med.* 2017;14(1):e1002221.
24. Wang G, Warrington N, Evans DM. *Personal communication.* 2020.

25. Lawlor D, Richmond R, Warrington N, *et al.* Using Mendelian randomization to determine causal effects of maternal pregnancy (intrauterine) exposures on offspring outcomes: Sources of bias and methods for assessing them. *Wellcome Open Research*. 2017;2.
26. Ebbeling CB, Pawlak DB, Ludwig DS. Childhood obesity: public-health crisis, common sense cure. *The Lancet*. 2002;360(9331):473-82.
27. Zhang G, Bacelis J, Lengyel C, *et al.* Assessing the causal relationship of maternal height on birth size and gestational age at birth: a mendelian randomization analysis. *PLoS Med*. 2015;12(8):e1001865.
28. Chen J, Bacelis J, Sole-Navais P, *et al.* Dissecting maternal and fetal genetic effects underlying the associations between maternal phenotypes, birth outcomes, and adult phenotypes: A mendelian-randomization and haplotype-based genetic score analysis in 10,734 mother–infant pairs. *PLoS Med*. 2020;17(8):e1003305.
29. Brion M-JA, Shakhbazov K, Visscher PM. Calculating statistical power in Mendelian randomization studies. *Int J Epidemiol*. 2012;42(5):1497-501.
30. Yengo L, Sidorenko J, Kemper KE, *et al.* Meta-analysis of genome-wide association studies for height and body mass index in ~700000 individuals of European ancestry. *Hum Mol Genet*. 2018;27(20):3641-9.
31. Khera AV, Chaffin M, Wade KH, *et al.* Polygenic prediction of weight and obesity trajectories from birth to adulthood. *Cell*. 2019;177(3):587-96. e9.
32. Evans DM, Brion MJA, Paternoster L, *et al.* Mining the human phenome using allelic scores that index biological intermediates. *PLoS Genetics*. 2013;9(10):e1003919.
33. Evans DM, Visscher PM, Wray NR. Harnessing the information contained within genome-wide association studies to improve individual prediction of complex disease risk. *Hum Mol Genet*. 2009;18(18):3525-31.
34. Tyrrell J, Richmond RC, Palmer TM, *et al.* Genetic evidence for causal relationships between maternal obesity-related traits and birth weight. *JAMA*. 2016;315(11):1129-40.
35. Fraser A, Macdonald-Wallis C, Tilling K, *et al.* Cohort profile: the Avon Longitudinal Study of Parents and Children: ALSPAC mothers cohort. *Int J Epidemiol*. 2012;42(1):97-110.
36. Boyd A, Golding J, Macleod J, *et al.* Cohort profile: the ‘children of the 90s’—the index offspring of the Avon Longitudinal Study of Parents and Children. *Int J Epidemiol*. 2013;42(1):111-27.
37. Wright J, Small N, Raynor P, *et al.* Cohort profile: the Born in Bradford multi-ethnic family cohort study. *Int J Epidemiol*. 2012;42(4):978-91.
38. West J, Santorelli G, Whincup PH, *et al.* Association of maternal exposures with adiposity at age 4/5 years in white British and Pakistani children: findings from the Born in Bradford study. *Diabetologia*. 2018;61(1):242-52.
39. McCarthy S, Das S, Kretzschmar W, *et al.* A reference panel of 64,976 haplotypes for genotype imputation. *Nat Genet*. 2016;48(10):1279.
40. The 1000 Genomes Project Consortium. A global reference for human genetic variation. *Nature*. 2015;526(7571):68.
41. Huang J, Howie B, McCarthy S, *et al.* Improved imputation of low-frequency and rare variants using the UK10K haplotype reference panel. *Nature Communications*. 2015;6:8111.
42. O’Connell J, Gurdasani D, Delaneau O, *et al.* A General Approach for Haplotype Phasing across the Full Spectrum of Relatedness. *PLOS Genetics*. 2014;10(4):e1004234.
43. Choi SW, O’Reilly PF. PRSice-2: Polygenic Risk Score software for biobank-scale data. *GigaScience*. 2019;8(7):giz082.
44. Vilhjálmsson BJ, Yang J, Finucane HK, *et al.* Modeling linkage disequilibrium increases accuracy of polygenic risk scores. *The American Journal of Human Genetics*. 2015;97(4):576-92.
45. Mak TSH, Porsch RM, Choi SW, Zhou X, Sham PC. Polygenic scores via penalized regression on summary statistics. *Genet Epidemiol*. 2017;41(6):469-80.
46. Mak TSH, Porsch RM. *lassosum: LASSO with summary statistics and a reference panel. R package version 0.4.4*. 2019.

47. Loh P-R, Kichaev G, Gazal S, Schoech AP, Price AL. Mixed-model association for biobank-scale datasets. *Nat Genet.* 2018;1.
48. Bycroft C, Freeman C, Petkova D, *et al.* Genome-wide genetic data on~ 500,000 UK Biobank participants. *BioRxiv.* 2017:166298.
49. Sudlow C, Gallacher J, Allen N, *et al.* UK biobank: an open access resource for identifying the causes of a wide range of complex diseases of middle and old age. *PLoS Med.* 2015;12(3):e1001779.
50. Locke AE, Kahali B, Berndt SI, *et al.* Genetic studies of body mass index yield new insights for obesity biology. *Nature.* 2015;518(7538):197-206.
51. Speliotes EK, Willer CJ, Berndt SI, *et al.* Association analyses of 249,796 individuals reveal 18 new loci associated with body mass index. *Nat Genet.* 2010;42(11):937.
52. Burgess S, Small DS, Thompson SG. A review of instrumental variable estimators for Mendelian randomization. *Stat Methods Med Res.* 2017;26(5):2333-55.
53. Thomas DC, Lawlor DA, Thompson JR. Re: Estimation of bias in nongenetic observational studies using "Mendelian triangulation" by Bautista *et al.* *Ann Epidemiol.* 2007;7(17):511-3.
54. Yang J, Zaitlen NA, Goddard ME, Visscher PM, Price AL. Advantages and pitfalls in the application of mixed-model association methods. *Nat Genet.* 2014;46(2):100.
55. Yang J, Lee SH, Goddard ME, Visscher PM. GCTA: a tool for genome-wide complex trait analysis. *Am J Hum Genet.* 2011;88(1):76-82.
56. Yang Q, Sanderson E, Tilling K, Borges MC, Lawlor DA. Exploring and mitigating potential bias when genetic instrumental variables are associated with multiple non-exposure traits in Mendelian randomization. *medRxiv.* 2019:19009605.
57. Hemani G, Bowden J, Davey Smith G. Evaluating the potential role of pleiotropy in Mendelian randomization studies. *Hum Mol Genet.* 2018;27(R2):R195-R208.
58. Bowden J, Davey Smith G, Burgess S. Mendelian randomization with invalid instruments: effect estimation and bias detection through Egger regression. *Int J Epidemiol.* 2015;44(2):512-25.
59. StataCorp. *Stata 13 Base Reference Manual.* College Station, TX: Stata Press; 2013.
60. R Core Team. *R: A language and environment for statistical computing.* Vienna, Austria: R Foundation for Statistical Computing; 2018.
61. Bond TA, Karhunen V, Wielscher M, *et al.* Exploring the role of genetic confounding in the association between maternal and offspring body mass index: evidence from three birth cohorts. *Int J Epidemiol.* 2019.
62. Smith J, Cianflone K, Biron S, *et al.* Effects of maternal surgical weight loss in mothers on intergenerational transmission of obesity. *The Journal of Clinical Endocrinology & Metabolism.* 2009;94(11):4275-83.
63. Kral JG, Biron S, Simard S, *et al.* Large maternal weight loss from obesity surgery prevents transmission of obesity to children who were followed for 2 to 18 years. *Pediatrics.* 2006;118(6):e1644-e9.
64. Willmer M, Berglind D, Sørensen TI, Näslund E, Tynelius P, Rasmussen F. Surgically induced interpregnancy weight loss and prevalence of overweight and obesity in offspring. *PLoS One.* 2013;8(12):e82247.
65. Barisione M, Carlini F, Gradaschi R, Camerini G, Adami GF. Body weight at developmental age in siblings born to mothers before and after surgically induced weight loss. *Surg Obes Relat Dis.* 2012;8(4):387-91.
66. Patel C, Karasouli E, Shuttlewood E, Meyer C. Food Parenting Practices among Parents with Overweight and Obesity: A Systematic Review. *Nutrients.* 2018;10(12):1966.
67. Lynch M, Walsh B. *Genetics and analysis of quantitative traits.* Sunderland, MA: Sinauer; 1998.
68. Kong A, Thorleifsson G, Frigge ML, *et al.* The nature of nurture: Effects of parental genotypes. *Science.* 2018;359(6374):424-8.
69. Brumpton B, Sanderson E, Heilbron K, *et al.* Avoiding dynastic, assortative mating, and population stratification biases in Mendelian randomization through within-family analyses. *Nature Communications.* 2020;11(1):3519.

70. Baud A, Mulligan MK, Casale FP, *et al.* Genetic variation in the social environment contributes to health and disease. *PLoS Genetics*. 2017;13(1):e1006498.
71. Schnurr TM, Morgen CS, Borisevich D, *et al.* The influence of transmitted and non-transmitted parental BMI-associated alleles on the risk of overweight in childhood. *Sci Rep*. 2020;10(1):1-10.
72. Richardson TG, Harrison S, Hemani G, Smith GD. An atlas of polygenic risk score associations to highlight putative causal relationships across the human phenotype. *Elife*. 2019;8:e43657.
73. Burgess S, Butterworth AS, Thompson JR. Beyond Mendelian randomization: how to interpret evidence of shared genetic predictors. *J Clin Epidemiol*. 2016;69:208-16.
74. Tubbs JD, Zhang YD, Sham PC. Intermediate confounding in trio relationships: The importance of complete data in effect size estimation. *Genet Epidemiol*. 2020;44(4):395-9.
75. Sharp GC, Lawlor DA, Richmond RC, *et al.* Maternal pre-pregnancy BMI and gestational weight gain, offspring DNA methylation and later offspring adiposity: findings from the Avon Longitudinal Study of Parents and Children. *Int J Epidemiol*. 2015;44(4):1288-304.
76. Lawlor DA, Benfield L, Logue J, *et al.* Association between general and central adiposity in childhood, and change in these, with cardiovascular risk factors in adolescence: prospective cohort study. *BMJ*. 2010;341:c6224.
77. Fox C, Massaro J, Hoffmann U, *et al.* Abdominal visceral and subcutaneous adipose tissue compartments: association with metabolic risk factors in the Framingham Heart Study. *Circulation*. 2007;116(1):39-48.
78. Eriksson B, Löf M, Forsum E. Body composition in full-term healthy infants measured with air displacement plethysmography at 1 and 12 weeks of age. *Acta Paediatr*. 2010;99(4):563-8.
79. Burgess S, Davies NM, Thompson SG, EPIC-InterAct Consortium. Instrumental variable analysis with a nonlinear exposure-outcome relationship. *Epidemiology*. 2014;25(6):877-85.
80. Munafò MR, Tilling K, Taylor AE, Evans DM, Davey Smith G. Collider scope: when selection bias can substantially influence observed associations. *Int J Epidemiol*. 2017.