Novel prognostic determinants of COVID-19-related mortality: a pilot study on severely-ill patients in Russia

Rubina K.^{4*}, Shmakova A.¹, Shabanov A.³, Andreev Yu.³, Borovkova N.³, Kulabukhov V.³, Evseev A.³, Popugaev K.³, Petrikov S.³, Semina E.^{1,2}

¹Laboratory of Molecular Endocrinology, Institute of Experimental Cardiology, Federal State Budgetary Organization National Cardiology Research Center Ministry of Health of the Russian Federation, Moscow, Russia

²Department of Biochemistry and Molecular Medicine, Faculty of Medicine, Lomonosov Moscow State University, Moscow, Russia

³Department of Intensive Care, Sklifosovsky Research Institute of Emergency Medicine of the Moscow Healthcare Department, Moscow, Russia

⁴Laboratory of Morphogenesis and Tissue Reparation, Faculty of Medicine, Lomonosov Moscow State University, Moscow, Russia

***Corresponding author: Rubina Kseniya kseniiarubina1971@gmail.com**

Abstract

COVID-19 pandemic has posed a severe healthcare challenge calling for an integrated approach in determining the clues for early non-invasive diagnostics of the potentially severe cases and efficient patient stratification.

Here we analyze the clinical, laboratory and CT scan characteristics associated with high risk of COVID-19-related death outcome in the cohort of severely-ill patients in Russia. The data obtained reveal that elevated dead lymphocyte counts, decreased early apoptotic lymphocytes, decreased CD14+/HLA-Dr+ monocytes, increased expression of JNK in PBMCs, elevated IL-17 and decreased PAI-1 serum levels are associated with a high risk of COVID-19-related mortality thus suggesting them to be new prognostic factors. This set of determinants could be used as early predictors of potentially severe course of COVID-19 for trials of prevention or timely treatment.

Keywords: COVID-19, SARS-CoV-2, patient prognosis, clinical outcomes, COVID-19-related death.

Funding: The reported study was funded by RFBR according to the research project № 20-24-60029.

The ongoing wave of COVID-19 infections resulting from an outbreak, which initially started in Wuhan in China, has reached alarming proportions across the globe, as reported by the World Health Organization (WHO) ¹. The unprecedented challenge with SARS-CoV-2 infection calls for an integrated approach, encompassing the development of effective vaccines to prevent the spread of the disease, search for new drugs aimed at lowering the viral load and preventing the complications, and quest for the clues to early diagnostics of a severe course of the disease.

The clinical spectrum of COVID-19 manifestations appears to be remarkably wide, ranging from asymptomatic infection, mild upper respiratory tract illness to severe viral pneumonia with respiratory failure, systemic inflammation, disseminated intravascular coagulation, multiorgan

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

failure and even death²⁻⁴. Older age, diabetes and cardiovascular comorbidities are among the highest risk factors influencing the disease severity and the clinical outcome of SARS-CoV-2 infection^{4,5}. However, severe COVID-19 has been registered not only in elderly patients or patients suffering from comorbidities, but also in apparently healthy young people^{6,7}. The development of new prognostic algorithms for differential diagnostics to predict and reduce the severity and complications of COVID-19 remains essential.

The data obtained in the present study reveal novel prognostic factors associated with high risk of COVID-19-related death overtime in patients with severe COVID-19. These include: the elevated dead lymphocyte counts, decreased early apoptosis of lymphocytes, decreased CD14+, HLA-Dr+ (human leukocyte antigen D related) monocytes, increased JNK (c-Jun N-terminal kinase) expression in PBMCs (peripheral blood mononuclear cell), increased IL-17 and decreased PAI-1 (plasminogen activator inhibitor-1) serum level. The prognostic value of each determinant can potentially impact decision-making on the level of healthcare and require more extensive follow-up strategies.

To the best our knowledge this is the first integrated study based on the Russian population aimed to reveal the original prognostic COVID-19-related death factors.

Material and methods

Study design and clinical workflow

The present study was a single-center retrospective study carried out on a cohort of 52 symptomatic adult patients with the confirmed SARS-CoV-2 infection, admitted to intensive care unit (ICU) of N.V. Sklifosovsky Research Institute for Emergency Medicine, Moscow from April to July 2020. Demographic, clinical, and laboratory data were recorded at admission. The study was approved by the ethics committee of Sklifosovsky Research Institute of Emergency Medicine of the Moscow Healthcare Department (protocol #5-120, issued on 01.04.2020). The need for written patient consent was waived by the ethics committee of the hospital because laboratory investigations were conducted according to the local standard. The local ethical committee approved the retrospective study.

Nasopharyngeal swabs were collected from all patients, followed by RT-PCR assay to confirm the SARS-CoV-2 infection. Philips Ingenuity Core 128 CT Scanner (Netherlands) was used for all examinations. In all cases, semi-quantitative CT scoring (severity ranging from 0 to 4) was evaluated according to the protocol recommended by Moscow Healthcare Department adapted from the International Protocols and enriched with local experience⁸.

Routine blood tests were carried out for all patients and the following parameters were evaluated: white blood cells (WBC), lymphocytes, monocytes, neutrophil, red blood cell (RBC), platelet counts, levels of hemoglobin, serum total protein, albumins, globulins, bilirubin, urea, creatinine, alanine aminotransferase (ALT), aspartate aminotransferase (AST), lactate dehydrogenase (LDH), creatine kinase (CK), prothrombin, fibrinogen, D-dimer levels, activated partial thromboplastin time (APTT) and the international normalized ratio (INR).

Biochemical parameters analysis in blood serum and PBMCs

In addition to blood tests routinely and periodically performed in hospital, we assessed the levels of plasminogen (PLG), plasminogen activator inhibitor 1 (PAI-1), IL-1 α , IL-17, transforming growth factor beta (TGF β), tumor necrosis factor alpha (TNF α), and adiponectin (ADP) in blood serum. Blood serum was collected and immediately frozen. Commercially available ELISA kits (SEA605Hu for ADP, SEA071Hu for IL-1 α , SEA063Hu for IL-17, SEA532Hu for PAI-1, SEB236Hu for PLG, SEA124Hu for TGF β , and SEA133Hu for TNF α (Cloud-Clone Corp., Wuhan, Hubei) were used to evaluate the corresponding analytes according

to the manufacturers' protocols. The 3,3', 5,5'-tetramethylbenzidine (TMB) was used as a chromogen and the maximum absorption wavelength of 450 nm was detected using a microplate reader Multiscan Ascent (Thermo Fisher). The experiments were performed in duplicate.

Whole blood specimens were collected into EDTA-containing tubes either directly labelled with antibodies or subjected to PBMC purification using Lympholyte-H (Cedarlane) density gradient centrifugation. Antibodies against the following surface markers were applied for whole blood sample labelling: CD95 (mouse monoclonal anti-hu CD95-FITC 1F-362-T100, EXBIO), CD14 (mouse monoclonal anti-hu CD14-PE Conjugated Antibody A07764, Beckman Coulter) and HLA-Dr (HLA-Dr-FITC Conjugated Antibody, IM1638U, Beckman Coulter). Isotypic non-immune IgG (IgG1 Mouse-FITC Isotype Control, A07795, Beckman Coulter) was used as a control. For lymphocyte death analysis, PBMCs were stained with anti-CD45, Annexin V, 7AAD (mouse monoclonal anti-hu CD45-FITC Conjugated Antibody A07782 Beckman Coulter; Annexin V-FITC Kit-AAD Kit, IM3614, Beckman Coulter; 7-AAD Viability Dye A07704, Beckman Coulter). Labelled cells were analysed by flow cytometry with CYTOMICS FC500 flow cytometer (Beckman Coulter) and CXP software. Lymphocyte and monocyte populations were gated on FSC vs. SSC plot. The following parameters were assessed: the absolute number per 1 μ l and the percentage of dead lymphocytes (CD45+/7AAD+), the percentage of early (Annexin V+/7AAD) and late apoptotic (Annexin V+/7AAD+) lymphocytes as well as the percentage of CD95+ lymphocytes and HLA-Dr+ monocytes (CD14+/HLA-Dr+). The gating strategy is shown in **Supplementary figure 1**.

PBMC lysates were prepared using protease inhibitor Gordox®, and the apoptotic marker expression, such as Akt, Bad, Bcl-2, Caspase-8, Caspase-9, JNK and p53 was evaluated using Milliplex®MAP (Early Apoptosis 7-plex Magnetic Bead Kit, cat#48-660MAG) assay and Luminex® (USA). The values for each parameter are presented as Median Fluorescence Intensity (MFI). The experiments were performed in duplicate.

Outcome and exposures

The outcome used in the survival analysis in the current study was COVID-19-related death, defined on the basis of International Classification of Diseases 10 code U07.1 (COVID-19, virus identified), recorded as either a primary underlying or secondary cause of death.

With respect to potential risk factors assessed as exposures, we obtained data for sex, age, body mass index (BMI), tobacco smoking status, chest CT severity score (the worst CT imaging data per patient presented in Table 1). Patients were grouped by age as: younger than 40 years, 40-49 years, 50-59 years, 60-69 years, 70-79 years, and 80 years or older. BMI was grouped as: less than 30 (no evidence of obesity), 30-34.9 (obese class I), 35-39.9 (obese class II), and 40 $\text{kg}\cdot\text{m}^{-2}$ or higher (obese class III). Tobacco smoking status was identified as smoker, non-smoker (not a current smoker), or missing data. Chest CT score was grouped as 1-2 or 3-4.

Table 1. Cohort description with COVID-19 deaths by patient characteristics. Data are presented as n (%). BMI = body mass index.

	All patients (n = 52)	Deaths (n = 11)
Sex		
Male	28 (54%)	5 (45%)
Female	24 (46%)	6 (55%)
Age, years		
18-39	3 (6%)	0 (0%)
40-49	8 (15%)	1 (9%)

50-59	12 (23%)	1 (9%)
60-69	11 (21%)	3 (27%)
70-79	7 (13%)	1 (9%)
80+	11 (21%)	5 (45%)
BMI, kg·m⁻²		
<30	31 (60%)	7 (64%)
30-34.9	16 (31%)	4 (36%)
35-39.9	2 (4%)	0 (0%)
40+	3 (6%)	0 (0%)
Tobacco smoking status		
Non-smoker	41 (79%)	6 (55%)
Smoker	2 (4%)	0 (0%)
Missing data	9 (17%)	5 (45%)
Chest CT severity score (the worst CT imaging data per patient)		
0-2	32 (62%)	2 (18%)
3-4	20 (38%)	9 (82%)

Routine blood test parameters were organized into three groups relying on the reference range: lower, within, or higher the reference range. The accepted reference values of the blood tests parameters are presented in Supplementary **Table 1**. In case the data of several measurements were available, the mean of all measurements was calculated and used for subsequent analysis. Biochemical serum and leukocyte parameters categorization are presented in **Supplementary table 2**.

Statistical analysis

Categorical variables are described as numbers and percentages. Continuous variables are reported as the median (interquartile range). To identify risk factors associated with COVID-19-related death, a survival analysis of the outcome was performed: time was defined as the time from hospital admission to the end of a follow-up (hospital discharge or death) and the event defined as death or hospital discharge after recovery. The Kaplan-Meier method was used to analyze cumulative survival curves and their difference between the groups categorized by type of exposure compared by log-rank tests. Exposures with difference between the groups $P < 0.05$ by log-rank tests were considered significant. We created univariable Cox proportional hazards survival analysis models for the evaluated parameters of each patient to test for their association with survival. We also created multivariable Cox proportional hazards survival analysis models to identify independent risk factors. Complete cases were considered. The variables included in the final models were based on clinical and scientific knowledge and the results of univariable analyses. Considering the small size of the cohort and to avoid overfitting in the model, we created three separate final multivariable Cox models, with variables grouped as clinical characteristics and hematological results; clinical biochemical routine blood test results; laboratory blood findings. The data were analyzed in R studio v.1.2.5042 with the survival and survminer packages.

Results

A total of 52 patients, aged 26-88, with confirmed SARS-CoV-2 infection were included in the study cohort (**Table 1**). The median age was 64 (54-76) years; the median BMI was 28.7 (25.8-31.1) kg·m⁻²; 28 patients (54%) were men. 1 patient was admitted to ICU with CT score 0; 13 patients with CT score 1; 18 patients with CT score 2; 13 patients with CT score 3, and 4 patients with CT score 4. Overall, 11 (21%) patients died.

The survival analysis was performed with time defined as a period from hospital admission to the end of the follow-up (hospital discharge or death) and event defined as death or hospital discharge after recovery (**Figure 1**). The Kaplan-Meier survival function analysis revealed that 20-day survival probability was 91.0% (95% CI 82.9-99.9%), 40-day survival probability was 64.3% (95% CI 46.1-89.5), 60-day survival probability was 48.2% (95% CI 28.7-81.0%).

Figure 1. Kaplan-Meier survival curve with associated 95% confidence intervals (shaded area) for patients admitted with COVID-19 infection to ICU of N.V. Sklifosovsky Research Institute for Emergency Medicine, Moscow from April to July 2020. The fraction of survival is expressed on the y-axis, while time (days) of the observation period is expressed on the x-axis. Vertical traits indicate censored data (hospital discharge).

We performed the Kaplan-Meier analysis to compare the survival over time between different groups of patients for each variable. The Kaplan-Meier survival plots for the clinical and routine blood test factors with statistically significant results are presented in **Figure 2**. We found, that the risk of COVID-19-related death over time was significantly increased in patients with CT score 3-4 at admission ($P = 0.042$), patients on pulmonary ventilation ($P = 0.00047$), patients with WBC count more than $11.8 \cdot 10^3/\mu\text{l}$ ($P = 0.001$), neutrophil count more than $8.2 \cdot 10^3/\mu\text{l}$ ($P = 0.00027$), lymphocyte count less than $1.1 \cdot 10^3/\mu\text{l}$ ($P = 0.018$), increased serum levels of globulins more than 30 g/l ($P = 0.046$), urea more than 7.2 mM ($P = 0.01$), CK more than 200 U/l ($P = 0.023$), prothrombin less than 78% ($P = 0.023$), INR more than 1.17 ($P = 0.02$) and D-dimer more than 2 mg/l ($P = 0.0078$). Gender, age, BMI, tobacco smoking status and other parameters of routine blood tests were not associated with statistically significant differences as revealed by Kaplan-Meier analysis (**Supplementary figure 2**).

Figure 2. Kaplan-Meier survival curves for patients grouped by clinical and routine blood test parameters. The fraction of survival is expressed on the y-axis, while time (days) of the observation period is expressed on the x-axis. Vertical traits indicate censored data (hospital discharge). P values of log-rank tests are indicated for each graph. Reference group is shown in red.

The Kaplan-Meier survival plots for the biochemical test factors being statistically significant are presented in **Figure 3**. We found, that the risk of COVID-19-related death over time was significantly increased in patients with more than 100 dead lymphocytes in 1 μ l ($P = 0.0026$), less than 5% of early apoptotic lymphocytes ($P = 0.0081$), less than 85% of CD14+/HLA-Dr+ monocytes ($P = 0.029$), increased expression of JNK in PBMCs (>200 MFI, $P = 0.034$), increased serum levels of IL-17 (>7 pg/ml, $P = 0.037$) and decreased PAI-1 serum levels (<40 pg/ml, $P = 0.00073$). Percentage of dead and late apoptotic lymphocytes, percentage of CD95+ lymphocytes, expression of Akt, Bad, Bcl-2, Caspase-8, Caspase-9, p53 and serum levels of PLG, IL-1 α , TGF β , TNF α , adiponectin were not associated with statistically significant differences in survival as revealed by Kaplan-Meier analysis (**Supplementary figure 3**).

Figure 3. Kaplan-Meier survival curves for patients grouped by biochemical test parameters. The fraction of survival is expressed on the y-axis, while time (days) of the observation period is expressed on the x-axis. Vertical traits indicate censored data (hospital discharge). P values of log-rank tests are indicated for each graph. Reference group is shown in red.

To identify the independent risk factors, we created multivariate Cox proportional hazards survival analysis models separately for clinical characteristics and hematology results; for clinical biochemical routine blood test results and for laboratory blood findings (**Table 2**). On multivariate Cox proportional hazards survival analysis of clinical and routine hematological test parameters, ventilation (HR 55.66 (2.2-1388.4), $P = 0.014$) and platelet count (HR 1.01 (1.002-1.027), $P = 0.017$) were found to be the independent prognostic factor for COVID-19 related death. On multivariate analysis of clinical biochemical routine blood test results, the serum levels of globulins (HR 1.32 (1.03-1.70), $P = 0.026$), CK (HR 1.02 (1.005-1.04), $P = 0.013$) and D-dimer (3.41 (1.11-10.50), $P = 0.033$) were found to be the independent prognostic factor for COVID-19 related death. On multivariate analysis of laboratory blood findings, the serum level of IL-17 (HR 1.07 (1.001-1.14), $P = 0.046$) and PAI-1 (HR 0.82 (0.69-0.98), $P = 0.026$) were found to be the independent prognostic factor for COVID-19 related death. The results of univariable Cox

proportional hazards survival analysis for each parameter are presented in **Supplementary table 3**.

Table 2. Multivariable Cox proportional hazards survival analysis models for 3 groups of prognostic factors. HR – hazard ratio, CI – confidence interval, Inf – infinite. * – $P < 0.05$.

Parameter	HR (95% CI)	P value
Clinical characteristics and hematology results		
CT	0-2: reference 3-4: 0.52 (0.03-7.85)	0.634
Ventilation	No: reference Yes: 55.66 (2.23-1388.40)	0.014 *
WBC, $10^3/\mu\text{l}$	0.58 (0.11-2.96)	0.509
Neutrophil count, $10^3/\mu\text{l}$	2.68 (0.57-12.48)	0.210
Lymphocyte count, $10^3/\mu\text{l}$	0.29 (0.01-6.93)	0.447
Monocyte count, $10^3/\mu\text{l}$	0.15 (0.00-1564.4)	0.692
Platelet count, $10^3/\mu\text{l}$	1.01 (1.002-1.027)	0.017 *
Clinical biochemical routine blood test results		
Globulins, g/l	1.32 (1.03-1.70)	0.026 *
Urea, mM	0.88 (0.62-1.20)	0.467
Creatinine, μM	1.02 (0.99-1.10)	0.244
ALT, U/l	0.99 (0.93-1.00)	0.638
AST, U/l	1.01 (0.98-1.00)	0.572
LDH, U/l	0.99 (0.985-1.00)	0.705
CK, U/l	1.02 (1.005-1.04)	0.013 *
D-dimer, mg/l	3.41 (1.11-10.50)	0.033 *
Laboratory blood findings		
Dead lymphocyte count	1.02 (1.00-1.04)	0.054
Early lymphocyte apoptosis	1.09 (0.77-1.55)	0.624
CD14+, HLA-Dr+ monocytes, %	0.96 (0.91-1.01)	0.108
IL-17, pg/ml	1.07 (1.001-1.14)	0.046 *
IL-1 α , pg/ml	1.09 (0.90-1.31)	0.385
PLG, $\mu\text{g/ml}$	0.66 (0.32-1.32)	0.238
PAI-1, pg/ml	0.82 (0.69-0.98)	0.026 *
TNF α , pg/ml	0.90 (0.68-1.19)	0.464
TGF β , pg/ml	1.00 (1.00-2.02)	0.069
ADP, $\mu\text{g/ml}$	0.56 (0.30-1.08)	0.083

Discussion

With the rapid spread of COVID-19 around the world since its outbreak in early 2020, it has stirred up an international concern and still remains so. Chest CT imaging has been recognized to play a pivotal role in monitoring COVID-19 disease progression and predicting adverse prognosis. We found that CT score 3-4 at admission and pulmonary ventilation significantly increase the risk of COVID-19-related death over time (**Figure 2**) that is in accordance with the previously published data ^{2,9-14}.

A compelling body of evidence indicate that clinical and routine blood test factors such as an increased WBC count ^{2,15-22}, elevated neutrophil count ^{2,16-23}, low lymphocyte count ^{2,4,9,17,18,20-22}, elevated urea ^{2,15,19,21-24}, upregulated CK ^{2,4,9,18,22} and increased level of serum globulins ²³ or

immunoglobulins²⁵ are statistically significant and strongly correlate with the adverse outcome in COVID-19 patients. Our data also confirm the recent papers (**Figure 2**). Previously, neutrophil-lymphocyte ratio (NLR) was shown to be a prognostic marker of systemic inflammation in various pathological conditions^{26,27}. Moreover, NLR was found to have a high prognostic value in COVID-19 patients with the increase in neutrophils and the decrease in the total number of lymphocytes in the peripheral blood. Generally, the COVID-19 patients with increased NLR have a poor prognosis and a high risk of death²⁸. Earlier, it has been demonstrated that COVID-19 patients with severe respiratory failure exhibit low HLA-Dr expression on CD14+ monocytes relative to that of mild COVID-19 patients or healthy volunteers^{29,30}. Xu and co-authors hypothesized that monocytes in severe COVID-19 displayed a phenotype similar to immunosuppressive monocytic myeloid-derived suppressor cells (CD14+/HLA-DR-/lo)³⁰. Interestingly, the expression of HLA-Dr on monocytes from healthy donors decreased following the cultivation in media with plasma from patients with COVID-19 who have immune dysregulation³¹. Our results also indicate that the risk of COVID-19-related death over time is significantly increased in patients with elevated neutrophil and decreased lymphocyte counts. Moreover, we have defined that the decrease in CD14+/HLA-Dr+ monocyte fraction are significant predictors of disease severity in COVID-19 patients (**Figure 3**).

The biological role of lymphocyte apoptosis in infectious disease is ambiguous; whether it operates as an intrinsic protective mechanism or mediates a severe course of the disease still remains unclear. Lymphocytes can undergo extensive and ostensibly out-of-control apoptosis during severe infections, potentially contributing to immunosuppression³². In the majority of viral infections, lymphocytes are subjected to apoptosis, which is considered to be a defense mechanism preventing the spread of infection without a local inflammatory reaction³³. Although lymphocyte count can be initially augmented at the onset of COVID-19, its subsequent decline in the course of infection may result in lymphocytopenia and immunodeficiency associated with an increased COVID-19 severity³⁴. The frequency of peripheral blood lymphocyte subpopulations, including T-CD4+, T-CD8+, NK, B cells, and monocytes, and their apoptosis pattern was assessed in Iranian COVID-19 patients. The authors reported a marked increase in apoptosis of mononuclear cells from COVID-19 patients as compared to control, however no association with mortality has been addressed in this study³⁵. Our data indicate that elevated dead lymphocyte counts are associated with COVID-19 mortality (**Figure 3**), however, early lymphocyte apoptosis is rather a protective mechanism reflecting the intensity of the adaptive cell immune response to an infectious agent, whereas a decrease in the early apoptosis of lymphocytes is an important predictor of COVID-19-related mortality (**Figure 3**).

Elevated plasma D-dimer is a well-established prognostic factor for adverse outcome in respiratory diseases and COVID-19^{28,36,37}. Our results go along with these previously published data and ascertain that the COVID-19-related death over time is significantly increased in patients with augmented D-dimer (**Figure 2**).

A potentially practical opportunity was proposed by the sustained elevated production of IL-6^{2,4,9,15,16,18,19,21-23,38}, IL-1, IL-2R, IL-8, IL-10^{2,9} and some other blood circulating factors, whose early evaluation anticipated disease progression and duration of hospital stay of COVID-19 patients³⁹. Such interleukins as IL-1 β , IL-6, IL-17A, TNF- α , and monocyte chemoattractant peptide (MCP)-1 were identified in patients with severe COVID-19⁴⁰. Huang et al.⁴¹ demonstrated the upregulated levels of IL-1 β , IL-6, IL-8, IL-17, interferon γ (IFN- γ), TNF- α in patients with COVID-19 as compared to healthy donors. The same study revealed increased plasma concentrations of MCP-1 and TNF- α in patients with COVID-19 admitted to the ICU. Similarly, Chen and co-authors⁶ reported the correlation between high concentrations of plasma IL-6 and TNF- α and the severe course of COVID-19. In contrast, in the study by Kang and co-authors these cytokines, including IL-1 β , IL-12p40, and IL-17 were undetectable in patients with severe COVID-19⁴². Several clinical trials investigating interleukin inhibitors (IL-1 inhibitor anakinra, and IL-6 inhibitors tocilizumab or sarilumab) revealed that IL-1 inhibition, but not IL-6

inhibition, was associated with a significant reduction of mortality in patients admitted to hospital with COVID-19, ARDS (acute respiratory distress syndrome), and hyperinflammation⁴³.

IL-17 is a promising target in COVID-19, since it operates upstream of both, IL-1 and IL-6⁴⁴. Although the pathogenic role of IL-17 has been suggested previously, this cytokine was not detected as a mortality risk factor in COVID-19 patients^{40,44}. MERS-CoV infection was reported to be associated with an increased serum level of IL-17, although its correlation with mortality was not demonstrated⁴⁵. IL-17 polymorphisms are associated with susceptibility and poor prognosis in acute respiratory distress syndrome⁴⁶. Meanwhile, in obese patients with confirmed COVID-19, IL-17 boosts neutrophil recruitment via induction of inflammatory cytokine production by respiratory epithelium, smooth muscle cells and fibroblasts resulting in respiratory dysfunction⁴⁰.

In the present study IL-1 α , TNF- α and TGF- β were not associated with the statistically significant difference in patient survival (**Supplementary figure 3**); however, IL-17 and PAI-1 were determined as independent prognostic factors for COVID-19 related death (**Figure 3**), underlying their potential significance as predictors of mortality for patients with COVID-19.

PAI-1 is a well-known inhibitor of the plasminogen activators (tissue plasminogen activator – tPA, and urokinase – uPA) and hence the overall fibrinolysis⁴⁷. Of note, PAI-1 levels are increased in hypertension, obesity, diabetes, cardiovascular diseases, and old age, which is consistent with the scenario of COVID-19 disease^{48,49}. Plasmin, and other proteases, may cleave the furin site in the S protein of SARS-CoV-2, thus increasing the virus virulence. Moreover, the dysregulation in fibrinolytic homeostasis in COVID-19 patients with excessive fibrin degradation may result in an unbalanced fibrinolysis inducing multiple organ hemorrhage^{50,51}. Hyperfibrinolysis resulting from excessive plasmin activity is also associated with elevated D-dimer in severely-ill COVID-19 patients.

While Umemura et al. reported an unchanged PAI-1 level in COVID-19 patients⁵², several other groups detected the increased levels of PAI-1, tPA and uPA in COVID-19 patients as compared to healthy controls^{53,54}. Zou and co-authors demonstrated that high levels of tPA and PAI-1 were associated with worse respiratory status in patients hospitalized with COVID-19⁴⁸. It has been found that overproduced PAI-1 binds to specific receptors on macrophages and upregulates the production of proinflammatory cytokines and chemokines, which further activate the innate immune cells within the infected lungs resulting in lung damage. In turn, high hypoxic environment additionally stimulates PAI-1 production creating a vicious cycle of cytokine storm generated during SARS virus pathogenesis⁵¹. In contrast, our data indicate that COVID-19-related mortality is associated with decreased PAI-1, establishing that PAI-1 being the constituent component of the fibrinolytic system plays an important role in COVID-19 pathogenesis. The explanation for the reduced PAI-1 content in the critically-ill patients may reside in compensation for the hyperactivation of plasminogen activator system, reflecting the exhaustion of plasmin inhibitors (**Figure 3**).

The C-Jun NH2-terminal kinase (JNK1/JNK2) pathway activation ultimately leads to various cellular effects such as inflammatory response, cell proliferation, survival or even apoptosis. JNK signaling pathway was suggested to play a prominent role in immune response to viral infection due to activation of several interleukins (IL-2, IL-4) and IFN- γ ⁵⁵. To uncover the mechanisms of JNK-mediated inflammation in COVID-19, Shirato and Kizaki carried out an *in vitro* study and analyzed the effects of SARS-CoV-2 spike protein S1 subunit on murine and human macrophages. Exposure to S1 subunit activated pro-inflammatory mediators (nuclear factor- κ B (NF- κ B) and JNK) signaling pathways via the activation of a toll-like receptor 4 (TLR4) on macrophage surface. Pro-inflammatory cytokine (TNF- α , IL-6, IL-1 β , and nitric oxide) production induced by S1 was abrogated by specific inhibitors of NF- κ B and JNK pathways⁵⁶.

In line with the suggested strategy for inhibition of JNK pathway in COVID-19 critically-ill patients as a promising therapeutic approach⁵⁷, here we present the first clinical study revealing the increased JNK expression in PBMCs to be a significant prognostic marker of COVID-19-related mortality (**Figure 3**). It is tempting to speculate that the elevated JNK, which has been

demonstrated to induce TLR4-mediated apoptosis in various types of cells^{56,58}, and the decreased level of lymphocyte early apoptosis in severe COVID-19 patients revealed in the present study are intertwined. However, this assumption needs further investigation.

In conclusion, this study unveiled an integrated set of immunological features and new prognostic markers of patients with severe COVID-19 that are significant predictors of the related mortality. The obtained results suggest a high risk of COVID-19-related death overtime in patients with elevated dead lymphocyte counts, decreased early apoptosis of lymphocytes, reduced fraction of CD14+/HLA-Dr+ monocytes, increased JNK expression in PBMCs, elevated IL-17 and decreased PAI-1 serum levels. Gender, age, BMI, tobacco smoking status and several parameters of routine blood tests (**Supplementary figure 2**) were not statistically significant.

Our analysis revealed novel determinants of a high COVID-19-related death risk overtime that can provide for patient stratification at admission to ICU or point to the potential requirement of a more extensive follow up treatment strategies.

Limitations

Our study has some limitations. First, the cohort might not be fully representative as the study was a single center one and only the patients admitted to ICU were included. Therefore, a relatively high mortality rate was detected, which cannot reflect the actual COVID-19 mortality in a population. Second, our results might be limited by the sample size. Third, the outcomes were evaluated at the end of the follow-up period instead of a fixed time period during the course of the disease. Fourth, not all laboratory tests were done in all patients, however, the data were missing completely at random. Hence, the interpretation of our results might be limited.

However, we believe that our study uncovers some important tendencies that should be further considered with the relation to COVID-19 mortality. As far as we are aware, this is the first retrospective cohort study among patients with COVID-19 in Russia that evaluated the biochemical parameters in blood serum different from those that are routinely used in healthcare practice. The obtained results may provide a rationale for testing these new parameters as novel markers of COVID-19 severity and point to potential targets for therapeutical intervention in quest to improve the outcomes.

Contribution

Rubina K. and Semina E. conceived and designed the study. Petrikov S. managed intensive care unit facility, resources and patients' admission. Shabanov A., Andreev Yu., Borovkova N., Kulabukhov V. collected and managed the patient's data and performed the biochemical tests. Shmakova A. performed the statistical analysis and described the results. Rubina K. and Semina E. contributed to the writing, review and editing of the manuscript. Evseev A., Popugaev K., Petrikov S. made a substantial contribution to the design of the study.

Supplementary tables and figures

Supplementary table 1. The accepted reference values of the routine blood tests parameters.

Parameter	Reference values	Units
RBC count	3.7-5.4	10 ¹² /l
Hemoglobin	120-140	g/l
WBC count	3.8-11.8	10 ³ /μl
Neutrophil count	1.9-8.2	10 ³ /μl

Neutrophils %	42.7-76.8	%
Lymphocyte count	1.1-3.8	10 ³ /μl
Lymphocytes %	16-45.9	%
Monocyte count	0.2-0.9	10 ³ /μl
Monocytes %	4.3-11.0	%
Platelet count	179-408	10 ³ /μl
Total protein	66-83	g/l
Albumins	35-52	g/l
Globulins	20-30	g/l
Total bilirubin	<19	μM
Urea	<7.2	mM
Creatinine	<96	μM
ALT	<34	U/l
AST	<31	U/l
LDH	<247	U/l
CK	<200	U/l
APTT	25.1-36.5	S
Prothrombin	78-142	%
INR	0.64-1.17	-
Fibrinogen	2.2-4.98	g/l
D-dimer	<0.5	mg/l

Supplementary table 2. Categorization of biochemical parameters. Reference group is highlighted in bold. MFI – Median Fluorescence Intensity.

Parameter	Groups			Units
Dead lymphocytes count	<100	>100		1/μl
Dead lymphocytes %	<1.5	>1.5		%
Early apoptotic lymphocytes	<5%	5-10%	>10%	%
Late apoptotic lymphocytes	<0.1%	>0.1%		%
CD95+ lymphocytes	<37	37-46	>46	%
CD14+, HLA-Dr+ monocytes	<85%	>85%		%
Akt	<107	>107		MFI
Bad	<101	>101		MFI
Bcl-2	<20.25	>20.25		MFI
Caspase-8	<81.25	>81.25		MFI
Caspase-9	<200	>200		MFI
JNK	<200	>200		MFI
p53	<72.25	>72.25		MFI
IL-17	<7	>7		pg/ml
PLG	<20	>20		μg/ml
IL-1α	<0.3	>0.3		pg/ml
PAI-1	<40	>40		pg/ml
TGFβ	<30	>30		pg/ml
TNFα	<11	>11		pg/ml
Adiponectin	<5	5-10	>10	μg/ml

382 **Supplementary table 3. Univariable Cox proportional hazards survival analysis**
 383 **models.** HR – hazard ratio, CI – confidence interval, Inf – infinite. * – $P < 0.05$, ** – $P < 0.01$.

Parameter	HR (95% CI)	P value
Demographic and clinical characteristics		
Gender	F: <i>reference</i> M: 1.13 (0.34-3.78)	0.838
Age, years	18-39: 0.00 (0.00-Inf)	0.999
	40-49: 1.18 (0.07-19.52)	0.909
	50-59: <i>reference</i>	
	60-69: 2.24 (0.23-21.81)	0.488
	70-79: 0.60 (0.03-10.57)	0.728
	80+: 2.53 (0.28-22.59)	0.406
BMI, kg·m ⁻²	<30: <i>reference</i>	
	30-34.9: 0.95 (0.28- 3.24)	0.930
	35-39.9: 0.00 (0.00-Inf)	0.999
	40+: 0.00 (0.00-Inf)	0.999
Smoking	Non-smoker: <i>reference</i> Smoker: 0.00 (0.00-Inf)	0.999
CT	0-2: <i>reference</i> 3-4: 4.40 (0.94-20.63)	0.06
Ventilation	No: <i>reference</i> Yes: 9.86 (2.10-46.23)	0.004 **
Hematology results		
RBC count, 10 ¹² /l	0.75 (0.3-1.86)	0.534
Hemoglobin, g/l	0.99 (0.95-1.02)	0.421
WBC count, 10 ³ /μl	1.16 (1.03-1.30)	0.014 *
Neutrophil count, 10 ³ /μl	1.17 (1.03-1.33)	0.013 *
Neutrophil, %	1.11 (1.01-1.22)	0.025 *
Lymphocyte count, 10 ³ /μl	0.22 (0.04-1.12)	0.068
Lymphocyte, %	0.86 (0.75-0.99)	0.036 *
Monocyte count, 10 ³ /μl	0.37 (0.01-19.28)	0.619
Monocyte, %	0.74 (0.54-1.01)	0.056
Platelet count, 10 ³ /μl	1.003 (0.996-1.009)	0.423
Clinical biochemical routine blood test results		
Total serum protein, g/l	1.01 (0.94-1.09)	0.807
Albumins, g/l	0.91 (0.82-1.01)	0.071
Globulins, g/l	1.15 (1.02-1.30)	0.021 *
Total bilirubin	0.99 (0.96-1.02)	0.534
Urea, mM	1.09 (1.04-1.15)	0.001 **
Creatinine, μM	1.005 (1.002-1.009)	0.006 **
ALT, U/l	1.002 (1.0004-1.003)	0.013 *

AST, U/l	1.001 (1.0002-1.001)	0.010 *
LDH, U/l	1.001 (1.0002-1.001)	0.005 **
CK, U/l	1.003 (1.001-1.005)	0.002 **
APTT, s	1.01 (0.998-1.022)	0.093
Prothrombin, %	0.99 (0.96-1.01)	0.391
INR	1.17 (0.66-2.07)	0.591
Fibrinogen, g/l	1.26 (0.95-1.67)	0.105
D-dimer, mg/l	1.26 (1.02-1.55)	0.032 *
Laboratory blood findings		
Dead lymphocytes, count/ μ l	1.009 (1.003-1.013)	0.001 **
Dead lymphocytes, %	2.45 (1.21-4.96)	0.013 *
Early lymphocyte apoptosis	0.91 (0.77-1.06)	0.216
Late lymphocyte apoptosis	3.08 (0.01-882.66)	0.696
CD95+ lymphocytes, %	0.98 (0.94-1.02)	0.370
CD14+/HLA-Dr+ monocytes, %	0.97 (0.95-0.99)	0.005 **
AKT	1.00 (1.00-1.00)	0.593
BAD	1.01 (1.00-1.01)	0.063
BCL2	0.90 (0.79-1.02)	0.107
Caspase 8	1.03 (0.99-1.07)	0.128
Caspase 9	1.00 (1.00-1.01)	0.258
JNK	1.00 (1.00-1.00)	0.510
P53	0.97 (0.94-1.02)	0.230
IL-17, pg/ml	1.03 (1.01-1.05)	0.001 **
IL-1 α , pg/ml	1.07 (1.02-1.13)	0.005 **
PLG, μ g/ml	1.01 (0.87-1.17)	0.916
PAI-1, pg/ml	0.96 (0.91-1.01)	0.104
TNF α , pg/ml	1.07 (0.95-1.20)	0.263
TGF β , pg/ml	1.00 (1.00-1.00)	0.159
ADP, μ g/ml	0.88 (0.73-1.06)	0.263

384

385

Supplementary figure 1. The flow cytometry analysis. A – Representative flow cytometry graphs showing the gating strategy used to assess lymphocyte apoptosis. Forward scatter (FSC) vs. side scatter (SSC) plot applied to gate the lymphocyte cell population and remove the debris. Annexin-FITC-channel vs. 7AAD-channel plot used to gate early (Annexin V+/7AAD-) and late (Annexin V+/7AAD+) apoptotic cells in lymphocyte population. B – Representative flow cytometry graphs showing the gating strategy used to evaluate the percentage of CD14+/HLA-Dr+ cells in monocyte cell population from the whole blood. Isotypic non-immune IgG (IgG1 Mouse-FITC Isotype Control, A07795, Beckman Coulter) was applied as a control.

Supplementary figure 2. Kaplan-Meier survival curves for demographic, other clinical and routine blood test parameters. The fraction of survival is expressed on the y-axis, while time (days) of the observation period is expressed on the x-axis. Vertical traits indicate censored data (hospital discharge). *P* values of log-rank tests are indicated for each graph. Reference group is shown in red.

Supplementary figure 3. Kaplan-Meier survival curves for other biochemical test parameters. The fraction of survival is expressed on the y-axis, while time (days) of the observation period is expressed on the x-axis. Vertical traits indicate censored data (hospital discharge). *P* values of log-rank tests are indicated for each graph. Reference group is shown in red.

References

1. WHO. Coronavirus Disease (COVID-19) Situation Reports. <https://www.who.int/emergencies/diseases/novel-coronavirus-2019/situation-reports> (2021).
2. Wang, Y. *et al.* Clinical course and outcomes of 344 intensive care patients with COVID-19. *American Journal of Respiratory and Critical Care Medicine* vol. 201 1430–1434 (2020).
3. Coleman, J. J., Manavi, K., Marson, E. J., Botkai, A. H. & Sapey, E. COVID-19: to be or not to be; that is the diagnostic question. *Postgraduate medical journal* vol. 96 392–398 (2020).
4. Zhou, F. *et al.* Clinical course and risk factors for mortality of adult inpatients with

- COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet* **395**, 1054–1062 (2020).
5. Williamson, E. J. *et al.* Factors associated with COVID-19-related death using OpenSAFELY. *Nature* **584**, 430–436 (2020).
6. Chen, G. *et al.* Clinical and immunological features of severe and moderate coronavirus disease 2019. *J. Clin. Invest.* **130**, 2620–2629 (2020).
7. Zhang, S. Y. *et al.* Severe COVID-19 in the young and healthy: monogenic inborn errors of immunity? *Nature Reviews Immunology* vol. 20 455–456 (2020).
8. Морозов, С. П. *et al.* Лучевая диагностика коронавирусной болезни (Covid-19): организация, методология, интерпретация результатов. Версия 2 (17.04.2020). vol. № ЦДТ – 2020 – II (Государственное бюджетное учреждение здравоохранения города Москвы «Научно-практический клинический центр диагностики и телемедицинских технологий Департамента здравоохранения города Москвы», 2020).
9. Zhang, X.-B. *et al.* Risk factors for mortality of coronavirus disease-2019 (COVID-19) patients in two centers of Hubei province, China: A retrospective analysis. *PLoS One* **16**, e0246030 (2021).
10. Hu, Y., Zhan, C., Chen, C., Ai, T. & Xia, L. Chest CT findings related to mortality of patients with COVID-19: A retrospective case-series study. *PLoS One* **15**, (2020).
11. Tabatabaei, S. M. H., Rahimi, H., Moghaddas, F. & Rajebi, H. Predictive value of CT in the short-term mortality of Coronavirus Disease 2019 (COVID-19) pneumonia in nonelderly patients: A case-control study. *Eur. J. Radiol.* **132**, 109298 (2020).
12. Lei, Q. *et al.* Correlation between CT findings and outcomes in 46 patients with coronavirus disease 2019. *Sci. Rep.* **11**, 1103 (2021).
13. Li, K. *et al.* Radiographic findings and other predictors in adults with Covid-19. *medRxiv* 2020.03.23.20041673 (2020) doi:10.1101/2020.03.23.20041673.
14. Patel, U. *et al.* Age-Adjusted Risk Factors Associated with Mortality and Mechanical Ventilation Utilization Amongst COVID-19 Hospitalizations—a Systematic Review and Meta-Analysis. *SN Compr. Clin. Med.* **2**, 1740–1749 (2020).
15. Ruan, Q., Yang, K., Wang, W., Jiang, L. & Song, J. Clinical predictors of mortality due to COVID-19 based on an analysis of data of 150 patients from Wuhan, China. *Intensive Care Medicine* vol. 46 846–848 (2020).
16. Li, M. *et al.* Analysis of the Risk Factors for Mortality in Adult COVID-19 Patients in Wuhan: A Multicenter Study. *Front. Med.* **7**, 545 (2020).
17. Izcovich, A. *et al.* Prognostic factors for severity and mortality in patients infected with COVID-19: A systematic review. *PLoS One* **15**, e0241955 (2020).
18. Huang, Y. *et al.* A cohort study of 676 patients indicates D-dimer is a critical risk factor for the mortality of COVID-19. *PLoS One* **15**, e0242045 (2020).
19. Martins-Filho, P. R., Tavares, C. S. S. & Santos, V. S. Factors associated with mortality in patients with COVID-19. A quantitative evidence synthesis of clinical and laboratory data. *European Journal of Internal Medicine* vol. 76 97–99 (2020).
20. Ioannou, G. N. *et al.* Risk Factors for Hospitalization, Mechanical Ventilation, or Death Among 10 131 US Veterans With SARS-CoV-2 Infection. *JAMA Netw. open* **3**, e2022310 (2020).
21. Mesas, A. E. *et al.* Predictors of in-hospital COVID-19 mortality: A comprehensive systematic review and meta-analysis exploring differences by age, sex and health conditions. *PLoS One* **15**, e0241742 (2020).
22. Wang, L. *et al.* Coronavirus disease 2019 in elderly patients: Characteristics and prognostic factors based on 4-week follow-up. *J. Infect.* **80**, 639–645 (2020).
23. Wu, C. *et al.* Risk Factors Associated with Acute Respiratory Distress Syndrome and Death in Patients with Coronavirus Disease 2019 Pneumonia in Wuhan, China. *JAMA Intern. Med.* **180**, 934–943 (2020).

- 472 24. Cheng, A. *et al.* Diagnostic performance of initial blood urea nitrogen combined with D-
473 dimer levels for predicting in-hospital mortality in COVID-19 patients. *Int. J. Antimicrob.*
474 *Agents* **56**, 106110 (2020).
- 475 25. Zhao, Y. *et al.* Abnormal immunity of non-survivors with COVID-19: Predictors for
476 mortality. *Infect. Dis. Poverty* **9**, 108 (2020).
- 477 26. Huang, H. *et al.* Preoperative neutrophil–lymphocyte and platelet–lymphocyte ratios as
478 independent predictors of T stages in hilar cholangiocarcinoma. *Cancer Manag. Res.*
479 **Volume 11**, 5157–5162 (2019).
- 480 27. Celikbilek, M. *et al.* Neutrophil-Lymphocyte Ratio as a Predictor of Disease Severity in
481 Ulcerative Colitis. *J. Clin. Lab. Anal.* **27**, 72–76 (2013).
- 482 28. Ye, W. *et al.* Dynamic changes of D-dimer and neutrophil-lymphocyte count ratio as
483 prognostic biomarkers in COVID-19. *Respir. Res.* **21**, 169 (2020).
- 484 29. Giamarellos-Bourboulis, E. J. *et al.* Complex Immune Dysregulation in COVID-19
485 Patients with Severe Respiratory Failure. *Cell Host Microbe* **27**, 992-1000.e3 (2020).
- 486 30. Xu, G. *et al.* The differential immune responses to COVID-19 in peripheral and lung
487 revealed by single-cell RNA sequencing. *Cell Discov.* **6**, 1–14 (2020).
- 488 31. Dempsey, L. A. Immune dysregulation. *Nat. Immunol.* **21**, 596 (2020).
- 489 32. Parrino, J., Hotchkiss, R. S. & Bray, M. Prevention of immune cell apoptosis as potential
490 therapeutic strategy for severe infections. *Emerging Infectious Diseases* vol. 13 191–198
491 (2007).
- 492 33. Somova, L. M., Besednova, N. N. & Plekhova, N. G. Apoptosis and infectious diseases.
493 *Russ. J. Infect. Immun.* **4**, 303–318 (2015).
- 494 34. Fathi, N. & Rezaei, N. Lymphopenia in COVID-19: Therapeutic opportunities. *Cell Biol.*
495 *Int.* **44**, 1792–1797 (2020).
- 496 35. Taghiloo, S. *et al.* Apoptosis and immunophenotyping of peripheral blood lymphocytes in
497 Iranian COVID-19 patients: Clinical and laboratory characteristics. *J. Med. Virol.* **93**,
498 1589–1598 (2021).
- 499 36. Ge, Y. L. *et al.* Elevated plasma D-dimer in adult community-acquired pneumonia
500 patients is associated with an increased inflammatory reaction and lower survival. *Clin.*
501 *Lab.* **65**, 131–138 (2019).
- 502 37. Fruchter, O., Yigla, M. & Kramer, M. R. D-dimer as a prognostic biomarker for mortality
503 in chronic obstructive pulmonary disease exacerbation. *Am. J. Med. Sci.* **349**, 29–35
504 (2015).
- 505 38. Mikami, T. *et al.* Risk Factors for Mortality in Patients with COVID-19 in New York
506 City. *J. Gen. Intern. Med.* **36**, 17–26 (2021).
- 507 39. Laing, A. G. *et al.* A dynamic COVID-19 immune signature includes associations with
508 poor prognosis. *Nat. Med.* **26**, 1623–1635 (2020).
- 509 40. Leija-Martínez, J. J. *et al.* IL-17A and TNF- α as potential biomarkers for acute respiratory
510 distress syndrome and mortality in patients with obesity and COVID-19. *Med. Hypotheses*
511 **144**, 109935 (2020).
- 512 41. Huang, C. *et al.* Clinical features of patients infected with 2019 novel coronavirus in
513 Wuhan, China. *Lancet* **395**, 497–506 (2020).
- 514 42. Kang, S. *et al.* IL-6 trans-signaling induces plasminogen activator inhibitor-1 from
515 vascular endothelial cells in cytokine release syndrome. *Proc. Natl. Acad. Sci. U. S. A.*
516 **117**, 22351–22356 (2020).
- 517 43. Cron, R. Q. COVID-19 cytokine storm: targeting the appropriate cytokine. *Lancet*
518 *Rheumatol.* **0**, (2021).
- 519 44. Pacha, O., Sallman, M. A. & Evans, S. E. COVID-19: a case for inhibiting IL-17? *Nature*
520 *Reviews Immunology* vol. 20 345–346 (2020).
- 521 45. Mahallawi, W. H., Khabour, O. F., Zhang, Q., Makhdoum, H. M. & Suliman, B. A.
522 MERS-CoV infection in humans is associated with a pro-inflammatory Th1 and Th17
523 cytokine profile. *Cytokine* **104**, 8–13 (2018).

46. Xie, M., Cheng, B., Ding, Y., Wang, C. & Chen, J. Correlations of IL-17 and NF- κ B gene polymorphisms with susceptibility and prognosis in acute respiratory distress syndrome in a Chinese population. *Biosci. Rep.* **39**, (2019).
47. Eden, G., Archinti, M., Furlan, F., Murphy, R. & Degryse, B. The urokinase receptor interactome. *Curr. Pharm. Des.* **17**, 1874–1889 (2011).
48. Zuo, Y. *et al.* Plasma tissue plasminogen activator and plasminogen activator inhibitor-1 in hospitalized COVID-19 patients. vol. 11 1580.
49. Cesari, M., Pahor, M. & Incalzi, R. A. REVIEW: Plasminogen Activator Inhibitor-1 (PAI-1): A Key Factor Linking Fibrinolysis and Age-Related Subclinical and Clinical Conditions. *Cardiovasc. Ther.* **28**, e72–e91 (2010).
50. Ji, H. L., Zhao, R., Matalon, S. & Matthay, M. A. Elevated plasmin(Ogen) as a common risk factor for COVID-19 susceptibility. *Physiol. Rev.* **100**, 1065–1075 (2020).
51. Matsuyama, T., Kubli, S. P., Yoshinaga, S. K., Pfeffer, K. & Mak, T. W. An aberrant STAT pathway is central to COVID-19. *Cell Death and Differentiation* vol. 27 3209–3225 (2020).
52. Umemura, Y. *et al.* Hematological Phenotype of COVID-19-Induced Coagulopathy: Far from Typical Sepsis-Induced Coagulopathy. *J. Clin. Med.* **9**, 2875 (2020).
53. Cabrera-Garcia, D. *et al.* High levels of plasminogen activator inhibitor-1, tissue plasminogen activator and fibrinogen in patients with severe COVID-19. *medRxiv* 2020.12.29.20248869 (2021) doi:10.1101/2020.12.29.20248869.
54. Cugno, M. *et al.* Complement activation and endothelial perturbation parallel COVID-19 severity and activity. *J. Autoimmun.* **116**, 102560 (2021).
55. Hemmat, N. *et al.* The roles of signaling pathways in SARS-CoV-2 infection; lessons learned from SARS-CoV and MERS-CoV. *Archives of Virology* vol. 166 675–696 (2021).
56. Shirato, K. & Kizaki, T. SARS-CoV-2 spike protein S1 subunit induces pro-inflammatory responses via toll-like receptor 4 signaling in murine and human macrophages. *Heliyon* **7**, e06187 (2021).
57. Solimani, F., Meier, K. & Ghoreschi, K. Janus kinase signaling as risk factor and therapeutic target for severe SARS-CoV-2 infection. *Eur. J. Immunol.* eji.202149173 (2021) doi:10.1002/eji.202149173.
58. Zhang, Y. & Bliska, J. B. Role of toll-like receptor signaling in the apoptotic response of macrophages to Yersinia infection. *Infect. Immun.* **71**, 1513–1519 (2003).