1 Original Article

2 **Title:** Industry and workplace characteristics associated with the use of a COVID-19
3 contact tracing app in Japan: a nation-wide employee survey

4

5 Tomohiro Ishimaru¹, Koki Ibayashi¹, Masako Nagata², Ayako Hino³, Seiichiro Tateishi⁴,
6 Mayumi Tsuji⁵, Akira Ogami⁶, Shinya Matsuda⁷, Yoshihisa Fujino¹ for the
7 CORoNaWork Project

8

9 **Author contributions:** Y.F. was the chairperson of the study group. T.I. conceived
10 the research questions. All the authors designed the research protocol and developed
11 the questionnaire. T.I. conducted the statistical analysis with Y.F.. T.I. drafted the
12 initial manuscript. All the authors revised and approved the final manuscript.

13

14 Authors' affiliations:

15 ¹Department of Environmental Epidemiology, Institute of Industrial Ecological
16 Sciences, University of Occupational and Environmental Health, Japan, Kitakyushu,
17 Japan

18 ²Department of Occupational Health Practice and Management, Institute of Industrial
19 Ecological Sciences, University of Occupational and Environmental Health, Japan,
20 Kitakyushu, Japan

21 ³Department of Mental Health, Institute of Industrial Ecological Sciences, University of
22 Occupational and Environmental Health, Japan, Kitakyushu, Japan

23 ⁴Department of Occupational Medicine, School of Medicine, University of
24 Occupational and Environmental Health, Japan, Kitakyushu, Japan

25 ⁵Department of Environmental Health, School of Medicine, University of Occupational

26 and Environmental Health, Japan, Kitakyushu, Japan

27 ⁶Department of Work Systems and Health, Institute of Industrial Ecological Sciences,
28 University of Occupational and Environmental Health, Japan, Kitakyushu, Japan

29 ⁷Department of Preventive Medicine and Community Health, School of Medicine,
30 University of Occupational and Environmental Health, Japan, Kitakyushu, Japan

31

32 **Corresponding author:**

33 Tomohiro Ishimaru, MD, MPH, PhD

34 Department of Environmental Epidemiology, Institute of Industrial Ecological Sciences,
35 University of Occupational and Environmental Health, 1-1 Iseigaoka, Yahata-nishi-ku,
36 Kitakyushu, Fukuoka 807-8555, Japan

37 Tel: +81-93-603-1611, Fax: +81-93-601-7324, e-mail: ishimaru@med.uoeh-u.ac.jp

38

39 Abstract

40 **Objectives:** To combat coronavirus disease 2019 (COVID-19), many countries have
41 used contact tracing apps, including Japan's voluntary-use contact-confirming
42 application (COCOA). The current study aimed to identify industry and workplace
43 characteristics associated with the use of this COVID-19 contact tracing app.

44 **Methods:** This cross-sectional study of full-time workers used an online survey.
45 Multiple logistic regression analysis was used to evaluate the associations of industry
46 and workplace characteristics with contact tracing app use.

47 **Results:** Of the 27,036 participants, 25.1% had downloaded the COCOA. Workers in
48 the public service (adjusted odds ratio [aOR] = 1.29, 95% confidence interval [CI]:
49 1.14–1.45) and information technology (aOR = 1.38, 95% CI: 1.20–1.58) industries
50 were more likely to use the app than were those in the manufacturing industry. In
51 contrast, app usage was less common among workers in the retail and wholesale
52 (aOR = 0.87, 95% CI: 0.76–0.99) and food/beverage (aOR = 0.81, 95% CI:
53 0.70–0.94) industries, but further adjustment for business size attenuated these
54 associations. Workers at larger companies were more likely to use the app.
55 Compared with permanent employees, the odds of using the app were higher for
56 managers and civil servants but lower for those who were self-employed.

57 **Conclusions:** One possible reason for the under-implementation of the contact
58 tracing app in the retail and wholesale and food/beverage industries is small business
59 size, as suggested by the fully adjusted model results. An awareness campaign
60 should be conducted to promote the widespread use of the contact tracing app in
61 these industries.

62
63 **Keywords:** Contact tracing, COVID-19, SARS-CoV-2, smartphone, worksite

64

65 **Introduction**

66 A number of countries have used digital contact tracing tools to combat coronavirus
 67 disease 2019 (COVID-19) by tracing possible infected cases to prevent the spread of
 68 the disease.^{1, 2} Japan's government released a contact tracing app for
 69 smartphones—the contact-confirming application (COCOA)—on June 19, 2020.³
 70 Downloading the COCOA is not mandatory but it is recommended by authorities. As
 71 of March 18, 2021, the COCOA had been downloaded by 20.8% of Japanese citizens
 72 (26 million downloads) and used for 2.5% of positive cases (11,513/452,863).³ The
 73 COCOA uses Bluetooth to collect information on close contacts—those who have
 74 been within 1 meter of an individual who has installed the app for a duration of at least
 75 15 minutes in the last 14 days. When a user is infected with COVID-19, they can alert
 76 others with whom they have been in close contact through the app. No personal
 77 information is stored in the central database. The notified person can then get tested
 78 at the nearest COVID-19 testing center.

79

80 High utilization is important for contact tracing apps because they are only effective
 81 when both those who are infected with COVID-19 and their close contacts have
 82 installed the app. According to a simulation study, 90% of the population needs to use
 83 the COCOA for effective infection control.⁴ In other countries, contact tracing apps,
 84 such as TraceTogether in Singapore, have frequently been used with mandatory
 85 installation and use.² In contrast, this practice is less frequent in countries where
 86 smartphones are not widely used.² In terms of using contact tracing apps, migrant
 87 workers, persons with low income, and older adults who may find the technology
 88 necessary to use the app difficult are all at a disadvantage.⁵ The acceptance rate of

89 contact tracing apps has been found to range from 42% to 70% in developed
90 countries.⁶⁻⁸ Several factors influence this acceptance: trust in the government, health
91 concerns, privacy, battery usage, and concerns about the effectiveness of the app.⁷

92

93 In Japan, employers have promoted the use the COCOA among workers for effective
94 contact tracing to prevent workplace COVID-19 clusters.⁹ Many workers are at risk of
95 COVID-19 infection because of their close contact with colleagues or customers.
96 However, there is little evidence about the associations of industry and workplace
97 characteristics with the use of contact tracing apps. Promoting the use of contact
98 tracing apps in the workplace is also important for preventing the spread of clusters
99 from the working-age population to high-risk populations, such as older adults. To
100 develop a strategy for the widespread use of contact tracing apps, it is necessary to
101 identify the industries or workplaces in which such apps are under-implemented. The
102 purpose of the current study was to identify which industry and workplace
103 characteristics were associated with the use of a COVID-19 contact tracing app.

104

105 **Subjects and Methods**

106 Study design and participants

107 The current study was conducted as part of the Collaborative Online Research on the
108 Novel-coronavirus and Work (CORoNaWork) Project. Details of the study protocol
109 have been published elsewhere.¹⁰ Briefly, the CORoNaWork Project was an online
110 cohort study in Japan. We extracted online self-administered questionnaire data from
111 the baseline dataset, collected December 22–26, 2020, for cross-sectional analysis.
112 Panelists who had registered with an online research company and who were currently
113 working full-time were invited to participate in the survey. Health care workers and

caregivers were not invited to participate. We selected 33,087 participants through cluster sampling stratified by sex, region, and job type. We excluded invalid responses, leaving data on 27,036 participants for analysis. When the baseline survey was conducted, the numbers of COVID-19 infections and deaths were much higher than they had been during the first and second waves of the disease in Japan; therefore, Japan was on maximum alert during this third wave. This study was approved by the Ethics Committee of the University of Occupational and Environmental Health, Japan (R2-079).

Questionnaire

This study used questionnaire data including variables on (a) demographic characteristics; (b) health behavior; (c) risk perception; (d) industry and workplace characteristics; and (e) the outcome (use of the contact tracing app). The demographic variables included sex, age, marital status, education, and annual household income. The health behavior variables were smoking and regular alcohol intake. Risk perception was assessed using a single question on anxiety about contracting COVID-19, with the response options of *yes* and *no*. Three variables were used to assess industry and workplace characteristics: type of industry, business size, and occupation. Type of industry followed the Japan Standard Industrial Classification, with industries accounting for less than 3% of the sample categorized as “other.”¹¹ Business size was classified as < 10, 10–49, 50–999, or ≥ 1,000 employees. Each participant reported their occupation as *permanent employee*, *manager*, *civil servant*, *dispatched or contract worker*, *self-employed*, or *other*. The outcome variable, the use of the contact tracing app, was assessed by asking “Have you downloaded the COCOA?” The response options were *yes* and *no*.

139

140 Data analysis

141 Univariate and multiple logistic regression analyses were used to evaluate the
 142 associations of occupational factors with the use of the contact tracing app. We
 143 assessed three occupational factors as explanatory variables: type of industry,
 144 business size, and occupation. Three models were evaluated. Model 1 was the
 145 univariate model. Model 2 was adjusted for demographic, health behavior, and risk
 146 perception variables (sex, age, marital status, education, annual household income,
 147 smoking, alcohol intake, and anxiety about contracting COVID-19). Model 3
 148 additionally adjusted for type of industry and business size. We did not include the
 149 occupation variables in Model 3 because of the high correlations between the
 150 occupation of civil servant and the public service industry and between self-employed
 151 occupation and business size of 1–9 employees. All *P*-values were two-sided, and
 152 statistical significance was set at $P < 0.05$. We used Stata/SE 16.1 (StataCorp,
 153 College Station, TX, USA) for all analyses.

154

155 Results

156 Table 1 shows the general characteristics of the study participants. Of the 27,036
 157 participants, 6,786 (25.1%) reported having downloaded the COCOA. Approximately
 158 half of the participants were women (48.9%), and about half were married (55.6%).
 159 The most frequently observed category for annual household income was 4–8 million
 160 JPY (44.1%; USD 1 = JPY 106.78 as of 2020).¹² Excluding the “other” industry
 161 category, manufacturing accounted for the largest percentage of the sample (17.0%),
 162 followed by medical and welfare (16.6%) and public service (7.0%). Establishments
 163 with 50–999 employees (35.9%) and permanent employee (46.5%) were the most

frequently observed business size and occupation categories, respectively.

Table 2 shows the associations of industry and workplace characteristics with the use of the COVID-19 contact tracing app. Participants in the public service (adjusted odds ratio [aOR] = 1.29, 95% confidence interval [CI]: 1.14–1.45, Model 3) and information technology (aOR = 1.38, 95% CI: 1.20–1.58, Model 3) industries were more likely to use the app than were those in the manufacturing industry. In contrast, app usage was less common among participants working in the retail and wholesale and food/beverage industries, and adjusting for demographic, health behavior, and risk perception variables did not alter these associations remarkably (retail and wholesale: aOR = 0.87, 95% CI: 0.76–0.99; food/beverage: aOR = 0.81, 95% CI: 0.70–0.94; Model 2); however, after further adjustment for business size in Model 3, these industry types showed no significant difference from the manufacturing industry.

Participants in larger companies were more likely to use the app (50–999 employees: aOR = 1.21, 95% CI: 1.12–1.32; ≥ 1,000 employees: aOR = 1.37, 95% CI: 1.24–1.52; Model 3). Regarding occupation, compared with permanent employees, the odds of app usage were higher for managers (aOR = 1.40, 95% CI: 1.28–1.53, Model 2) and civil servants (aOR = 1.22, 95% CI: 1.11–1.34, Model 2) but lower for those who were self-employed (aOR = 0.76; 95% CI: 0.69–0.85, Model 2).

Discussion

The current study identified that working in the public service sector, in the information technology industry, or for a business with 50 or more employees, as well as holding the position of manager or civil servant, were factors facilitating the use of the contact

tracing app—over 30% of each of these groups had downloaded the app. In contrast, working in the retail and wholesale and food/beverage industries and being self-employed were associated with a lower likelihood of using the app—around 20% of these groups had downloaded the app. These findings give insight needed for the development of a strategy for the widespread use of COVID-19 contact tracing apps.

An important finding of this study was that the contact tracing app was under-implemented in the retail and wholesale and food/beverage industries. This finding did not change substantially after adjusting for demographic, health behavior, and risk perception variables. However, further adjustment for business size attenuated these associations. One possible reason for this change in results is that most workplaces in these types of industries are small-scale, which may be the cause of insufficient use of the app. This idea is consistent with a previous study showing that small companies were relatively unlikely to implement workplace measures against COVID-19.¹³ This may be because of the limited opportunities to obtain information and the lack of human, time, and financial resources to take countermeasures against health threats at small companies.¹⁴ However, workers in these industries have particularly a strong need to use the contact tracing app because they are in contact with an unspecified number of people, such as in restaurants and bars, where customers eat and drink without masks.¹⁵ An awareness campaign should be conducted for the retail and wholesale and food/beverage industries to promote the use of the contact tracing app.¹⁶

We found that the contact tracing app was frequently used by civil servants and managers. This finding implies that civil servants and managers might follow requests

to use the app made through organized channels because workers were encouraged to use the COCOA by the government and by employer associations.⁹ A previous study conducted in Japan reported that the public service sector most frequently received announcements regarding measures to prevent COVID-19 in the early stage of the pandemic¹⁷—this also suggests that the public service sector provides COVID-19 advisory information to civil servants. These results indicate that app uptake may increase when workers, including managers, receive messages through organized channels, even if the requests are voluntary. For non-managers, a sharing function allowing users to invite coworkers and friends to download the app may increase the rate of installation of the app.⁷

Workers in the information technology industry, such as system engineers, have the advantage of expertise in digital technology and know how to use smartphone apps. The current study revealed that information technology industry workers used the contact tracing app more frequently, compared with workers in other industries. This finding is in line with a previous study showing that high information technology literacy eliminated technical obstacles to using contact tracing apps.¹⁸ In addition, the information technology industry actively implements workplace measures against COVID-19, such as teleworking, and these practices may also relate to the finding.¹⁷ These results suggest that a technical support service for those who have difficulty using the contact tracing app may improve app uptake.

This study has several limitations. First, the study recruited panelists who registered with an online research company. Therefore, the participants may not represent general workers. For example, online panelists may be particularly willing to use

online tools or more familiar than others with using apps. Consequently, the results may underestimate negative factors for use of the app. Second, we evaluated the use of the contact tracing app by asking participants about whether they had downloaded it. Therefore, we did not confirm whether the application was installed. However, we think that most people who downloaded the app also installed it. Despite these limitations, to the best of our knowledge, this study represents the first study in Japan to examine current use of the COCOA with a large sample.

In conclusion, the present study evaluated the associations of industry and workplace characteristics with the use of a COVID-19 contact tracing app in a large-scale online survey of Japanese workers. Those working in the public service sector or in information technology, as well as managers, were frequently found to use the contact tracing app, whereas those working in the retail and wholesale and food/beverage industries were less likely to use it. One possible reason for the under-implementation of the contact tracing app in the retail and wholesale and food/beverage industries may be the small size of businesses in these types of industries. An awareness campaign should be conducted for workers in these industries to promote the widespread use of the contact tracing app to help these workers trace possible infected cases, preventing the spread of the disease.

Acknowledgments

This study was supported by a research fund from the University of Occupational and Environmental Health, Japan; General Incorporated Foundation (Anshin Zaidan): The Development of Educational Materials on Mental Health Measures for Managers at Small-sized Enterprises; Health, Labour and Welfare Sciences Research Grants:

Comprehensive Research for Women's Healthcare (H30-josei-ippan-002); Research for the Establishment of an Occupational Health System in Times of Disaster (H30-roudou-ippan-007), and scholarship donations from Chugai Pharmaceutical Co., Ltd.

The current members of the CORoNaWork Project, in alphabetical order, are as follows: Dr. Yoshihisa Fujino (present chairperson of the study group), Dr. Akira Ogami, Dr. Arisa Harada, Dr. Ayako Hino, Dr. Chimed-Ochir Odgerel, Dr. Hajime Ando, Dr. Hisashi Eguchi, Dr. Kazunori Ikegami, Dr. Keiji Muramatsu, Dr. Koji Mori, Dr. Kyoko Kitagawa, Dr. Masako Nagata, Dr. Mayumi Tsuji, Dr. Rie Tanaka, Dr. Ryutaro Matsugaki, Dr. Seiishiro Tateishi, Dr. Shinya Matsuda, Dr. Tomohiro Ishimaru, Dr. Tomohisa Nagata, Dr. Yosuke Mafune, and Ms. Ning Liu. All members are affiliated with the University of Occupational and Environmental Health, Japan.

Disclosure

Approval of the research protocol: This study was approved by the Ethics Committee of the University of Occupational and Environmental Health, Japan (R2-079).

Informed Consent: Informed consent was obtained from all participants. *Registry and the Registration No. of the study/trial:* N/A. *Animal Studies:* N/A. *Conflict of Interest:* N/A.

References

1. Salathe M, Althaus C, Anderegg N, *et al.* Early evidence of effectiveness of digital contact tracing for SARS-CoV-2 in Switzerland. *Swiss Med Wkly* 2020;150:w20457.

- 289 2. Elkhodr M, Mubin O, Iftikhar Z, *et al.* Technology, Privacy, and User Opinions of
290 COVID-19 Mobile Apps for Contact Tracing: Systematic Search and Content
291 Analysis. *J Med Internet Res* 2021;23:e23467.
- 292 3. Ministry of Health Labour and Welfare. COVID-19 Contact-Confirming Application.
293 https://www.mhlw.go.jp/stf/seisakunitsuite/bunya/cocoa_00138.html. Published
294 2021. Accessed March 18, 2021.
- 295 4. Nakamoto I, Jiang M, Zhang J, *et al.* Evaluation of the Design and Implementation
296 of a Peer-To-Peer COVID-19 Contact Tracing Mobile App (COCOA) in Japan.
297 *JMIR Mhealth Uhealth* 2020;8:e22098.
- 298 5. Anglemyer A, Moore TH, Parker L, *et al.* Digital contact tracing technologies in
299 epidemics: a rapid review. *Cochrane Database Syst Rev* 2020;8:CD013699.
- 300 6. Lewandowsky S, Dennis S, Perfors A, *et al.* Public acceptance of
301 privacy-encroaching policies to address the COVID-19 pandemic in the United
302 Kingdom. *PLoS One* 2021;16:e0245740.
- 303 7. Garrett PM, White JP, Lewandowsky S, *et al.* The acceptability and uptake of
304 smartphone tracking for COVID-19 in Australia. *PLoS One* 2021;16:e0244827.
- 305 8. Garrett PM, Wang Y, White JP, *et al.* Young Adults View Smartphone Tracking
306 Technologies for COVID-19 as Acceptable: The Case of Taiwan. *Int J Environ*
307 *Res Public Health* 2021;18.
- 308 9. Japan Federation of Economic Organizations. The NEW growth strategy.
309 https://www.keidanren.or.jp/en/policy/2020/108_proposal.html. Published 2020.
310 Accessed March 18, 2021.
- 311 10. Fujino Y, Ishimaru T, Eguchi H, *et al.* Protocol for a nationwide Internet-based
312 health survey in workers during the COVID-19 pandemic in 2020. *JUOEH*
313 2021:in press.

- 314 11. Ministry of Internal Affairs and Communications. Japan Standard Industrial
315 Classification.
316 https://www.soumu.go.jp/toukei_toukatsu/index/seido/sangyo/H25index.htm.
317 Published 2014. Accessed March 18, 2021.
- 318 12. Organisation for Economic Co-operation and Development. Exchange rates.
319 [https://data.oecd.org/conversion/exchange-rates.htm?fbclid=IwAR2Bn6JIF8Wf](https://data.oecd.org/conversion/exchange-rates.htm?fbclid=IwAR2Bn6JIF8WfAbVhwDEiKFhG0_rsTu-iLq1PW47o1ujebwc7CWF0e0wUx_A)
320 [AbVhwDEiKFhG0_rsTu-iLq1PW47o1ujebwc7CWF0e0wUx_A](https://data.oecd.org/conversion/exchange-rates.htm?fbclid=IwAR2Bn6JIF8WfAbVhwDEiKFhG0_rsTu-iLq1PW47o1ujebwc7CWF0e0wUx_A). Published 2021.
321 Accessed March 18, 2021.
- 322 13. Ishimaru T, Nagata M, Hino A, *et al*. Workplace measures against COVID-19
323 during the winter third wave in Japan: company size-based differences.
324 *medRxiv* 2021:2021.2002.2014.21251716.
- 325 14. Hasle P, Limborg HJ. A review of the literature on preventive occupational health
326 and safety activities in small enterprises. *Ind Health* 2006;44:6-12.
- 327 15. Kim J, Kim J, Wang Y. Uncertainty risks and strategic reaction of restaurant firms
328 amid COVID-19: Evidence from China. *Int J Hosp Manag*
329 2021;92:102752-102752.
- 330 16. Rodriguez P, Grana S, Alvarez-Leon EE, *et al*. A population-based controlled
331 experiment assessing the epidemiological impact of digital contact tracing. *Nat*
332 *Commun* 2021;12:587.
- 333 17. Sasaki N, Kuroda R, Tsuno K, Kawakami N. Workplace responses to COVID-19
334 and their association with company size and industry in an early stage of the
335 epidemic in Japan. *Environmental and Occupational Health Practice*
336 2020;2:eohp.2020-0007-OA.
- 337 18. von Wyl V, Hoglinger M, Sieber C, *et al*. Drivers of Acceptance of COVID-19
338 Proximity Tracing Apps in Switzerland: Panel Survey Analysis. *JMIR Public*

339 *Health Surveill* 2021;7:e25701.

340

341 Table 1. General characteristics of the study participants

	Total <i>N</i> = 27,036 <i>n</i> (%)	COVID-19 contact tracing app users <i>n</i> = 6,786 (25.1%)	
		<i>n</i> (%)	Utilization rate (%)
Sex			
Female	13,222 (48.9)	3,181 (46.9)	24.1
Male	13,814 (51.1)	3,605 (53.1)	26.1
Age			
20–39 years	6,763 (25.0)	1,669 (24.6)	24.7
40–49 years	8,011 (29.6)	1,886 (27.8)	23.5
50–59 years	9,012 (33.4)	2,359 (34.8)	26.2
60–65 years	3,250 (12.0)	872 (12.8)	26.8
Marital status			
Single	9,164 (33.9)	1,991 (29.4)	21.7
Divorced or widowed	2,843 (10.5)	722 (10.6)	25.4
Married	15,029 (55.6)	4,073 (60.0)	27.1
Education			
Junior high or high school	7,321 (27.1)	1,647 (24.3)	22.5
Vocational school or college	6,544 (24.2)	1,552 (22.9)	23.7
University or graduate school	13,171 (48.7)	3,587 (52.8)	27.2
Annual household income			
< 2,000,000 Japanese yen	1,709 (6.3)	277 (4.1)	16.2
2,000,000–3,999,999 Japanese yen	5,548 (20.5)	1,188 (17.5)	21.4
4,000,000–7,999,999 Japanese yen	11,927 (44.1)	2,925 (43.1)	24.5
≥ 8,000,000 Japanese yen	7,852 (29.1)	2,396 (35.3)	30.5
Current smoking	7,004 (25.9)	1,740 (25.6)	24.8
Regular alcohol intake	11,017 (40.8)	2,901 (42.7)	26.3
Anxiety about contracting COVID-19 (yes)	20,017 (74.0)	5,737 (84.5)	28.7
Industry			
Manufacturing	4,591 (17.0)	1,165 (17.2)	25.4
Public service	1,891 (7.0)	614 (9.0)	32.5
Information technology	1,331 (4.9)	429 (6.3)	32.2
Retail and wholesale	1,797 (6.6)	394 (5.8)	21.9
Food/beverage	1,319 (4.9)	269 (4.0)	20.4
Medical and welfare	4,500 (16.6)	1,071 (15.8)	23.8
Finance	1,226 (4.5)	364 (5.4)	29.7
Construction	944 (3.5)	240 (3.5)	25.4
Other	9,437 (34.9)	2,240 (33.0)	23.7
Business size			
1–9 employees	6,165 (22.8)	1,272 (18.7)	20.6
10–49 employees	4,390 (16.2)	957 (14.1)	21.8
50–999 employees	9,703 (35.9)	2,491 (36.7)	25.7
≥ 1,000 employees	6,778 (25.1)	2,066 (30.4)	30.5
Occupation			
Permanent employee	12,575 (46.5)	3,006 (44.3)	23.9
Manager	3,403 (12.6)	1,160 (17.1)	34.1
Civil servant	2,810 (10.4)	840 (12.4)	29.9
Dispatched or contract worker	2,894 (10.7)	671 (9.9)	23.2
Self-employed	2,819 (10.4)	525 (7.7)	18.6
Other	2,535 (9.4)	584 (8.6)	23.0

343 Table 2. Associations between occupational factors and the use of the COVID-19 contact tracing app

	Model 1			Model 2			Model 3		
	OR	(95% CI)	p value	OR	(95% CI)	P-value	OR	(95% CI)	P-value
Industry									
Manufacturing	1.00	-	-	1.00	-	-	1.00	-	-
Public service	1.41	(1.25–1.58)	< 0.001	1.32	(1.17–1.49)	< 0.001	1.29	(1.14–1.45)	< 0.001
Information technology	1.39	(1.21–1.58)	< 0.001	1.36	(1.18–1.55)	< 0.001	1.38	(1.20–1.58)	< 0.001
Retail and wholesale	0.83	(0.73–0.94)	0.004	0.87	(0.76–0.99)	0.034	0.92	(0.81–1.05)	0.223
Food/beverage	0.76	(0.66–0.89)	< 0.001	0.81	(0.70–0.94)	0.007	0.89	(0.76–1.03)	0.125
Medical and welfare	0.94	(0.85–1.03)	0.201	0.86	(0.78–0.94)	0.004	0.91	(0.82–1.01)	0.068
Finance	1.26	(1.09–1.45)	0.001	1.19	(1.03–1.38)	0.016	1.15	(0.99–1.32)	0.061
Construction	1.00	(0.85–1.18)	0.999	0.98	(0.84–1.16)	0.845	1.10	(0.93–1.30)	0.271
Other	0.92	(0.84–0.99)	0.034	0.94	(0.87–1.03)	0.175	1.01	(0.93–1.10)	0.807
Business size									
1–9 employees	1.00	-	-	1.00	-	-	1.00	-	-
10–49 employees	1.09	(0.99–1.20)	0.063	1.04	(0.94–1.14)	0.453	1.03	(0.97–1.11)	0.329
50–999 employees	1.36	(1.25–1.46)	< 0.001	1.24	(1.14–1.34)	< 0.001	1.22	(1.13–1.33)	< 0.001
≥ 1,000 employees	1.71	(1.57–1.85)	< 0.001	1.52	(1.39–1.65)	< 0.001	1.44	(1.32–1.57)	< 0.001
Occupation*									
Permanent employee	1.00	-	-	1.00	-	-			
Manager	1.62	(1.48–1.76)	< 0.001	1.40	(1.28–1.53)	< 0.001			
Civil servant	1.34	(1.23–1.47)	< 0.001	1.22	(1.11–1.34)	< 0.001			
Dispatched or contact worker	0.95	(0.86–1.04)	0.277	1.02	(0.92–1.12)	0.769			
Self-employed	0.72	(0.64–0.80)	< 0.001	0.76	(0.69–0.85)	< 0.001			
Other	0.95	(0.86–1.05)	0.290	0.89	(0.80–0.99)	0.033			

344 COVID-19: coronavirus disease 2019; OR: odds ratio; CI: confidence interval

345 Model 1: Adjusted for sex and age

346 Model 2: Adjusted for sex, age, income, marital status, education, smoking, alcohol intake, and anxiety about contracting COVID-19

347 Model 3: Adjusted for sex, age, income, marital status, education, smoking, alcohol intake, anxiety about contracting COVID-19, type of industry,
348 and business size

349 * The occupation variables were not included in Model 3 because of the high correlations between the occupation of civil servant and the public
350 service industry and between self-employed occupation and business size of 1–9 employees.