Examining changes in sleep duration associated with the onset of the COVID-19 pandemic: Who is sleeping and who is not?

Salma Batool-Anwar, M.D., M.P.H.^{1,2}

Rebecca Robbins, Ph.D.^{1,2}

Shahmir H. Ali, B.A.³

Ariadna Capasso, M.F.A.³

Joshua Foreman, Ph.D.^{5,3}

Abbey M. Jones, M.P.H.⁶

Yesim Tozan, Ph.D.⁴

Ralph J. DiClemente, Ph.D.³

Stuart F. Quan, M.D.^{1,2}

¹Department of Medicine, Division of Sleep and Circadian Disorders, Brigham & Women's Hospital, Boston, MA

²Division of Sleep Medicine, Harvard Medical School, Boston MA

³Department of Social and Behavioral Sciences, School of Global Public Health, New York University, New York, NY

⁴Global Health Program, School of Global Public Health, New York University, New York, NY

⁵Ophthalmology, Department of Surgery, University of Melbourne, Melbourne, Australia

⁶Department of Epidemiology, School of Global Public Health, New York University, New York, NY

Running Head: Changes in Sleep with COVID-19

Correspondent: Stuart F. Quan, M.D.

Division of Sleep and Circadian Disorders, Brigham and Women's Hospital

221 Longwood Ave.

Boston, MA 02115

Email: Stuart_Quan@hms.harvard.edu

Attestation: All authors have seen and approved this manuscript.

Declarations:

Financial: This work was supported by K01HL150339 from the National Heart, Lung and Blood Institute.

Conflicts of Interest: Dr. Robbins has received personal fees from Denihan Hospitality, Rituals Cosmetics, byNight, and SleepCycle. Dr. Quan serves as a consultant to Jazz Pharmaceuticals, Best Doctors and Whispersom. Drs. Batool-Anwar, Foreman, Tozan, DiClemente, and Mr. Ali, and Mmes. Capasso and Jones have no conflicts of interest.

Word Counts: Abstract	Brief Summary	Manuscript
250		3745
Tables: 4	Figures: 0	

Changes in Sleep with COVID-19

Abstract

Introduction

The COVID-19 pandemic has resulted in social isolation and reports of insomnia. However, reports of changes in sleep duration and associated factors are few.

Methods

Data were from an online survey of adults recruited via social media that included a question asking whether the respondent slept less or more after the onset of the pandemic. Analyses determined the association between changes in sleep duration and self reported sociodemographic and occupational information; beliefs about COVID-19; changes in sleep patterns; and responses pertaining to loneliness, anxiety, and depression.

Results

There were 5,175 respondents; 53.9% had a change in sleep duration. 17.1% slept less and 36.7% slept more. Sleeping more was related to greater education, being single/divorced/separated, unemployed or a student. Being retired, divorced/separated or a homemaker, and living in the Mountain or Central time zones were associated with less sleep. Beliefs that COVID-19 would result in personal adverse consequences was associated with both more and less sleep. However, the strongest associations with both more and less sleep were seen with depression, anxiety, and loneliness with adjusted odds ratios ranging from 1.92 (95% CI 1.67-2.21) for sleeping more and loneliness to 5.29 (95% CI 4.1-6.7) for sleeping less and anxiety.

Conclusions

Changes in sleep duration since the start of the COVID-19 pandemic were highly prevalent among social media users and were associated with several sociodemographic factors and beliefs that

Changes in Sleep with COVID-19

COVID-19 would have adverse personal impacts. However, the strongest associations occurred with worse mental health suggesting that improvements may occur with better sleep.

Keywords

COVID-19, sleep duration, mental health, depression, anxiety, psychosocial beliefs

Changes in Sleep with COVID-19

Introduction

It is widely recognized that sleep is a vital psychophysiologic process for mental health. In addition to increased prevalence of mental health conditions among people with sleep disorders,¹ a bidirectional relationship also exists between sleep and psychiatric disorders.² This association is further strengthened during times of crisis from natural disasters and pandemics. Increased rates of insomnia among adolescents (38.3%) were seen after the Wenchuan earthquake of 2008³, and these sleep disturbances were associated with depression and post traumatic stress disorder.⁴ Similar experiences of a negative effect on mental health were seen during previous pandemics.⁵ After the MERS pandemic of 2015, high rates (47.2%) of anxiety symptoms were seen as many as six months after the isolation ended.⁶ Individuals infected with SARS and MERS also reported greater anxiety, depression, insomnia, and post-traumatic stress disorder (PTSD) during and in the post-illness stage.⁷

These experiences from past epidemics and natural disasters have raised concerns for imminent psychosocial impact following the current COVID-19 pandemic. Although SARS-CoV-2 mainly affects the respiratory system, recent reports have demonstrated neurologic symptoms related to COVID-19.⁸ There has been increased focus on neuropsychiatric effects of the virus and its relationship with impaired sleep.^{7,9}

The current pandemic has caused increased mortality, economic instability, and social isolation. The devastation of this magnitude is generating considerable burden in terms of population-wide mental health.^{5,10} Besides the direct effects of SARS-CoV-2 on the brain, the detrimental effects of lockdown and quarantine on psychosocial health also have been reported globally.¹¹⁻¹³ Increased anxiety and stress have been reported among people during isolation and quarantine.⁵ Increased cases of stigmatization and discrimination during these periods of social

Changes in Sleep with COVID-19

isolation have been shown to further contribute to psychosocial stress.¹⁴ Additionally, the effects of home confinement on light exposure and more access to use of digital tools has been shown to have a negative impact on sleep quality.¹⁵

Worsening psychosocial as well as sleep health can be seen among vulnerable populations. Healthcare workers including physicians, nurses, laboratory workers, and technicians delivering care at the front line are considered the most vulnerable during a pandemic situation. A recent meta-analysis demonstrated increased rates of anxiety, depression, PTSD, and insomnia among healthcare workers affecting their well-being.¹⁶ They are put in an unprecedented situation by having to make difficult decisions and work under extreme pressure, both resulting in moral injury.¹⁷

Although several studies have reported increased depression, irritability, PTSD, and insomnia among COVID-19 survivors and healthcare workers, the data on the impact of COVID-19 on changes in sleep duration among the general population are limited. The objective of this study was to extend the results of previous studies by examining the effect of the COVID-19 pandemic on self-reported changes in sleep duration and the sociodemographic and mental health factors associated with these changes among the general population of the United States (US).

Methods

Participants

Recruitment for this survey has been described extensively elsewhere.^{18,19} Briefly, during April 16-21, 2020, a social-media advertisement campaign was launched on Facebook and affiliated platforms (i.e., Instagram, Facebook Messenger) to attract participation in a survey of COVID-19 knowledge, beliefs and behaviors. Previous research has shown social media to be a valid method of recruitment for health surveys, including those focused on COVID-19.^{13,20} The

Changes in Sleep with COVID-19

campaign was targeted to persons who were at least 18 years of age and residing in the United States. Interested social media users were directed to an anonymous web-based survey administered by Qualtrics (Provo, UT). Eligibility was confirmed by brief screening questions. Participants were not compensated for survey completion. All study material and procedures underwent Human Subjects Research review by the New York University Institutional Review Board and were determined to be exempt, and the need for explicit written or oral consent was also waived (IRB-FY2020–4285).

Questionnaire

Survey development has been outlined previously.^{18,19} In summary, the survey was based on principles expressed in the Health Belief Model,²¹ a paradigm used in surveys related to other viral outbreaks.²²⁻²⁴ In addition to demographic information, the survey included a variety of questions related to the participant's behaviors, attitudes and beliefs towards COVID-19 as well as their current mental health. Items were adapted from surveys utilized during previous infectious disease outbreaks so that they were pertinent to the COVID-19 pandemic.²⁵ The survey also contained a single question regarding sleep: ["Since hearing about the Coronavirus outbreak, how have the below behaviors changed for you? Sleeping"]. Responses were elicited using a 5-point Likert scale: "Much less," "Little less," "Not changed," "Little more," and "Much more" with an additional Not Applicable response. Participants who selected Not Applicable were removed from the analysis. These responses were aggregated into 3 categories "sleeping less than usual" (combination of "Much less" and "Little less sleep"), "sleeping more than usual" (combination of "Little more" and "Much more") and "no change in sleep." Besides sociodemographic characteristics of the cohort, we chose additional survey items to be included in these analyses based on our *a priori* assessment that their content would be relevant to changes in sleep behavior.

Changes in Sleep with COVID-19

Beliefs about COVID-19 assessed included perceived risk [“What do you think is your risk of getting infected with Coronavirus?”] and perceived severity [“If you were infected with Coronavirus, how severe do you think it would be?”], which were measured on a 10-point scale from “not at all likely” to “extremely high risk” and “extremely severe,” respectively. Responses to perceived risk and severity were re-coded so responses of “not at all” to the scale midpoint (5) represented “low” beliefs about risk and severity, while responses above the scale mid-point (6) to 10 (Very Severe) were re-coded to represent “high” beliefs about risk and severity. Responses to financial ability to quarantine [“I can financially afford to self-quarantine”], beliefs about discrimination [“Since the Coronavirus outbreak, I feel discriminated against”], beliefs that COVID-19 is deadlier than influenza [“Coronavirus is more deadly than the seasonal flu”], beliefs COVID-19 is not as big a problem as the media suggest [“Coronavirus is not as big of a problem as the media suggests”], and beliefs that COVID-19 is a bigger problem than the government suggests [“Coronavirus is a bigger problem than the government suggests”] were captured on scales from 0 “strongly disagree” to 3 “strongly agree.” Responses were dichotomized so that “strongly disagree” and “disagree” were re-coded to indicate “low” beliefs or concerns, while “disagree” or “strongly disagree” were re-coded to represent “high” beliefs or concerns, with the exception of financial ability to quarantine, which was re-coded so that “strongly agree” and “agree” were represented low financial worries and “strongly disagree” and “disagree” represented high financial worries.

Mental health was assessed using the Patient Health Questionnaire-4 (PHQ-4),²⁶ an ultra-brief screening scale for depression and anxiety, which was included in the survey and modified to refer to feelings about the coronavirus outbreak over the past 7 days. Anxiety was considered

Changes in Sleep with COVID-19

present if there was a score of 3 or greater on the first 2 items of the PHQ-4. Similarly, depression was determined to be present if there was a score of 3 or greater on the last 2 items of the PHQ-4. In addition to anxiety and depression, the construct of “loneliness” was assessed using responses to the following 3 questions: “In the last 3 months, how often have you felt: 1) left out; 2) isolated; 3) that you lacked companionship.” Possible responses for all 3 questions were the following: 0) Never; 1) Often; 2) Rarely; 3) Sometimes. A cumulative score of 6 or higher was considered as indicative of “loneliness.”²⁷ Additional details regarding the wording of the survey items and possible responses are provided in the supplement.

Statistical Analysis

We estimated differences in sociodemographic characteristics, COVID-19 beliefs, and mental health by reported change in sleep (i.e., no change, versus either sleeping more than usual or less than usual). We utilized Pearson χ^2 to determine if associations existed between sociodemographic factors and the reported change in sleep. Next, to examine sleep changes by sociodemographic factors, we conducted multivariable binary logistic regressions to characterize the sociodemographic profile of those who reported sleeping more than usual and less than usual. In both cases, models used reports of “no change in sleep” as the reference group. To examine COVID-19 beliefs by sleep changes, we ran adjusted multivariable binary logistic regressions to examine reports of sleeping more and sleeping less and their relationship to COVID-19 beliefs, after adjusting for confounders (i.e., age, sex, race, presence of a child at home, urban residence, marital status, education, employment, and time zone). Finally, to examine mental health by sleep changes, we conducted adjusted binary logistic regression to examine reports of sleeping more and sleeping less and their relationship to loneliness, anxiety, and depression, controlling for the

Changes in Sleep with COVID-19

aforementioned confounders. The level of statistical significance for all models was set at 0.05.

Analyses were conducted in SPSS (Version 21; Armonk, NY).

Results

There were 5,175 participants who responded to the survey question pertaining to change in sleep. Their demographic information disaggregated by whether they reported a change in sleep or not, is shown in Table 1. Irrespective of whether they reported a change in their sleep duration, most respondents were between the ages of 50 and 69 years (26-34%), predominantly of white race (92.7%-93.5%), and more likely to be female (54.6-62.2%). A majority also reported being married (68.7%-74.1%), residing in an urban area (51.6%-55%) in the Eastern time zone (54%-57.8%), not having children in the household (71.9%-76.4%), having gone to college (88.6%-91.7%), and being employed (53.7%-58.9%).

Overall, 2,787 (53.9%) participants experienced a change in their sleep duration as a consequence of the COVID-19 pandemic. As shown in Table 1, younger persons were more impacted by the pandemic than older individuals. In addition, more women (62.2%) than men (37.8%) altered their sleep duration. Other demographic factors associated with a large (>3% absolute difference) change in sleep duration were living in an urban area (55.0% [changed] vs. 51.6% [unchanged]), having a child at home (28.1% [changed] vs. 20.3% [unchanged]), being single (17.9% [changed] vs. 13.9% [unchanged]), or being employed (58.9% [changed] vs 53.7% [unchanged]). Conversely, living in a rural area (28.8% [changed] vs. 34.6% [unchanged]) and being retired (20.4% [changed] vs. 31.9% [unchanged]) appeared to have a large protective effect on change in sleep duration.

Associations of sociodemographic factors with sleeping less since hearing about the COVID-19 outbreak are displayed in Table 2. There were 884 (17.1%) participants who slept less

Changes in Sleep with COVID-19

than usual. Significant associated factors were younger age (<60 years), female sex, urban and suburban residential area and being divorced or separated.

Table 2 also shows associations of sociodemographic variables with sleeping more since hearing about the COVID-19 outbreak. There were 1,903 (36.7%) participants who slept more than usual. Age was an important associated factor, and analogous to sleeping less than usual, the odds of sleeping more appeared to become progressively greater with decreasing age. Similar to sleeping less, sleeping more was positively associated with female sex and being divorced or separated (vs. married). However, sleeping more also was related to being single in comparison to being married. Having a college education or higher was associated with more sleep compared to those with only a high school education. The impact of work status was mixed. In comparison to those who were employed, those who were unemployed or students slept more. In contrast, those who were retired or were homemakers were less likely to sleep more. The impact of time zone showed that those residing in the Mountain or Central time zones were less likely to sleep more. Although those whose race was other than Black, Hispanic or Asian appeared to be less likely to sleep more than Whites, the overall number of respondents to the survey in this demographic group was small.

Table 3 displays the association between COVID-19 beliefs and the likelihood of sleeping more or less since hearing about COVID-19. Sleeping less was associated with a higher perceived risk of COVID-19 infection, belief that COVID-19 was deadlier than the flu and belief that the problem was greater than what the government suggests. In contrast, more financial worries and disagreement with the statement that “COVID-19 is not as big a problem as the media suggest” was associated with lower odds of sleeping less. Sleeping more than usual was associated with the same pattern of responses as sleeping less than usual with one exception; there was a greater

Changes in Sleep with COVID-19

likelihood of sleeping more related to the belief that infection with COVID-19 would be associated with severe symptoms.

Table 4 shows the relationship between mental health constructs and changes in sleep duration. Both sleeping less than usual and sleeping more than usual were strongly associated with anxiety, depression, and loneliness in unadjusted and fully adjusted models.

Discussion

This study examined changes in sleep duration among a national sample of social media users since hearing about the COVID-19 pandemic as well as factors associated with these changes. We demonstrated that a large proportion of the United States adult population suffered a change in sleeping patterns during this pandemic. Importantly, these changes disproportionately impacted women, younger to middle aged adults, and those living in urban settings. Differential associations were further noted as a consequence of marital status, employment, and education. Being single and employed was associated with sleeping less than usual. However, the strongest associations were observed with symptoms of mental health impairment.

One of the most important findings in our analyses was that 53.9% of our cohort experienced a change in their sleep duration as a result of the COVID-19 pandemic with the largest proportion (36.7%) sleeping more than usual. This latter finding is consistent with recent data from a global internet survey which demonstrated an average 26 minute increase in sleep duration and 50 minute delay in mid-sleep time on workdays²⁸ and from another survey using data downloaded from a widely used sleep app showing a mean increase of 14 minutes in 5 large American, European and Asian cities,²⁹ as well as an internet survey in Americans.³⁰ The ability to work from home without “social time pressure” is the most likely explanation.²⁸ Moreover, given the

Changes in Sleep with COVID-19

multitude of stressors during this pandemic, fatigue and depression may also be contributing to increased sleep time.

Notwithstanding the large percentage of those who increased their sleep duration, there also were 17.1% of participants who were sleeping less than usual. This finding of worsening sleep during the pandemic is consistent with previous studies demonstrating increased rates of insomnia. In mainland China, an online survey of participants between the ages of 18-76 years found that the prevalence of insomnia rose from 26.2% before to 33.7% during the pandemic.³¹ Similarly, a recent meta-analysis also demonstrated a high prevalence of insomnia (23.8%) with a greater impact among healthcare workers in comparison to the general population.³² In another meta-analysis,³³ researchers reported an overall pooled prevalence of insomnia of over 37% with higher rates among healthcare workers compared to the general population (47.3% vs 18.2%). Other studies have reported poor sleep among healthcare workers during previous SARS and MERS pandemics.^{34,35} However, most of these studies are from China and Europe where the devastation from this pandemic was felt initially and without prior knowledge of the overall impact of this virus. The uncertainty surrounding the pandemic likely contributed to stress,³⁶ a major contributor to insomnia.

Our study found that women were more likely to report both sleeping less and sleeping more in response to the current pandemic. Our results are consistent with those reported in a Greek population.³⁷ Women maybe more vulnerable to stress, and in general report more somatic symptoms and poor sleep than men.³³ As evidenced in our study, sleep duration is also impacted more by childcare needs which has a greater consequence for women. Additionally, most studies demonstrating higher prevalence of insomnia among healthcare workers have a higher proportion of women in their cohorts; insomnia generally is more prevalent in women.³⁸

Changes in Sleep with COVID-19

The other major demographic factor associated with a change in sleep duration was age. For both sleeping more and for sleeping less, we found that the impact of the pandemic was greater in younger persons. To our knowledge this is the first study demonstrating increased sleep among older participants during this crisis. This observation is in contradiction to the known association of age with the greater likelihood of having a sleep disturbance.³⁹ It is possible that financial security among the elderly lead to less worries and anxiety and hence less risk of sleep disturbance despite greater morbidity and mortality risk from COVID-19. Furthermore, in a previous influenza epidemic, high stress levels were observed disproportionately in younger adults.⁴⁰ It is possible that younger persons are more vulnerable and have lesser coping skills than older participants. Consistent with this explanation, Huang et al. have reported elevated levels of anxiety among young adults during the pandemic.⁴¹

We found that those living in urban and suburban settings were more likely to experience a change in sleep duration. Our findings are consistent with a study demonstrating greater sleep disturbances among urban Greek residents during the pandemic compared to those living in rural settings.³⁷ The fact that the first phase of the COVID-19 pandemic primarily affected the more densely populated urban and suburban areas of the United States is the most likely explanation for this finding in our study. It also is consistent with previous observations demonstrating that sleep quality generally is worse among urban residents.⁴² Rural living also may be protective by providing more opportunities for safe physical activity and exposure to natural environments resulting in less stress.³⁷

Several other sociodemographic factors appeared to play a role in altering sleep duration. Some categories of living alone were a risk factor for less sleep or more sleep suggesting that the presence of a partner mitigated some of the pandemic-related social isolation. Education was

Changes in Sleep with COVID-19

associated with sleeping more. It is likely that higher levels of education provided more opportunity to work from home.⁴³ Being unemployed or a student increased the likelihood of sleeping more; absence of “social time pressure” is a likely explanation. Being a homemaker or retired was not associated with sleeping more. The former may reflect the realities of the high workload of a homemaker. The latter may be either financially secure or are less likely to be affected by changes in income / loss of income and employment due to COVID-19. An interesting finding was the impact of time zone with participants residing in the Mountain or Central time zones less likely to sleep more. A previous study has demonstrated that Mountain or Central time zone residents sleep on average 15 minutes more per night possibly related to scheduling of television network programming (1 hour earlier in these time zones).⁴⁴ A pre-existing longer sleep duration may have made these participants less vulnerable to increasing their sleep duration further.

We observed that both sleeping more and sleeping less were highly associated with beliefs related to the impact of COVID-19 on personal health or financial status. Individuals who felt that COVID-19 would negatively impact them were more likely to have a change in sleep duration. Our results are consistent with those reported from the United Kingdom with respect to increased risk of being infected with COVID-19.⁹ The likely explanation for these observations is greater stress levels and their negative impact on sleep. However, the finding that greater worry regarding the ability to quarantine was associated with a lower association with less sleep seems counterintuitive and without an obvious explanation. Interestingly, despite reports of adverse racially motivated actions perpetrated against Asians, there was no impact of high levels of perceived discrimination on sleep duration. However, the number of Asians in the study cohort was very low, precluding us from drawing conclusions with regards to race and ethnicity.

Changes in Sleep with COVID-19

Mental health constructs appeared to be the most highly associated factors with both more sleep and less sleep. The association between depression and anxiety, and insomnia during the COVID-19 pandemic is well-established in both health care workers and the general population.^{16,33} We extend these findings by demonstrating that both depression and anxiety are related to both more sleep and less sleep. Surveys during the pandemic have also documented that loneliness occurs commonly and is related to the social isolation imposed by COVID-19 mitigation strategies.^{45,46} Our findings documenting the association between loneliness and changes in sleep duration highlight the impact of social isolation on sleep. It is possible that interventions to improve mental health during the pandemic will result in better sleep health.

A major strength of this study is that it utilized a dataset containing substantial information pertaining to sociodemographic and mental health status that might explain change in sleep duration. However, we do acknowledge some limitations. First, the study recruitment was comprised mainly of a convenience sample of online social media users. Although 70% of Americans use Facebook,⁴⁷ certain demographic groups may have been underrepresented. Nevertheless, a recent systematic review has identified Facebook as a useful tool in shortening recruitment periods, reducing costs, and enhancing representation of target populations.²⁰ Second, people with low socioeconomic status and poor access to computer or internet connection (11.2%-19.6%)⁴⁸ at home also may have been underrepresented in this study. Although we tried to recruit racial and ethnic minorities,¹⁸ we do acknowledge that our sample was predominantly white and not representative of the United States population. However, a mitigating factor is that we had respondents from every state, age group and sex. Unfortunately, given social distancing regulations caused by the COVID-19 pandemic, online data collection is one of the few platforms that is available. Third, our data regarding sleep was limited only to a single self-reported question

Changes in Sleep with COVID-19

on changes in sleep duration. It is therefore relatively non-specific and carries the risk of recall bias. In addition, we lack information on COVID-19 infection status for study participants; this maybe important information since sleep disturbances can be a consequence of viral infection. Lastly, our analyses were cross-sectional, and thus causality cannot be inferred.

In summary, we have presented data showing that during the COVID-19 pandemic, the majority of adults in the United States experience either an increase or a decrease in their sleep and that changes in sleep duration are related to a number of sociodemographic factors and beliefs regarding susceptibility to COVID-19 infection. Importantly, mental health constructs have the strongest association with sleep suggesting that in addition to prevention and treatment of COVID-19, recognizing psychosocial aspects during stressful times is crucial. Future studies are needed to examine this association between COVID-19 and neuropsychiatric problems, and to develop interventions to address these challenges across populations resulting in better sleep health.

Changes in Sleep with COVID-19

Acknowledgments

The authors respectfully acknowledge the contribution of Ms. Kristen Monten at an earlier stage of this manuscript. This work was supported in part by the NIH grant numbers: K01HL150339 (RR)

Changes in Sleep with COVID-19

References

1. Baglioni C, Nanovska S, Regen W, et al. Sleep and mental disorders: A meta-analysis of polysomnographic research. *Psychol Bull* 2016;142:969-90.
2. Scott AJ, Webb TL, Rowse G. Does improving sleep lead to better mental health? A protocol for a meta-analytic review of randomised controlled trials. *BMJ Open* 2017;7:e016873,2017-016873.
3. Morganstein JC, Ursano RJ. Ecological Disasters and Mental Health: Causes, Consequences, and Interventions. *Front Psychiatry* 2020;11:1.
4. Fan F, Zhou Y, Liu X. Sleep Disturbance Predicts Posttraumatic Stress Disorder and Depressive Symptoms: A Cohort Study of Chinese Adolescents. *J Clin Psychiatry* 2017;78:882-8.
5. Brooks SK, Webster RK, Smith LE, et al. The psychological impact of quarantine and how to reduce it: rapid review of the evidence. *Lancet* 2020;395:912-20.
6. Jeong H, Yim HW, Song YJ, et al. Mental health status of people isolated due to Middle East Respiratory Syndrome. *Epidemiol Health* 2016;38:e2016048.
7. Rogers JP, Chesney E, Oliver D, et al. Psychiatric and neuropsychiatric presentations associated with severe coronavirus infections: a systematic review and meta-analysis with comparison to the COVID-19 pandemic. *Lancet Psychiatry* 2020;7:611-27.
8. Nath A. Neurologic complications of coronavirus infections. *Neurology* 2020;94:809-10.
9. Wright L, Steptoe A, Fancourt D. Are adversities and worries during the COVID-19 pandemic related to sleep quality? Longitudinal analyses of 48,000 UK adults. *PLoS One* 2021;16:e0248919.

Changes in Sleep with COVID-19

10. Lades LK, Laffan K, Daly M, Delaney L. Daily emotional well-being during the COVID-19 pandemic. *Br J Health Psychol* 2020;25:902-11.
11. Gualano MR, Lo Moro G, Voglino G, Bert F, Siliquini R. Effects of Covid-19 Lockdown on Mental Health and Sleep Disturbances in Italy. *Int J Environ Res Public Health* 2020;17:10.3390/ijerph17134779.
12. Janati Idrissi A, Lamkaddem A, Benouajjit A, et al. Sleep quality and mental health in the context of COVID-19 pandemic and lockdown in Morocco. *Sleep Med* 2020;74:248-53.
13. Li DJ, Ko NY, Chen YL, et al. COVID-19-Related Factors Associated with Sleep Disturbance and Suicidal Thoughts among the Taiwanese Public: A Facebook Survey. *Int J Environ Res Public Health* 2020;17:10.3390/ijerph17124479.
14. Dubey S, Biswas P, Ghosh R, et al. Psychosocial impact of COVID-19. *Diabetes Metab Syndr* 2020;14:779-88.
15. Altena E, Baglioni C, Espie CA, et al. Dealing with sleep problems during home confinement due to the COVID-19 outbreak: Practical recommendations from a task force of the European CBT-I Academy. *J Sleep Res* 2020;29:e13052.
16. Pappa S, Ntella V, Giannakas T, Giannakoulis VG, Papoutsis E, Katsaounou P. Prevalence of depression, anxiety, and insomnia among healthcare workers during the COVID-19 pandemic: A systematic review and meta-analysis. *Brain Behav Immun* 2020;88:901-7.
17. Greenberg N, Gnanapragasam S, Wessely S. Managing mental health challenges faced by healthcare workers during covid-19 pandemic. *BMJ* 2020;368:m1211.

Changes in Sleep with COVID-19

18. Ali SH, Foreman J, Capasso A, Jones AM, Tozan Y, DiClemente RJ. Social media as a recruitment platform for a nationwide online survey of COVID-19 knowledge, beliefs, and practices in the United States: methodology and feasibility analysis. *BMC Med Res Methodol* 2020;20:116,020-01011-0.
19. Ali SH, Foreman J, Tozan Y, Capasso A, Jones AM, DiClemente RJ. Trends and Predictors of COVID-19 Information Sources and Their Relationship with Knowledge and Beliefs Related to the Pandemic: Nationwide Cross-Sectional Study. *JMIR Public Health Surveill* 2020;6:e21071.
20. Whitaker C, Stevelink S, Fear N. The Use of Facebook in Recruiting Participants for Health Research Purposes: A Systematic Review. *J Med Internet Res* 2017;19:e290.
21. Janz NK, Becker MH. The Health Belief Model: a decade later. *Health Educ Q* 1984;11:1-47.
22. Painter JE, DiClemente RJ, von Fricken ME. Interest in an Ebola vaccine among a U.S. national sample during the height of the 2014-2016 Ebola outbreak in West Africa. *Vaccine* 2017;35:508-12.
23. Hasan F, Khan MO, Ali M. Swine Flu: Knowledge, Attitude, and Practices Survey of Medical and Dental Students of Karachi. *Cureus* 2018;10:e2048.
24. Al-Rabiaah A, Temsah MH, Al-Eyadhy AA, et al. Middle East Respiratory Syndrome-Corona Virus (MERS-CoV) associated stress among medical students at a university teaching hospital in Saudi Arabia. *J Infect Public Health* 2020;13:687-91.
25. World Health Organization. Regional Office for Europe. Survey tool and guidance: rapid, simple, flexible behavioral insights on COVID-19: 29 July 2020. 2020:40 p.

Changes in Sleep with COVID-19

26. Kroenke K, Spitzer RL, Williams JB, Lowe B. An ultra-brief screening scale for anxiety and depression: the PHQ-4. *Psychosomatics* 2009;50:613-21.
27. Hughes ME, Waite LJ, Hawkley LC, Cacioppo JT. A Short Scale for Measuring Loneliness in Large Surveys: Results from Two Population-Based Studies. *Res Aging* 2004;26:655-72.
28. Korman M, Tkachev V, Reis C, et al. COVID-19-mandated social restrictions unveil the impact of social time pressure on sleep and body clock. *Sci Rep* 2020;10:22225,020-79299-7.
29. Robbins R, Affouf M, Weaver MD, et al. Examining nocturnal sleep episode duration before and during COVID-19 in five major metropolitan areas on three different continents. *JMIR Preprints* 21/05/2020:20546, doi: 10.2196/preprints.20546
30. Sleep Standards. Sleep Habits Post Lockdown in The U.S. (2020). Publication Date: 5/2020. <https://sleepstandards.com/sleep-habits-post-quarantine/#Demographics>. Accessed: 1/11, 2021.
31. Li Y, Qin Q, Sun Q, Sanford LD, Vgontzas AN, Tang X. Insomnia and psychological reactions during the COVID-19 outbreak in China. *J Clin Sleep Med* 2020;16:1417-8.
32. Cenat JM, Blais-Rochette C, Kokou-Kpolou CK, et al. Prevalence of symptoms of depression, anxiety, insomnia, posttraumatic stress disorder, and psychological distress among populations affected by the COVID-19 pandemic: A systematic review and meta-analysis. *Psychiatry Res* 2020;295:113599.
33. Wu T, Jia X, Shi H, et al. Prevalence of mental health problems during the COVID-19 pandemic: A systematic review and meta-analysis. *J Affect Disord* 2020;281:91-8.

Changes in Sleep with COVID-19

34. Bai YM, Lin CC, Lin CY, Chen JY, Chue CM, Chou P. Survey of stress reactions among health care workers involved with the SARS outbreak. . *Psychiatr Serv* 2004;55:1055-7.
35. Su TP, Lien TC, Yang CY, et al. Prevalence of psychiatric morbidity and psychological adaptation of the nurses in a structured SARS caring unit during outbreak: a prospective and periodic assessment study in Taiwan. *J Psychiatr Res* 2007;41:119-30.
36. McGinty EE, Presskreischer R, Anderson KE, Han H, Barry CL. Psychological Distress and COVID-19-Related Stressors Reported in a Longitudinal Cohort of US Adults in April and July 2020. *JAMA* 2020;324:2555-7.
37. Voitsidis P, Gliatas I, Bairachtari V, et al. Insomnia during the COVID-19 pandemic in a Greek population. *Psychiatry Res* 2020;289:113076.
38. Jackson CL, Powell-Wiley TM, Gaston SA, Andrews MR, Tamura K, Ramos A. Racial/Ethnic Disparities in Sleep Health and Potential Interventions Among Women in the United States. *J Womens Health (Larchmt)* 2020;29:435-42.
39. Miner B, Kryger MH. Sleep in the Aging Population. *Sleep Med Clin* 2020;15:311-8.
40. Taylor MR, Agho KE, Stevens GJ, Raphael B. Factors influencing psychological distress during a disease epidemic: data from Australia's first outbreak of equine influenza. *BMC Public Health* 2008;8:347,2458-8-347.
41. Huang Y, Zhao N. Generalized anxiety disorder, depressive symptoms and sleep quality during COVID-19 outbreak in China: a web-based cross-sectional survey. *Psychiatry Res* 2020;288:112954.

Changes in Sleep with COVID-19

42. Arrona-Palacios A, Gradisar M. Self-reported sleep duration, sleep quality and sleep problems in Mexicans adults: Results of the 2016 Mexican National Halfway Health and Nutrition Survey. *Sleep Health* 2020.

43. Dey M, Frazis H, Loewenstein MA, Sun H. Ability to work from home: evidence from two surveys and implications for the labor market in the COVID-19 pandemic. *Monthly Labor Rev* 2020.

44. Silva GE, Goodwin JL, Sherrill DL, et al. Relationship between reported and measured sleep times: the sleep heart health study (SHHS). *J Clin Sleep Med* 2007;3:622-30.

45. Perez S, Masegoso A, Hernandez-Espeso N. Levels and variables associated with psychological distress during confinement due to the coronavirus pandemic in a community sample of Spanish adults. *Clin Psychol Psychother* 2020.

46. Robb CE, de Jager CA, Ahmadi-Abhari S, et al. Associations of Social Isolation with Anxiety and Depression During the Early COVID-19 Pandemic: A Survey of Older Adults in London, UK. *Front Psychiatry* 2020;11:591120.

47. Perrin A, Anderson M. Share of US adults using social media, including Facebook, is mostly unchanged since 2018. Pew Research Center. 2019. URL: <https://www.pewresearch.org/fact-tank/2019/04/10/>

48. U.S. Census Bureau, 2019. Quick Facts. <https://www.census.gov/quickfacts/fact/table/US/AGE135219>. AGE135219. Accessed November 3, 2020.

Changes in Sleep with COVID-19

Table 1

Demographic characteristics of the sample by report of any change in sleep (i.e., either reports of sleeping more or sleeping less than usual versus no change in sleep, N=5,175)

		<u>No Change in Sleep</u>		<u>Any Change (Slept More or Slept Less)</u>		Chi-Square	P-Value
Variables		N	%	N	%		
Age	>70	313	13.1%	201	7.2%	120.2	0.000
	60-69	811	34.0%	736	26.4%		
	50-59	624	26.1%	825	29.6%		
	40-49	335	14.0%	487	17.5%		
	30-39	227	9.5%	376	13.5%		
	18-29	78	3.3%	162	5.8%		
Sex	Male	1,078	45.4%	1,048	37.8%	30.8	0.000
	Female	1,294	54.6%	1,725	62.2%		
Race	White	2,214	92.7%	2,606	93.5%	9.8	0.007
	Black, Hispanic, Asian	73	3.1%	104	3.7%		
	Other	101	4.2%	77	2.8%		
Child	No children at home	1,825	76.4%	2,004	71.9%	13.5	0.000
	Child at home	563	20.3%	783	28.1%		
Urban/Rural	Rural	786	34.6%	772	28.8%	19.9	0.000
	Urban	1173	51.6%	1473	55.0%		
	Suburban	315	13.9%	433	16.2%		
Marital Status	Married/cohabiting	1,686	74.1%	1,839	68.7%	27.3	0.000
	Single	316	13.9%	480	17.9%		
	Widowed	106	4.7%	102	3.8%		
	Divorced/separated	166	7.3%	257	9.6%		
Education	High School Diploma or GED	257	11.4%	218	8.3%	14.1	0.000
	Some College or Higher	1,989	88.6%	2,424	91.7%		
Work	Employed	1,222	53.7%	1,577	58.9%	121.6	0.000
	Unemployed	123	5.4%	270	10.1%		
	Student	14	0.6%	47	1.8%		
	Retired	726	31.9%	547	20.4%		
	Unable to work	92	4.0%	142	5.3%		
	Housework	97	4.3%	95	3.5%		
Time Zone	Pacific	251	11.1%	327	12.3%	14.2	0.002
	Mountain	185	8.2%	178	6.7%		
	Central	606	26.8%	620	23.3%		
	Eastern	1221	54.0%	1539	57.8%		

Changes in Sleep with COVID-19

Table 2
Binary Logistic Regression Models Examining Sociodemographic Predictors of Sleeping Less than Usual and Sleeping More Than Usual (N=5,175).

Sociodemographic Variables		<i>Sleeping Less Than Usual (N=884)</i>				<i>Sleeping More Than Usual (N=1903)</i>			
		<i>P-</i>				<i>P-</i>			
		<i>OR</i>	<i>Value</i>	<i>Lower</i>	<i>Upper</i>	<i>OR</i>	<i>Value</i>	<i>Lower</i>	<i>Upper</i>
Age	>70	<i>Reference</i>							
	60-69	1.33	0.117	0.93	1.89	1.23	0.103	0.96	1.57*
	50-59	2.05	0.000	1.39	3.02	1.48	0.006	1.12	1.96
	40-49	2.24	0.000	1.45	3.46	1.58	0.005	1.15	2.17
	30-39	2.92	0.000	1.86	4.58	1.62	0.005	1.15	2.27
	18-29	2.10	0.013	1.17	3.78	2.18	0.000	1.42	3.36
Sex	Male	<i>Reference</i>							
	Female	1.86	0.000	1.56	2.22	1.20	0.007	1.05	1.37
Race	White	<i>Reference</i>							
	Black, Hispanic,								
	Asian	1.15	0.551	0.73	1.79	1.01	0.974	0.70	1.45
	Other	0.72	0.154	0.45	1.13	0.55	0.002	0.37	0.80
Child	No children	<i>Reference</i>							
	Child at home	1.06	0.605	0.85	1.32	1.02	0.822	0.86	1.22
Urban/Rural	Rural	<i>Reference</i>							
	Urban	1.32	0.002	1.11	1.63	1.16	0.049	1.00	1.34
	Suburban	1.35	0.027	1.03	1.77	1.24	0.038	1.01	1.52
Marital Status	Married/cohabiting	<i>Reference</i>							
	Single	1.09	0.468	0.86	1.40	1.27	0.012	1.05	1.53
	Widowed	1.18	0.440	0.77	1.81	1.05	0.781	0.75	1.46
	Divorced/separated	1.39	0.028	1.04	1.87	1.46	0.002	1.16	1.85
Education	High School	<i>Reference</i>							
	Diploma or GED								
	College or Higher	0.69	0.072	0.46	1.03	1.51	0.000	1.21	1.88
Work	Employed	<i>Reference</i>							
	Unemployed	1.21	0.250	0.87	1.68	1.80	0.000	1.41	2.30
	Student	1.55	0.356	0.61	3.96	2.67	0.008	1.29	5.55
	Retired	0.82	0.128	0.63	1.06	0.77	0.008	0.63	0.93
	Unable to work	1.37	0.100	0.94	2.01	1.11	0.524	0.81	1.52
	Housework	0.69	0.072	0.46	1.03	0.61	0.007	0.43	0.87
Time Zone	Pacific	<i>Reference</i>							
	Mountain	0.88	0.196	0.72	1.07	0.80	0.004	0.68	0.93
	Central	0.89	0.481	0.65	1.23	0.72	0.012	0.56	0.93
	Eastern	1.03	0.829	0.79	1.35	1.03	0.798	0.84	1.26

* χ^2 for trend=73.9, $p<0.0001$

Table 3

Binary Regression Examining the Relationships Between Sleeping Less or Sleeping More Than Usual and COVID-19 Beliefs in Adjusted and Unadjusted Models (N=5,175).

COVID-19 Beliefs [†]	<i>Sleeping Less than Usual (N=884)</i>							
	<i>Unadjusted</i>				<i>Adjusted</i>			
	<i>OR</i>	<i>P-</i>	<i>Lower</i>	<i>Upper</i>	<i>OR</i>	<i>P-</i>	<i>Lower</i>	<i>Upper</i>
		<i>Value</i>				<i>Value</i>		
High perceived risk for COVID-19 (Reference: low perceived risk)	1.41	0.000	1.20	1.66	1.22	0.021	1.03	1.45
High perceived severity of COVID-19 symptoms (Reference: low perceived severity)	1.09	0.281	0.93	1.28	1.11	0.221	0.94	1.33
High COVID-19 financial concerns regarding ability to quarantine (Reference: low financial concerns)	0.71	0.000	0.59	0.84	0.80	0.021	0.66	0.97
High perceived discrimination due to COVID-19 (Reference: low perceived discrimination)	1.10	0.391	0.88	1.38	1.15	0.246	0.91	1.47
Beliefs COVID-19 is deadlier than the flu (Reference: high disagreement with this belief)	2.19	0.000	1.78	2.69	1.94	0.000	1.56	2.42
Beliefs COVID-19 is not as big a problem as the media suggest (Reference: high disagreement with this belief)	0.46	0.000	0.39	0.55	0.51	0.000	0.42	0.62
Beliefs COVID-19 is a bigger problem than the government suggests (Reference: high disagreement with this belief)	2.12	0.000	1.80	2.49	2.00	0.000	1.68	2.38
COVID-19 Beliefs	<i>Sleeping More than Usual (N=1903)</i>							
	<i>Unadjusted</i>				<i>Adjusted</i>			
	<i>OR</i>	<i>P-</i>	<i>Lower</i>	<i>Upper</i>	<i>OR</i>	<i>P-</i>	<i>Lower</i>	<i>Upper</i>
		<i>Value</i>				<i>Value</i>		
High perceived risk for COVID-19 (Reference: low perceived risk)	1.28	0.000	1.13	1.45	1.17	0.018	1.03	1.33
High perceived severity of COVID-19 symptoms (Reference: low perceived severity)	1.21	0.003	1.06	1.37	1.27	0.001	1.11	1.45
High COVID-19 financial concerns regarding ability to quarantine (Reference: low worries)	0.80	0.002	0.70	0.92	0.87	0.069	0.75	1.01

Changes in Sleep with COVID-19

High perceived discrimination due to COVID-19 (Reference: low perceived discrimination)	0.90	0.277	0.75	1.08	0.94	0.541	0.78	1.14
Beliefs COVID-19 is deadlier than the flu (Reference: high disagreement with this belief)	1.94	0.000	1.67	2.26	1.90	0.000	1.62	2.23
Beliefs COVID-19 is not as big a problem as the media suggest (Reference: high disagreement with this belief)	0.64	0.000	0.56	0.73	0.66	0.000	0.58	0.76
Beliefs COVID-19 is a bigger problem than the government suggests (Reference: high disagreement with this belief)	1.72	0.000	1.52	1.94	1.72	0.000	1.51	1.96

Notes.

Adjusted models control for sociodemographic factors age, sex, education, marital status, work status, and time zone.

Bold indicates significance at the <.05 level.

†Responses to perceived risk [“On a scale from 0-10, what do you think is your risk of getting infected with Coronavirus?”] and perceived severity [“On a scale from 0-10, If you were infected with Coronavirus, how severe do you think it would be?”] were recoded so responses from 0 (Not Severe) to the scale midpoint (5) represented “low risk” and “low severity” responses and those responses from above the scale mid-point (6) to 10 (Very Severe) were re-coded to represent “high risk” and “high severity.” Responses to financial ability to quarantine [“I can financially afford to self-quarantine”], beliefs about discrimination [“Since the Coronavirus outbreak, I feel discriminated against”], beliefs that COVID-19 is deadlier than the flu [“Coronavirus is more deadly than the seasonal flu”], beliefs COVID-19 is not as big a problem as the media suggest [“Coronavirus is not as big of a problem as the media suggests”], and beliefs that COVID-19 is a bigger problem than the government suggests [“Coronavirus is a bigger problem than the government suggests”] were captured on scales from 0 “strongly disagree” to 3 “strongly agree.” Responses were dichotomized so that “strongly disagree” and “disagree” were re-coded to 0 “low” and responses of “disagree” or “strongly disagree” were re-coded to “high,” with the exception of financial ability to quarantine, which was re-coded so that “strongly agree” and “agree” were” represented low financial worries and “strongly disagree” and “disagree” represented high financial worries.

Table 4
Binary Regression Variables Examining the Relationships Between Sleeping Less or Sleeping More Than Usual and Mental Health in Adjusted and Unadjusted Models.

		<i>Sleeping Less than Usual</i>							
		<i>Unadjusted</i>				<i>Adjusted</i>			
Mental Health		<i>OR</i>	<i>P-Value</i>	<i>Lower</i>	<i>Upper</i>	<i>OR</i>	<i>P-Value</i>	<i>Lower</i>	<i>Upper</i>
Anxiety		5.29	0.000	4.17	6.70	4.52	0.000	3.52	5.79
<i>(Reference: no anxiety)</i>									
Depression		4.23	0.000	3.51	5.09	3.86	0.000	3.17	4.70
<i>(Reference: no depression)</i>									
Loneliness		3.18	0.000	2.63	3.84	2.67	0.000	2.18	3.27
<i>(Reference: no loneliness)</i>									
		<i>Sleeping More Than Usual</i>							
		<i>Unadjusted</i>				<i>Adjusted</i>			
Mental Health		<i>OR</i>	<i>P-Value</i>	<i>Lower</i>	<i>Upper</i>	<i>OR</i>	<i>P-Value</i>	<i>Lower</i>	<i>Upper</i>
Anxiety		2.80	0.000	2.42	3.24	2.63	0.000	2.25	3.07
<i>(Reference: no anxiety)</i>									
Depression		2.60	0.000	2.29	2.96	2.48	0.000	2.16	2.84
<i>(Reference: no depression)</i>									
Loneliness		2.16	0.000	1.89	2.46	1.92	0.000	1.67	2.21
<i>(Reference: no loneliness)</i>									

Supplement

1. Sex
 - a. Female
 - b. Male
 - c. Other
 - d. Prefer not to disclose
2. Age group
 - a. 18-29 years old
 - b. 30-39 years old
 - c. 40-49 years old
 - d. 50-59 years old
 - e. 60-69 years old
 - f. 70-79 years old
 - g. 80+ years old
3. What's your race/ethnicity?
 - a. Black, Non-Hispanic
 - b. White, Non-Hispanic
 - c. Asian/Pacific Islander
 - d. Hispanic/Latinx
 - e. Native American or American Indian
 - f. Interracial, Mixed race, or Other
4. What is your marital or cohabitation status?
 - a. Married/cohabitating
 - b. Single
 - c. Widowed
 - d. Divorced/separated
5. How would you describe where you live?
 - a. Urban
 - b. Suburban
 - c. Rural
6. What is your current working status?
 - a. Employed, full time
 - b. Employed, part time
 - c. Unpaid work (e.g., homemaker, eldercare, childcare)
 - d. Self-employed
 - e. Out of work and looking for work
 - f. Out of work but not currently looking for work
 - g. Student
 - h. Military personnel
 - i. Retired

Changes in Sleep with COVID-19

- j. Unable to work
7. What is your highest level of education?
- Less than a High School diploma
 - High School diploma or GED
 - Some college, no degree
 - Associate's degree
 - Bachelor's degree
 - Masters/Professional degree or above
8. In which state do you currently reside?
[Drop-down menu of all 50 states, District of Columbia, Puerto Rico]
9. What do you think is your risk of getting infected with the Coronavirus?
Not at all likely											Extremely likely
0	1	2	3	4	5	6	7	8	9	10	
10. If you were infected with the Coronavirus, how severe do you think it would be?
Not Severe											Very Severe
0	1	2	3	4	5	6	7	8	9	10	
11. I can financially afford to self-quarantine.
- Strongly Agree
 - Agree
 - Disagree
 - Strongly Disagree
12. Since the Coronavirus outbreak, I feel discriminated against.
- Strongly Agree
 - Agree
 - Disagree
 - Strongly Disagree
13. Coronavirus is more deadly than the seasonal flu.
- Strongly Agree
 - Agree
 - Disagree
 - Strongly Disagree
14. Coronavirus is not as big of a problem as the media suggests.
- Strongly Agree
 - Agree
 - Disagree
 - Strongly Disagree

Changes in Sleep with COVID-19

15. Coronavirus is a bigger problem than the government suggests.

- a. Strongly Agree
- b. Agree
- c. Disagree
- d. Strongly Disagree

In the last 3 months, how often have you felt:

16. Left out

- a. Never
- b. Rarely
- c. Sometimes
- d. Often

17. Isolated from others

- a. Never
- b. Rarely
- c. Sometimes
- d. Often

18. That you lack companionship

- a. Never
- b. Rarely
- c. Sometimes
- d. Often

Over the last 7 days, how often have you been bothered by any of the following problems because of THE CORONAVIRUS OUTBREAK?

19. Feeling nervous, anxious, or on edge?

- a. Not at all
- b. Several days
- c. More than half the days
- d. Nearly everyday

20. Not being able to stop or control worrying?

- a. Not at all
- b. Several days
- c. More than half the days
- d. Nearly everyday

21. Feeling down, depressed, or hopeless?

- a. Not at all
- b. Several days
- c. More than half the days
- d. Nearly everyday

Changes in Sleep with COVID-19

22. Little interest or pleasure in doing things (that I used to enjoy)?

- a. Not at all
- b. Several days
- c. More than half the days
- d. Nearly everyday

Since hearing about the Coronavirus outbreak, how have the below behaviors changed for you?

23. Sleeping

- a. Much more
- b. Little more
- c. Not changed
- d. Little less
- e. Much less
- f. Not applicable