Beneath the surface: mental health, and harassment and abuse of athletes participating in the FINA (Aquatics) World Championships, 2019

Margo Mountjoy^{1,2,3}, Astrid Junge^{4,5}, Christer Magnusson², Farhad Moradi Shahpar², Edgar Ivan Ortiz Lizcano², Josip Varvodic², Xinzhai Wang², Mohamed Yahia Cherif², Lee Hill¹, Jim Miller^{2,6}

From:

1 McMaster University Hamilton, Hamilton, Canada
2 Federation Internationale de Natation (FINA), Lausanne, Switzerland
3 International Olympic Committee (IOC), Games Group and Working Group on the Prevention of Harassment and Abuse in Sport Lausanne, Switzerland
4 Medical School University of Hamburg, Hamburg, Germany
5 Swiss Concussion Centre at Schulthess Clinic Zürich, Switzerland
6 Medicine Department, University of Virginia, Charlottesville, and Virginia Commonwealth University, Richmond, Virginia, USA

Contact for corresponding author:

Dr. Margo Mountjoy
McMaster Medical School
10-B Victoria Street South
Kitchener, ON, Canada N2G 1C5
Email: mmsportdoc@mcmaster.ca
Phone: +1 226 971 2940

Word Count: 3130

Abstract word count 249

Number of Tables and Figures: 3

Original Research: Sport Medicine + Sport Psychology

Patient and Public Involvement:

Athlete representatives and members of the FINA Athletes Commission were involved in determining the research priority of mental health and safeguarding, as

well as actively recruiting participants and promoting the research project during the FINA World Championships. We will invite athletes and the entourage to help with dissemination and application of the results.

Scientific contributions of each author:

Margo Mountjoy, MD, PhD: Substantial contributions to the conception and design of the study, collection and interpretation of data, drafting, writing, and revising of the manuscript, and approval of the final version to be published.

Astrid Junge, PhD: Substantial contributions to the conception and design of the study, analysis and interpretation of data, drafting, writing, and revising of the manuscript, and approval of the final version to be published.

Christer Magnusson, MD, Farhad Moradi Shapar, MD, PhD, Edgar Ivan Ortiz Lizcano, MD, Josip Varvodic, MD, Xinzhai Wang, MD, Mohamed Yahia Cherif MD: Contributions to data collection and approval of the final version to be published.

Lee Hill, PhD: Substantial contributions to data collection and approval of the final version to be published.

Jim Miller, MD, FAAFP/Sports Medicine: Substantial contributions to conception of the study, as well as data collection and approval of the final version to be published.

All authors agree to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

Funding: The data collection was funded by the Federation Internationale de Natation (FINA), Lausanne, Switzerland. FINA had no role in the study design, the collection, analysis, and interpretation of the data, writing the report, or the decision

61 to submit the article for publication. No other funding was received for the present
62 study.

63

64 **Competing Interest:** None declared.

65

66 **Ethics approval:** The present study had ethic approval from McMaster University -
67 Hamilton Integrated Research Ethics Board, Canada.

68

69 **No Additional data:** No supplementary data.

70

71 **Abbreviations:**

72 AS: Artistic swimming

73 BEDA-Q: Brief Eating Disorders in Athletes Questionnaire

74 CES-D-10: Center for Epidemiological Studies Depression Scale revised

75 DV: Diving

76 FINA: Federation Internationale de Natation

77 HD: High Diving

78 IOC: International Olympic Committee

79 OWS: Open water swimming

80 SW: Swimming

81 WC: World Championships

82 WP: Water polo

83

84 ABSTRACT

85

86 **Objective:** To assess the mental health and experience of sport-related harassment
87 and abuse of athletes participating in the FINA World Championships 2019, and to
88 analyse it in relation to gender and the aquatic disciplines.

89 **Design:** Cross-sectional study using an anonymous questionnaire.

90 **Methods:** During the Championships, registered athletes (swimmers, divers, high
91 divers, water polo players, artistic swimmers, open-water swimmers) completed a
92 survey including the main outcome measures of depression (CES-D-10), eating
93 disorders (BEDA-Q), the subjective need for psychotherapeutic support, and the
94 experience of harassment and/or abuse in their sports environment.

95 **Results:** A quarter of the athletes (n=62, 24.6%) were classified as depressed and
96 more than a third (n=111, 35.0%) as having an eating disorder. More than 40% of the
97 athletes stated that they wanted or needed psychotherapeutic support for mental
98 health problems. Fifty-one athletes (14.9%) had experienced harassment/abuse in
99 sport themselves, and 31 (9%) had witnessed it in another athlete. The experiences
100 of harassment and abuse ranged from unwanted comments about body or
101 appearance (40.2%) to rewards in sport for sexual favours (2.5%) and rape (0.3%).
102 Athletes who had experienced harassment/abuse in sport themselves had higher
103 average scores for depression and eating disorders, and more of them needed
104 psychotherapeutic support. Up to a third would not talk to anybody if they saw or
105 experienced harassment/abuse, and less than 20% would talk to an official for help.

106 **Conclusion:** Targeted initiatives are required to address the burden of mental health
107 issues and harassment and abuse in sport in the FINA aquatic disciplines.

108

109 **Keywords:** aquatics, elite sports, mental health, depression, harassment + abuse

INTRODUCTION

"I didn't want to be in the sport anymore ... I didn't want to be alive anymore." Michael Phelps 2018¹

The sport community was shocked when Michael Phelps, the most decorated Olympian of all time, first spoke about his struggles with mental health issues in 2016.² A systematic review and meta-analysis of mental health of elite athletes revealed a prevalence of 33.6% for anxiety/depression, 26.4% sleep disturbance, 19.6% symptoms of distress, 9% for alcohol misuse, and 1%–28% for adverse eating habits or eating disorder.³ There is a well-established correlation between mental health symptomatology and experience with harassment/abuse both inside and outside of sport.⁴ In athletics, a cross-sectional study of 198 elite Swedish athletes, suicidal ideation of the female athletes was associated with having been sexually abused combined with signs of psychological vulnerability.⁵ A cohort study on Canadian national team athletes demonstrates statistically significant correlations between harassment/abuse in sport and mental health outcomes defined as self-harm, disordered eating and eating disorders, suicidal thoughts and seeking help for mental health issues.⁶

While the original IOC publication on sexual harassment/abuse in sport was published in 2007,⁷ the IOC first implemented safeguarding against harassment/abuse at the 2016 Olympic Games,⁸ and published an updated consensus statement.⁹ A research project was conducted at the Youth Olympic Games 2018 to assess the participating elite youth athletes' understanding and perception of harassment/abuse in their respective sports. About a third stated that it 'very likely' or 'likely' occurred in their sport.¹⁰ This study is the only publication on harassment/abuse in a large cohort of international multi-sport elite athletes.

While there has been a recent increase in scientific publications on mental health and harassment/abuse in sport, there remains a dearth of scientific evidence of the elite aquatic athletes' experience in these domains. In order to address this gap in knowledge, the objectives of this study were to assess the athletes' mental health problems and their experience of sport-related harassment/abuse, and to analyse these findings in relation to gender and the aquatic disciplines.

MATERIALS AND METHODS

An anonymized survey for accredited athletes was performed during the FINA World Championships (WC), 2019 using the same recruitment and data collection procedures as in previous studies.¹¹⁻¹³

The **questionnaire** started with questions on gender, age, and aquatic discipline, followed by two questions on the extent of health complaints and their effect on performance or participation in the four weeks prior to the WC (based on the Oslo Trauma Research Centre Questionnaire¹⁴).

Mental health was assessed using the 10-item version of the Center for Epidemiological Studies Depression Scale revised (CESD-R-10),^{15,16} the Brief Eating Disorders in Athletes Questionnaire (BEDA-Q)¹⁷ and the question "Have you ever wanted or needed support from a psychotherapist for personal or mental health problems?" (no/ yes, previously / yes, currently).¹⁸

The section on **harassment and abuse** was newly developed based on the review of the literature.^{10,19} It consisted of two introductory questions „Do you think harassment and/or abuse occurs in your sport?" (no/not likely/not sure/likely/very likely) and „Have you ever witnessed or experienced yourself any forms of harassment or abuse in your sports environment?" (no/yes, experienced myself/yes, witnessed in another athlete) followed by a table with 13 situations, for each the

athlete should report whether or not they had witnessed or experienced it themselves, and if so, who did it (coach/another athlete/others). For details see Table 1.

The data were processed Excel (Microsoft Office 2011) and SPSS (V.23, IBM). The results are reported using means with standard deviation or frequencies with percentage, and in accordance with STROBE guidelines. We applied ANOVA, t- and Chi²-test statistical methods with an accepted significance at $p < 0.05$.

Support for travel to the FINA WC for data collection was funded by the FINA. Although FINA approved the project, it had no oversight or influence over the data collection, analysis, interpretation of results or drafting of the manuscript. Ethical design followed the principles of the Declaration of Helsinki for experiments involving humans, including informed consent and observation of privacy rights; ethical approval was granted by the Hamilton Integrated Research Ethics Board.

RESULTS

A total of 377 athletes participating in the 2019 FINA WC answered the questionnaire (response rate: 14.8%). The 229 (61.1%) female and 146 (38.9%) male athletes were on average 22.4 years old (SD=5.37; range: 14-56), with female participants (21.4, sd=4.71) being significantly younger than male (23.9; sd=5.93; $t=4.48$, $p<.001$). The majority of participants were swimmers (SW, $n=132$, 36.2%), water polo players (WP, $n=82$, 22.5%) and artistic swimmers (AS, $n=79$, 21.6%) followed by divers (DV, $n=43$, 11.8%), high divers (HD, $n=21$, 5.8%) and open water swimmers (OWS, $n=8$, 2.2%).

In the 4 weeks prior to the WC, half of the athletes had no symptoms/complaints ($n=181$), nor any ***injury, illness or other health*** problems that affected their performance or participation ($n=187$; 51.8%). About a third reported

minor health problems (n=125; 34.5%), 40 (11.0%) moderate and 16 (4.4%) severe. More than a quarter of the athletes (n=100; 27.7%) reported minor, 51 (14.2%) moderate and 16 (4.4%) major effects of health problems on their performance; seven (1.9%) could not participate at all.

Based on their answers to the CES-D-10 (n=268, mean=7.38, sd=4.75, range 0-25), a quarter of the athletes (n=62, 24.6%) were classified as **depressed** without significant differences between genders. The depression score correlated significantly with the extent of symptoms/health complaints ($r=.30$, $p<.001$) and their effect on performance in the four weeks prior to the WC ($r=.28$, $p<.001$).

Based on their answers to the six BEDA-Q questions (n=317, mean=0.67, sd=0.60, range 0-3), more than a third of the athletes (n=111, 35.0%) had an **eating disorder**. The proportion of athletes with an eating disorder differed significantly between females and males (40.2% vs 27.3%; $\chi^2=5.5$, $p<.05$). Athletes with an eating disorder had a significantly higher depression score ($F=4.8$; $p<.05$). Thirty athletes (11.6%) were classified to have both depression and eating disorder. About a quarter of the athletes (n=90, 25.1%) stated that they were *trying to lose weight now* (i.e. during the WC), with significantly more females than males (31.7% vs 14.8%; $\chi^2=12.6$, $p<.001$). Sixty athletes (17.4%) had tried to lose weight more than five times.

More than 40% of the athletes stated that they “wanted or needed **psychotherapeutic support** for personal or mental health problems” previously (n=103, 28.8%) or currently (n=57, 15.9%) without differences across genders or ages. Athletes with and without a subjective need for psychotherapeutic support differed significantly ($F=9.9$, $p<.001$) in their average depression score. About half of the athletes (53.3%) who currently wanted or needed psychotherapeutic support for

personal or mental health problems were classified as depressed. No differences were observed regarding eating disorders.

About a third of the athletes thought that **harassment/abuse** occurs in their sport 'likely' (n=83, 23.9%) or 'very likely' (n=42, 12.1%). Seventy-five athletes (21.6%) were 'unsure', 43 (12.4%) regarded this as 'unlikely', and 105 athletes (30.2%) thought that 'no' harassment/abuse occurs in their sport. The answers differed significantly between genders ($\chi^2=17.9$, $p<.01$), with males regarding it more likely than females. Almost 80% of the athletes (n=273, 79.6%) stated that they have never witnessed or experienced themselves any forms of harassment/abuse in their sport environment. Fifty-one athletes (14.9%) had experienced it themselves, and 31 (9.0%) had witnessed it in another athlete (percentages include 12 athletes who had both) without difference between genders. Athletes with different experiences of harassment/abuse differed significantly ($F=13.4$, $p<.001$) on how likely they think it occurs in their sport. Athletes who had not witnessed nor experienced harassment/abuse regarded it as less likely than athletes who had witnessed it in another athlete or experienced it themselves, and those who had both experienced and witnessed it regarded it as most likely. Athletes with different experiences of harassment/abuse differed significantly in their average score of depression ($F=4.6$, $p<.01$, see Figure 1) and eating disorder screening tools ($F=5.0$, $p<.01$, see Figure 1) as well as on the subjective need for psychotherapeutic support ($\chi^2=15.4$, $p<.05$). Athletes who experienced harassment/abuse themselves *and* witnessed it in another athlete had average lower values of depression and eating disorders than those who exclusively experienced it themselves (see Figure 1).

Insert Figure 1 here

240 From a list of 13 *situations of harassment/abuse in the sports environment*
 241 (see Table 1) 'unwanted comments about body or appearance' was the most
 242 frequently (40.2%) and 'rewards in sport for sexual favours' (2.5%) and 'rape'
 243 (reported as 'others' n=1, 0.3%) the least frequently witnessed or experienced
 244 situation. In general, situations were more frequently witnessed in another athlete
 245 than experienced themselves, except for 'inappropriate training', 'unsafe training
 246 conditions', and 'forced dieting, limited or no access to needed water or food'. About
 247 10% of the athletes had experience these situations, which were reported to have
 248 been perpetrated most often by the coach. On average, one in twenty athletes (5.0%)
 249 had experienced themselves 'unwanted and unnecessary touching of the body'
 250 mostly by the coach or another athlete. Three athletes (0.8%) had experienced
 251 'rewards in sport for sexual favours' themselves, and a further six (1.6%) had
 252 witnessed this in another athletes.

253

254 Insert Table 1 here

255

256 Significant more female than male athletes reported to have experienced
 257 harassment/abuse themselves with regard to 'unwanted comments about body or
 258 appearance' (38.5% vs 15.1%, $\chi^2=19.1$, $p<.001$), 'forced dieting, limited or no
 259 access to needed water or food' (11.8% vs 1.5%, $\chi^2=12.3$, $p<.001$), 'bullying /
 260 cyber bullying' (16.7% vs 5.3%, $\chi^2=9.4$, $p<.01$), 'unwanted and unnecessary
 261 touching of the body' (7.3% vs 2.2%, $\chi^2=4.2$, $p<.01$). About half of the athletes
 262 (n=181, 53.9%) reported they would talk to friends or family if they saw or
 263 experienced a situation like the ones listed in Table 1, but less than one in five (n=61,
 264 18.8%) would talk to an official contact to seek help. About one in eight athletes
 265 (n=43, 12.8%) would not talk to anybody, and 68 (20.2%) were not certain whether or

not to share the experience. The answers differed significantly between genders ($F=22.7$, $p<.001$). More than twice as many males than females (19.7% vs 8.7%) would not talk about it, and more females than males would talk to their friends and family (57.0%vs 33.9%).

Specific analysis combining gender and discipline were conducted for SW, WP, DV/HD (DV and HD combined) and female AS (see Table 2), since just three female and five male OWS and two male AS responded the survey.

Insert Table 2 here

Gender differences within a discipline, did not reach statistical significance due to the small subgroups, except for 'trying to lose weight now' in DV/HD ($\chi^2=6.6$, $p<.05$), need for psychotherapy in WP ($\chi^2=11.8$, $p<.01$), 'talk about experience/witnessed harassment/abuse' in SW ($\chi^2=20.6$, $p<.001$), and 'experienced themselves forced dieting, limited or no access to needed water or food' in SW ($\chi^2=10.1$, $p\leq.001$). However, in several variables, a substantially higher proportion of female than male athletes was observed, e.g. athletes with an eating disorder in WP and DV/HD, 'try to lose weight now' in SW, experienced harassment/abuse themselves in SW and HD (see Table 2). Further, it is interesting to note that a similar proportion of female and male SW reported a current need for psychotherapy, and that in WP more males than females had experienced harassment/abuse themselves. The latter concerns specifically 'inappropriate training' and 'intentional hitting, beating, or kicking (not fouls or rough play)'.

Comparing *female athletes* competing in AS, SW, DV/HD and WP, significant differences were observed with regard to eating disorders ($\chi^2=8.4$, $p<.05$), 'try to lose weight now' ($\chi^2=11.0$, $p<.05$) and the need for psychotherapeutic support

($\chi^2=15.6$, $p<.05$). Also, age ($F=8.1$, $p<.001$) and the perception of the occurrence of harassment/abuse in their sport ($F=5.8$, $p<.01$) differed significantly between the four disciplines (see Table 2). Significant differences between disciplines were observed for female athletes who experiences themselves 'unwanted comments about body or appearance' ($\chi^2=27.8$, $p<.001$) ranging from AS ($n=39$, 55.7%) to WP ($n=7$; 14.9%), and 'inappropriate training' ($\chi^2=8.1$, $p<.05$) ranging from AS ($n=13$, 18.8%) to DV/HD ($n=0$). 'Forced dieting, limited or no access to needed water or food' ranged from AS ($n=13$, 17.1%) to DV/HD ($n=1$, 3.8%) but the difference did not reach statistical significance ($\chi^2=6.5$, $p<.09$) (see Table 2). In five of the 13 situations of harassment/abuse, female AS had the highest percentage of affected athletes, and in four female SW.

Comparing *male athletes* between three disciplines, age ($F=6.6$, $p<.01$), the perception of the occurrence of harassment/abuse in their sport ($F=4.6$, $p<.05$) and the proportion of athletes who had witnessed or experienced harassment/abuse ($\chi^2=13.5$, $p<.05$) differed significantly (see Table 2). Further differences, such as more SW than WP or DV/HD were classified as depressed (31.9% vs 12.5% resp 15.4%) and needed psychotherapeutic support currently (23.3% vs 15.2% resp. 9.7%), were not statistically significant. Also differences in the 13 situations of harassment/abuse did not reach significance between disciplines in male athletes, most probably because of small numbers (see Table 2).

DISCUSSION

This study is the first to evaluate mental health and harassment/abuse in elite aquatic athletes. About a quarter of athletes were classified as **depressed**, which is consistent with previous publications.²⁰⁻²² In the general population and other studies on athletes,^{21,23} more female than male athletes reported symptoms of depression.

This also applies to the present study, except for swimmers. The prevalence of depression in male swimmers was about twice as high as in male athletes of the other aquatic disciplines. Our study demonstrated a significant correlation between the depression score and the athletes *i)* symptoms/health complaints and *ii)* their effect on performance in the four weeks prior to the WC. The intersection between injury and mental health is multi-faceted as elite sport can have stressors that increase the risk of both mental health disorders and injury. In addition, injury may trigger mental health symptoms while conversely, mental health disorders may prolong recovery from injury.⁴

More than a third of the athletes were screened as having an **eating disorder**, with a higher proportion of female (40%) than male athletes (27%), and highest in AS (54.1%). This finding is not surprising as AS is an aesthetic judged sport, where a quintessential lean body composition is implicitly valued in the culture of the sport.²⁴ Interestingly, in SW, there were equal proportions of female and males (28.1% vs 26.4%) with eating disorders which could be related to the gender-neutral importance of body composition on performance (decreasing drag), rather than the more common culturally-driven body image reasons that more often influence female than male athletes. Overall, a quarter of the athletes stated that they were trying to lose weight during the WC, however this applied to more than 40% of females in the two judged disciplines (AS, DV/HD). This finding is disturbing with almost half of the competing athletes in the highest competition level in the world engaging in weight loss behaviours that could affect their nutritional fuelling of performance. It also has significant medical implications for team physicians with the associated risks of weight loss during highly competitive sport. Interestingly, in WP, the male and female reporting of current weight loss behaviours was similar (about 20%), which is high for males and low for females. The estimated prevalence of eating disorders and/or

disordered eating among athletes in general, ranges from 0% to 19% in males and from 6% to 45% in females.²⁵

In the present study, 45% of participating athletes wanted or needed **psychotherapeutic support** for personal or mental health problems either previously (29%) or currently (16%). Similar results were reported in a survey on elite female football players (24% previous; 16% currently).¹⁸ Consistent with the high prevalence of depression in male swimmers, approximately 25% of both female and male swimmers were *currently* wanting or needing professional mental health support. These findings underscore the need for team physicians of elite athletes to have the core clinical competency to diagnose and treat mental health disorders²⁶, and for organizing committees and International Federations to incorporate mental health professionals into the medical care services for large multi-sport events.²⁷

Over one-third (36%) of athletes participating in the present study stated that **harassment/abuse** was either 'very likely' or 'likely' to occur in their sport. Similar percentages were found in youth athletes participating in the Youth Olympic Games 2018.¹⁰ In the present study 15% of the athletes reported to have experienced harassment/abuse in their sports environment themselves. For specific situations, the frequency ranged from 'unwanted comments about body or appearance' (40.2%) to 'rewards in sport for sexual favours' (2.5%) and 'rape' (0.3%). A study of German and Belgian-Dutch elite athletes found a prevalence of psychological (72%), physical (25%) and sexual harassment/abuse (31%) with significant overlap of types.²⁸ A cross-sectional study of 198 elite Swedish athletics athletes reported a sexual abuse prevalence of 11% (females 16% and males 4%) and physical abuse of 18% (females 14% and males 23%).²⁹ A survey of 1001 elite Canadian athletes from 64 sports reported a prevalence of 59% for emotional harm, 67% for neglect, 20% for sexual harm and 12% for physical harm.⁶ In the present study, athletes who have

experienced harassment or abuse in their sports environment themselves had higher average scores of the depression and of eating disorders, and more of them needed psychotherapeutic support than those who have not witnessed nor experienced it. While our findings are consistent with reports in the literature, they remain alarming. Validated prevention strategies to decrease harassment/abuse in the aquatic sports are urgently required.

In this survey, only 19% stated that they would **officially report harassment/abuse**, while 13% would say nothing, and 20% were unsure what to do. These numbers are consistent with the Canadian study who found that those athletes who had experienced harassment/abuse, 56% did not report their experience, and 84% did not submit a formal complaint.⁶ These findings demonstrate the need *i)* to encourage athletes to report their abuse, *ii)* to develop safe reporting mechanisms, and *iii)* to ensure experienced athlete support systems are accessible.

There are **limitations** to this study inherent in the retrospective design including selection and recall bias. In addition, the response rate was low which could result in an over- or underestimation of the prevalence.³⁰ However, we believe that a selection bias is unlikely, since responses to the questions on health complaints in the 4 weeks prior to the WC were similar to the athlete surveys in FINA WC 2013.¹¹⁻

¹³

CONCLUSIONS

This novel study has uncovered the stark realities that lie beneath the surface of the water for elite aquatic athletes. Attention is needed to improve understanding through targeted research regarding differences between genders and disciplines, as well as to develop educational initiatives to decrease the stigma of mental health in sport. Mechanisms are required to improve athlete help-seeking, to facilitate reporting of

harassment/abuse, and to enhance athlete support systems. Prevention strategies should be implemented and evaluated to address both athlete mental health challenges and harassment/abuse in the aquatic sports. The results of this study should serve as an alarm to engage and empower athletes and their entourage to work together with FINA to address athlete mental well-being and Safe Sport in the daily training environments of the swimming pools around the world.

„There are times where I feel absolutely worthless, where I completely shut down but have this bubbling anger that is through the roof. If I'm being honest, more than once I've just screamed out loud, "I wish I wasn't me!... The more you understand that it's ok, to not be ok – it's so powerful" Michael Phelps (2020)³¹

PRACTICAL IMPLICATIONS

- Sport organizations should develop strategies to decrease stigma related to athlete mental health problems.
- Team physicians should be educated to have the clinical competency to screen, diagnose and treat mental health problems in elite aquatic athletes regarding differences between genders and disciplines.
- Prevention strategies should be implemented to decrease the burden of mental health issues and harassment/abuse in sport.
- Reporting mechanisms and athlete support systems should be made available and accessible for athletes experiencing harassment/abuse in sport.

Acknowledgement: The authors greatly appreciate the cooperation of the athletes attending the FINA World Aquatic Championships who volunteered their time to provide data for this project, and to FINA for supporting the concept and logistics of the study.

REFERENCES

1. Scutti S. "Michael Phelps: I am extremely thankful that I did not take my life" CNN 20 Jan 2018. Available: <https://www.cnn.com/2018/01/19/health/michael-phelps-depression/index.html> [Accessed 22 Sep 2020]
2. Drehs W. Michael Phelps' final turn. 2016. EPSN the Magazine. Available: https://www.espn.com/espn/feature/story/_/id/16425548/michael-phelps-prepares-life-2016-rio-olympics [Accessed 22 Sep 2020]
3. Gouttebarga V, Castaldelli-Maia JM, Gorczynski P, et al. Occurrence of mental health symptoms and disorders in current and former elite athletes: a systematic review and meta-analysis. *Br J Sports Med*. 2019;53:700-6. <http://dx.doi.org.libaccess.lib.mcmaster.ca/10.1136/bjsports-2019-100671>
4. Reardon C, Hainline B, Aron CM, et al. Mental health in elite athletes: International Olympic Committee consensus statement (2019). *Br J Sports Med* 2019;53(11):667-99. <http://dx.doi.org/10.1136/bjsports-2019-100715>
5. Timpka T, Spreco A, Dahlstrom et al. Suicidal thoughts (ideation) among elite athletics (track and field) athletes: cross-sectional study of associations with sexual and physical abuse victimization, aspects of sports participation, and psychological and behavioural resourcefulness. *Br J Sports Med* 2020 <https://doi:10.1136/bjsports-2019-101386>
6. Kerr G, Willson E, Stirling A. Prevalence of maltreatment among current and former national team athletes. University of Toronto in partnership with Athletes CAN 20 April 2019. Available: http://athletescan.com/sites/default/files/images/prevalence_of_maltreatment_reporteng.pdf [Accessed 22 Sep 2020]
7. International Olympic Committee. Consensus Statement on "Sexual harassment and abuse in sport". February 2007. Available: <https://www.olympic.org/news/ioc-adopts-consensus-statement-on-sexual-harassment-and-abuse-in-sport> [Accessed 22 Sep 2020]

8. International Olympic Committee Safeguarding Framework, Rio de Janeiro Olympic Games 2016. Available: https://cdn.dosb.de/alter_Datenbestand/Bilder_allgemein/Veranstaltungen/Rio_2016/IOC_Framework_for_safeguarding_athletes.pdf [Accessed 22 Sep 2020]
9. Mountjoy M, Brackenridge C, Arrington M, et al. The IOC Consensus Statement: Harassment and abuse (non-accidental violence) in sport. *Br J Sports Med* 2016; (17):1019-29. <https://doi.org/10.1136/bjsports-2016-096121>
10. Mountjoy M, Vertommen T, Burrows K, et al., #SafeSport: Safeguarding initiatives at the Youth Olympic Games 2018. *Br J Sports Med* 2020;54:176-82. <https://doi.org/10.1136/bjsports-2019-101461>
11. Mountjoy M, Junge A, Benjamin S, et al. Competing with injuries: injuries prior to and during the 15th FINA World Championships 2013 (aquatics). *Br J Sports Med*. 2015;49(1):37-43. <http://dx.doi.org/10.1136/bjsports-2014-093991>
12. Prien A, Mountjoy M, Miller J, et al. Injury and illness in aquatic sport – how is the risk? A comparison of results from three FINA World Championships. *Br J Sports Med* 2017;51(4):277-82. <http://dx.doi.org/10.1136/bjsports-2016-096075>
13. Mountjoy M, Junge A, Slys J, et al. An uneven playing field: Athlete injury, illness, load, and daily training environment in the year prior to the FINA World Championships, 2017. [published online ahead of print, 2019 Dec 12]. *Clin J Sport Med* 2019;. <https://doi:10.1097/JSM.0000000000000814>
14. Clarsen B, Rønsen O, Myklebust G, et al. The Oslo Sports Trauma Research Center questionnaire on health problems: a new approach to prospective monitoring of illness and injury in elite athletes. *Br J Sports Med* 2014;48:754-60. <http://dx.doi.org/10.1136/bjsports-2012-092087>
15. Björgvinsson T, Kertz SJ, Bigda-Peyton JS, et al. Psychometric properties of the CES-D-10 in a psychiatric sample. *Assessment* 2013, 20, 429-36. <https://doi.org/10.1177/1073191113481998>
16. Zhang W, O'Brien N, Forrest JI, et al. Validating a shortened depression scale (10 item CES-D) among HIV-positive people in British Columbia, Canada. *PLoS One*. 2012;7(7):e40793. <https://doi.org/10.1371/journal.pone.0040793>

17. Martinsen M, Holme I, Pensgaard AM, et al. The development of the Brief Eating Disorder in Athletes Questionnaire (BEDA-Q). *Med Sci Sport Exer* 2014;46(8):1666–75. <https://doi.org/10.1249/mss.0000000000000276>
18. Junge A, Prinz B. Depression and anxiety symptoms in 17 teams of female football players including 10 German first league teams. *Br J Sports Med* 2019;53:471–7. <http://dx.doi.org/10.1136/bjsports-2017-098033>
19. Timpka T, Janson S, Jacobsson J. et al. Protocol design for large-scale cross-sectional studies of sexual abuse and associated factors in individual sports: Feasibility study in Swedish athletics. *J Sport Sci Med* 2015;14:179-87. PMID: 25729306; PMCID: PMC4306771
20. Armstrong S, Oomen-Early J. Social connectedness, self-esteem, and depression symptomatology among collegiate athletes versus non-athletes. *J Am Coll Health* 2009;57:521–6. <https://doi.org/10.3200/JACH.57.5.521-526>
21. Gulliver A, Griffiths KM, Mackinnon A, et al. The mental health of Australian elite athletes. *J Sci Med Sport* 2015;18:255–61. <https://doi.org/10.1016/j.jsams.2014.04.006>
22. Yang J, Peek-Asa C, Corlette JD, et al. Prevalence of and risk factors associated with symptoms of depression in Competitive Collegiate Student Athletes. *Clin J Sport Med* 2007;17:481–7. <https://doi:10.1097/JSM.0b013e31815aed6b>
23. Junge A, Feddermann-Demont N. Prevalence of depression and anxiety in top-level male and female football players. *BMJ Open Sport Exerc Med* 2016;2:e000087. <http://dx.doi.org/10.1136/bmjsem-2015-000087>
24. Ferrand C, Magnan C, Rouveix M, et al. Disordered eating, perfectionism and body-esteem of elite synchronized swimmers. *Eur J Sport Sci* 2007: 223–30. <http://doi:10.1080/17461390701722168>
25. Bratland-Sanda S, Sundgot-Borgen J. Eating disorders in athletes: overview of prevalence, risk factors and recommendations for prevention and treatment. *Eur J Sport Sci* 2013;13:499–508. <https://doi.org/10.1080/17461391.2012.740504>
26. Gouttebarga V, Bindra A, Blauwet C, et al. International Olympic Committee (IOC) Sport Mental Health Assessment Tool 1 (SMHAT-1) and Sport Mental

- Health Recognition Tool 1 (SMHRT-1): towards better support of athletes' mental health. *Br J Sports Med*. 2020;bjsports-2020-102411. <http://doi:10.1136/bjsports-2020-102411>
27. Mountjoy M, Moran J, Amed H, et al. Athlete health and safety at large sport events: the development of consensus-driven guidelines. *Br J Sports Med* 2020. [in publication]
28. Ohlert J, Vertommen T, Rulofs B, et al., Elite athletes' experiences of interpersonal violence in organised sport in Germany, the Netherlands, and Belgium. *Europ J Sport Sci* 2020; 1-10. [published online ahead of print, 2020 Jul 7]. <https://doi:10.1080/17461391.2020.1781266>
29. Timpka T, Janson S, Jacobsson J et al. Lifetime sexual and physical abuse among elite athletics athletes: A cross-sectional study of prevalence and correlates with athletics injury. *Br J Sports Med* 2014;48:667. <http://dx.doi.org/10.1136/bjsports-2014-093494.285>
30. Kawakami N, Yasuma N, Watanabe K, et al. Association of response rate and prevalence estimates of common mental disorders across 129 areas in a nationally representative survey of adults in Japan. *Soc Psychiatry Psychiatr Epidemiol*. 2020: <https://doi.org/10.1007/s00127-020-01847-3>
31. Phelps M. as told to Drehs W. „Michael Phelps: ,Thi is the most overwhelmed I've ever felt'. *ESPN* 18 May 2020. Available: https://www.espn.com/olympics/story/_/id/29186389/michael-phelps-most-overwhelmed-ever-felt [Accessed 22 Sep 2020]

FIGURES and TABLES

Figure 1: Average score on the depression (CESD-R-10, dark blue) and the eating disorder (BEDA-Q, light blue) scale in athletes with different experiences of harassment/abuse.

610 TABLES

611 **Table 1:** Athlete responses to the question: “Have you ever witnessed or
612 experienced yourself any of the following situations in your sports environment? If
613 yes, please answer both questions.”
614

		who was the victim?*		who did it?*		
	no	myself	another athlete	coach	another athlete	other
Unwanted comments about body or appearance (n=361)	216 59.8%	91 47+44	98 54+44	95	63	27
Exclusion, isolation or intentional denial of attention and/or support (n=356)	276 77.5%	44 27+17	53 36+17	26	51	16
Bullying / cyber bullying (n=359)	286 79.7%	39 22+17	52 34+17	20	46	20
Comments, jokes, messages or gestures of a sexual nature (n=361)	296 82.0%	30 9+21	56 35+21	29	42	13
Negative comments on race, gender, religion or country of origin (n=361)	302 83.7%	22 12+10	47 37+10	17	39	23
Inappropriate training, e.g. forced physical exertion beyond capabilities (n=357)	306 85.7%	39 22+17	29 12+17	45	2	1
Intentional hitting, beating or kicking (not fouls or rough play) (n=360)	317 88.1%	18 8+10	35 25+10	16	22	7
Forced dieting, limited or no access to needed water or food (n=361)	323 89.5%	27 13+14	25 11+14	28	2	4
Unsafe training conditions (n=358)	322 89.9%	30 14+16	22 6+16	27	2	4
Unwanted and unnecessary touching of the body (n=360)	324 90.0%	18 6+12	30 18+12	18	16	9
Unwanted touching of breast, buttock, genitals or attempt to kiss (n=361)	336 93.1%	11 0+11	25 14+11	11	14	3
Forced use of supplements / medication (n=362)	350 96.7%	7 2+5	10 5+5	9	4	4
Rewards in sport for sexual favours (n=362)	353 97.5%	3 2+1	7 6+1	7	1	0
other, please specify (n=327)	323 98.8%	2 1+1	3 2+1	1	1	

615 *Multiple answers are possible; + number of athletes who selected both “myself” and
616 “another athlete”
617

Table 2. Characteristics of all elite aquatic athletes, and of male and female athletes from different selected disciplines (i.e. results from open water swimmers and male artistic swimmers not presented separately due to small numbers but included in total) and number (%) of athletes who experienced different types of harassment/abuse themselves with regard to gender and discipline.

	Swimming		Water Polo		Diving / High Diving		Artistic Swimming	Total
Gender	female	male	female	male	female	male	female	both
Number of responding athletes	65	67	48	34	29	34	77	375
Age								
Mean (sd)	21.4 (4.9)	22.3 (5.7)	23.1 (3.9)	24.8 (4.1)	23.3 (5.9)	26.7 (7.4)	19.6 (3.8)	22.4 (5.4)
Depression (CES-D-10)								
Athletes with depression	12 (27.3%)	15 (31.9%)	9 (27.3%)	3 (12.5%)	6 (23.1%)	4 (15.4%)	15 (28.3%)	70 (26.1%)
Eating disorder (BEDA-Q)								
Athletes with eating disorder	16 (28.1%)	14 (26.4%)	16 (38.1%)	8 (27.6%)	11 (42.3%)	8 (27.6%)	33 (54.1%)	111 (35.2%)
Lose weight now	15 (24.2%)	9 (14.8%)	9 (19.6%)	6 (18.8%)	12 (41.4%)	4 (12.5%)	33 (44.0%)	90 (25.3%)
Need for psychotherapy								
No expressed need	28 (45.2%)	33 (55.0%)	20 (42.6%)	24 (72.7%)	17 (58.6%)	22 (71.0%)	46 (61.3%)	198 (55.6%)
Currently receiving	16 (25.8%)	14 (23.3%)	4 (8.5%)	5 (15.2%)	5 (17.2%)	3 (9.7%)	7 (9.3%)	57 (16.0%)
Previously received	18 (29.0%)	13 (21.7%)	23 (48.9%)	4 (12.1%)	7 (24.1%)	6 (19.4%)	22 (29.3%)	101 (28.4%)
Harassment /abuse								
Likely in the athlete's sport ^(a)	3.0 (1.4)	3.2 (1.4)	3.0 (1.4)	3.3 (1.3)	2.2 (1.4)	2.4 (1.3)	2.2 (1.2)	2.8 (1.4)
Not experienced nor witnessed	47 (75.8%)	51 (87.9%)	35 (77.8%)	18 (58.1%)	22 (84.6%)	27 (87.1%)	57 (81.4%)	271 (79.5%)
Experienced themselves	9 (14.5%)	4 (6.9%)	3 (6.7%)	5 (16.1%)	4 (15.4%)	2 (6.5%)	9 (12.9%)	39 (11.4%)
Witnessed in another athlete	3 (4.8%)	2 (3.4%)	5 (11.1%)	5 (16.1%)	0	1 (3.2%)	2 (2.9%)	19 (5.6%)

Experienced and witnessed	3 (4.8%)	1 (1.7%)	2 (4.4%)	3 (9.7%)	0	1 (3.2%)	2 (2.9%)	
Seek assistance								
No	6 (10.0%)	15 (25.9%)	2 (4.7%)	5 (17.2%)	2 (8.0%)	2 (6.7%)	7 (9.9%)	43 (12.9%)
Maybe	10 (16.7%)	16 (27.6%)	8 (18.6%)	7 (24.1%)	5 (20.0%)	8 (26.7%)	13 (18.3%)	68 (20.4%)
Friends or family	35 (58.3%)	11 (19.0%)	25 (58.1%)	14 (48.3%)	12 (48.0%)	13 (43.3%)	41 (57.7%)	161 (48.2%)
Official contact to seek help	5 (8.3%)	12 (20.7%)	7 (16.3%)	3 (10.3%)	3 (12.0%)	6 (20.0%)	5 (7.0%)	43 (12.9%)
Friends/family and official contact	4 (6.7%)	4 (6.9%)	1 (2.3%)	0	3 (12.0%)	1 (3.3%)	5 (7.0%)	19 (5.7%)
Athletes who experienced the specified type of harassment/abuse themselves								
Unwanted comments about body or appearance	13 (20.6%)	8 (11.9%)	7 (24.9%)	4 (12.1%)	8 (32.0%)	5 (15.2%)	39 (55.7%)	91 (25.2%) of 361
Exclusion, isolation, or intentional denial of attention and/or support	13 (21.0%)	8 (12.1%)	4 (8.5%)	3 (9.1%)	1 (3.8%)	2 (6.1%)	11 (16.4%)	44 (12.4%) of 356
Bullying / cyber bullying	10 (16.1%)	4 (6.0%)	4 (8.7%)	1 (3.0%)	3 (11.5%)	1 (3.0%)	14 (20.3%)	39 (10.8%) of 359
Comments, jokes, messages, or gestures of a sexual nature	9 (14.8%)	5 (7.6%)	4 (8.5%)	3 (9.1%)	2 (7.7%)	1 (3.0%)	4 (5.6%)	30 (8.3%) of 361
Negative comments on race, gender, religion, or country of origin	3 (4.8%)	2 (3.0%)	3 (6.4%)	2 (6.3%)	2 (7.7%)	0	7 (9.9%)	22 (6.1%) of 361
Inappropriate training, e.g. forced physical exertion beyond capabilities	9 (14.8%)	6 (9.1%)	3 (6.5%)	4 (12.1%)	0	0	13 (18.8%)	39 (10.9%) of 357
Intentional hitting, beating, or kicking (not fouls or rough	2 (3.2%)	5 (7.5%)	3 (6.4%)	3 (9.1%)	0	0	4 (5.9%)	18 (5.0%) of 360

play)								
Forced dieting, limited or no access to needed water or food	9 (14.3%)	0	2 (4.3%)	1 (3.0%)	1 (3.8%)	0	12 (17.1%)	27 (7.5%) of 361
Unsafe training conditions	7 (11.3%)	4 (6.2%)	4 (8.7%)	1 (3.0%)	5 (19.2%)	2 (6.1%)	5 (7.1%)	30 (8.4%) of 358
Unwanted and unnecessary touching of the body	8 (12.9%)	3 (4.5%)	4 (8.5%)	0	0	0	3 (4.3%)	18 (5.0%) of 360
Unwanted touching of breast, buttock, genitals or attempt to kiss	5 (7.9%)	2 (3.0%)	1 (2.1%)	0	0	0	2 (2.9%)	11 (3.0%) of 361
Forced use of supplements / medication	3 (4.8%)	1 (1.5%)	1 (2.1%)	1 (3.0%)	0	0	0	7 (1.9%) of 362
Rewards in sport for sexual favours	2 (3.2%)	1 (1.5%)	0	0	0	0	0	3 (0.8%) of 362
Other, please specify	1 (1.9%)	0	0	0	0	0	0	2 (0.6%) of 327

Numbers do not add up to responding athletes due to missing values in gender (n=2), discipline (n=12) and the respective question.

(a) no =1; not likely =2; not sure =3; likely =4; very likely =5.

DO NOT COPY