Variant Creutzfeldt-Jakob Disease: The Challenge of Preventing a Rare but Potentially Devastating Exposure

Ritambhara Pandey MD, Devesh Rai MD, Julie Giles NP, Maryrose Laguio-Vila MD, Emil
Lesho DO

Rochester Regional Health, Rochester, NY, USA

Abstract

Although rare, patients with variant Creutzfeldt-Jakob Disease (vCJD) in their differential
diagnosis of progressive dementia and movement disorder could continue to present to hospitals
for care. However, U.S.-based infection control guidelines do not fully address the possibility of
vCJD. After near-misses involving increasing numbers of patients with clinical findings and
epidemiologic risks compatible with vCJD, or exposures to chronic wasting disease, we sought
to improve recognition and prevention of iatrogenic spread of these prion-related diseases.

21 **Introduction**

22 Transmissible spongiform encephalopathies (TSEs) are progressively fatal
23 neurodegenerative diseases caused by infectious, misshapen prion proteins (PrP^{Sc}). In humans,
24 they include variant and classic (or sporadic) Creutzfeldt-Jakob Disease (vCJD/sCJD), the latter
25 being far more common in the U.S. Animal forms of prion disease include bovine spongiform
26 encephalopathy (BSE) and chronic wasting disease in cervids and some rodents.

27 TSEs present substantial diagnostic, infection control, and patient notification challenges
28 detailed in the discussion section below. Because prions resist usual standard disinfection and
29 sterilization protocols, iatrogenic exposures can occur via contaminated instruments or
30 equipment. Iatrogenic TSE can also be transmitted by percutaneous exposure to certain types of
31 tissue (mostly neural for sCJD), while vCJD can also be transmitted by exposure to infected
32 lymphoid tissue and possibly blood transfusions. This is why people from regions where BSE
33 was prevalent are still precluded from donating blood.¹⁻³

34 However, U.S.-based cleaning and disinfection and control guidelines and updates do not
35 address the possibility of vCJD.^{4,5} One of the reasons it is not included in general procedures is
36 that it is a rare occurrence and a special consideration. vCJD involves separate procedures that
37 are not applied across all reprocessing procedures, and different types of tissue that are
38 potentially infectious (not only neural).

39 Those guidelines do, however, recommend heightened awareness and policy
40 modification as events dictate.^{4,5} Additionally, experts have noted persistent risk of TSEs other
41 than sCJD in unsuspected geographic locations and a need for heightened suspicion and
42 surveillance.^{6,7} Furthermore, surveys of lymphoid tissue where PrP^{Sc} is found in cases of vCJD,

suggest the prevalence of asymptomatic disease can be as high as 1 in 2,000 in persons born between 1941 and 1985.⁸

Two Illustrative Incidents

Two incidents typified the challenges we faced in caring for these and similar patients and provided the impetus for this approach. The first incident involved a patient who presented with memory problems, apathy and depression leading to job loss. These symptoms were followed by confusion, impaired gait, occasional jerking movements and eventually dementia. The patient lived in England and France from 1985 to 1993 and ate canned dog food occasionally. The patient also required urgent pelvic and abdominal surgery immediately after admission. MRI showed restricted diffusion in the frontal, parietal, and occipital cortexes. 14-3-3, and Tau protein were negative. Since immune mediated encephalitis was also a leading consideration on the list of possible diagnoses, the patient underwent plasma exchange. 13 days later, the RTQuIC result became available and was positive. Contact tracing revealed that in the meantime, three other patients had undergone pheresis on the same machine. The Red Cross was contacted and recommended decommissioning the machine until further notice. The dilemma of patient notification was debated by the ethics, risk management, infection control team and the hospital leaders (Table 1).

The second incident involved a middle-aged patient who presented with psychiatric symptoms and cognitive decline. The patient, who was an avid hunter and consumed venison and squirrel brains, also developed impaired gait. An MRI showed bilateral pulvinar signs. The EEG did not reveal periodic sharp waves, but showed nonspecific abnormalities including intermittent rhythmic delta activity. The patient required several diagnostic and therapeutic procedures including endoscopies and endotracheal intubation that entailed exposure to

lymphoid tissue. Approximately 2 weeks after admission, 14-3-3, Tau protein, and RTQuIC results became available and were positive, creating a dilemma for reprocessing critical and semi-critical equipment. After several months in the hospital, the patient was discharged to a long-term care facility, and died more than 11 months after symptom onset. vCJD could not be definitively excluded until autopsy results became available several months later and revealed sCJD.

Against this backdrop, we sought to enhance our efforts to detect and prevent iatrogenic spread of TSE, including vCJD. We also aimed to increase awareness among all providers, prevent over-reaction or unnecessarily discarding expensive equipment, while simultaneously not missing an opportunity to prevent avoidable exposures.

Methods

Rochester Regional Health System consists of 8 hospitals and 9 urgent care centers in St Lawrence County, and the 9 counties comprising the Finger Lakes region of NY. The system serves diverse populations including refugee communities and permanent or temporarily relocated citizens or expatriates from many countries. In 2018, requests for CSF analysis for suspected TSE increased 20% compared to the preceding four years.

The two-part intervention consisted of an educational campaign nearly identical to the one successfully used for device-related infections and described previously.⁹ The second component entailed an extensive revision of our existing CJD policy, incorporating guidelines from the UK Department of Health and Social Care Advisory Committee and the World Health Organization regarding vCJD.^{10, 11} Briefly, the educational campaign included one-on-one engagements between infection preventionists (IPs), the hospital epidemiologist, the neurologist champion and frontline nurses and providers. Modes of delivery included individual

89 face-to-face meetings, team huddles, committee meetings, group lectures and grand rounds.

90 Relevant guidelines were posted on hospital websites and intranet.

91 One goal of the policy revision was early notification of all teams involved in the care
 92 including physicians, nursing staff including laboratory and sterile processing. Another goal was
 93 a simplified classification scheme of potentially infected tissue. Tissues were divided into three
 94 categories depending on their risk of infectivity: high, low, and no detectable infectivity (Table
 95 2). Only single use or disposable instruments are recommended for suspected cases who undergo
 96 surgery or invasive procedures involving contact with other than “no-risk” tissues. Non-
 97 disposable instruments or durable medical equipment that contact other than no risk tissues
 98 should be quarantined for possible decommissioning with appropriate labeling. All instruments
 99 are considered potentially contaminated when used on patients with conditions not clearly
 100 diagnosed before the procedure, and those instruments are either reprocessed according to prion
 101 guidelines or quarantined until the diagnosis is confirmed.

102 **Results**

103 During the 2-year follow up period, there were no similar incidents or other near misses.
 104 In response to queries from other IPs and epidemiologists, the policy was shared with hospitals
 105 outside of the network. Feedback was uniformly positive.

106 **Discussion**

107 TSEs present diagnostic, infection control and patient notification challenges. Most
 108 clinicians are unfamiliar with TSEs due to the rarity of those diseases. vCJD and CJD can be
 109 hard to distinguish clinically. No gold standard laboratory test exists for differentiating types of
 110 TSEs. The turn-around-time for RT-QiC can be up to 14 days or longer. In some cases, definitive

diagnosis requires brain biopsy or only becomes available after autopsy. Last, the National Prion Center typically only releases autopsy results to public health departments, furthering diagnostic delay.

This approach could be criticized for not being warranted or justifiable due to the fact vCJD is extraordinarily uncommon. Even though sCJD is far more common than vCJD in the U.S., patients with suspected vCJD have been increasingly encountered in our health system. The suspicion was based on epidemiologic exposures and symptoms or signs consistent with vCJD. For example, having lived in the United Kingdom or France for several years or decades, or occupational exposure as taxidermists or abattoirs, or regular consumption of animal brains or cervid meat products. Furthermore, available data indicate that the incidence of CJD in cervids is increasing, and the potential exists for transmission to humans and subsequent human disease, and the species barrier is not static.⁷ These occurrences are unlikely to be unique to our healthcare system. To that end, we make our policy to those interested. It is available from the corresponding author upon request.

Acknowledgements: None

Funding: There was no funding.

Conflicts of Interest: There are no conflicts of interest.

References:

1. American Red Cross. Eligibility reference material.
<https://www.redcrossblood.org/donate-blood/how-to-donate/eligibility-requirements/eligibility-criteria-alphabetical/eligibility-reference-material.html>.
Accessed January 19, 2021.
2. Belay ED, Schonberger LB. The public health impact of prion diseases. *Annu Rev Public Health*. 2005;26:191-212.
3. Ricketts MN, Cashman NR, Stratton EE, El Saadany S. Is Creutzfeldt-Jakob disease transmitted in blood? *Emerg Infect Dis*. 1997;3(2):155-163.
4. Rutala WA, Weber DJ. Creutzfeldt-Jakob disease: recommendations for disinfection and sterilization. *Clin Infect Dis*. 2001;32(9):1348-1356.
5. Rutala WA, Weber DJ; Society for Healthcare Epidemiology of America. Guideline for disinfection and sterilization of prion-contaminated medical instruments. *Infect Control Hosp Epidemiol*. 2010;3(2)1:107-117.
6. Maheshwari A, Fischer M, Gambetti P, et al. Recent US case of variant Creutzfeldt-Jakob Disease - global implications. *Emerg Infect Dis*. 2015;21(5):50-759.
7. Osterholm MT, Anderson CJ, Zabel MD, Scheftel JM, Moore KA, Appleby BS. Chronic wasting disease in cervids: implications for prion transmission to humans and other animal species. *mBio*. 2019;10(4):e01091-19.
8. Diack AB, Will RG, Manson JC. Public health risks from subclinical variant CJD. *PLoS Pathog*. 2017;13(11):e1006642.
9. Lesho EP, Clifford R, Vore K, et al. Sustainably reducing device utilization and device-related infections with DeCATHlons, device alternatives, and decision support. *Infect Control Hosp Epidemiol*. 2020;41(11):1344-1347.

10. Advisory Committee on Dangerous Pathogens TSE Subgroup. ACDP's August 2016 Updated position statement on occurrence of vCJD and prevalence of infection in the UK. <https://www.gov.uk/government/groups/advisory-committee-on-dangerous-pathogens>. Accessed March 14, 2021
11. World Health Organization. WHO infection control guidelines for transmissible spongiform encephalopathies. Report of a WHO consultation, Geneva, Switzerland, 23-26 March 1999. https://www.who.int/csr/resources/publications/bse/WHO_CDS_CSRAPH_2000_3/en/. Accessed March 14, 2021.

176 **Table 1: Diagnostic, Infection Control, and Notification Challenges Associated with**
 177 **Transmissible Spongiform Encephalopathies**

Diagnostic Challenge
Most clinicians are unfamiliar with TSEs due to the rarity of those diseases.
vCJD and CJD can be hard to distinguish clinically.
No gold standard laboratory test exists for differentiating types of TSEs.
Turn-around-time for RT-QiC can be up to 14 days or longer.
In some cases, definitive diagnosis requires brain biopsy or only becomes available after autopsy.
The National Prion Center typically only releases autopsy results to public health departments, furthering delay.
Infection Control Challenges
Prions resist usual disinfection and sterilization methods.
US-based disinfection and sterilization guidelines do not address possibility of vCJD contaminated equipment.
Notification Challenges or Considerations
Should potentially exposed patients be informed given the below considerations?
i. Potential negative impact of informing patients about a fatal, untreatable brain disease.
ii. No effective screening test is available for TSE.
iii. No post exposure prophylaxis is available.
iv. Contaminated instruments often become mixed during re-processing making identification and linkage to specific patients nearly impossible.
v. Autoclaving is generally superior to chemical reprocessing for prions.
vi. If equipment was re-processed more than twice, prions would be less likely to survive.

178 **Table 2: Infectivity of Specific Types of Tissue**

	Classic/ Sporadic CJD	Variant CJD
High Infectivity Tissue NO Reprocessing Allowed*	Brain (including dura mater) Spinal cord Posterior eye Pituitary tissue	Brain (including dura mater) Spinal cord, spinal ganglia Posterior eye Pituitary tissue Trigeminal ganglia
Low Infectivity Tissue NO Reprocessing Allowed*	Cerebrospinal fluid (CSF) Liver Lymph node Kidney Lung (Bronchoscope) Spleen Placenta Olfactory epithelium	Adrenal gland Blood Tonsils (Laryngoscope) Cerebrospinal fluid (CSF) Intestine (Endoscope) Bone marrow Liver Lymph node Kidney Lung (Bronchoscope) Spleen Skeletal muscle Placenta Olfactory epithelium Peripheral Nerves Rectum (Endoscope)
Tissue with no detectable infectivity OK to Reprocess*	Peripheral nerve Intestine Bone marrow Whole blood, leukocytes, serum, plasma Thyroid gland Adrenal gland Heart Skeletal muscle Adipose tissue Gingiva Prostate and testis Tears, sweat, saliva, and sputum Urine, and feces Semen, vaginal secretions, Milk	Thyroid gland Heart Adipose tissue Gingiva Prostate and testis Tears, sweat, saliva, and sputum Urine, and feces Semen, vaginal secretions, Breast milk

180 *Applies to surgical instruments and certain invasive equipment such as endoscopes and
181 laryngoscope blades
182