Clinical malaria incidence following an outbreak in Ecuador was predominantly associated with *Plasmodium falciparum* with recombinant variant antigen gene repertoires

Shazia Ruybal-Pesántez^{1§}, Fabian E. Sáenz², Samantha Deed^{1,3}, Erik K. Johnson⁴, Daniel B. Larremore^{5,6},
Claudia A. Vera-Arias^{2#}, Kathryn E. Tiedje^{1,3}, Karen P. Day^{1,3*}

¹ School of BioSciences/Bio21 Institute, University of Melbourne, Australia

² Centro de Investigación para la Salud en América Latina, Facultad de Ciencias Exactas y Naturales, Pontificia Universidad Católica del Ecuador, Av. 12 de octubre 1076, Apartado: 17-01-2184, Quito, Ecuador

³ Department of Microbiology and Immunology, University of Melbourne, Bio21 Institute, Melbourne, Australia

⁴ Department of Applied Mathematics, University of Colorado Boulder, Boulder, Colorado, USA

⁵ Department of Computer Science, University of Colorado Boulder, Boulder, Colorado, USA

⁶ BioFrontiers Institute, University of Colorado Boulder, Boulder, Colorado, USA

[§] *Current affiliations:* Population Health and Immunity Division, Walter and Eliza Hall Institute of Medical Research, Melbourne, Australia; Department of Medical Biology, University of Melbourne, Australia; Burnet Institute, Australia

[#] *Current affiliation:* Eck Institute for Global Health, University of Notre Dame, Notre Dame, IN, 46556, USA

**Corresponding author:* Karen P. Day; karen.day@unimelb.edu.au

Abstract

To better understand the factors underlying the continued incidence of clinical episodes of falciparum malaria in E-2020 countries targeting elimination, we have characterised *Plasmodium falciparum* disease transmission dynamics after a clonal outbreak on the northwest coast of Ecuador over a period of two years. We apply a novel, high-resolution genotyping method, the “varcode” based on a single PCR to fingerprint the DBLα region of the 40-60 members of the variant surface antigen-encoding *var* multigene family. *Var* genes are highly polymorphic within and between genomes, with *var* repertoires rapidly evolving by outcrossing during the obligatory sexual phase of *P. falciparum* in the mosquito. The continued incidence of clinical malaria after the outbreak in Ecuador provided a unique opportunity to use varcodes to document parasite microevolution and explore signatures of local disease transmission on the time scale of months to two years post-outbreak. We identified nine genetic varcodes circulating locally with spatiotemporal parasite genetic relatedness networks revealing that diversification of the

clonal outbreak parasites by sexual recombination was associated with increased incidence of clinical episodes of malaria. Whether this was due to chance, immune selection or sexual recombination per se is discussed. Comparative analyses to other South American parasite populations where *P. falciparum* transmission remains endemic elucidated the possible origins of Ecuadorian varcodes. This analysis demonstrated that the majority of clinical cases were due to local transmission and not importation. Nonetheless, some of the varcodes that were unrelated to the outbreak varcode were found to be genetically related to other South American parasites. Our findings demonstrate the utility of the varcode as a high-resolution surveillance tool to spatiotemporally track disease outbreaks using variant surface antigen genes and resolve signatures of recombination in an E-2020 setting nearing elimination.

Introduction

In 2016 the WHO identified 21 countries, known as the E-2020 countries, having the potential to eliminate local transmission of malaria by 2020¹, with seven of them in Latin America at the time (Paraguay and El Salvador have since been declared malaria-free). Malaria transmission in many of these E-2020 countries is epidemic/unstable with risks of outbreaks and resurgent malaria²⁻⁴. The elimination target requires a local reduction to zero incidence of indigenous cases. As progress is made towards elimination in these countries, it becomes critical to understand risk factors for disease transmission while maximizing limited funds and resources to examine individual cases. Ecuador is one of the five remaining E-2020 countries in Latin America. National malaria elimination efforts have largely focused on tropical areas, specifically the northwest coast and the Amazon region. These areas border non E-2020 countries, Colombia and Peru, that still have endemic transmission. Thus, rather than low transmission, Ecuador has epidemic/unstable *P. falciparum* transmission with *P. vivax* the dominant species causing clinical malaria infections.

Clinical cases caused by *P. falciparum* in Ecuador are mostly concentrated in the northwest coast, where a steady decrease in annual reported clinical cases was observed from 2009 (161 cases) until late 2012 (44 cases)⁵⁻⁷. A localized outbreak from November 2012 until November 2013 in Esmeraldas City documented 151 *P. falciparum* cases, the majority of cases (76%) in the northwest coast and Amazon region during this time period⁸. Population genetic analyses using microsatellite markers demonstrated that this outbreak was caused by the clonal expansion (i.e., single-source) of a residual parasite lineage previously reported on the north coast of Peru and on the coast of Ecuador⁸. Immediately after the outbreak, 31 cases were reported in 2014 and 189 cases in 2015^{5,9,10}. It was unclear whether the clonal

outbreak had contributed to *P. falciparum* clinical cases since 2013, with the factors underlying sustained disease transmission in Ecuador poorly understood. Moreover, Ecuador was categorized as “off-track” in the most recent E-2020 progress report and has not met the elimination target due to malaria resurgence^{11,12}. This points to an urgent need to better understand disease transmission patterns and whether clinical cases are due to imported or locally-acquired parasites.

Molecular surveillance of malaria in Latin America has focused on non E-2020 countries, e.g. Colombia, Brazil, Peru and Panama where malaria is endemic and transmission is moderate to low^{8,13–19}. This surveillance has relied on genotyping neutral molecular markers such as microsatellites or single-nucleotide polymorphisms. However, neutral markers, considered relatively slowly evolving, may not provide enough discriminatory resolution to define parasite evolution in relevant epidemiological time (1–2 years) in areas of low or epidemic and unstable transmission^{20,21}. Genes under selection can provide an alternative but complementary view of microevolution and they are key to examine population adaptation²².

In this context, the microbiological paradigm for the surveillance of diverse pathogens is to study transmission dynamics using the genes encoding the major surface antigen to reflect pathogen transmission dynamics in relation to immune selection^{23,24}. The failure to observe this paradigm in the malaria field has stemmed from both the misconception that neutral markers track transmission dynamics in relation to immunity and the fact that the genetics of the *var* genes encoding the major surface antigen of the blood stages (PfEMP1) are considered “too complex”. Single copy antigen genes under much less selection have become an alternative²⁵. Instead, we have embraced the complexity of the *var* system to understand the population genetics of *var* genes^{23,26–31} in multiple locations in Africa, South America, and globally to present here a way to explore transmission dynamics through the lens of these immune evasion genes.

Each genomic *var* repertoire of *P. falciparum* consists of 40 to 60 members of the *var* multigene family with evidence that South American parasites may have fewer *var* genes, e.g. 42 *var* genes (excluding *var2csa*) reported in the Honduran laboratory strain HB3 after whole genome sequencing and assembly^{32,33}. Our analysis of a South American dataset with a global dataset also showed fewer *var* genes in South American isolates as did a Brazilian data set^{31,34}. As *var* genes lie on almost all of the 14 chromosomes of the haploid *P. falciparum* genome³⁵ they undergo independent assortment during conventional meiosis

to create new *var* repertoires as recombinants of parental *var* repertoires. During infection, differential switching between these genes in a repertoire enables immune evasion in the human host^{36,37} thereby regulating parasite density and risk of clinical infection³⁸. Indeed, several serologic studies examining anti-PfEMP1 antibody responses in naturally-infected children have shown that the acquisition of anti-PfEMP1 protective immunity is variant-specific and susceptibility to clinical episodes is dependent on gaps in the individual's pre-existing antibody repertoire³⁸⁻⁴⁷.

A body of work defining the population genomics of the *var* multigene family by ourselves and others⁴⁸ has demonstrated that *var* genes provide a promising biomarker for surveillance over epidemiological time scales (1-10 years). *Var* DBL α sequences can resolve population structure at global, continental, regional and local scales in areas of varying endemicity in Africa, Oceania, and South America^{23,26-31,49}. Characterization of these major surface antigen genes also provides information about circulating repertoires of variants to enable measurement of immune responses. Translation of *var* population genetics in different epidemiological settings leads us to propose a new surveillance approach we call *var*coding. This is essentially a fingerprinting method to describe the diversity of these genes within and between human hosts as well as repertoire diversification by sexual recombination, with Bayesian statistics to account for missing data (Fig. 1).

Defining the 2012-2013 clinical disease outbreak in Ecuador as a baseline, we applied the *var*code methodology to characterize the transmission dynamics of clinical *P. falciparum* cases during and up to two years after the outbreak. Reconstruction of parasite genetic relatedness networks in space and time revealed that parasites with recombinant *var* repertoires were predominantly associated with the continued incidence of clinical cases after the outbreak. Further comparative analyses to published data from South American isolates²⁸ elucidated possible origins of Ecuadorian parasites, demonstrating that the majority of clinical cases were due to local transmission and not importation. Our findings also demonstrate the translational application of the *var*code as a high-resolution tool to detect, spatiotemporally track immune evasion genes as well as resolve signatures of recombination in E-2020 or low transmission settings.

Figure 1. Schematic diagram of the varcoding approach. For more details about each step, see Methods.

Results

Varcoding was completed for 58 *P. falciparum* isolates that were collected between 2013 and 2015 in endemic areas of the northwest coast and Amazon region of Ecuador (Fig. 2) from individuals of all ages presenting with malaria symptoms and confirmed *P. falciparum*-positive by microscopy or rapid diagnostic tests. These isolates represent 21% of the total cases reported in 2013, 61% in 2014 and 3% in 2015 (60% of the cases reported in January, 7% in May and 8% in November). For more details on the sampling scheme and bioinformatics workflow see Methods and Supplementary Fig. S1. Overall, we identified 195 unique *var* DBLα variants or types from 2,141 DBLα sequences from the 58 isolates, representative of the diversity circulating in Ecuadorian *P. falciparum* populations, as indicated by accumulation curves approaching a plateau (Fig. S2).

Figure 2. Sampling of *P. falciparum* isolates across endemic areas of Ecuador and over a period of three years (2013 – 2015). (a) Map of Ecuador depicting the sampling locations during the study. (b) Bar plot showing the number of *P. falciparum* positive samples collected in each year and their respective sampling locations. Every location is indicated with a different color.

Defining varcodes circulating in Ecuador

To define distinct varcodes circulating locally in Ecuador, we estimated parasite genetic relatedness by calculating the similarity index, Pairwise Type Sharing (P_{TS}) and a Bayesian version of the same (Bayesian pairwise type sharing; BP_{TS} , see Methods). BP_{TS} estimates include statistical uncertainty due to differences in isolate repertoire size (i.e. the number of DBL α types identified in each isolate repertoire), enabling a rigorous examination of relatedness. The median repertoire size in Ecuadorian isolates was 37 DBL α types (range: 11-43 types, Table S1), which is in line with the expected number of var genes in the Honduran laboratory reference strain HB3 ($n=42$ var genes excluding *var2csa*,³³). Nonetheless, we used a uniform prior distribution for repertoire sizes between 40 and 50 types, which accounted for the possibility of missing data in total isolate repertoire sizes using degenerate primers for var DBL α PCR. For a further analysis to validate our PCR and sequencing methodology, including a further sub-classification of DBL α types into upsA and non-upsA groups and their respective isolate genomic proportions, see Methods and Supplementary Text 1.

We constructed genetic relatedness networks at a threshold of $P_{TS} \geq 0.90$ to identify putatively identical genomes within the margin of error of detection of all DBL α types in an isolate. We confirmed this definition by comparing varcodes derived from P_{TS} to those derived from posterior mean BP_{TS} estimates (Supplementary Fig. S3), and by examining the lower and upper bounds of the 95% highest density posterior intervals (HDPIs). This revealed nine genetically distinct varcodes in this study with varcode4, varcode5, varcode9 seen only once and at the other extreme 36 isolates had varcode1 (Fig. 3a and

Supplementary Fig. S4a). We were unable to confirm whether any of the Ecuadorian *var*codes were circulating pre-outbreak as samples were not available. As expected, the outbreak *var*code1 was clonal (indicated by HDPIs that included 1; Fig. S5d), as previously demonstrated by microsatellite genotyping⁸. It is worth noting that even though we only sampled 30 *P. falciparum* isolates in Esmeraldas City from 140 (21%) outbreak cases in 2013, the fact that the disease outbreak was clonal and represented 91% of the 154 total cases on the northwest coast and Amazon region during 2013, means that we characterized a representative sample of the cases from the outbreak and during that year.

For each pairwise comparison, we calculated the HDPI width as an estimate of uncertainty (analogous to calculating the width of a frequentist confidence interval) and found there was high confidence (HDPI width ≤ 0.2) for 94.6% (632/668) of pairwise BP_{TS} estimates (Supplementary Fig. S4b). Low confidence pairwise BP_{TS} estimates (HDPI width > 0.2 ; max = 0.33) were found only between the two *var*code1 isolates with the smallest repertoire sizes (11 and 19 DBL α types; 36/36 comparisons). This lower statistical confidence follows naturally from under sampling of those isolates' DBL α types. After aggregating the within-*var*code pairwise comparisons, there was high confidence in the definition of all *var*codes (Supplementary Fig. S5a-b).

For comparison of relatedness by microsatellite genotyping, we calculated Pairwise Allele Sharing (P_{AS}) using previously published microsatellite data for the same isolates^{8,50}. Not surprisingly, 7 microsatellites did not resolve the nine *var*codes but only identified four, three of which were single parasite isolates, at the threshold of $P_{AS} \geq 0.90$ (Supplementary Fig. S6a). Because two of the seven microsatellite markers were fixed in the population^{8,50}, we also generated the relatedness networks at the threshold of $P_{AS} \geq 0.80$, which allowed for an error of detection of 1 microsatellite allele. However, the majority of isolates clustered at this threshold and only three *var*codes were resolved (Supplementary Fig. S6b). There were several pairwise comparisons (14%, 72/541 by P_{TS} and 6%, 31/541 by BP_{TS}) where *P. falciparum* isolates were found to be identical by microsatellites ($P_{AS} = 1$) but much less related by *var* (P_{TS} or BP_{TS} ≤ 0.50 , Supplementary Fig. S7). Nevertheless, the relationship between P_{TS} and P_{AS} scores was positively correlated (Pearson's correlation coefficient = 0.757, $p < 0.001$, Supplementary Fig. S7a). Similarly, the relationship between BP_{TS} and P_{AS} estimates was also positively correlated (Pearson's correlation coefficient = 0.779, $p < 0.001$, Supplementary Fig. S7b).

Persistent association of clinical cases with the same *var*codes over time and large distances

We next explored whether parasites with the same *var*codes caused clinical episodes after the outbreak. Persistent disease transmission over time (Fig. 3b) and large distances (Fig. 3e-f) was found as the same *var*codes were identified post outbreak and sometimes in different sampling locations. The median time between first and last identification of the same *var*codes with any clinical case was 216 days or approximately 7 months (range = 190 – 823 days) during the study period. For example, the outbreak *var*code1 identified in Esmeraldas City was also identified in San Lorenzo, Esmeraldas (~150km from Esmeraldas city) in 2013, then Cascales, Sucumbios (>300km away) in 2014, and then in Tobar Donoso, Carchi (~150km away) in 2015 (Fig. 3b). This is in line with an earlier study using microsatellites that showed that *P. falciparum* isolates from the outbreak were genetically related to an Ecuadorian parasite isolate collected around 1990⁸. This result was also consistent with reports in Peru⁵¹ and Colombia^{18,52} where identical parasites were shown to be circulating up to five and eight years after first identification, respectively.

Figure 3. Reconstructing the disease transmission dynamics of *P. falciparum* reveals signatures of local transmission and sexual recombination after the outbreak. (a) A network visualization of the genetic relatedness of *P. falciparum* isolates at the threshold of $P_{TS} \geq 0.90$ to define varcodes (see Methods). Every node represents a *P. falciparum* isolate and an edge represents the P_{TS} value between two particular nodes/isolates. Isolates that cluster together (i.e., connected by edges) are considered to have the same varcode. Each color represents a different varcode. Two *P. falciparum* isolates belonging to varcode1 appear as outliers in the network due to undersampling of their DBL α types. N refers to the number of isolates. For comparison to BP_{TS} estimates see Supplementary Fig. S3-5 and S8. (b) The number of *P. falciparum* isolates within each varcode (i.e. size of circle) in each year and their persistence over time. The three varcodes (varcode1, varcode2, varcode3) identified in 2013 were identified again in 2014, and varcode1 was identified again in 2015. Two varcodes identified in 2014 (varcode6, varcode7) were also identified in 2015. (c) A network visualization of the genetic relatedness of varcodes at the threshold of $P_{TS} \geq 0.50$ to discriminate between recombinant varcodes and genetically distinct varcodes. Isolates/varcodes that cluster together (i.e., connected by edges) represent recombinants. (d) A clustered heatmap showing the genetic profiles of each *P. falciparum* isolate with rows representing each isolate and columns representing each DBL α type. Black and white denote the presence and absence of each type, respectively. Isolates that clustered together were more genetically similar (i.e., the same DBL α types were present). Similarly, varcodes that clustered together were more genetically similar (e.g. recombinants). (e) The spatial distribution of varcodes in each year, with the size of the circle representing the number of *P. falciparum* isolates sampled in a given location in each year and the pie chart depicting the proportion of each varcode identified. (f) Spatiotemporal genetic relatedness network of varcodes between 2013-2015. Every node represents a sampling location, the size of the circle represents the number of *P. falciparum* isolates

sampling in each location, the pie chart depicts the proportion of each *varcode* identified, and the weighted edges show genetically-related *varcodes* ($P_{TS} \geq 0.50$). (Online version: GIF animation depicts the spatiotemporal transmission dynamics chronologically during and after the outbreak).

Spatiotemporal genetic relatedness networks reveal signatures of recombination

We next reconstructed spatiotemporal genetic relatedness networks to explore signatures of local disease transmission and parasite microevolution after the outbreak. We applied a P_{TS} threshold of ≥ 0.50 to detect recombinant and/or genetically-related *varcodes*. By using the date and location where each isolate was collected, we animated the network chronologically to examine spatiotemporal disease transmission dynamics during the outbreak in 2013 and post-outbreak in 2014 and 2015 (Fig. 3c, 3f, GIF). This analysis resolved parasite microevolution relevant to disease transmission after the outbreak and revealed that four *varcodes* (*varcode3*, *varcode4*, *varcode6*, *varcode7*) were recombinants of the outbreak *varcode1* (Fig. 3c). The remaining *varcodes* did not cluster in the network and were genetically distinct. These patterns were confirmed using BP_{TS} estimates based on posterior means (Supplementary Fig. S8a) and their corresponding 95% HDPIs (Supplementary Fig. S5c-f and S8b-c). For the *varcodes* that were recombinants and/or genetically-related (i.e., cluster in the networks in Fig. 3c and Supplementary Fig. S8a), 70.9% (433/611) of the HDPIs were confidently above the threshold of $BP_{TS} \geq 0.50$ (Supplementary Fig. S5d). For *varcodes* that were genetically-distinct (i.e., do not cluster in the networks), 98.4% (368/374) of the HDPIs were confidently below the threshold of $BP_{TS} \leq 0.50$ (Supplementary Fig. S5f). As expected, the relatedness networks based on the 7 microsatellites did not resolve these patterns since all isolates were connected at the threshold of $P_{AS} \geq 0.50$ and genetic relatedness was overestimated (Supplementary Fig. S9a). Indeed, we were not always able to discriminate recombinants from genetically distinct parasites since several pairwise comparisons (25%, 307/1232 by P_{TS} and 16%, 196/1232 by BP_{TS}) that resulted in $P_{AS} \geq 0.50$ were found to be less related by *var* (P_{TS} or $BP_{TS} \leq 0.50$) (Supplementary Fig. S7). Furthermore, the majority of recombinants were classified as essentially clonal since they clustered together at the threshold of $P_{AS} \geq 0.80$ (allowing for differences in one microsatellite allele, Supplementary Fig. S6b).

We were able to detect that *varcode3* was in fact a recombinant of the outbreak *varcode1*, even though it was sampled in San Lorenzo during the same time period as the ongoing outbreak in 2013 (Fig. 3c, 3e). With the available data we were unable to confirm whether *varcode3* was ‘imported’ to San Lorenzo from Esmeraldas City. The most plausible explanation is that the recombination event occurred in San Lorenzo during the outbreak. Parasites with recombinant *varcode3* caused clinical episodes in multiple

locations in 2014 (Fig. 3e). The other recombinant *var*codes (*varcode4*, *varcode6*, *varcode7*) were identified after the outbreak in 2014 or 2015. Whether the recombinant *var*codes resulted from sexual recombination events between outbreak *varcode1* and genetically distinct parasites that were already circulating at low levels and/or in asymptomatic reservoirs in Ecuador (e.g. ⁵) or those that were previously imported could not be ascertained from the current study population. Interestingly, all the recombinant *var*codes were identified in San Lorenzo indicating this area is a transmission hot spot and a reservoir of parasites with diverse *var* repertoires (Fig. 3e).

To further visualize the genetic profiles of *var*codes and confirm the observed recombination signatures, we constructed a clustered heatmap based on the presence and absence of the 195 DBL α types across all isolates, such that isolates with similar genetic profiles cluster together (Fig. 3d). This analysis confirmed that in the case of isolates identified as recombinants of the outbreak *varcode1* (i.e., *varcode3*, *varcode4*, *varcode6*, *varcode7*), a proportion of outbreak DBL α types as well as different DBL α types were present. This is consistent with the generation of a new *varcode* through outcrossing between two genomes with different *var* repertoires by conventional meiosis (i.e., approximately $\geq 50\%$ sharing of DBL α types). By contrast, the genetic profiles of *varcodes* 2, 5, 8, and 9 had different DBL α types, leading to the definition as parasites with genetically distinct *var*codes. When using 7 microsatellites, these recombination signatures in the genetic profiles were not identified (Supplementary Fig. S9b).

Parasites with recombinant *var*codes most frequently caused *P. falciparum* clinical episodes following the 2012-13 outbreak

Prior to the outbreak there had been a steady decline in clinical cases, however, increased incidence of disease occurred after the outbreak. Therefore, we analyzed trends in the epidemiology of *P. falciparum* cases that occurred post-outbreak to understand if there were any key risk factors. Of the 25 individuals in our study that had a clinical *P. falciparum* episode after the outbreak in 2014 or 2015, we had age data for 18 individuals (72%, Table 1). There was no significant difference (Kruskal-Wallis test, $p = 0.65$) in the median age of individuals experiencing clinical episodes caused by parasites with the outbreak *varcode1* (19 years, range = 19 – 57 years, $N = 3$ patients), a recombinant of *varcode1* (i.e., *varcodes* 3, 4, 6, or 7) (25 years, range = 17 – 58 years, $N = 13$ patients), or by a different *varcode* (34.5 years, range = 32 – 37 years, $N = 2$ patients). A greater diversity of *var*codes (1-9 inclusive) was associated with incidence of clinical malaria post-outbreak than during the 2013 outbreak (*varcodes* 1,2,3). Indeed in 2014, 79% of the

cases sampled were caused by either parasites with the outbreak *varcode1* (21%) or with any of the four recombinants of *varcode1* (58%). The trend was similar in 2015 with 83% of the cases sampled caused by either parasites with the outbreak *varcode1* (33%) or with any of the four recombinants of *varcode1* (50%), although our sampling of reported cases in 2015 after January was limited. Importantly, overall we found that 80% of the cases we sampled after the outbreak were caused by either parasites with the outbreak *varcode1* (24%) or with any of the four recombinants of *varcode1* (56%), especially with *varcode7*. Thus, disease transmission after the outbreak in 2014 and into 2015 was predominantly associated with parasites with recombinant *var* repertoires of the outbreak clone.

Comparative analyses to South America: elucidating possible origins of Ecuadorian *varcodes*

We next examined the possible origins of the *varcodes* circulating locally in Ecuador by comparing to published data²⁸ from 128 *P. falciparum* isolates collected from neighboring South American *P. falciparum* populations (Fig. 4a). When we combined the sequences from Ecuadorian isolates with those previously sequenced, we identified 543 unique *var* DBL α types in South America (compared to 458 in ²⁸). This is largely representative of the *var* diversity circulating in South American *P. falciparum* populations, as indicated by sampling accumulation curves approaching saturation (Supplementary Fig. S10). *Var*coding resolved a total of 97 *varcodes* in South America ($P_{TS} \geq 0.90$), with those identified in each country ranging from 9 in Ecuador and Venezuela, to 56 *varcodes* in French Guiana (Supplementary Fig. S11). The number of *var* DBL α types per isolate repertoire was low in all countries and indicative of only one *P. falciparum* genome infecting an individual (i.e. repertoire size ≤ 60 , Fig. 4b, Supplementary Table S1). We found that the median number of *var* DBL α types in each isolate ranged from 36.5 in Venezuela to 48.0 in French Guiana and the maximum number of *var* DBL α types per isolate was 42 or 43 in all countries except French Guiana that appeared to have more multi-genome infections (max = 92). The median repertoire size was in the range of the number of *var* genes seen in the genome of the Honduran laboratory reference strain HB3 (N=42 excluding *var2csa*, ^{32,33}). An unrooted phylogenetic neighbor-joining tree revealed distinct clusters of genetically-related isolates in South America generally clustering by country (Fig. 4c), consistent with previous analyses in the region demonstrating geographic population structure²⁸.

To examine whether any of the parasites with Ecuadorian *varcodes* were genetically related to other South American parasites, we constructed regional spatiotemporal relatedness networks and applied our threshold of $P_{TS} \geq 0.50$ (Fig. 4d). We identified a genetically-related Peruvian *P. falciparum* isolate that clustered with the outbreak and recombinant *varcodes* ($P_{TS} = 0.66-0.75$ with *varcode1* isolates). This is consistent with previous analyses using microsatellites showing the outbreak source was possibly a residual parasite lineage circulating in Peru in 1999-2000 and in Ecuador in the early 1990s^{8,13}. The fact that these isolates did not cluster with anything else suggests the outbreak may have been caused locally due to circulating parasites in unidentified reservoirs. *Varcode5*, which was sampled in the Amazon had a very different genetic profile to the other *varcodes* ($\geq 83\%$ unique types, Fig. 3d). It is worth noting that the parasite isolate with *varcode5* also had a different genetic background with respect to its microsatellite haplotype as compared to the rest of the Ecuadorian isolates (3 of 7 unique microsatellite alleles). The epidemiological data for the putative infection location was recorded as possibly imported from Peru (EC40, Table 1). However, our analysis would suggest that this *varcode* is more related to *P. falciparum* populations from French Guiana and Venezuela. Nonetheless, Peruvian isolates also clustered with *P. falciparum* isolates from French Guiana even though they did not cluster with *varcode5* and we found that 53% of the types identified in *varcode5* were also identified in Peruvian parasites (Fig. 4e). We were able to determine a putative importation of a Colombian parasite (*varcode9*, EC53, Table 1, Fig. 4d). The *varcodes* 2 and 8 did not cluster with any other South American isolate suggesting parasites with these *varcodes* may be local to Ecuador. The epidemiological data for the putative infection location for *varcode8* was recorded as Colombia (EC49 and EC50, Table 1). Although it did not cluster with Colombian *P. falciparum* isolates in our network analysis, we found that 67% of the types identified in *varcode8* (collapsing all 3 isolates) were also identified in Colombian parasites (Fig. 4e). For *varcode2*, we found that 44% of the types identified in *varcode2* were also identified in Colombia (Fig. 4e), providing evidence that parasites with *varcode2* and *varcode8* may represent residual parasites previously or recently imported from Colombia.

Figure 4. Relatedness networks in South America provide insights into the origins of the Ecuadorian varcodes and diversity patterns in South America. (a) A map showing the study sites across South America. The dates of sample collection in these countries occurred from 2002 to 2008, around five to thirteen years prior to the sample collection of the Ecuadorian isolates. N refers to the number of isolates. (b) Repertoire sizes are shown for each *P. falciparum* isolate, from which multiplicity of infection can be inferred (i.e., ~40-60 var DBLα types per *P. falciparum* genome). (c) An unrooted neighbor-joining tree shows the relatedness patterns among South American *P. falciparum* isolates. (d) A network visualization of the genetic relatedness of South American varcodes at the threshold of $P_{TS} \geq 0.50$. Due to the differences in time points and sample collections, for this analysis a genetically-related parasite did not necessarily reflect a recent transmission event. Every node represents a *P. falciparum* isolate and an edge represents the P_{TS} value between two particular nodes/isolates. Isolates/varcodes that cluster together (i.e., connected by edges) are genetically-related. (e) Relatedness of Ecuadorian varcodes to *P. falciparum* populations from South America. The relatedness of each varcode was measured by first concatenating all possible var DBLα types that were identified in the *P. falciparum* isolates comprising each varcode as well as concatenating all var DBLα types that were identified in each country. Then P_{TS} was calculated between each varcode and each country. The color gradient denotes the P_{TS} value for a particular comparison, with darker shades showing higher relatedness and the different colors corresponding to the different country comparisons. N refers to the number of var DBLα types. (f) A clustered heatmap showing the patterns of diversity across South America, with a row representing all the types identified in a particular country, and columns representing each of the 543 DBLα types. Black and white denote the presence and absence of each type, respectively. The total number of unique var DBLα types identified in each country ranged from 112 in Colombia to 249 in French Guiana. Countries that cluster together share more var DBLα types.

Conservation of individual DBL α types in space and time

Of significance, we found the same *var* DBL α types (56% of types identified in Ecuador, N=110/195, Fig. 4f) and *var*codes that were genetically related to other South American parasites and sampled up to 13 years before (Fig. 4d-e) were contributing to local transmission in Ecuador. Indeed, conservation of the DBL α types in all *var*codes was observed in space and time to varying degrees (Fig. 4e). We also found conservation of DBL α types on a continental scale with the number of the same types identified in any two or three countries being 124 (23%) and 59 (11%), respectively (Fig. 4f). Moreover, even DBL α types that were identified in relatively low frequencies in Ecuador (i.e., identified in a few *P. falciparum* isolates) were also identified in the South America dataset, often at relatively high frequencies in other countries (Supplementary Fig. S12). These patterns are consistent with some parasite *var*codes being putative importations and/or highly-related to parasites in other countries, as described previously.

No evidence of recombination in any DBL α type over the course of the outbreak

The fact that the outbreak *var*code1 was clonal provided a unique opportunity to examine the conservation of its 47 individual DBL α types after transmission to multiple human hosts over the course of the outbreak and up to two years after (N=36 infections). DBL α types are very variable, with the average pairwise identity of the sequences encoding this domain being approximately 42%³³. The cut-off of 96% sequence identity in our pipeline has been routinely used to identify the same DBL α types within the limits of sequencing errors and PCR artifacts (^{27,53}, Methods). Our data show that the same DBL α types were conserved in isolates with *var*code1 over the course of the outbreak (N=30 infections) and up to two years after the outbreak in the other six cases with *var*code1 (Supplementary Fig. S13a). In total, we report 1,283 independent observations of the same 41 DBL α types identified in 36 cases with *var*code1 assessed over 2 to 3 years (any type was identified in a median of 34 infections, ranging from 1 to all 36 infections) (Supplementary Fig. S13a). Given our threshold of $P_{TS} \geq 0.90$ to identify putatively identical *var* DBL α repertoires, only 10 of the 47 DBL α types found in the outbreak clone were not found in the majority of the isolates with *var*code1 (median frequency = 1, max = 6, Supplementary Fig. S13a). Four of these were identified in the cases with recombinant *var*codes (median frequency = 17.5, range = 6-19), two of which were also identified in the Peruvian isolate that clustered with the outbreak/recombinants in Fig. 4d, suggesting they are also conserved. The other 6/10 were singletons in the entire dataset (including the South America data) and could be a consequence of the PCR with degenerate primers and/or low-quality DNA.

Among the 15 cases with recombinant *var*codes (*var*codes 3,4,6, and 7), we report 339 independent observations of 39 of the 47 outbreak DBL α types. There were 26-27 outbreak types shared between *var*code1 and the recombinant *var*codes 3,4, and 7, and 17 types between *var*code1 and *var*code6 (Supplementary Fig. S13b). These findings highlight the conservation of DBL α types of the outbreak clone in space and time, a pattern further supported by the fact that 31 of 47 outbreak DBL α types were found among previously recorded South American isolates (in 2002 to 2008²⁸; see Supplementary Fig. S13c). This evidence points to conservation of these types despite the many transmission events that must have occurred between 2002 and 2013-15 across South America. We found that the highest number of outbreak DBL α types were identified in Peru (N=28, 60%) followed by Colombia (N=14, 30%) and Venezuela (N=8, 17%), and the lowest in French Guiana (N= 2, 4%, Supplementary Fig. S9c).

Discussion

Our investigation of the spatiotemporal incidence of clinical episodes of *P. falciparum* in Ecuador by *var*coding supports the view that disease transmission after the 2012-2013 outbreak was sustained predominantly by Ecuadorian parasites. We observed persistence of the outbreak clonal lineage (identified with the same *var*codes) and outcrossing of this lineage with other locally circulating parasites causing clinical cases, rather than recent importation of clonally transmitted parasites. Our results point to the need for resources to be focused locally in Ecuador to uncover the circulating asymptomatic reservoirs of infection identified by *var*code recombination events. A role for human mobility must also be considered in the spread of *P. falciparum* as parasites with the same *var*codes were also observed across large distances (~150-300km).

San Lorenzo was found to be a transmission hotspot likely due to mining activities and occupation-related travel in these areas, as well as its proximity to the Ecuador-Colombia border^{3,50,54-56}. These findings point to San Lorenzo for both genetic surveillance and targeted interventions. Importantly, our results provide strong evidence for ongoing transmission in Ecuador and provide the first baseline characterization of *P. falciparum* antigenic diversity and parasite *var*codes circulating in the northwest coast and Amazon regions of Ecuador. It is worth noting that several recent outbreaks have occurred in the same areas of Ecuador (in 2016¹⁰, 2018⁵⁷, 2019⁶ and 2020⁵⁸) with malaria transmission remaining unstable rather than endemic.

We demonstrate that 58% of clinical cases sampled in 2014 immediately after the outbreak in Ecuador were caused by parasites with recombinant *var*codes. The same trend was observed in 2015, although our sampling was limited relative to the number of reported cases for the months sampled. We ask is this a chance finding or a consequence of immune evasion? We hypothesize that these parasites with recombinant *var*codes have new combinations of *var* DBL α types that may provide an immunological advantage in a population with variable levels of immunity, as would be expected during elimination campaigns⁵⁹.

Network analyses and computational models combining evolution and epidemiology point to variant-specific immune selection defining *var* population structure^{60,61}. In scenarios where transmission is low and there is limited *var* diversity, the chance of recombination is also relatively low and thus immune selection is weaker. Selection in the context of an E-2020 country like Ecuador is likely variable due to the recent transition from moderate to epidemic transmission over the last 15 years¹ with declining population immunity as a consequence of approaching elimination. Low population immunity would select for circulating clones leading to high similarity between *var* repertoires. Here, the signatures of recently recombined genomes with the outbreak clone that we detect in four *var*codes were clearly associated with the continued incidence of clinical disease after the outbreak. Thus parasites with recombinant *var*codes may present a risk factor for disease. We hypothesize that they may be better able to evade variant-specific immunity to PfEMP1 variants expressed by the outbreak clone by sharing 50% or less of the DBL α types of the outbreak clone. The availability of *var* DBL α sequences provides the potential to test this hypothesis by serological methods that detect variant-specific immunity. In this context, inferences based on neutral markers do not give information about immune evasion genes and variant-specific immunity, whereas *var*codes identify circulating variants to which individuals can become immune. e.g. as was done in serologic studies in Papua New Guinea⁶² and the Brazilian Amazon⁶³.

A previous study that included some of the same *P. falciparum* isolates from Esmeraldas City and San Lorenzo in the northwest coast described three main genetic clusters based on microsatellite genotyping⁵⁰. In contrast, when considering the same isolates, *var*coding resolved six *var*codes circulating in the northwest coast. In this study, as expected comparative analyses to “gold-standard” microsatellite genotyping demonstrated that *var*coding provided higher discriminatory resolution to detect fine-scale genetic variation and describe parasite microevolution related to disease transmission in epidemiological time. These findings were in accordance with the rapidly evolving *var* multigene family due to immune

selection^{30,60,64}, the overall number of microsatellite loci examined and the fact that two loci were fixed in the population^{8,50} decreasing the denominator of polymorphic microsatellite markers from seven to five. Similar observations have been described in an earlier study in Venezuela where sympatric parasites identical at neutral microsatellite loci were shown to have very different *var* repertoires⁶⁵. These South American genetic epidemiology studies can be compared to those of a peri-urban area in Senegal where reduced transmission has resulted from intense vector and antimalarial control. In Senegal, highly related genotypic clusters of *P. falciparum* isolates were resolved by both *var* and use of a 24-SNP barcode as neutral loci rather than microsatellite loci⁶⁶. Genotyping *var* loci under immune selection in conjunction with neutral loci proves to be informative to validate clonality and avoids the need for whole genome sequencing as a potential rapid response to an outbreak investigation. Even in areas where WGS is routinely conducted, *var*coding has the potential to identify relevant samples for downstream WGS as part of a larger and complementary genomic surveillance framework.

In contrast to the microevolution of parasite *var*codes, we documented conservation of individual *var* DBL α types over space and time. This is similar to Africa^{27,29} where we see highly variable but many conserved DBL α types yet rapidly evolving *var* repertoires or *var*codes. Here we observed conservation of the outbreak *var*code1 clone DBL α types after transmission within and between multiple human hosts on the scale of months to two years in Ecuador (1,283 independent observations of the same types when examining 36 cases with *var*code1). Moreover, 110 of the 195 (56.4%) Ecuadorian DBL α types were identified across South America over 5 to 13 years (2,130 independent observations of the Ecuadorian DBL α types in 128 different infections from²⁸). These observations are consistent with balancing selection⁶⁷ and in line with other studies demonstrating high levels of *var* DBL α sequence conservation on the time scale of up to 23 years in Brazil^{34,68}. Another independent example of conservation of DBL α sequences over time is found in a peri-urban area of Senegal over 7 years⁶⁶. This pattern of conservation of *var* DBL α types observed in diverse sites confirms the potential of this genetic marker for malaria surveillance.

Of note, the observed conservation of DBL α types in natural infections in humans in Africa^{27,29,69} and South America would not be predicted by data from *in vitro* passage of long-term cloned lines of *P. falciparum* showing high rates of mitotic recombination within *var* genes^{70–72}. Indeed, Claessens et al⁷¹ reported a very high rate of $2-8 \times 10^{-3}$ mitotic recombinants created per erythrocytic cycle for three of four long-term *in vitro* cultured lines, “with DBL α domains showing the most recombinations”. The clone

tree of HB3, selected for passage through mosquitos, did not show any recombinants suggesting that the high rates observed may relate to passage *in vitro*. While we identified conservation rather than diversification of *var* DBL α types in space and time, it is possible that mitotic recombinants are generated on the observed time scale but not transmitted. Such is the case in HIV (e.g. reviewed in ⁷³). Further exploration of within and between host diversification of *var* genes in natural infections requires a more detailed genomic approach analyzing recently adapted parasite lines.

This is the first case study demonstrating the translational application of the *var*code for outbreak surveillance in epidemic/unstable malaria transmission such as E-2020 settings. Specifically, data presented here have shown that *var*coding was able to resolve the population genetics of sustained disease transmission after an outbreak in Ecuador. We conclude that *var*coding requiring a single PCR with degenerate primers has great potential as a simple, cost effective and high-resolution approach to examine *P. falciparum* antigenic diversity in relation to population immunity as well as genome diversification by recombination. This will prove particularly useful in the context of changing patterns of human mobility and gene flow in the Americas where there is high demand for such a diagnostic surveillance method. The tool will also be useful in other epidemic or low transmission settings targeting elimination across the globe.

Here we show that *var* repertoire evolution via sexual recombination was associated with sustained transmission resulting in clinical infections. This is in line with our previous predictions of epidemic disease transmission in South America with parasites with novel *var* repertoires, or *var*codes²⁸, and serves as a warning of what may come in terms of monitoring the complexity of malaria resurgence in Latin America with both importation and microevolution of *P. falciparum*. Surveillance with neutral markers alone cannot investigate this potential risk related to recombination of variant antigen gene repertoires. The widespread migration of Venezuelans has already led to massive increases in *P. falciparum* cases in Brazil and Colombia⁷⁴, as well as in the Peru-Ecuador border³. Such migration points to the possibility of importation of diverse parasites with antigenically novel *var* genes as we predicted ²⁸, and is now also worrisome in the context of increased transmission of malaria in tropical regions due to the diversion of healthcare resources to the COVID-19 pandemic, as recently warned by the WHO⁷⁵. Going the distance to elimination must be supported by appropriate molecular surveillance.

Methods

Ethics statement

The *P. falciparum* isolates collected from individuals of all ages presenting with uncomplicated malaria cases were obtained from the malaria surveillance protocol approved by the Ethical Review Committee of Pontificia Universidad Católica del Ecuador. Ethical approval for this study was obtained from Pontificia Universidad Católica del Ecuador (Quito, Ecuador) and University of Melbourne (Melbourne, Australia). Written informed consent was provided by study participants and/or their legal guardians.

Study design, study area and sample collection

In this molecular epidemiological study, we examined samples that were collected passively from 2013 to 2015 in endemic areas of Ecuador, namely the northwest coast and the Amazon region, as part of ongoing malaria surveillance by the Ecuadorian Ministry of Health and patients showing at the local clinics. During 2013, the majority of samples were collected from an ongoing outbreak that occurred in Esmeraldas city, Esmeraldas province. Details of the sample collection are described in Sáenz et al⁸. In 2014 and 2015, the “post-outbreak” samples were mostly collected from across the northwest coast with few from the Amazon region, reflective of the epidemiology of clinical *P. falciparum* cases in Ecuador (Northwest coast: 20 and 173 cases; Amazon: 11 and 16 cases in 2014 and 2015, respectively). Details of the sample collection are described in Vera-Arias et al⁵⁰. It is worth noting that although cases were reported throughout 2015, samples were not available for many of them due to logistical challenges. Briefly, subjects who were diagnosed with falciparum malaria either by light microscopy or a rapid diagnostic test were asked if they wanted to participate in the study and signed an informed consent form. The participants had to: 1) live in the areas where the samples were taken, 2) be over 2 years old, 3) agree to participate, 4) sign an informed consent form, 5) give a blood sample, 6) be able to understand the informed consent. From each participant a blood sample (either venous blood or dried blood spot) was collected as well as their responses to a basic demographic questionnaire (including places recently travelled and their address). The positivity of all samples was confirmed in the laboratory by microscopy and PET-PCR⁷⁶.

DNA extraction

Genomic DNA was extracted from venous blood or dried blood spot samples using a QIAamp DNA MINI KIT (QIAGEN, Germantown, MD, USA) using the protocol recommended by the manufacturer.

Microsatellite genotyping

The same *P. falciparum* isolates in this study were previously genotyped for seven putatively neutral microsatellite markers (TA1, POLYα, PfpK2, TA109, 2490, C2M34, C3M39) as described in^{8,50}. These previously published microsatellite genotyping data^{8,48,75} were used in the present analyses.

Var genotyping and related data processing

For *var* genotyping, the DBLα domains of antigen-encoding *var* genes were amplified and sequenced on a MiSeq 2x300bp paired-end Illumina platform as described in^{29,60}. We obtained 6,929,470 raw sequence reads, which we cleaned and processed using the DBLαCleaner pipeline^(60, <http://github.com/Unimelb-Day-Lab/DBLαCleaner>). Our customized bioinformatic pipeline has been described in detail in⁶⁰. Briefly, we de-multiplexed and merged the reads as well as removed low-quality reads and chimeras using several filtering parameters (see Supplementary Fig. S1 for details). This resulted in 2,141 cleaned DBLα sequences (Supplementary Fig. S1). To identify distinct or unique DBLα types (i.e., unique genetic variants), we clustered the DBLα sequences from Ecuadorian *P. falciparum* isolates with 5,699 previously published²⁸ DBLα sequences from other South American countries (Colombia (N=21 isolates), French Guiana (N=76 isolates), Peru (N=21 isolates), and Venezuela (N=10 isolates) at the standard 96% sequence identity²⁷ using the clusterDBLalpha pipeline (<http://github.com/Unimelb-Day-Lab/clusterDBLalpha>). All the cleaned DBLα sequences in this study have been submitted to the DDBJ/ENA/GenBank (BioProject Number: PRJNA642683).

We further curated our dataset by translating the DBLα types into amino acid sequences using the classifyDBLalpha pipeline (<http://github.com/Unimelb-Day-Lab/classifyDBLalpha>) and removing any DBLα types that were non-translatable (N=4). In addition, any *P. falciparum* isolate with low sequencing quality (< 10 DBLα types) was removed from the analysis. *Var*coding was completed for a total of 70 *P. falciparum* Ecuadorian isolates, however, 12 *P. falciparum* isolates with low DNA quality and/or poor sequencing quality were removed and we obtained *var* DBLα data for 58 isolates (82.9%). A total of 543 unique DBLα types identified in the 186 South American *P. falciparum* isolates (N = 58 Ecuadorian *P. falciparum* isolates from this study, N = 128 South American *P. falciparum* isolates previously published in²⁸) were used for subsequent *var* analyses at the regional-level, and only the 195 unique DBLα types identified in Ecuador were used for Ecuador-specific analyses.

A further validation of our sequencing and genotyping methodology was undertaken by analyzing the isolate genomic proportions of DBL α types classified as upsA and non-upsA using our classifyDBLalpha pipeline²⁹ and are described in Supplementary Text 1.

Genetic relatedness analyses

Measuring pairwise type or allele sharing

To estimate genetic relatedness among isolates, we calculated the similarity index Pairwise Type Sharing (P_{TS})²⁶, as adapted by He et al⁶⁰ to account for differences in *var* DBL α sampling across isolates in the case of *var*, and Pairwise Allele Sharing (P_{AS})⁷⁷ in the case of microsatellites and constructed similarity matrices. Because *P. falciparum* is haploid we refer to *var* repertoires and microsatellite haplotypes in the same manner, i.e. a *var* repertoire of DBL α types or a multilocus haplotype of microsatellite alleles transmitted together. Every parasite isolate was compared to every other parasite isolate in the population to determine the proportion of shared loci (i.e. *var* types or microsatellites), but only “retrospective” pairwise comparisons were analyzed as a conservative approach to account for time order for the Ecuador-specific analyses. In other words, since we were interested in using the outbreak as a baseline this approach allowed us to look at genetic relatedness of an isolate with everything else circulating before its identification. For the comparative analyses with South American data we included all pairwise comparisons and did not correct for time order due to the differences in sampling time points (five to thirteen years before the Ecuadorian isolates were collected).

Measuring Bayesian pairwise type sharing

Genetic relatedness (or repertoire overlap) can also be explored more rigorously with unbiased Bayesian pairwise type sharing (BP_{TS} , Johnson et al *in preparation*), which accounts for uncertainty in P_{TS} estimates due to differences in isolate repertoire size. This approach uses Bayesian inference methods, which estimate repertoire overlap and uncertainty⁷⁸, and uses them in a subsequent P_{TS} calculation, carrying that uncertainty forward. The prior distribution for repertoire size, used in inference, was informed by observations as follows. First, the median observed repertoire size in Ecuadorian isolates was 37 types, ranging from 11 to 43 (Table S1). Second, the number of expected *var* genes with DBL α domains from whole genome sequencing data of the Honduran laboratory reference strain HB3 was 42³³ and 50 *var* genes based on long-read PacBio sequencing of HB3⁷⁹. And third, based on our sequencing data of 37

technical replicates of HB3, the median repertoire size or number of DBL α types per isolate was 39 (range 36-41 types), with 40 types consistently identified in the majority of replicates (range 21-37 replicates) (Supplementary Text 1). We therefore used a uniform prior on repertoire sizes between 40 and 50 types, combined with the general Bayesian repertoire overlap framework⁷⁸ to produce unbiased estimates (posterior means). These were used to confirm our P_{TS} estimates. As expected, the P_{TS} and BP_{TS} estimates were positively correlated (Pearson's correlation coefficient = 0.919, $p < 0.001$, Supplementary Fig. S3). To measure uncertainty in central estimates, we computed a 95% highest density posterior interval (HDPI), a Bayesian version of confidence intervals, for each pairwise estimate. Like a frequentist confidence interval, the width of the HDPI provides a measure of uncertainty of each pairwise comparison. All posteriors were sampled using MCMC.

Interpretation of genetic relatedness measures

Since *P. falciparum* undergoes sexual recombination (i.e. meiosis) in the mosquito, conventional genetic interpretations can be applied to P_{TS} and P_{AS} . Therefore, pairwise isolate comparisons resulting in a P_{TS} or P_{AS} of 0 (0% shared loci), 0.5 (50% shared loci), and 1 (100% shared loci) would indicate genetically distinct isolates, recombinant isolates, and clones or genetically-identical isolates, respectively. We applied this approach to define “*var*codes” as groups of isolates sharing $\geq 90\%$ of their DBL α types ($P_{TS} \geq 0.90$) to identify putatively identical genomes within the margin of error of detection of 1-5 DBL α types in an isolate (i.e., within the margin of error of detection of all DBL α sequences in an isolate using degenerate primers for *var* DBL α PCR). We confirmed our interpretations of *var*codes, recombinants and genetically distinct isolates by comparing to unbiased BP_{TS} estimates (posterior means) and examining the lower and upper bounds of the 95% HDPIs as a measure of the statistical uncertainty of each pairwise comparison.

In the case of microsatellite genotyping, we applied a threshold of $P_{AS} \geq 0.90$ (sharing $\geq 90\%$ of their microsatellite alleles) to define clones and also applied a threshold of $P_{AS} \geq 0.80$ to identify putatively identical genomes within the margin of error of detection of 1 microsatellite allele in an isolate (for comparative purposes to the threshold for *var*, Supplementary Fig. S2). It is important to note that 2 out of the 7 microsatellite markers genotyped were fixed in the population⁸.

Visualization of genetic relatedness networks

To visualize the genetic relatedness among isolates as determined by P_{TS} or P_{AS} , we constructed networks using the R packages *ggraph*⁸⁰ and *tidygraph*⁸¹ where isolates are depicted as nodes and edges as the P_{TS}

or P_{AS} values at a given threshold. The R package *ggspatial*⁸² was used to plot spatial networks using latitude/longitude coordinates for sampling location. To visualize genetic relatedness of parasites over time we used the R package *gganimate*⁸³ to construct spatiotemporal relatedness networks. We generated a clustered heatmap based on the presence/absence matrix of DBL α types to visualize the genetic profiles of each isolate in Ecuador and each country in South America using the R package *pheatmap*⁸⁴ and the “complete” clustering method. Unrooted neighbor-joining phylogenetic trees based on pairwise genetic distance (calculated as $1 - P_{TS}$) were constructed using the R packages *ape*⁸⁵ and *ggtree*^{86,87}.

Statistical analysis

We used R version 3.5.2 for all analyses⁸⁸. The package *dplyr*⁸⁹ was used for further data curating. We used base R and the R package *epiR*⁹⁰ to perform chi-squared tests for univariate analyses of categorical variables to compare proportions and for non-parametric tests to compare distributions of continuous variables between two groups (Mann-Whitney U test) or among k groups (Kruskal-Wallis test) with a Bonferroni correction for multiple comparisons. To evaluate how well we sampled the true pool of *var* DBL α diversity (i.e. the true number of genetic variants circulating) in Ecuador and in South America, we used the R package *vegan*⁹¹ to generate species accumulation curves.

Data availability

The cleaned DBL α sequences generated in this study from Ecuadorian *P. falciparum* isolates have been submitted to the DDBJ/ENA/GenBank (BioProject Number: PRJNA642683). The python code for the DBL α Cleaner pipeline is available on GitHub at <https://github.com/Unimelb-Day-Lab/DBLaCleaner>. The python code for the clusterDBLalpha pipeline is available on GitHub at <https://github.com/Unimelb-Day-Lab/DBLalpha>. The python code for the classifyDBLalpha pipeline is available on GitHub at <https://github.com/Unimelb-Day-Lab/classifyDBLalpha>. All other deidentified data and analysis code are available on the open-source GitHub repository at <https://github.com/shaziaruybal/varcode-ecuador>.

References

1. Organization, W. H. *Eliminating Malaria*. <https://www.who.int/malaria/publications/atoz/eliminating-malaria/en/> (2016).

- 677 2. Espinoza, J. Malaria Resurgence in the Americas: An Underestimated Threat. *Pathogens* **8**, 11 (2019).
- 678 3. Jaramillo-Ochoa, R. *et al.* Effects of Political Instability in Venezuela on Malaria Resurgence at Ecuador–
679 Peru Border, 2018. *Emerg Infect Dis* **25**, 834–836 (2019).
- 680 4. Ferreira, M. U. & Castro, M. C. Malaria Situation in Latin America and the Caribbean: Residual and
681 Resurgent Transmission and Challenges for Control and Elimination. *Methods Mol Biology Clifton N J*
682 **2013**, 57–70 (2019).
- 683 5. Sáenz, F. E. *et al.* Malaria epidemiology in low-endemicity areas of the northern coast of Ecuador: high
684 prevalence of asymptomatic infections. *Malaria J* **16**, 300 (2017).
- 685 6. MSP. *Gaceta Epidemiológica Semanal No. 52.* (2019).
- 686 7. PAHO. *Report on the Situation of Malaria in the Americas 2012.* (2013).
- 687 8. Sáenz, F. E. *et al.* Clonal population expansion in an outbreak of Plasmodium falciparum on the
688 northwest coast of Ecuador. *Malaria J* **14**, 497 (2015).
- 689 9. MSP. *Gaceta Epidemiológica Semanal No. 52.* (2016).
- 690 10. MSP. *Gaceta Epidemiológica Semanal No. 52.* (2017).
- 691 11. Organization, W. H. *World Malaria Report 2019.* (2019).
- 692 12. Organization, W. H. THE E-2020 INITIATIVE OF 21 MALARIA-ELIMINATING COUNTRIES: 2019 progress
693 report. (2019).
- 694 13. Griffing, S. M. *et al.* South American Plasmodium falciparum after the Malaria Eradication Era: Clonal
695 Population Expansion and Survival of the Fittest Hybrids. *Plos One* **6**, e23486 (2011).
- 696 14. Larrañaga, N. *et al.* Genetic structure of Plasmodium falciparum populations across the Honduras–
697 Nicaragua border. *Malaria J* **12**, 354 (2013).
- 698 15. Obaldia, N. *et al.* Clonal outbreak of Plasmodium falciparum infection in eastern Panama. *J Infect Dis*
699 **211**, 1087–96 (2014).
- 700 16. Okoth, S. A. *et al.* Molecular Investigation into a Malaria Outbreak in Cusco, Peru: Plasmodium
701 falciparum BV1 Lineage is Linked to a Second Outbreak in Recent Times. *Am J Tropical Medicine Hyg* **94**,
702 128–31 (2015).
- 703 17. Baldeviano, G. C. *et al.* Molecular Epidemiology of Plasmodium falciparum Malaria Outbreak, Tumbes,
704 Peru, 2010–2012. *Emerg Infect Dis* **21**, 797–803 (2015).
- 705 18. Chenet, S. M., Taylor, J. E., Blair, S., Zuluaga, L. & Escalante, A. A. Longitudinal analysis of Plasmodium
706 falciparum genetic variation in Turbo, Colombia: implications for malaria control and elimination. *Malaria*
707 *J* **14**, 363 (2015).

708 19. Carter, T. E. *et al.* Genetic Diversity of *Plasmodium falciparum* in Haiti: Insights from Microsatellite
709 Markers. *Plos One* **10**, e0140416 (2015).

710 20. Redmond, S. N. *et al.* De Novo Mutations Resolve Disease Transmission Pathways in Clonal Malaria.
711 *Mol Biol Evol* **35**, 1678–1689 (2018).

712 21. Chenet, S. M., Schneider, K. A., Villegas, L. & Escalante, A. A. Local population structure of
713 *Plasmodium*: impact on malaria control and elimination. *Malaria J* **11**, 412 (2012).

714 22. Barton, N. Understanding Adaptation in Large Populations. *Plos Genet* **6**, e1000987 (2010).

715 23. Chen, D. S. *et al.* A Molecular Epidemiological Study of var Gene Diversity to Characterize the
716 Reservoir of *Plasmodium falciparum* in Humans in Africa. *Plos One* **6**, e16629 (2011).

717 24. Kirk, H. & Freeland, J. R. Applications and Implications of Neutral versus Non-neutral Markers in
718 Molecular Ecology. *Int J Mol Sci* **12**, 3966–3988 (2011).

719 25. Nelson, C. S. *et al.* High-resolution micro-epidemiology of parasite spatial and temporal dynamics in a
720 high malaria transmission setting in Kenya. *Nat Commun* **10**, 5615 (2019).

721 26. Barry, A. E. *et al.* Population Genomics of the Immune Evasion (var) Genes of *Plasmodium falciparum*.
722 *Plos Pathog* **3**, e34 (2007).

723 27. Day, K. P. *et al.* Evidence of strain structure in *Plasmodium falciparum* var gene repertoires in children
724 from Gabon, West Africa. *Proc National Acad Sci* **114**, E4103–E4111 (2017).

725 28. Rougeron, V. *et al.* Evolutionary structure of *Plasmodium falciparum* major variant surface antigen
726 genes in South America: Implications for epidemic transmission and surveillance. *Ecol Evol* **7**, 9376–9390
727 (2017).

728 29. Ruybal-Pesántez, S. *et al.* Population genomics of virulence genes of *Plasmodium falciparum* in clinical
729 isolates from Uganda. *Sci Rep-uk* **7**, 11810 (2017).

730 30. Rorick, M. M. *et al.* Signatures of competition and strain structure within the major blood-stage
731 antigen of *Plasmodium falciparum* in a local community in Ghana. *Ecol Evol* **8**, 3574–3588 (2018).

732 31. Tonkin-Hill, G. *et al.* Evolutionary analyses of the major variant surface antigen-encoding genes reveal
733 population structure of *Plasmodium falciparum* within and between continents. *Plos Genet* **17**, e1009269
734 (2021).

735 32. Kraemer, S. M. *et al.* Patterns of gene recombination shape var gene repertoires in *Plasmodium*
736 *falciparum*: comparisons of geographically diverse isolates. *Bmc Genomics* **8**, 45 (2007).

737 33. Rask, T. S., Hansen, D. A., Theander, T. G., Pedersen, A. G. & Lavstsen, T. *Plasmodium falciparum*
738 Erythrocyte Membrane Protein 1 Diversity in Seven Genomes – Divide and Conquer. *Plos Comput Biol* **6**,
739 e1000933 (2010).

740 34. Albrecht, L. *et al.* The South American Plasmodium falciparum var gene repertoire is limited, highly
741 shared and possibly lacks several antigenic types. *Gene* **453**, 37–44 (2010).

742 35. Gardner, M. J. *et al.* Genome sequence of the human malaria parasite Plasmodium falciparum. *Nature*
743 **419**, 498–511 (2002).

744 36. Biggs, B. A. *et al.* Antigenic variation in Plasmodium falciparum. *Proc National Acad Sci* **88**, 9171–9174
745 (1991).

746 37. Scherf, A., Lopez-Rubio, J. J. & Riviere, L. Antigenic variation in Plasmodium falciparum. *Annu Rev*
747 *Microbiol* **62**, 445–70 (2008).

748 38. Bull, P. C. *et al.* Parasite antigens on the infected red cell surface are targets for naturally acquired
749 immunity to malaria. *Nat Med* **4**, 358–360 (1998).

750 39. Marsh, K. & Howard, R. Antigens induced on erythrocytes by P. falciparum: expression of diverse and
751 conserved determinants. *Science* **231**, 150–153 (1986).

752 40. Forsyth, K. P. *et al.* Diversity of Antigens Expressed on the Surface of Erythrocytes Infected with
753 Mature Plasmodium Falciparum Parasites in Papua New Guinea. *Am J Tropical Medicine Hyg* **41**, 259–265
754 (1989).

755 41. Newbold, C. I., Pinches, R., Roberts, D. J. & Marsh, K. Plasmodium falciparum: The human
756 agglutinating antibody response to the infected red cell surface is predominantly variant specific. *Exp*
757 *Parasitol* **75**, 281–292 (1992).

758 42. IQBAL, J., PERLMANN, P., GREENWOOD, B. M. & BERZINS, K. Seroreactivity with the Plasmodium
759 falciparum blood stage antigen Pf332 in adults and children from malaria-endemic regions. *Clin Exp*
760 *Immunol* **94**, 68–74 (1993).

761 43. Bull, P. C., Lowe, B. S., Kortok, M. & Marsh, K. Antibody Recognition of Plasmodium falciparum
762 Erythrocyte Surface Antigens in Kenya: Evidence for Rare and Prevalent Variants. *Infect Immun* **67**, 733–
763 739 (1999).

764 44. Giha, H. A. *et al.* Antibodies to variable Plasmodium falciparum-infected erythrocyte surface antigens
765 are associated with protection from novel malaria infections. *Immunol Lett* **71**, 117–126 (2000).

766 45. Ofori, M. F. *et al.* Malaria-Induced Acquisition of Antibodies to Plasmodium falciparum Variant Surface
767 Antigens. *Infect Immun* **70**, 2982–2988 (2002).

768 46. Chattopadhyay, R. *et al.* Plasmodium falciparum Infection Elicits Both Variant-Specific and Cross-
769 Reactive Antibodies against Variant Surface Antigens. *Infect Immun* **71**, 597–604 (2003).

770 47. Mackintosh, C. L. *et al.* Failure to respond to the surface of Plasmodium falciparum infected
771 erythrocytes predicts susceptibility to clinical malaria amongst African children. *Int J Parasitol* **38**, 1445–
772 1454 (2008).

- 773 48. Otto, T. D. *et al.* Evolutionary analysis of the most polymorphic gene family in falciparum malaria.
774 *Wellcome Open Res* **4**, 193 (2019).
- 775 49. Tessema, S. K. *et al.* Phylogeography of var gene repertoires reveals fine-scale geospatial clustering of
776 *Plasmodium falciparum* populations in a highly endemic area. *Mol Ecol* **24**, 484–97 (2015).
- 777 50. Vera-Arias, C. A., Castro, L. E., Gómez-Obando, J. & Sáenz, F. E. Diverse origin of *Plasmodium*
778 *falciparum* in northwest Ecuador. *Malaria J* **18**, 251 (2019).
- 779 51. Branch, O. H. *et al.* *Plasmodium falciparum* genetic diversity maintained and amplified over 5 years of
780 a low transmission endemic in the Peruvian Amazon. *Mol Biol Evol* **28**, 1973–86 (2010).
- 781 52. Echeverry, D. F. *et al.* Long term persistence of clonal malaria parasite *Plasmodium falciparum*
782 lineages in the Colombian Pacific region. *Bmc Genet* **14**, 2 (2013).
- 783 53. Rask, T. S., Petersen, B., Chen, D. S., Day, K. P. & Pedersen, A. G. Using expected sequence features to
784 improve basecalling accuracy of amplicon pyrosequencing data. *Bmc Bioinformatics* **17**, 176 (2016).
- 785 54. Douine, M. *et al.* Illegal gold miners in French Guiana: a neglected population with poor health. *Bmc*
786 *Public Health* **18**, 23 (2017).
- 787 55. Douine, M. *et al.* Investigation of a possible malaria epidemic in an illegal gold mine in French Guiana:
788 an original approach in the remote Amazonian forest. *Malaria J* **18**, 91 (2019).
- 789 56. Mosquera-Romero, M., Zuluaga-Idárraga, L. & Tobón-Castaño, A. Challenges for the diagnosis and
790 treatment of malaria in low transmission settings in San Lorenzo, Esmeraldas, Ecuador. *Malaria J* **17**, 440
791 (2018).
- 792 57. MSP. *Gaceta Epidemiológica Semanal No. 52*. (2018).
- 793 58. MSP. Personal communication. (2020).
- 794 59. Deloron, P. & Chougnet, C. Is Immunity to malaria really short-lived? *Parasitol Today* **8**, 375–378
795 (1992).
- 796 60. He, Q. *et al.* Networks of genetic similarity reveal non-neutral processes shape strain structure in
797 *Plasmodium falciparum*. *Nat Commun* **9**, 1817 (2018).
- 798 61. Pilosof, S. *et al.* Competition for hosts modulates vast antigenic diversity to generate persistent strain
799 structure in *Plasmodium falciparum*. *Plos Biol* **17**, e3000336 (2019).
- 800 62. Barry, A. E. *et al.* The Stability and Complexity of Antibody Responses to the Major Surface Antigen of
801 *Plasmodium falciparum* Are Associated with Age in a Malaria Endemic Area. *Mol Cell Proteomics* **10**,
802 M111.008326 (2011).
- 803 63. Carlos, B. C. *et al.* Expressed var gene repertoire and variant surface antigen diversity in a shrinking
804 *Plasmodium falciparum* population. *Exp Parasitol* **170**, 90–99 (2016).

805 64. Artzy-Randrup, Y. *et al.* Population structuring of multi-copy, antigen-encoding genes in *Plasmodium*
806 *falciparum*. *Elife* **1**, e00093 (2012).

807 65. Tami, A., Ord, R., Targett, G. A. & Sutherland, C. J. Sympatric *Plasmodium falciparum* isolates from
808 Venezuela have structured var gene repertoires. *Malaria J* **2**, 7 (2003).

809 66. Bei, A. K. *et al.* *Plasmodium falciparum* population genetic complexity influences transcriptional profile
810 and immune recognition of highly related genotypic clusters. *Biorxiv* 2020.01.03.894220 (2020)
811 doi:10.1101/2020.01.03.894220.

812 67. Zilversmit, M. M. *et al.* Hypervariable antigen genes in malaria have ancient roots. *Bmc Evol Biol* **13**,
813 110 (2013).

814 68. Albrecht, L. *et al.* Extense variant gene family repertoire overlap in Western Amazon *Plasmodium*
815 *falciparum* isolates. *Mol Biochem Parasit* **150**, 157–165 (2006).

816 69. Bei, A. K. *et al.* Immune Characterization of *Plasmodium falciparum* Parasites with a Shared Genetic
817 Signature in a Region of Decreasing Transmission. *Infect Immun* **83**, 276–285 (2014).

818 70. Duffy, M. F., Byrne, T. J., Carret, C., Ivens, A. & Brown, G. V. Ectopic Recombination of a Malaria var
819 Gene during Mitosis Associated with an Altered var Switch Rate. *J Mol Biol* **389**, 453–469 (2009).

820 71. Claessens, A. *et al.* Generation of Antigenic Diversity in *Plasmodium falciparum* by Structured
821 Rearrangement of Var Genes During Mitosis. *Plos Genet* **10**, e1004812 (2014).

822 72. Zhang, X. *et al.* Rapid antigen diversification through mitotic recombination in the human malaria
823 parasite *Plasmodium falciparum*. *Plos Biol* **17**, e3000271 (2019).

824 73. Theys, K. *et al.* The impact of HIV-1 within-host evolution on transmission dynamics. *Curr Opin Virol*
825 **28**, 92–101 (2018).

826 74. Grillet, M. E. *et al.* Venezuela’s humanitarian crisis, resurgence of vector-borne diseases, and
827 implications for spillover in the region. *Lancet Infect Dis* **19**, e149–e161 (2019).

828 75. Organization, W. H. The potential impact of health service disruptions on the burden of malaria:
829 (2020).

830 76. Lucchi, N. W. *et al.* Molecular Diagnosis of Malaria by Photo-Induced Electron Transfer Fluorogenic
831 Primers: PET-PCR. *Plos One* **8**, e56677 (2013).

832 77. Ruybal-Pesántez, S. *et al.* Lack of Geospatial Population Structure Yet Significant Linkage
833 Disequilibrium in the Reservoir of *Plasmodium falciparum* in Bongo District, Ghana. *Am J Tropical*
834 *Medicine Hyg* **97**, 1180–1189 (2017).

835 78. Larremore, D. B. Bayes-optimal estimation of overlap between populations of fixed size. *Plos Comput*
836 *Biol* **15**, e1006898 (2019).

837 79. Otto, T. D. *et al.* Long read assemblies of geographically dispersed Plasmodium falciparum isolates
838 reveal highly structured subtelomeres. *Wellcome Open Res* **3**, 52 (2018).

839 80. Pedersen, T. L. *ggraph: An Implementation of Grammar of Graphics for Graphs and Networks*. (2020).

840 81. Pedersen, T. L. *tidygraph: A Tidy API for Graph Manipulation*. (2019).

841 82. Dunnington, D. *ggspatial: Spatial Data Framework for ggplot2*. (2018).

842 83. Pedersen, T. L. & Robinson, D. *gganimate: A Grammar of Animated Graphics*. (2020).

843 84. Kolde, R. *pheatmap: Pretty Heatmaps*. (2019).

844 85. Paradis, E. & Schliep, K. ape 5.0: an environment for modern phylogenetics and evolutionary analyses
845 in R. *Bioinformatics* **35**, 526–528 (2018).

846 86. Yu, G., Smith, D. K., Zhu, H., Guan, Y. & Lam, T. T. ggtree : an r package for visualization and annotation
847 of phylogenetic trees with their covariates and other associated data. *Methods Ecol Evol* **8**, 28–36 (2016).

848 87. Yu, G., Lam, T. T.-Y., Zhu, H. & Guan, Y. Two Methods for Mapping and Visualizing Associated Data on
849 Phylogeny Using Ggtree. *Mol Biol Evol* **35**, 3041–3043 (2018).

850 88. Team, R. C. *R: A language and environment for statistical computing*. (R Foundation for Statistical
851 Computing, 2019).

852 89. Wickham, H., François, R., Henry, L. & Müller, K. *dplyr: A Grammar of Data Manipulation*. (2020).

853 90. Stevenson, M. *epiR: Tools for the Analysis of Epidemiological Data*. (2020).

854 91. Oksanen, J. *et al.* *vegan: Community Ecology Package*. (2019).

855

Acknowledgements

The authors would like to thank the patients who contributed samples and the field teams who were involved in sample collections. We thank the Ecuadorian Ministry of Health, especially Luis E Castro, Julio Valencia and Javier Gomez Obando. We appreciate the support of the Australian Genome Research Facility for illumina sequencing of the samples. This research was financially supported by the Pontificia Universidad Católica del Ecuador (grant numbers: L13058, L13248, M13416, N13416, O13087[QINV0084] to F.E.S), the Fogarty International Center at the National Institutes of Health [Program on the Ecology and Evolution of Infectious Diseases (EEID), Grant number: R01-TW009670 to K.P.D.], and the National Institutes of Allergy and Infectious Diseases, National Institutes of Health (grant number: R01-AI084156 to K.P.D.). S.R-P. was supported by a Melbourne International Engagement Award from the University of Melbourne and gratefully acknowledges the J.D. Smyth Travel Award from the Australian Society for Parasitology that enabled her to travel to Ecuador to establish this collaborative project.

Author contributions

S.R-P. and K.P.D. conceived and designed the study; F.E.S. and C.A.V-A. carried out field work to obtain *P. falciparum* isolates and epidemiological metadata; S.R-P. and K.E.T. developed the varcoding methodology; S.R-P. and S.D. performed the molecular experiments and sequencing; S.R-P., F.E.S. and C.A.V-A. curated the clinical and epidemiological metadata; S.R-P. performed the formal data analysis and visualization; E.K.J. and D.B.L. designed and conducted the Bayesian statistical analysis; S.R-P., F.E.S., K.E.T. and K.P.D. contributed to the interpretation of the data; K.P.D. supervised the work; S.R-P., F.E.S. and K.P.D. acquired funding; S.R-P. and K.P.D. wrote the paper with contributions from all authors.

877 Table 1. Epidemiological characteristics of study participants.

Case	Month collected		Age range, years	Sex	Sampling location (County, Province)	Self-reported infection location*	varcode	Putative origin**
EC1	Jan 2013	Outbreak	NA	NA	Esmeraldas, Esmeraldas		varcode1	Local
EC2	Jan 2013	Outbreak	NA	NA	Esmeraldas, Esmeraldas		varcode1	Local
EC3	May 2013	Outbreak	>30-40	M	Esmeraldas, Esmeraldas		varcode1	Local
EC4	May 2013	Outbreak	>30-40	M	Esmeraldas, Esmeraldas		varcode1	Local
EC5	May 2013	Outbreak	>50-60	F	Esmeraldas, Esmeraldas		varcode1	Local
EC6	May 2013	Outbreak	>30-40	M	Esmeraldas, Esmeraldas		varcode1	Local
EC7	May 2013	Outbreak	>50-60	F	Esmeraldas, Esmeraldas		varcode1	Local
EC8	May 2013	Outbreak	>10-20	F	Esmeraldas, Esmeraldas		varcode1	Local
EC9	May 2013	Outbreak	>10-20	F	Esmeraldas, Esmeraldas		varcode1	Local
EC10	May 2013	Outbreak	>10-20	M	Esmeraldas, Esmeraldas		varcode1	Local
EC11	May 2013	Outbreak	>20-30	F	Esmeraldas, Esmeraldas		varcode1	Local
EC12	Jun 2013	Outbreak	>10-20	F	Esmeraldas, Esmeraldas		varcode1	Local
EC13	Jun 2013	Outbreak	>10-20	M	Esmeraldas, Esmeraldas		varcode1	Local
EC14	Jun 2013	Outbreak	>40-50	M	Esmeraldas, Esmeraldas		varcode1	Local
EC15	Jun 2013	Outbreak	>40-50	M	Esmeraldas, Esmeraldas		varcode1	Local
EC16	Jun 2013	Outbreak	>10-20	F	Esmeraldas, Esmeraldas		varcode1	Local
EC17	Jun 2013	Outbreak	>10-20	F	Esmeraldas, Esmeraldas		varcode1	Local
EC18	Jun 2013	Outbreak	>20-30	M	Esmeraldas, Esmeraldas		varcode1	Local
EC19	Jun 2013	Outbreak	>40-50	M	Esmeraldas, Esmeraldas		varcode1	Local
EC20	Jul 2013		>20-30	M	San Lorenzo, Esmeraldas		varcode2	Local
EC21	Jul 2013		>10-20	F	San Lorenzo, Esmeraldas		varcode2	Local
EC22	Jul 2013	Outbreak	>10-20	M	Esmeraldas, Esmeraldas		varcode1	Local
EC23	Jul 2013	Outbreak	>0-10	M	Esmeraldas, Esmeraldas		varcode1	Local
EC24	Jul 2013	Outbreak	NA	F	Esmeraldas, Esmeraldas		varcode1	Local
EC25	Sep 2013	Outbreak	>30-40	M	Esmeraldas, Esmeraldas		varcode1	Local
EC26	Oct 2013	Outbreak	>20-30	F	Esmeraldas, Esmeraldas		varcode1	Local
EC27	Oct 2013		>10-20	F	San Lorenzo, Esmeraldas		varcode3	Local
EC28	Oct 2013	Outbreak	>20-30	M	Esmeraldas, Esmeraldas		varcode1	Local
EC29	Nov 2013	Outbreak	>10-20	M	Esmeraldas, Esmeraldas		varcode1	Local
EC30	Nov 2013	Outbreak	>10-20	M	Esmeraldas, Esmeraldas		varcode1	Local
EC31	Nov 2013	Outbreak	>10-20	F	Esmeraldas, Esmeraldas		varcode1	Local
EC32	Nov 2013	Outbreak	>30-40	M	Esmeraldas, Esmeraldas		varcode1	Local
EC33	Nov 2013	Outbreak	>30-40	M	Esmeraldas, Esmeraldas		varcode1	Local
EC34	Jan 2014		NA	NA	Jaramijo, Manabi		varcode3	Local
EC35	Jan 2014		>50-60	F	San Lorenzo, Esmeraldas		varcode1	Local
EC36	Jan 2014		>10-20	M	San Lorenzo, Esmeraldas		varcode1	Local
EC37	Jan 2014		>30-40	M	San Lorenzo, Esmeraldas		varcode2	Local
EC38	Jan 2014		>10-20	M	San Lorenzo, Esmeraldas		varcode4	Local
EC39	Jan 2014		>10-20	M	San Lorenzo, Esmeraldas		varcode1	Local
EC40	Jan 2014		>30-40	F	Aguarico, Orellana	Peru	varcode5	French Guiana/Peru/Venezuela
EC41	Mar 2014		>20-30	M	San Lorenzo, Esmeraldas		varcode6	Local
EC42	May 2014		>50-60	F	Esmeraldas, Esmeraldas	Colombia	varcode3	Local
EC43	May 2014		NA	NA	Cascales, Sucumbios	San Lorenzo, Esmeraldas	varcode1	Local
EC44	Jul 2014		>30-40	F	San Lorenzo, Esmeraldas		varcode7	Local
EC45	Aug 2014		>20-30	F	San Lorenzo, Esmeraldas		varcode7	Local
EC46	Aug 2014		>20-30	M	San Lorenzo, Esmeraldas		varcode7	Local
EC47	Aug 2014		>50-60	M	San Lorenzo, Esmeraldas		varcode7	Local
EC48	Aug 2014		>20-30	M	San Lorenzo, Esmeraldas		varcode7	Local
EC49	Sep 2014		NA	F	Lago Agrio, Sucumbios	Colombia	varcode8	Local
EC50	Oct 2014		NA	NA	Lago Agrio, Sucumbios	Colombia	varcode8	Local
EC51	Oct 2014		>30-40	M	San Lorenzo, Esmeraldas	Colombia	varcode6	Local
EC52	Dec 2014		>20-30	M	San Lorenzo, Esmeraldas		varcode7	Local
EC53	Jan 2015		NA	NA	Lumbaqui, Sucumbios	Colombia	varcode9	Colombia
EC54	Jan 2015		>20-30	M	San Lorenzo, Esmeraldas		varcode7	Local
EC55	Jan 2015		>10-20	M	San Lorenzo, Esmeraldas		varcode7	Local
EC56	Nov 2015		>10-20	F	San Lorenzo, Esmeraldas		varcode6	Local
EC57	May 2015		NA	NA	Tobar Donoso, Carchi		varcode1	Local
EC58	May 2015		NA	NA	Tobar Donoso, Carchi		varcode1	Local

NA refers to no data collected for that particular epidemiological variable.

*The self-reported infection location was recorded at the time of sample collection based on the answers provided by the study participants.

**The putative origin as determined by varcoding.

Supplementary Information

Supplementary Figures

Figure S1. Bioinformatic sequence data processing flowchart. The flowchart shows the bioinformatic process to clean the raw illumina sequence reads with details on filter parameters for each step. This custom bioinformatic pipeline was developed to de-multiplex and remove PCR and sequencing artefacts and is described in detail in He et al ⁶⁰.

Figure S2. Sampling depth of *var* DBLα types in Ecuador. Accumulation curves depict the number of observed DBLα types plotted as a function of the number of DBLα sequences sampled.

Figure S3. Relatedness of isolates based on observed and unbiased Bayesian pairwise type sharing estimates. There was a positive correlation between the two relatedness measures (correlation coefficient = 0.919, $p < 0.001$). Circles colored in light red correspond to pairwise comparisons involving the two *P. falciparum* isolates with the smallest repertoire sizes (11 and 19 DBL α types), many of which had lower statistical confidence and the largest discrepancies between observed and unbiased estimates.

Figure S4. Genetic relatedness networks based on Bayesian pairwise type sharing estimates to define varcodes. A network visualization of the genetic relatedness of *P. falciparum* isolates at the threshold of (a) $BP_{TS} \geq 0.90$ to define clusters of genetically-related isolates based the mean posterior Bayesian PTS (BP_{TS}). Every node represents a *P. falciparum* isolate and an edge represents the BP_{TS} value between two particular nodes/isolates. Isolates that cluster together (i.e., connected by edges) are considered to be genetically identical (i.e., clones). Every color represents a different varcode. (b) The proportion of pairwise comparisons within a given 95% highest density posterior interval width, which provides a measure of the uncertainty of each pairwise estimate. Only varcodes with >1 *P. falciparum* isolate are shown. For the full distribution of within-varcode BP_{TS} estimates see Fig. S4b.

Figure S5. Distribution of observed and Bayesian pairwise type sharing estimates used to define varcodes, as well as recombinant, genetically-related, and genetically distinct varcodes. For observed P_{TS} estimates (a, c, e), the boxplots (in blue) show the median and interquartile range of the observed P_{TS} estimates (circles) for all pairwise comparisons among *P. falciparum* isolates for a given within-varcode or between-varcode comparison. For Bayesian PTS (BP_{TS}) estimates (b, d, f), the boxplots (in blue) show the median and interquartile range of the posterior mean BP_{TS} estimates. The circles show the BP_{TS} estimates for the lower and upper bound of the 95% highest density posterior intervals (HDPis). Circles showing all of the posterior mean BP_{TS} estimates are omitted for clarity. Dashed lines indicate the threshold of (a-b) P_{TS} or $BP_{TS} \geq 0.90$ and (c-f) P_{TS} or $BP_{TS} \geq 0.50$. For comparison to the genetic relatedness network edges see Figures 2a, 2c, S3a, S7a. In all panels, light grey circles correspond to pairwise comparisons involving the two *P. falciparum* isolates with the smallest repertoire sizes (11 and 19 DBL α types), many of which had lower statistical confidence and the largest discrepancies between observed and unbiased estimates. Black circles denote all other estimates. Only varcodes with >1 *P. falciparum* isolate are shown.

Figure S6. Genetic relatedness networks by microsatellite genotyping. A network visualization of the genetic relatedness of *P. falciparum* isolates at the threshold of (a) $P_{AS} \geq 0.90$ and (b) $P_{AS} \geq 0.80$ to define clusters of genetically-related isolates based on microsatellite genotyping. Every node represents a *P. falciparum* isolate and an edge represents the P_{AS} value between two particular nodes/isolates. Isolates that cluster together (i.e., connected by edges) are considered to be genetically identical (i.e., clones) at the threshold of $P_{AS} \geq 0.90$ in (a) but can differ by 1 allele at the threshold of $P_{AS} \geq 0.80$ in (b). Every color represents a different varcode and are colored based on the definition of varcodes by var for comparative purposes. It is important to note that 2 out of the 7 microsatellite markers genotyped were fixed in the population⁸.

Figure S7. Relatedness of isolates based on var and microsatellite genotyping. There was a positive correlation between (a) P_{TS} and P_{AS} relatedness measures (correlation coefficient = 0.757, $p < 0.001$) as well as (b) unbiased Bayesian P_{TS} and P_{AS} relatedness measures (correlation coefficient = 0.779, $p < 0.001$).

Figure S8. Genetic relatedness networks based on unbiased Bayesian pairwise type sharing estimates to identify recombinants. A network visualization of the genetic relatedness of *P. falciparum* isolates at the threshold of (a) $BP_{TS} \geq 0.50$ to define clusters of genetically-related isolates based on the posterior mean BP_{TS} estimates. Every node represents a *P. falciparum* isolate and an edge represents the BP_{TS} value between two particular nodes/isolates. Isolates that cluster together (i.e., connected by edges) are considered to be genetically identical (i.e., clones). Every color represents a different varcode. The (b) number and (c) proportion of pairwise comparisons within a given 95% highest density posterior interval width, which provides a measure of the uncertainty of each pairwise estimate. For the full distribution of within-varcode BP_{TS} estimates see Fig. S4d-f.

Figure S9. Relatedness networks based on microsatellite genotyping. (a) A network visualization of the genetic relatedness of *P. falciparum* isolates at the threshold of $P_{AS} \geq 0.50$ colored by varcode for comparative purposes. Every node represents a *P. falciparum* isolate and an edge represents the P_{AS} value between two particular nodes/isolates. (b) A clustered heatmap showing the genetic profiles of each *P. falciparum* isolate with rows representing each isolate and columns representing each microsatellite allele. Black and white denote the presence and absence of each allele, respectively. Isolates that clustered together were more genetically similar (i.e., the same types were present).

Figure S10. Sampling depth of var DBLα types in South America. Accumulation curves depict the number of observed DBLα types plotted as a function of the number of DBLα sequences sampled.

Figure S11. South American varcodes. A network visualization of the genetic relatedness of *P. falciparum* isolates at the threshold of $P_{TS} \geq 0.90$ to define clusters of genetically-related isolates, or *varcodes* in South America. Every node represents a *P. falciparum* isolate and an edge represents the P_{TS} value between two particular nodes/isolates. Isolates that cluster together (i.e., connected by edges) are considered to have the same *varcode*. The colors depict the country of origin/sampling location of each *P. falciparum* isolate.

Figure S12. Frequency distribution of the 195 DBLα types identified in Ecuador. Each point corresponds to an individual DBLα type and is colored by its frequency (i.e., the number of Ecuadorian *P. falciparum* isolates it was identified in). The y-axis depicts the frequency at which each Ecuadorian DBLα type was identified in the 128 *P. falciparum* isolates from South America. Of the 56 “singleton” DBLα types (i.e., frequency of 1) in the Ecuadorian dataset, 32 (57.1%) were identified in other South American countries and 24 (42.9%) were not identified anywhere else in South America.

Figure S13. Identification of the outbreak varcode1 DBLα types. Each column represents one of the 47 DBLα types and black and white denote the presence and absence of each type, respectively, in (a) the 36 infections with varcode1, (b) in the 15 infections with recombinant varcodes and (c) in South American *P. falciparum* populations. (a) The rows represent case numbers and are not clustered such that from top to bottom they are representative of sampling date (see Table 1 for details on sampling dates and whether they correspond to during or post-outbreak), and the columns are ordered from left to right in decreasing frequency of each DBLα type. *Note:* EC2 and EC7 were under sampled as compared to the other *P. falciparum* isolates (11 and 19 DBLα types per isolate, respectively). (b-c) Clustered heatmaps show the identification of the 47 DBLα types from the outbreak varcode1 *P. falciparum* isolates in the other *P. falciparum* isolates from South America. A total of 31 (66%) of the outbreak types were identified in one or more countries. The highest number of types identified in another country was 28 (60%) in isolates collected in 2003-2004 from Peru, followed by 14 (30%) in isolates collected from 2002-2006 from Colombia, followed by 8 (17%) in isolates collected from 2003-2007 from Venezuela and only 2 (4%) in isolates collected from 2006-2008 from French Guiana. The isolate collection dates are reported in²⁸.

Supplementary Tables

Table S1. Summary statistics for total number of *var* DBL α types per isolate (i.e., repertoire size) in South American *P. falciparum* populations.

Location	min*	med	mean	max
Colombia	19.0	40.0	38.4	43.0
Ecuador	11.0	37.0	36.6	43.0
French Guiana	13.0	48.0	50.5	92.0
Peru	19.0	36.0	33.4	42.0
Venezuela	28.0	36.5	35.2	43.0

*Note: all isolates with <10 types were removed due to low sequencing quality (see Methods)

Supplementary Text 1

As described in the Methods, the degenerate primers we use amplify the DBL α domain of *var* genes. We performed an additional validation of our PCR (using these degenerate DBL α primers) and sequencing methodology to understand the margin of error of detection of all DBL α types in an isolate. DBL α types were translated into amino acid sequences and classified as upsA or non-upsA, using the classifyDBLalpha pipeline (Ruybal-Pesántez et al. 2017) to examine whether the expected genomic proportions of upsA/non-upsA were obtained in each isolate.

Given our study was conducted in South America, we used the Honduran laboratory reference strain HB3 as a benchmark for the expected number of total DBL α types (i.e. repertoire size), as well as the proportion of upsA/non-upsA. Whole genome sequencing of HB3 has identified 44 *var* genes, with 8 upsA DBL α types (defined by DBL α domain 1), 34 non-upsA DBL α types (DBL α domains 0 and 2) and 2 upsE (*var2csa* genes, defined by DBLpam domains and do not have DBL α domains) (Rask et al. 2010). We independently verified this using 37 technical replicates of *var* DBL α PCR amplification and illumina sequencing of our HB3 laboratory isolate. It is worth noting that since HB3 has two *var2csa* genes that do not have DBL α domains, these will not be amplified by our degenerate primers, so we expect that only 42 of the *var* genes of HB3 could be amplified.

HB3 technical replicates

From the data obtained from 37 HB3 technical replicates, we identified the expected repertoire sizes with a median of 39 DBL α types (range: 36-41). A median of 7 upsA DBL α types (range: 6-8) and a median of 33 non-upsA DBL α types (range: 30-34) were identified (Figure 1). The median genomic proportion of upsA DBL α types was 17.5% (range: 15.4-19.5%) and 82.5% (range: 80.5-84.6%) for non-upsA DBL α types (Figure 2). We found that of the 46 identified in the 37 technical replicates, 40 of them were consistently identified in the majority of replicate isolates (range 21 to 37 replicates). Of these 40 types, 7 were ups-A and 33 were non-upsA. All of these findings are in line with what is expected from whole genome sequencing data (Rask et al. 2010).

Figure 1: The distribution of the number of upsA and non-upsA types identified in each HB3 isolate repertoire.

Figure 2: The proportion of upsA and non-upsA types identified in each HB3 isolate repertoire.

Field isolate genomic proportions of upsA and non-upsA DBL α types

The 543 unique DBL α types identified in the 186 South American *P. falciparum* isolates (N = 58 Ecuadorian *P. falciparum* isolates from this study, N = 128 previously published *P. falciparum* isolates from Colombia, French Guiana, Peru and Venezuela) were translated into amino acid sequences and classified as upsA or non-upsA, using the classifyDBLalpha pipeline (Ruybal-Pesántez et al. 2017). There were 79 upsA and 464 non-upsA types.

Looking first at the 195 types (26 upsA and 169 non-upsA) identified in Ecuadorian isolates, we obtained a median genomic proportion of upsA of 10.8% (range: 5.1-18.2%) and 89.2% (range: 81.8-94.9%) of non-upsA types for all isolate *varcodes* (Figure 3).

Figure 3: The proportion of upsA and non-upsA types in each isolate repertoire. The case numbers in the x-axis can be used to identify the clinical information of the participant in Table 1.

To confirm the patterns we observed in Ecuadorian isolates, we also compared them to the other South American isolates. In the other South American isolate *varcodes* the genomic proportions of upsA/non-upsA were similar, with a median proportion of upsA of 9-14% and 86-91% for non-upsA types (Table 1).

With regards to the number of upsA identified in all the South American isolate *varcodes*, we identified a median of 4-5 upsA types and 31-42.5 non-upsA types (Table 2).

Table 1: The proportion of upsA types in South American isolates

country	min	median	mean	max
Colombia	0.00	0.09	0.09	0.14
Ecuador	0.05	0.11	0.11	0.18
French Guiana	0.04	0.11	0.11	0.18
Peru	0.08	0.14	0.14	0.20
Venezuela	0.05	0.11	0.12	0.21

Table 2: The number of upsA types identified in South American isolates

country	min	median	mean	max
Colombia	0	4	4	6
Ecuador	2	4	4	6
French Guiana	2	5	6	14
Peru	2	5	5	7
Venezuela	2	4	4	9

Figure 4: The proportion of upsA and non-upsA types in each isolate repertoire stratified by country.

Inheritance of DBLa types in the recombinant *var*codes

For the outbreak *varcode*1, its 47 DBLa types were classified as 5 upsA and 42 non-upsA. In Figure 5 we look specifically at the “inheritance” of upsA (blues) vs non-upsA (greens) types from the parental outbreak *varcode*1 in the case of the parasites with recombinant *var*codes (*varcodes*3,4,6,7). This provides a proxy to examine inheritance of types with regards to their chromosomal location. The proportion of the outbreak types that were inherited in the recombinant parasite *var*codes is indicated in the darker shades of blue or green, showing that the proportion of inherited types varied both by *varcode* and upsA/non-upsA. Overall, the DBLa type sharing patterns in parasites with recombinant *var*codes are consistent with inheritance of 50% types, with a higher proportion of non-upsA inherited types (~40-70%, i.e. 17 to 30 of the 42 types) vs upsA (~20-50%, i.e. 1 to 3 of the 5 upsA types). The exception was *varcode*7 where 60-80% of upsA were inherited, i.e. 3-4 of the 5 upsA types. The lighter shades correspond to those types that were not inherited from the outbreak clone but from the other parent.

Figure 5: The proportion of the outbreak types that were inherited in the recombinant parasite *var*codes is indicated in the darker shades of blue or green. The proportion of the outbreak types that were inherited in the recombinant parasite *var*codes is indicated in the darker shades of blue or green and the lighter shades correspond to those types that were not inherited from the outbreak clone but from the other parent.

References

Rask, Thomas S., Daniel A. Hansen, Thor G. Theander, Anders Gorm Pedersen, and Thomas Lavstsen. 2010. “Plasmodium Falciparum Erythrocyte Membrane Protein 1 Diversity in Seven Genomes Divide and Conquer.” Edited by Jonathan A. Eisen. *PLoS Computational Biology* 6 (9): e1000933. <https://doi.org/10.1371/journal.pcbi.1000933>.

- 58 Ruybal-Pesántez, Shazia, Kathryn E. Tiedje, Gerry Tonkin-Hill, Thomas S. Rask, Moses R. Kamya, Bryan
59 Greenhouse, Grant Dorsey, Michael F. Duffy, and Karen P. Day. 2017. “Population Genomics of
60 Virulence Genes of Plasmodium Falciparum in Clinical Isolates from Uganda.” *Scientific Reports* 7 (1).
61 <https://doi.org/10.1038/s41598-017-11814-9>.