

Supplement 1: Cohort Selection Diagram



Supplement 2 : ICD 10 Codes and Text descriptors Used for Study Outcomes

Condition	ICD-10 Diagnosis Code	ICD-10 Text Descriptors			
Anxiety					
Symptom Level	R45.2	Unhappiness (worries NOS)			
	R45.82	Worries			
	R45.89	Feeling worried, anxious appearance, feeling anxious			
Disorder Level	F06.4,F10.180,F10.280,F12.9 80,F13.180,F13.280,F13.980, F14.180,F14.280,F14.980,F15 .180,F15.280,F15.980,F16.18 0,F16.280,F16.980,F18.180,F 18.280,F18.980,F19.180,F19. 280,F19.980,F31.3,F40.0,F40. 00,F40.01,F40.02,F40.10,F40. 11,F40.210,F40.218,F40.220, F40.228,F40.23,F40.230,F40. 231,F40.233,F40.240,F40.241 ,F40.242,F40.243,F40.248,F4 0.291,F40.298,F40.8,F40.9,F4 1.0,F41.1,F41.3,F41.8,F41.9,F 93.0,F94.0	All			
Depression					
Symptom Level	R45.7	State of emotional stress or shock			
	R45.81	Low self esteem			
	R45.84	Anhedonia			
	R45.851	Suicidal ideations			
	R45.86	Emotional lability			
	R45.89	Sad, sad mood, feeling sad, sadness, feeling of sadness, at risk for self harm, depressed affect, difficulty coping, dysphoric mood, feelings of worthlessness, feels depressed, flat affect, non-suicidal depressed mood, non-suicidal self harm, self-esteem disturbance, suicidal risk, tearfulness, thoughts of self harm			
Disorder Level	F06.30,F06.31,F06.32,F06.34, F10.14,F10.24,F10.94,F11.14, F11.24,F11.94,F13.14,F13.24, F13.94,F14.14,F14.24,F14.94, F15.14,F15.24,F15.94,F16.14, F16.24,F16.94,F18.14,F18.24, F18.94,F19.14,F19.24,F19.94, F32.0,F32.1,F32.2,F32.3,F32. 4,F32.5,F32.8,F32.81,F32.89, F32.9,F33.0,F33.1,F33.2,F33. 3,F33.40,F33.41,F33.42,F33.8 ,F33.9,F34.1,F34.81,F34.89,F 34.9	All			



Supplement 3: Medication Prevalence By Age

Ethnicity		Race			
Hispanic	Non- Hispanic	Asian	Black	White	Other
30/265	178/1409	17/241	7/72	160/1011	37/427
(11.3%)	(12.6%)	(7.1%)	(9.7%)	(15.8%)	(8.7%)
24/175	141/756	22/157	6/45	117/527	34/268
(13.7%)	(18.7%)	(14%)	(13.3%)	(22.2%)	(12.7%)
34/83	155/356	14/50	11/23	134/266	52/139
(41.0%)	(43.5%)	(28%)	(47.%)	(50.4%)	(37.4%)
	Hispanic 30/265 (11.3%) 24/175 (13.7%) 34/83	Hispanic Non- Hispanic 30/265 178/1409 (11.3%) (12.6%) 24/175 141/756 (13.7%) (18.7%) 34/83 155/356	Hispanic Non- Hispanic Asian 30/265 178/1409 17/241 (11.3%) (12.6%) (7.1%) 24/175 141/756 22/157 (13.7%) (18.7%) (14%) 34/83 155/356 14/50	Hispanic Non- Hispanic Asian Black 30/265 178/1409 17/241 7/72 (11.3%) (12.6%) (7.1%) (9.7%) 24/175 141/756 22/157 6/45 (13.7%) (18.7%) (14%) (13.3%) 34/83 155/356 14/50 11/23	Hispanic Non- Hispanic Asian Black White 30/265 178/1409 17/241 7/72 160/1011 (11.3%) (12.6%) (7.1%) (9.7%) (15.8%) 24/175 141/756 22/157 6/45 117/527 (13.7%) (18.7%) (14%) (13.3%) (22.2%) 34/83 155/356 14/50 11/23 134/266

Supplement 4: Medication Prevalence by Race and Ethnicity

Supplemental Figure 5: Medication Prevalence by Patient Factor for 6-18 year old patients

	Anxiety (N=2385)		Depressio	on (N=1390)	Anxiety+Depression (N=723)		
	Medicated	Nonmedicated	Medicated	Nonmedicated	Medicated	Nonmedicated	
Age							
6-12	125/1185	1060/1185	40/325	285/325	61/162	101/162	
	(10.54%)	(89.45%)	(12.30%)	(87.70%)	(37.65%)	(62.35%)	
3-18	232/1200	968/1200	245/1065	820/1065	282/561	279/561	
	(19.33%)	(80.67%)	(23.00%)	(77.00%)	(50.26%)	(49.74%)	
Sex							
Female	184/1352	1168/1352	166/833	667/833	218/462	244/462	
	(13.60%)	(86.40%)	(19.92%)	(80.08%)	(47.18%)	(52.82%)	
Male	173/1033	860/1033	119/557	438/557	125/261	136/261	
	(16.74%)	(83.26%)	(21.36%)	(78.64%)	(47.89%)	(52.11%)	
Insurance							
Private	280/1869	1589/1869	212/1004	792/1004	263/530	267/530	
	(14.98%)	(85.02%)	(21.12%)	(78.88%)	(49.62%)	(50.38%)	
Public	69/425	356/425	59/343	284/343	64/159	95/159	
	(16.23%)	(83.77%)	(17.21%)	(82.79%)	(40.25%)	(59.75%)	
Military	6/65	59/65	11/33	22/33	13/28	15/28	
,	(9.23%)	(90.77%)	(33.33%)	(66.67%)	(46.42%)	(53.57%)	
NA	2/26	24/26	3/10	7/10	3/6	3/6	
	(7.70%)	(92.30%)	(30.00%)	(70.00%)	(50.00%)	(50.00%)	
Number of Co- morbidities							
0	170/1570	1400/1570	160/924	764/924	152/366	214/366	
	(10.83%)	(89.17%)	(17.32%)	(82.68%)	(41.53%)	(58.47%)	
1	133/637	504/637	92/369	277/369	123/244	121/244	
	(20.88%)	(79.12%)	(24.94%)	(75.06%)	(50.40%)	(49.60%)	
2+	54/178	124/178	33/97	64/97	68/113	45/113	
	(30.34%)	(69.66%)	(34.03%)	(65.97%)	(60.18%)	(39.82%)	
Comorbid Diagnoses							
ADHD	116/452	336/452	69/220	151/220	93/170	77/170	
	(25.66%)	(74.34%)	(31.37%)	(68.63%)	(54.71%)	(45.29%)	
ASD	40/104	64/104	4/30	26/30	15/19	4/19	
	(38.46%)	(61.54%)	(13.33%)	(86.67%)	(78.94%)	(21.06%)	
Sleep	38/227	189/227	27/118	91/118	60/117	57/117	
	(16.74%)	(83.26%)	(22.88%)	(77.12%)	(51.28%)	(48.72%)	
Trauma	10/82	72/82	16/74	58/74	30/57	27/57	
and	(12.20%)	(87.8%)	(21.62%)	(78.38%)	(52.63%)	(47.37%)	
Stressor Related	,		, ,		, , , , , , , , , , , , , , , , , , ,		
Disorders							
Referral to	43/232	189/232	48/208	160/208	63/143	80/143	
DPB or Psychiatry	(18.54%)	(81.46%)	(23.08%)	(76.92%)	(44.05%)	(55.95%)	