Reference intervals of amino acid for newborns in North Asia: Diagnosing Inborn Errors of Metabolism Using Liquid Chromatography Tandem Mass Spectrometry (Amino acid reference in North Asia)

TA. Bairova, AV. Belskikh, AV. Atalyan, OV. Bugun,
NV. Nemchinova, LV. Rychkova

Scientific Centre for Family Health and Human Reproduction Problems

Location: 16 Timiryazev str., Irkutsk city, 664003 Irkutsk region

AUTORS

T.A. Bairova, Laboratory of Personalized Medicine, Scientific Centre for Family Health and Human Reproduction Problems, Irkutsk, Russia. Roles: Conceptualization, Supervision, Methodology, Formal analysis, Writing – original draft, Writing – review & editing, Data curation, Methodology. <http://orcid.org/0000-0003-3704-830X>

A.V. Belskikh, Laboratory of Personalized Medicine, Scientific Centre for Family Health and Human Reproduction Problems, Irkutsk, Russia. Roles: Methodology, Investigation, Data curation, Visualization, Writing – original draft. <http://orcid.org/0000-0003-3678-7274>

A.V. Atalyan, Functional Group of Information Systems and Biostatistics, Scientific Centre for Family Health and Human Reproduction Problems, Irkutsk, Russia. Roles: Methodology, Data analysis. <http://orcid.org/0000-0002-3407-9365>

O.V. Bugun, chief medical officer, Scientific Centre for Family Health and Human Reproduction Problems, Irkutsk, Russia. Roles: Project administration. <https://orcid.org/0000-0002-2162-3683>

N.V. Nemchinova, laboratory assistant, Scientific Centre for Family Health and Human Reproduction Problems, Irkutsk, Russia. Roles: Investigation, Data curation. <https://orcid.org/0000-0002-9720-8750>

25 L.V. Rychkova, director, Scientific Centre for Family Health and Human Reproduction Prob-
26 lems, Irkutsk, Russia. Roles: Project administration, Funding acquisition.
27 <https://orcid.org/0000-0002-0117-2563>

28 Corresponding author: Tatiana Bairova (tbairova38@mail.ru)

29 ABSTRACT

30 Aim. To establish reference intervals (RI) for blood amino acids (AA) in healthy newborns of
31 North Asia measured by liquid chromatography tandem mass spectrometry (LC-MS/MS) and
32 evaluate their differences from respective reference values for newborns from other popula-
33 tions.

34 Objectives. A cross-sectional study of 381 healthy newborns was conducted. De-identified
35 dried blood spots annotated by age, birth-weight, and sex were obtained from 381 healthy
36 newborns aged 0–7 days. Data was collected from April to May of 2020.

37 Methods. Dried blood spots collected from filtered paper were used to analyze and measure
38 of 13 derivatized amino acids using LC-MS/MS method. Nonparametric statistical approach-
39 es were used to generate 2.5th–97.5th percentile distributions for newborns in North Asia in
40 accordance with CLSI EP28-A3c.

41 Results. Reference intervals (RI) for phenylalanine, tyrosine, citrulline, alanine, ornithine,
42 proline in North Asian newborns differ slightly from those of newborns in other countries
43 around the world. This allows for the use of universal RI in the diagnosis of congenital meta-
44 bolic disorders of the indicated amino acids (AA) in different populations around the world.
45 The RI for branched-chain essential AA (valine, leucine, and isoleucine are metabolic criteria
46 for maple syrup disease) are higher in North Asian infants as compared to infants in other
47 populations. In addition, the RI for arginine, aspartic acid, and glutamic acid in North Asian
48 newborns are higher than in newborns in other countries around the world. In our study, the
49 RI for methionine in newborns were lower than in many countries worldwide [McHugh D,
50 2011]. For optimal clinical practice, RI for certain AA in newborns (valine, leucine, isoleu-

51 cine, methionine, arginine, aspartic acid, glutamic acid) should be determined for specific
52 populations- further augmenting the diagnosis of inborn errors of metabolism.

53 Keywords. Inborn errors of metabolism; newborn screening; tandem mass spectrometry.

54 INTRODUCTION

55 Inborn errors of metabolism (IEM) are ‘rare genetic disorders which invariably lead to meta-
56 bolic defects accompanied by an accumulation of toxic intermediate substances. The absence
57 of specific clinical manifestation(s) determines the complexity of diagnostics. IEM as a dis-
58 ease can begin at any age, but often manifests right after birth. The consequences are danger-
59 ous and without appropriate therapy and nutritional limitations is oftentimes fatal.

60 Currently, more than 2000 cases of IEM have been identified and described. [1] Factors that
61 influence the prevalence of the disorder(s) vary greatly amongst countries and social groups:
62 1:667 in Saudi Arabia [2], 1:784 in UK [3], 1:1200 in Canada [4], from 1:4100 [5] to 1:2900
63 [6] in Germany, from 1:1178 to 1:8304 [7] in China, 1:1944 in Egypt [8], 1:2396 in Portugal
64 [9], 1:2800 in South Korea [10], 1:2916 in Malaysia [11], 1:3165 in Singapore [12], 1:5882
65 for Taiwan [13], 1:9000 in Japan [14]. Key factors include variations in analytical techniques
66 [15], total number of IEMs analyzed, age differences at onset of disorder manifestation, and
67 ethnicity and race of studied population(s) [16, 17]. Reference AA concentration ranges for 0-
68 5 days of life were established in the R4S project, and included 45 countries [18]. This trial
69 included 47 US states, Australia, several countries of the European Union and Africa. Japan,
70 South Korea, Malaysia, and Taiwan represented the Asian territories involved in this project.
71 There were no representative regions from Northern Asia. The possibility of using reference
72 ranges, established for Europe, in the evaluation of AA levels in Northern Asia is not clear –
73 herein lies the rationale for conducting the current study.

74 The aim of this study was to establish reference intervals for AA levels by LC-MS/MS, which
75 have not been conducted in Northern Asia. Our secondary objective was to compare Refer-
76 ence Intervals (RI) with other parts of the world.

77 RI of 13 amino acids in newborns from North Asia were determined. The RI of the essential
78 AA (valine, leucine, isoleucine, methionine, arginine) and nonessential AA (aspartic acid,
79 glutamic acid) in newborns from North Asia differed from those in many nations around the
80 world [McHugh D, 2011].

81 2. MATERIALS AND METHODS

82 2.1. Study design. Cross-sectional descriptive study.

83 2.2. Study population. 400 dried de-identified blood spot (DBS) samples were collected in the
84 regional perinatal center. Data for 19 samples were excluded due to extended collection date
85 i.e., collection at 7 or further days of life, unusual bodyweight or unexpected concentrations.
86 A possible reason for unexpected AA concentration levels could be explained by contamina-
87 tion or IEM. 381 DBS samples from healthy children were included in the analysis. Boys
88 203(53.28 %), girls–178 (46.72 %).

89 In accordance with the World Health Organization's (WHO) criteria, «newborns» are categor-
90 ically defined from the moment of birth to the 28th day of life [19]. The period from birth to
91 the 7th day is considered «Early Neonatal». Newborn screening was conducted during this
92 period for the purposes of early IEM detection.

93 Inclusion criteria:

- 94 1. Healthy male and female newborns;
- 95 2. Gestational age, 37-42 weeks;
- 96 3. Birth weight 2500 – 4000 g.;
- 97 4. APGAR score greater than 7 at 10;
- 98 5. Age 4-6 days.
- 99 6. Breastfeeding to the moment of sample collection.

100 Exclusion criteria:

- 101 1. Gestation less than 36 weeks;
- 102 2. Inborn errors of metabolism;

- 103 3. Positive family history with metabolic disorders;
- 104 4. Lethargy;
- 105 5. Irritation;
- 106 6. Poor feeding;
- 107 7. Tachypnea;
- 108 8. Persistent vomiting;
- 109 9. Abnormal movement;
- 110 10. Spasms;
- 111 11. Malformations;
- 112 12. Diagnosed disorders of the liver, kidneys, heart, skeletal muscles;
- 113 13. Prescribed drug therapy;
- 114 14. Mother's diabetes mellitus disease.

115 The average age of infants that participated in our study was median (Me) (25th; 75th per-
116 centiles) 3 (3; 4) days. Mean birth body mass was M(25th;75th percentiles) 3408 (3095;3700)g,
117 for boys M(25th;75th percentiles)3480 (3150;3730) g, for girls- 3310 (3037,5;3637,5) g.

118 The study protocol was reviewed and approved by the local Ethic Committee of Scientific
119 Centre for Family Health and Human Reproduction Problems (number 4.2; date of registra-
120 tion 12th of May 2020). All procedures were conducted in accordance with the Declaration of
121 Helsinki. Prior to the collection of blood specimens, written consent was obtained from the
122 parents of the newborns. Each newborn was assigned a unique identification (ID) number.
123 Documented data collected from the mother and infants' medical records included date of
124 birth, sex, weight, height/length, and mother's health.

125 2.3. Description of the study area. All mothers lived in the Irkutsk Region during pregnancy.
126 The Irkutsk Region, located in the Asian part of Russia is 774 846 km² (4,52 % of Russia).
127 The territory's continental climate type is characterized as arctic and/or sub-arctic (Fig. 1).

128 Summer is cool. Winter is cold (down to -40 C). Average temperatures are below -10°C
129 throughout the year.

130 2. 39 million people live in the Irkutsk Region (actual for the 2020 year), with more than 30,
131 000 births registered annually. The population is multinational: Russians – 88%, Ukrainians –
132 1.2%, Tatars – 0.94%. Indigenous population consist of Buryats – 3.2%, evenks-0.05%,
133 Yakuts – 0.04% and Tofalars – 0.03% (2010).

134 Fig 1. World map indicating the region in which the research was performed

135 2.4. Sample Collection.

136 Samples were collected, April-May of 2020. Blood samples were taken on the 4th day after
137 birth in the regional perinatal center of Irkutsk Region, in accordance with the local newborn
138 screening program. Blood specimens were collected by heel-stick and spotted on filter paper
139 cards (Whatman 905, USA) a total of 5 spots, approximately 40 mkl of blood each. Filter pa-
140 per cards were dried (DBSs) for 4 hours at room temperature on a horizontal, non-humid and
141 nonabsorbent surface and then stored frozen at -20° C with a drying agent for 1-3 months.

142 2.5. Analysis of Amino Acids.

143 Concentrations of 13 AA were evaluated: acid aspartic (Asp), acid glutamic (Glu), alanine
144 (Ala), arginine (Arg), citrulline (Cyt), glycine (Gly), leucine (Leu) / isoleucine (Ile),
145 methionine (Met), ornithine (Orn), phenylalanine (Phe), proline (Pro), tyrosine (Tyr), valine
146 (Val). AA spectrum defined by reagent kit for AA level measurement "Amino Acid and
147 Acylcarnitines from dried blood" (Chromsystems, Munich/Germany).

148 4 mm dried blood spot disks were punched out of the filter card and placed on to a 96-well
149 microtiter plate. 200 mkl of the reconstituted internal standard was added. The plate was
150 evaporated at 60°C and 600 rpm until dry. 60 mkl of derivatization reagent (butan-1-ol, n-
151 butyl acetate, and hydrogen chloride) was added, the microtiter plate was sealed with a pro-
152 tective sheet, incubated for 15 min at 60°C and 600 rpm, and evaporated at 60°C and 600 rpm

153 until dry. 100 mkl reconstitution buffer was added and agitated for 1 min at 600 rpm. 10 mkl
154 was used for analysis with an optimized tandem mass spectrometer HPLC-MS/MS method.
155 The analysis was performed with Shimadzu LCMS-8060 triple quadrupole mass spectrometer
156 with Shimadzu LC-30AD HPLC system (Shimadzu, Kyoto, Japan). Optimization of the
157 method was performed in accordance with the certified manufacturer's specifications [20].
158 Optimal operating conditions included a 1.7-minute run time and mobile phase gradient with
159 a flow rate of 20–200 mkl/min

160 2.6. Statistical Analysis. Statistical analysis was performed using STATISTICA (data analysis
161 software system), version 10 (StatSoft, Inc.), and R – a free software environment for statisti-
162 cal computing and graphics (<https://www.r-project.org/>).

163 AA values, and characteristics of the newborns (gender, age, and weight) were described us-
164 ing descriptive statistics. Results were reported as median (Me) and 25, 75 percentiles for
165 quantitative variables and percentages for the categorical variables. The nonparametric
166 Mann–Whitney U test was used to analyze for differences in skewed distributed continuous
167 variables in two gender groups. Empirical distribution were tested by the Kolmogorov-
168 Smirnov test. Two-tailed $\alpha < 0.05$ were considered significant.

169 Moreover, 1, 2.5, 5, 10, 25, 75, 90, 97.5, 99, 99.5, 99.9 percentiles and the 95% confidence
170 intervals were calculated for 13 AA. Bootstrap resampling was used to estimate confidence
171 intervals (R packages: `boot()`, `boot.ci()`).

172 Calculation of sampling sufficiency was not done in accordance with requirements from
173 Clinical and Laboratory Standards Institute (CLSI CA28-3, 2018). Normally, for RI estima-
174 tion no less than 120 individuals are required.

175 3. RESULTS AND DISCUSSION

176 In this study, we calculated blood levels for 13 AA from DBS of healthy boys and girls in the
177 early neonatal period (from birth to the 7th day of life) within North Asia (Table 1).

178 Amino acids (AA) were divided into two groups: essential and non-essential. Essential
179 AA cannot be synthesized by a human organism and therefore must be obtained from food
180 sources. In our trial, essential AA included valine, leucine, isoleucine, methionine, and phe-
181 nylalanine. For newborns, arginine was also considered an essential AA.

182 Table 1. Comparison of the medians Me (25; 75 percentiles) of AA in the blood of newborn
183 boys and girls

№	AA	All (n=381)	Boys (n=203)	Gerls (n=178)	p
		Essential AA			
1	Leu/Ile	151.70 (113.93; 239.46)	157.92 (112.10;256.69)	148.75 (117.36;235.99)	0.905
2	Val	89.53 (66.84; 116.38)	89.95 (69.03;118.94)	89.06 (65.82;114.36)	0.322
3	Met	11.89 (9.23; 16.66)	11.73 (9.08;16.95)	12.06 (9.57;16.20)	0.805
4	Phe	43.65 (34.80; 54.29)	44.22 (35.25;54.83)	42.97 (34.02;53.00)	0.314
5	Arg	15.71(12.06; 22.41)	16.36 (11.79;23.35)	15.45 (12.08;22.18)	0.500
		No-essential AA			
6	Ala	223.79 (144.10; 333.69)	215.58 (130.37;333.69)	225.66 (162.62;330.77)	0.301
7	Asp	185.88 (120.33; 395.48)	184.35 (122.76;397.55)	186.51 (114.66;393.78)	0.574
8	Glu	409.23 (337.50; 483.73)	408.18 (320.99;485.87)	410.48 (345.39;479.54)	0.703
9	Tyr	62.75 (47.02; 86.18)	62.61(47.65;86.25)	62.80 (45.72;86.18)	0.728
10	Cyt	9.13 (6.73; 11.56)	9.38 (6.73;12.03)	9.03 (6.73;11.29)	0.628
11	Gly	346.00 (224.90; 412.30)	343.64 (232.02;412.29)	347.74 (221.66;411.24)	0.583
12	Orn	77.33(58.34; 104.86)	77.33 (58.34;107.63)	77.44 (57.38;103.33)	0.976
13	Pro	168.38 (127.09; 210.84)	165.13 (124.99;210.20)	171.99 (128.33;211.61)	0.566

184 When comparing AA levels, no significant differences were found between boys and girls.
185 Similar results, without significant differences between sexes for newborns, were shown by
186 Dietzen, D. J. (2016) within the US population [21], Jing Liu (2016) in the Chinese popula-
187 tion [22] and Jaraspong Uaariyapanichkul (2018) in the Thai language [23]. Gender influ-
188 ences AA levels in older children and adults as opposed to newborns [24, 25]. Men have
189 higher concentrations of essential AAs as compared to women [26].
190 A key factor in making a sound clinical diagnosis involves qualified interpretation of labora-
191 tory data, most notably the use of reference values for levels of AAs.. A few studies have ex-
192 amined AA reference intervals in newborns; however, in our study some tests were excluded
193 from consideration due to the use of parametric statistical analysis, since our data did not
194 show a normal distribution of values. Differences in statistical analysis were difficult to com-
195 pare neonatal RIs across studies. Some experts considered the 1st percentile as the bottom
196 range [18, 27, 30]. The upper bounds represented values from the 90 to 99.9 percentiles. This
197 is why we calculated different percentiles for each AA (Table 2).

198 Table 2. Amino acid percentiles in neonatal dried blood spots 381 newborns, analyzed by tan-
199 dem mass spectrometry (MS/MS)

AA marker	Percentiles (mmol/L) and 95% CI										
	1	2,5	5	10	25	75	90	97,5	99	99,5	99,9
Essential AA											
Leu/Ile	68.1 (64.8, 77.5)	77.3 (69.2, 84.8)	85.1 (79.34, 89.46)	91.6 (89.3, 96.9)	112.4 (108.1, 118.6)	227.8 (210.0, 261.0)	357.6 (330.7, 394.2)	495.0 (444.6, 577.9)	594.9 (502.5, 725.5)	654.1 (544.0, 923.5)	844.0 (601.0, 923.5)
Val	31.9 (25.7, 40.0)	40.6 (36.6, 42.7)	43.2 (40.9, 45.3)	50.1 (45.6, 53.66)	66.9 (64.8, 71.3)	116.3 (111.9, 123.8)	183.4 (160.6, 202.8)	257.0 (232.6, 305.9)	322.4 (265.7, 409.4)	365.9 (280.2, 536.1)	488.9 (314.4, 536.1)
Met	4.7 (3.4, 5.6)	5.6 (4.9, 6.3)	6.4 (5.7, 7.0)	7.24 (6.9, 7.6)	9.2 (8.9, 9.6)	16.3 (15.4, 17.1)	21.9 (20.1, 24.6)	30.0 (26.3, 34.0)	34.2 (30.2, 38.4)	36.3 (31.4, 39.0)	38.8 (34.7, 39.0)
Phe	19.1 (18.7, 22.6)	22.6 (19.5, 24.8)	25.1 (22.9, 26.7)	28.4 (26.1, 29.8)	35.2 (33.4, 36.9)	54.3 (51.6, 56.3)	63.9 (61.3, 67.0)	82.4 (72.4, 91.4)	92.6 (83.5, 121.2)	98.0 (84.6, 127.9)	124.4 (92.5, 127.9)
Arg	4.9 (4.1, 5.7)	5.7 (5.1, 7.1)	7.2 (6.0, 7.9)	8.7 (8.0, 10.0)	12.2 (11.0, 12.8)	22.5 (21.6, 24.0)	30.8 (27.4, 35.3)	46.8 (41.8, 53.6)	56.4 (48.5, 69.8)	66.0 (53.3, 76.2)	73. (59.7, 76.2)
No-essential AA											
Ala	55.4 (44.3, 61.1)	61.1 (58.4, 69.9)	71.9 (63.0, 83.0)	96.3 (76.7, 109.1)	143.2 (130.3, 158.6)	335.3 (310.9, 362.0)	416.3 (403.4, 433.5)	483.0 (455.3, 525.6)	540.3 (485.8, 1094.8)	647.5 (504.7, 1436.6)	1309.8 (548.0, 1437.0)
Asp	69.5 (65.1, 75.6)	75.5 (70.3, 79.6)	81.2 (76.3, 85.6)	90.7 (85.2, 95.7)	119.9 (110.6, 125.9)	383.8 (336.9, 431.8)	553.8 (512.1, 604.5)	755.8 (703.3, 880.8)	902.3 (772.8, 1012.8)	1007.5 (824, 1182.0)	1119.4 (918.0, 1182.1)
Glu	177.9 (150.7, 211.4)	211.5 (187.1, 230.7)	245.2 (215.9, 259.9)	278.6 (258.5, 292.2)	337.6 (316.3, 346.2)	485.6 (465.9, 505.9)	627.2 (565.9, 699.6)	900.0 (767.7, 1083.2)	1122.7 (932.0, 1460.1)	1249.13 (1009.1, 1549.2)	1516.14 (1157.0, 1549.0)
Tyr	27.4 (24.1, 31.3)	31.4 (27.8, 33.7)	34.5 (32.0, 36.5)	38.7 (35.9, 40.4)	47.6 (45.0, 49.3)	87.0 (80.1, 93.1)	120.3 (105.2, 134.5)	176.3 (161.6, 192.4)	196.7 (179.6, 329.7)	244.2 (185.7, 348.7)	338.8 (201.2, 348.7)
Cyt	3.9 (3.5, 4.4)	4.4 (4.0, 4.6)	4.6 (4.4, 5.0)	5.4 (5.0, 5.6)	6.7 (6.3, 7.2)	11.5 (11.0, 12.1)	14.2 (13.4, 15.2)	18.4 (16.9, 20.2)	20.4 (18.5, 27.4)	20.9 (19.2, 43.8)	35.7 (20.3, 43.9)
Gly	97.6 (90.3, 130.7)	132.5 (99.5, 145.7)	150.9 (136.8, 159.0)	176.8 (157.7, 186.3)	224.7 (212.8, 243.0)	411.9 (399.5, 438.1)	492.1 (479.4, 521.7)	593.9 (562.2, 663.6)	665.4 (599.2, 771.3)	697.5 (635.2, 807.6)	794.1 (665.5, 807.6)
Orn	32.2 (28.9, 37.0)	36.9 (32.5, 39.5)	40.3 (37.3, 43.7)	46.6 (43.3, 49.1)	58.5 (55.8, 61.1)	104.3 (96.6, 113.5)	136.4 (130.4, 149.6)	200.0 (173.3, 243.9)	249.9 (201.8, 276.6)	266.1 (210.9, 406.6)	358.3 (252.4, 406.6)
Pro	64.1 (51.3, 70.9)	70.7 (65.3, 85.3)	85.9 (73.6, 92.5)	97.2 (91.4, 103.0)	125.9 (116.5, 131.4)	208.4 (200.4, 220.7)	271.2 (248.0, 289.6)	356.6 (333.9, 429.9)	441.1 (361.0, 476.9)	449.6 (378.9, 520.1)	497.53 (433.8, 520.1)

200 At the next stage, we compared the RI of AA in newborns from North Asia with similar indi-
201 cators in newborns among other populations. All studies presented used DBS samples and
202 HPLC-MS / MS. Percentiles of the reference values from other studies were compared with
203 95% confidence intervals of the same percentiles calculated in our study sample of newborns.
204 The decision on the comparability of the reference values was based on whether the value of
205 the compared percentile was included in the calculated 95% confidence interval of our refer-
206 ences (Table 3).

207 Table 3. Comparison of the results of our studies of the AA content in the blood of newborns
208 (0-7 days, N = 381) by tandem MS / MS mass spectrometry with the data of other researchers

Country	45 countries [18]	USA [21]	Germany [20]	Austria [20]	Colombia [27]	Banglade sh [28]	Thailand [23]	Korea [29]	Tibet [30]
Age, days after birth	1-5	0-4	0-28	0-28	1-18	1-7	0-4	New-born	Newborn

N, a бс.	25-30 million	310	1500	2500	891	120	85	15000	15029
Cutt-off	1:99ile	2,5:97,5ile	99,5ile	99,5ile	1:99ile	97,5ile	2,5:97,5ile	99,9ile	1:99ile
AA marker	Essential AA								
Leu/Ile	-	31*-130*	270*	226*	-	-	-	236*	-
Val	57 [#] :212*	46 [#] :224	198*	191*	59,18 [#] :204*	188.83*	89,1 [#] :179,2*	211*	55,19 [#] :326,0 [#]
Met	11*:44 [#]	10 [#] :39 [#]	22*	45 [#]	11 [#] :101,91 [#]	42.38 [#]	10,7 [#] :17,7 [#]	60 [#]	2,96*:32,5
Phe	33 [#] :97	30 [#] :97 [#]	92	92	23,92 [#] :78,94*	85.45	50,3 [#] :70,1*	99	21,23:103,41 [#]
Arg	2,3*:32*	<1*-36*	35*	47*	4,4:52,07	21.39*	5,6:14,7*	40*	1,71*: 61,38 [#]
Non-essential AA									
Ala	117 [#] :507	104 [#] :394*	743	561	91,43 [#] :348,82*	-	424 [#] :633	694	132,25 [#] :905,0
Asp	-	-	270*	147*	-	-	102*:216*	-	-
Glu	158:551*	193-566*	710*	800*	-	-	485 [#] :687,4 [#]	850*	-
Tyr	34 [#] :207	34 [#] :151*	225	197	37,03:201,25	199.96*	74,1:114,4	270	31,78:268,92
Cyt	6,0 [#] :28,0*	5 [#] :23 [#]	33 [#]	34	7,81 [#] :27,52	17.17	13,8 [#] :23 [#]	40	5,86 [#] :97,65 [#]
Gly	24:117*	182 [#] :637	717	834 [#]	-	-	300 [#] :414*	1000 [#]	5,86*:97,65*
Orn	-	19*:105*	350	258	-	112.77*	124 [#] :185*	300	20,37*:540,01 [#]
Pro	-	94*:245*	-	441	-	-	97 [#] :150*	354*	70,63:468,79

209 * our RI are higher than RI in a comparable study

210 # our RI are lower than RI in comparable study

211 Results from a comparative analysis of the AA reference values in newborns indicate the var-

212 iability of the AA spectrum in newborns in different countries around the world. The RI anal-

213 ysis of essential AAs indicates that the RI of phenylalanine in newborns from North Asia is

214 comparable with newborns from other countries. Despite the statistical significance of some

215 of the differences, these differences are not large enough to indicate the use of different refer-

216 ence intervals for the diagnosis of phenylketonuria in clinical practice.

217 Differences in references for other essential amino acids were revealed. Our data indicate high

218 concentrations of branched-chain AA (valine, leucine, and isoleucine) in newborns from

219 North Asia as compared to other populations. An exception were the newborns from Tibet,

220 whose concentrations of branched-chain AA were higher than those of the newborns in our

221 study (Ch. Zhang, 2019) [30]. Disturbances in the metabolism of leucine, isoleucine and

222 valine have been reported in maple syrup urine disease (MSUD). This is a rare (orphan) dis-

223 ease. The prevalence is 1: 110,000 newborns in Sweden [31] to 1: 1,148,000 in Korea [32].

224 MSUD diagnostics includes a quantitative assessment of the levels of leucine, isoleucine and

225 valine in the blood with an assessment of the leucine / alanine ratio and genetic diagnostics

226 with the detection of mutations in the branched-chain AA dehydrogenase genes: *BCKDHA*,

227 *BCKDHB*, *DBT*, *DLD*. Moreover, the spectrum of mutations in these genes in Russian pa-

228 tients with MSUD differs from other populations [33].

229 The same pattern was found for arginine, aspartic acid, glutamic acid in which levels were
 230 higher than mentioned in other epidemiological trials [18, 20, 21, 23, 28, 29].

231 We found low levels of the essential amino acid methionine in North Asian babies as com-
 232 pared to babies in other populations around the world. It is well-known and widely accepted
 233 that essential AAs cannot be synthesized in humans, including newborns. Levels of AA are
 234 determined by placental circulation and breast milk [34]. In our trial, data specific to protein
 235 consumption thru breastfeeding was not collected. Yet, earlier trials show that high protein
 236 consumption can define AA metabolism for the first 4 months of life [35]. Respectively, AA
 237 blood levels in newborns directly correlate with AA metabolism in their respective mothers.

238 As aforementioned, essential AA is not synthesized by the human body but rather sourced
 239 through food-intake; therefore, AA levels are influenced by nutritional patterns and thereby
 240 depend on cultural traditions, the availability and accessibility to food, presence of specific
 241 genes, (which assist in the metabolism of certain food products), and technological achieve-
 242 ments.

243 North Asia is a territory of temperate to cold climate. Nutrition is a very important aspect in
 244 response to adaptation to the climate of certain geographical locations. Quantity and quality of
 245 meals should meet biological needs. Cold stress drives the formation of livestock economies,
 246 leading to high-caloric diets with a high content of animal-origin proteins and fats. Nutrition
 247 patterns define the metabolism of proteins and AA. Meat-eaters have 47% greater AA con-
 248 centrations as compared with vegans (a diet that consist of little to no animal products and/or
 249 by-products); with significant differences in lysine, methionine, tryptophan, alanine glycine,
 250 and tyrosine levels [36, 37]. Furthermore, it has been demonstrated that diets high in carbo-
 251 hydrates did not lead to significant changes in AA levels; however, diets high in fat lead to
 252 increased levels of branched-chain AA: valine, leucine, and isoleucine [38]. Yet AA levels for
 253 North Asian newborns were higher than in other populations and races, excluding Tibet. J.A.
 254 Schmidt et al discovered another pattern during their clinical trial of 65,000 UK volunteers.

255 They concluded that only a weak correlation existed between nutrition patterns and AA levels
256 for meat-eaters, vegetarians, vegans, and fish-eaters i.e., pescatarians. However, repeated AA
257 level evaluation shows differences in alanine and glycine levels - these levels were lower for
258 meat-eaters as compared with other groups.

259 Analysis of the RI of non-essential AA (tyrosine, citrulline, alanine, ornithine, proline) shows
260 that the RI in newborns from North Asia is comparable to newborns from other countries. De-
261 spite the statistical significance of some of the differences, these differences were not large
262 enough to indicate the use of different RI for diagnosis within clinical practice. Essential AA
263 level differences should be noted for Caucasian newborns from North Asia and newborns
264 from Thai, due to higher levels in the Caucasian group. In 2006, Tan IK, et.al. noted such dif-
265 ferences between Caucasian and Asian newborns [16]. At the same time, we found compara-
266 bility of RI for essential AA for Caucasian newborns from North Asia and newborns from
267 Tibet – there were no significant differences between groups in levels of valine, methionine,
268 phenylalanine, and arginine.

269 4. CONCLUSION.

270 Our work evaluated RI for 13 AA in newborns from North Asia, considering 2.5 and 97.5
271 percentiles as lower and upper limits of reference intervals. RI for AA are variable in different
272 populations. The RI for phenylalanine, tyrosine, citrulline, alanine, ornithine, proline in North
273 Asian newborns differ slightly from the RI for newborns in other countries around the world.
274 This allows for the use of universal RI in the diagnosis of congenital metabolic disorders of
275 the indicated AA in different populations around the world. The RI for branched-chain essen-
276 tial AA (valine, leucine, and isoleucine are metabolic criteria for maple syrup urine disease)
277 are higher in North Asian infants than in infants in other populations around the world. In ad-
278 dition, the RI for arginine, aspartic acid, glutamic acid in North Asian newborns were higher
279 than in newborns in other countries around the world. The RI for methionine in newborns
280 within our study were lower than in many countries around the world. For optimal clinical

281 practice, RI for certain AA in newborns (valine, leucine, isoleucine, methionine, arginine, as-
282 partic acid, glutamic acid) should be determined for specific populations. This will help diag-
283 nose inborn errors of metabolism.

284 In an effort to increase the effectiveness of newborn screening for IEM expedient a unified
285 approach for RI estimation, in accordance with «Defining, Establishing, and Verifying Refer-
286 ence Intervals in the Clinical Laboratory, Approved Guideline – Third Edition» (2010), where
287 RI are considered at intervals between 2.5th and 97.5th percentiles.

288 A differential approach for AA RI is necessary. It allows for the optimization of analysis in
289 IEM diagnosis, formation of a solution attraction algorithm, and harmonizes approaches to the
290 formulation of regional IEM lists for each world population, with specific consideration given
291 to race and ethnic differences.

292 Acknowledgement

293 The authors gratefully acknowledge the staff of the Regional Perinatal Center (head MD,
294 Ph.D. NV. Protopopova). Department of Pathology for help in collecting material.

295 Conflict of Interest

296 The authors declare that the research was conducted in the absence of any commercial or fi-
297 nancial relationships that could be construed as a potential conflict of interest.

298 REFERENCE

- 299 1. Arnold GL. Inborn errors of metabolism in the 21st century: past to present. *Ann*
300 *Transl Med.* 2018;6(24):467. pmid: 30740398. doi: 10.21037/atm.2018.11.36.
- 301 2. Moammar H, Cheriyan G, Mathew R, Al-Sannaa N. Incidence and patterns of inborn
302 errors of metabolism in the Eastern Province of Saudi Arabia, 1983-2008. *Ann Saudi*
303 *Med.* 2010;30(4):271-7. pmid: 20622343. doi: 10.4103/0256-4947.65254.
- 304 3. Sanderson S, Green A, Preece MA, Burton H. The incidence of inherited metabolic
305 disorders in the West Midlands, UK. *Arch Dis Child.* 2006;91(11):896-9. pmid:
306 16690699. doi: 10.1136/adc.2005.091637.

- 307 4. Applegarth DA, Toone JR, Lowry RB. Incidence of inborn errors of metabolism in
308 British Columbia, 1969-1996. *Pediatrics*. 2000;105(1):e10. pmid: 10617747. doi:
309 10.1542/peds.105.1.e10.
- 310 5. Schulze A, Lindner M, Kohlmüller D, Olgemöller K, Mayatepek E, Hoffmann GF.
311 Expanded newborn screening for inborn errors of metabolism by electrospray ioniza-
312 tion-tandem mass spectrometry: results, outcome, and implications. *Pediatrics*.
313 2003;111(6):1399-406. pmid: 12777559. doi: 10.1542/peds.111.6.1399.
- 314 6. Lindner M, Gramer G, Haegi G, Fang-Hoffmann J, Schwab KO, Tacke U, et al. Effi-
315 cacy and outcome of expanded newborn screening for metabolic diseases--report of 10
316 years from South-West Germany. *Orphanet J Rare Dis*. 2011;6:44. pmid: 21689452.
317 doi: 10.1186/1750-1172-6-44.
- 318 7. Lu H, Zheng J, Yao Y, Yang M, Yao S, Shi N, et al. A retrospective analysis of tan-
319 dem mass spectrometry screening for neonatal genetic metabolic diseases in Yancheng
320 area. *J. Birth Health Hered*. 2016;24:83-85. doi: 10.13404/j.cnki.cjbhh.2016.03.036
- 321 8. Hassan FA, El-Mougy F, Sharaf SA, Mandour I, Morgan MF, Selim LA, et al. Inborn
322 errors of metabolism detectable by tandem mass spectrometry in Egypt: The first
323 newborn screening pilot study. *J Med Screen*. 2016;23(3):124-9. pmid: 26790708. doi:
324 10.1177/0969141315618229.
- 325 9. Vilarinho L, Rocha H, Sousa C, Marcão A, Fonseca H, Bogas M, et al. Four years of
326 expanded newborn screening in Portugal with tandem mass spectrometry. *J Inherit*
327 *Metab Dis*. 2010;33:133-8. pmid: 20177789. doi: 10.1007/s10545-010-9048-z.
- 328 10. Yoon HR, Lee KR, Kang S, Lee DH, Yoo HW, Min WK, et al. Screening of new-
329 borns and high-risk group of children for inborn metabolic disorders using tandem
330 mass spectrometry in South Korea: a three-year report. *Clin Chim Acta*. 2005;354(1-
331 2):167-80. pmid: 15748614. doi: 10.1016/j.cccn.2004.11.032.

- 332 11. Yunus ZM, Rahman SA, Choy YS, Keng WT, Ngu LH. Pilot study of newborn
333 screening of inborn error of metabolism using tandem mass spectrometry in Malaysia:
334 outcome and challenges. *J Pediatr Endocrinol Metab.* 2016;29(9):1031-9. pmid:
335 27544719. doi: 10.1515/jpem-2016-0028.
- 336 12. Lim JS, Tan ES, John CM, Poh S, Yeo SJ, Ang JS, et al. Inborn Error of Metabolism
337 (IEM) screening in Singapore by electrospray ionization-tandem mass spectrometry
338 (ESI/MS/MS): An 8 year journey from pilot to current program. *Mol Genet Metab.*
339 2014;113(1-2):53-61. pmid: 25102806. doi: 10.1016/j.ymgme.2014.07.018.
- 340 13. Niu DM, Chien YH, Chiang CC, Ho HC, Hwu WL, Kao SM, et al. Nationwide survey
341 of extended newborn screening by tandem mass spectrometry in Taiwan. *J Inherit*
342 *Metab Dis.* 2010;33:295-305. pmid: 20567911. doi: 10.1007/s10545-010-9129-z.
- 343 14. Yamaguchi S. Newborn screening in Japan: restructuring for the new era. *Ann Acad*
344 *Med Singap.* 2008;37(12):13-5. pmid: 19904475.
- 345 15. Hoffman GL. *Newborn Screening by Tandem Mass Spectrometry: Approved Guide-*
346 *line.* Clinical and Laboratory Standards Institute. 2010.
- 347 16. Tan IK, Gajra B. Plasma and urine amino acid profiles in a healthy adult population of
348 Singapore. *Ann Acad Med Singap.* 2006;35(7):468-75. pmid: 16902722.
- 349 17. Jeanmonod R, Asuka E, Jeanmonod D. Inborn Errors Of Metabolism. 2020 Aug 23
350 [cited 13 October 2020]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls
351 Publishing; 2020 Jan-. Available from:
352 <https://www.ncbi.nlm.nih.gov/books/NBK459183/>
- 353 18. McHugh D, Cameron CA, Abdenur JE, Abdulrahman M, Adair O, Al Nuaimi SA, et
354 al. Clinical validation of cutoff target ranges in newborn screening of metabolic disor-
355 ders by tandem mass spectrometry: a worldwide collaborative project. *Genet Med.*
356 2011;13(3):230-54. pmid: 21325949. doi: 10.1097/GIM.0b013e31820d5e67.

- 357 19. Child Development: Positive Parenting Tips: Infants (0-1 Year of Age). 2018 May 23
358 [cited 14 November 2020]. In: *Centers for Disease Control and Prevention (CDC)*
359 [Internet]. U.S. Department of Health and Human Services. Available from:
360 <https://www.cdc.gov/ncbddd/childdevelopment/positiveparenting/infants.html>
- 361 20. Chromsystems Instruments & Chemicals GmbH. Instruction Manual for LC-MS/MS
362 Analysis. MassChrom® Amino Acids and Acylcarnitines from Dried Blood, order No
363 55000. 2020.
- 364 21. Dietzen DJ, Bennett MJ, Lo SF, Grey VL, Jones PM. Dried Blood Spot Reference In-
365 tervals for Steroids and Amino Acids in a Neonatal Cohort of the National Children's
366 Study. *Clin Chem*. 2016;62(12):1658-1667. pmid: 27784706. doi:
367 10.1373/clinchem.2016.263434.
- 368 22. Liu J, Chen XX, Li XW, Fu W, Zhang WQ. Metabolomic Research on Newborn In-
369 fants With Intrauterine Growth Restriction. *Medicine (Baltimore)*. 2016;95(17):e3564.
370 pmid: 27124067. doi: 10.1097/MD.0000000000003564.
- 371 23. Uaariyapanichkul J, Chomtho S, Suphapeetiporn K, Shotelersuk V, Punnahitananda S,
372 Chinjarernpan P, et al. Age-Related Reference Intervals for Blood Amino Acids in
373 Thai Pediatric Population Measured by Liquid Chromatography Tandem Mass Spec-
374 trometry. *J Nutr Metab*. 2018;2018:5124035. pmid: 29854440. doi:
375 10.1155/2018/5124035.
- 376 24. Yi P, Liu L, Mei H, Zeng F, Huang Z, Niu H. Establishment of reference range of
377 plasma amino acids for younger Chinese children by reverse phase HPLC. *J Pediatr*
378 *Endocrinol Metab*. 2011;24(9-10):733-8. pmid: 22145465. doi:
379 10.1515/jpem.2011.270.
- 380 25. Sharer JD, De Biase I, Matern D, Young S, Bennett MJ, Tolun AA. Laboratory analy-
381 sis of amino acids, 2018 revision: a technical standard of the American College of

382 Medical Genetics and Genomics (ACMG). *Genet Med.* 2018;20(12):1499-1507. pmid:
383 30459394. doi: 10.1038/s41436-018-0328-6.

384 26. Yamamoto H, Kondo K, Tanaka T, Muramatsu T, Yoshida H, Imaizumi A, et al. Ref-
385 erence intervals for plasma-free amino acid in a Japanese population. *Ann Clin*
386 *Biochem.* 2016;53(3):357-64. pmid: 25829462. doi: 10.1177/0004563215583360.

387 27. Céspedes N, Valencia A, Echeverry CA, Arce-Plata MI, Colón C, Castiñeiras DE, et
388 al. Reference values of amino acids, acylcarnitines and succinylacetone by tandem
389 mass spectrometry for use in newborn screening in southwest Colombia. *Colomb Med*
390 *(Cali).* 2017;48(3):113-119. pmid: 29213153. doi: 10.25100/cm.v48i3.2180.

391 28. Sarker SK, Islam MT, Biswas A, Bhuyan GS, Sultana R, Sultana N, et al. Age-
392 Specific Cut-off Values of Amino Acids and Acylcarnitines for Diagnosis of Inborn
393 Errors of Metabolism Using Liquid Chromatography Tandem Mass Spectrometry. *Bi-*
394 *omed Res Int.* 2019;2019:3460902. pmid: 30723736. doi: 10.1155/2019/3460902.

395 29. Cho SE, Park EJ, Seo DH, Lee IB, Lee HJ, Cho DY, et al. Neonatal Screening Tests
396 for Inherited Metabolic Disorders using Tandem Mass Spectrometry: Experience of a
397 Clinical Laboratory in Korea. *Laboratory Medicine Online.* 2015;5(4):196-203.
398 doi:10.3343/lmo.2015.5.4.196.

399 30. Zhang Ch, Dha D, Cheng Y, Ma Y, Meng Y, Tse D, et al. An investigation of amino
400 acid and acylcarnitine levels in neonates from the Tibet Autonomous. Research
401 Square: PPR132791 [Preprint]. 2019 [cited 2020 November 17]. Available from:
402 <https://www.researchsquare.com/article/rs-9158/v1.pdf>

403 31. Sörensen L, von Döbeln U, Åhlman H, Ohlsson A, Engvall M, Naess K, et al. Ex-
404 panded Screening of One Million Swedish Babies with R4S and CLIR for Post-
405 Analytical Evaluation of Data. *Int J Neonatal Screen.* 2020;6(2):42. pmid: 33073033.
406 doi: 10.3390/ijns6020042.

407 32. Shibata N, Hasegawa Y, Yamada K, Kobayashi H, Purevsuren J, Yang Y, et al. Diver-
408 sity in the incidence and spectrum of organic acidemias, fatty acid oxidation disorders,
409 and amino acid disorders in Asian countries: selective screening vs. expanded new-
410 born screening. *Mol Genet Metab Rep*. 2018;16:5-10. pmid: 29946514. doi:
411 10.1016/j.ymgmr.2018.05.003.

412 33. Kurkina MV., Baydakova GV., Zakharova EYu. High frequency of homozygous mu-
413 tations in genes *BCKDHA*, *BCKDHB* in diseases with the smell of urine maple syrup
414 in Russian patients. *Russian Bulletin of perinatology and pediatrics*. 2016;(4):198.

415 34. Vaughan OR, Rosario FJ, Powell TL, Jansson T. Regulation of Placental Amino Acid
416 Transport and Fetal Growth. *Prog Mol Biol Transl Sci*. 2017;145:217-251. pmid:
417 28110752. doi: 10.1016/bs.pmbts.2016.12.008.

418 35. Haschke-Becher E, Kainz A, Bachmann C. Reference values of amino acids and of
419 common clinical chemistry in plasma of healthy infants aged 1 and 4 months. *J Inherit*
420 *Metab Dis*. 2016;39(1):25-37. pmid: 26227325. doi: 10.1007/s10545-015-9870-4.

421 36. Schmidt JA, Rinaldi S, Scalbert A, Ferrari P, Achaintre D, Gunter MJ, et al. Plasma
422 concentrations and intakes of amino acids in male meat-eaters, fish-eaters, vegetarians
423 and vegans: a cross-sectional analysis in the EPIC-Oxford cohort. *Eur J Clin Nutr*.
424 2016;70(3):306-12. pmid: 26395436. doi: 10.1038/ejcn.2015.144.

425 37. Bairova TA, Dolgikh VV, Kolesnikova LI, Pervushina OA. Nutritciogenetics and risk
426 factors of cardiovascular disease: associated research in eastern Siberia populations.
427 *Acta Biomedica Scientifica*. 2013; (4):92.

428 38. Swendseid ME, Yamada C, Vinyard E, Figueroa WG, Drenick EJ. Plasma amino acid
429 levels in subjects fed isonitrogenous diets containing different proportions of fat and
430 carbohydrate. *Am J Clin Nutr*. 1967;20(1):52-5. pmid: 6017009. doi:
431 10.1093/ajcn/20.1.52.

Therritory of Irkutsk Region