

Supplement 6: The sensitivity analysis results.

EXCLUDED TRIALS	Overall effect	General Approach	Efficacious intervention characteristics
<p>Trials with the education comparator (Boyne et al. 2014; Meng et al. 2016; Freedland et al. 2015; Brodie and Inoue 2005) (Ajiboye et al. 2015)</p>	<p>SMD = 0.31, 95%CI [0.21; 0.40] ***</p>	<p>Exercise: SMD = 0.34 , 95%CI [0.18; 0.51]; Remote communication and treatment: SMD = 0.42, 95%CI [0.24; 0.60]; Exercise and behaviour change: SMD = 0.21, 95%CI [0.004; 0.41]; Disease Management: SMD = 0.21, 95%CI [-0.03; 0.45]; Self-Management: SMD = 0.23, 95%CI [-0.17; 0.62].</p>	<p>No change in the significance of the effects associated with individual intervention or participant characteristics</p>

EXCLUDED TRIALS	Overall effect	General Approach	Efficacious intervention characteristics
<p>HF-ACTION trial (O'Connor et al. 2009)</p>	<p>SMD = 0.37, 95% CI: [0.10; 0.63]*</p>	<p>Exercise: SMD = 0.56, 95% CI [0.18; 0.94]; Remote communication and treatment: SMD =0.41, 95% CI [-0.29; 1.11]; MI: SMD = 0.03, 95% CI [-0.89; 0.94]; Exercise and behaviour change: SMD = 0.21, 95% CI [-0.59; 1.01]; CBT: SMD = 0.21, 95% CI [-0.91; 1.33]; Disease Management: SMD =0.23, 95% CI [-0.95; 1.41].</p>	<p>Behavioural Practice and Rehearsal * Graded task * Group-based * Centre-based *</p>

<p>High Risk of Bias trials (Ajiboye et al. 2015; Tomita et al. 2008)</p>	<p>SMD = 0.4578 95%CI [0.0903 0.8252 * Exercise: SMD =0.36, 95%CI [-0.2; 0.91]; Remote communication and treatment: SMD = 0.56, 95%CI [-0.57; 1.7]; MI: SMD = 0.03, 95%CI [-1.16; 1.21]; Exercise and behaviour change: SMD = 1.27, 95%CI [0.47; 2.07]***; CBT: SMD = 0.21, 95%CI [-1.38; 1.8]; Disease Management: SMD =0.23, 95%CI [-1.39 1.85].</p>	<p>Social Support (emotional) * Social Support (practical) ** TCS score * Monitoring of behaviour by others w/o feedback* Credible source*** Adding objects to the environment *** Self-monitoring of outcome(s) of behaviour* Self-monitoring of behaviour * Information about health consequences * Information on how to perform behaviour * Graded tasks * Action Planning * Goal setting (behaviour) * Behavioural Practice and Rehearsal**</p>
---	---	--

EXCLUDED TRIALS	Overall effect	General Approach	Efficacious intervention characteristics
HF-ACTION trial (O'Connor et al. 2009) and High Risk of Bias trials	SMD = 0.2670 95%CI [0.1549 0.3790 ***	Exercise: SMD=0.35, 95% CI [0.15; 0.54]*; Remote communication and treatment: SMD=0.47, 95%CI [0.24;0.70]*; MI: SMD=0.03, 95%CI [-0.46;0.52]; Exercise and behaviour change: SMD= 0.21; 95%CI:[-0.07; 0.49]; CBT SMD= 0.21, 95%CI:[-0.15; 0.57] Disease Management SMD= 0.23, 95%CI:[-0.25; 0.71].	Nurse *** Telehealth** Biological Feedback* Credible source * Self-monitoring of outcomes of behaviour* Self-monitoring of behaviour ** Demonstration of the behaviour ** Information about health consequences ** Graded tasks ** Action Planning ** Goal setting (behaviour)*** Behavioural Practice and Rehearsal***

Note: *p<0.05; **p<0.01, *** p<0.001. The changes in the findings, identified by the sensitivity analysis (corresponding exclusion of the trials) are in bold.

