Choosing the right tool for the job: A comprehensive assessment of serological assays for SARS-CoV-2 as surrogates for authentic virus neutralization

Nicholas Wohlgemuth¹, Kendall Whitt², Sean Cherry¹, Ericka Kirkpatrick Roubidoux¹, Chun-Yang Lin³, Kim J. Allison¹, Ashleigh Gowen¹, Pamela Freiden¹, E. Kaitlynn Allen³, St. Jude Investigative Team, Aditya H. Gaur¹, Jeremie H. Estep^{4,5}, Li Tang⁶, Tomi Mori⁶, Diego R. Hijano¹, Hana Hakim¹, Maureen A. McGargill³, Florian Krammer⁷, Michael A. Whitt², Joshua Wolf¹, Paul G. Thomas³, Stacey Schultz-Cherry^{1*}

St. Jude Investigative Team

David C. Brice³, Ashley Castellaw³, David E. Whittman⁸, Jason Hodges⁴, Ronald H. Dallas¹, Valerie Cortez¹, Ana Vazquez-Pagan¹, Richard J. Webby¹, Resha Bajracharya³, Brandi L Clark³, Lee-Ann Van de Velde³, Walid Awad³, Taylor L Wilson³, Allison M. Kirk³, Elaine I. Tuomanen¹, Randall T. Hayden⁹, James Hoffman¹⁰, Jamie Russell-Bell¹, James Sparks⁵

¹Department of Infectious Diseases, St Jude Children's Research Hospital, Memphis, TN, USA

²Department of Microbiology, Immunology and Biochemistry, University of Tennessee Health Science Center, Memphis, TN, USA

³Department of Immunology, St Jude Children's Research Hospital, Memphis, TN, USA

⁴Department of Hematology, St Jude Children's Research Hospital, Memphis, TN, USA

⁵Department of Global Pediatric Medicine, St Jude Children's Research Hospital, Memphis, TN, USA

⁶Department of Biostatistics, St Jude Children's Research Hospital, Memphis, TN, USA

⁷Department of Microbiology, Icahn School of Medicine at Mount Sinai, New York, NY, USA

⁸Office of Quality and Patient Care St Jude Children's Research Hospital, Memphis, TN, USA

⁹Department of Pathology, St Jude Children's Research Hospital, Memphis, TN, USA

27 ¹⁰Department of Pharmaceutical Sciences, St Jude Children's Research Hospital, Memphis, TN,

28 USA

29 *Corresponding author: Stacey-Schulz-Cherry stacey.schultz-cherry@stjude.org

Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) emerged in late 2019 and has since caused a global pandemic resulting in millions of cases and deaths¹⁻⁴. Diagnostic tools and serological assays are critical for controlling the outbreak⁵⁻⁷, especially assays designed to quantitate neutralizing antibody levels, considered the best correlate of protection⁸⁻¹¹. As vaccines become increasingly available¹², it is important to identify reliable methods for measuring neutralizing antibody responses that correlate with authentic virus neutralization but can be performed outside of biosafety level 3 (BSL3) laboratories. While many neutralizing assays using pseudotyped virus have been developed, there have been few studies comparing the different assays to each other as surrogates for authentic virus neutralization^{9,10,13,14}. Here we characterized three enzyme-linked immunosorbent assays (ELISAs) and three pseudotyped VSV virus neutralization assays and assessed their concordance with authentic virus neutralization. The most accurate assays for predicting authentic virus neutralization were luciferase and secreted embryonic alkaline phosphatase (SEAP) expressing pseudotyped virus neutralizations, followed by GFP expressing pseudotyped virus neutralization, and then the ELISAs.

Numerous serological assays have been developed to quantitate severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) antibody responses to determine total antibody concentration and neutralization activity. Neutralization assays can be performed with authentic virus or a pseudotyped virus expressing the SARS-CoV-2 spike (S) protein on its surface and a marker to measure infection of cells. The clear advantage of a pseudotyped virus is safety, as these studies can be performed in standard BSL2 laboratories. Another advantage is that results using pseudotyped virus can be obtained sooner, typically less than 24 hours whereas with authentic virus, plaque reduction-based neutralization assays take 2-3 days. A third advantage of using pseudotypes is flexibility. Pseudotypes expressing spike variants can be generated easily once the sequence is known since all that is needed is a plasmid that

expresses the variant of interest. One disadvantage of the pseudotyped virus neutralization assay is the pseudotyped viruses lack all but the spike protein from SARS-CoV-2, meaning they can only be neutralized by spike specific antibodies, and the clustering of proteins may not be representative of authentic virus particles. Yet, few studies have demonstrated whether the neutralization dose 50% (ND₅₀), the dilution at which 50% of virus will be neutralized, differs between pseudotyped virus detection platforms and, importantly, how they compare to authentic virus. To fill this gap in knowledge, we compared SARS-CoV-2 antibody titers in plasma from 24 PCR-positive individuals and 10 PCR negative individuals by enzyme-linked immunosorbent assay (ELISA), pseudotyped virus neutralization assay, and authentic virus neutralization. Adult participants were enrolled in the prospective, adaptive cohort study of St. Jude Children's Research Hospital employees "St. Jude Tracking of Viral and Host Factors Associated with COVID-19" (SJTRC, [clinicaltrials.gov #NCT04362995](https://clinicaltrials.gov/ct2/show/study/NCT04362995)) beginning in April of 2020. SJTRC was approved by the St. Jude Internal Review Board, and all participants provided written informed consent in a manner consistent with institutional policies. Cohort characteristics are provided in Supplemental Table 1. Samples were collected from PCR-positive individuals a median 33 days following diagnosis (interquartile range: 25.75-48.25) between April and August 2020.

The ELISAs included in the comparison detect antibodies to SARS-CoV-2 spike protein, nucleocapsid protein (N), or the receptor-binding domain (RBD) of the spike protein as described¹⁵. Briefly, plasma samples were diluted 1:50 for RBD and N ELISAs and results expressed as the ratio of the optical density (OD) from the sample over the negative control (a known negative, pre-pandemic plasma sample), which is common practice. To determine spike titers, plasma was diluted 1:100 to 1:8100 and an area under the curve (AUC) analysis performed. The pseudotyped virus platform was a vesicular stomatitis virus (VSV) glycoprotein (G) knockout VSV expressing full-length SARS-CoV-2 spike protein (VSV-ΔG-S) from the Wuhan-Hu-1 strain with three different reporter genes: green fluorescence protein (GFP), luciferase (Luci.) and secreted alkaline phosphatase (SEAP). Authentic virus neutralization

studies were performed under BSL3+ conditions with the 2019n-CoV/USA_WA1/2020 strain obtained from BEI Resources.

All PCR positive participants had ELISA titers to RBD, N, and spike, although the titers differed (Table 1). The average RBD ratio for the positive participants was 16.45 (95% CI: 14.53 – 18.36) and 1.62 (95% CI: 1.33 – 1.91) for negative participants while the average N ratios were 9.20 (95% CI: 7.69 – 11.11) for positive participants and 1.40 (95% CI: 0.79 – 2.01) for negative participants. Spike AUC average was 6.44 (95% CI: 5.04 – 7.85) for the positive samples (the spike ELISA was not performed on negative samples).

To quantitate neutralization titers, plasma was diluted 1:100 to 1:900 and tested by authentic virus and VSV-ΔG-S GFP, Luci., and SEAP pseudotyped viruses. AUC and ND₅₀ were calculated (Table 2). The average AUC values for the authentic virus neutralization assay were 51.14 (95% CI: 43.01 – 62.25) for positive participants and 10.96 (95% CI: 6.29 – 15.63) for negative participants. The GFP, Luci, and SEAP pseudotyped virus neutralization assays gave average AUC values of 71.07 (95% CI: 65.47 – 76.67), 54.20 (95% CI: 46.67 – 61.73), and 56.14 (95% CI: 48.26 – 64.01) respectively, for positive participants and 9.33 (95% CI: 3.62 – 15.04), 0 (95% CI: 0 – 0), and 1.21 (95% CI: 0 – 2.79) respectively for negative participants (Table 2). The geometric average ND₅₀ value for the authentic virus neutralization assay was 254.7 (95% CI: 92.97 – 697.6) for positive participants and 13.56 (95% CI: 5.08 – 36.14) for negative participants compared to 1305 (95% CI: 763.5 – 2232), 404.1 (95% CI: 225.1 – 725.5), and 474.3 (95% CI: 255.7–879.7), for positive participants and 12.12 (95% CI: 3.562 – 41.27), 1 (95% CI: 1 – 1), and 1.772 (95% CI: 0.7476 – 4.202) for negative participants respectively for the GFP, Luci and SEAP pseudotyped viruses. All neutralization platforms differentiated average negative and positive samples (Figure 1). While the AUC and ND₅₀ values were significantly higher for the GFP pseudotyped virus compared to the authentic virus or Luci. pseudotyped virus, suggesting that VSV-ΔG-S-GFP could be a more sensitive assay, it is balanced by increased AUC and ND₅₀ values in negative participants. Only the Luci. and SEAP

pseudotyped viruses showed no background in samples from PCR-negative participants. A Bland-Altman methods comparison test shows that there is systematic bias between the different pseudotyped virus neutralization assays and the authentic virus neutralization assay, leading to higher variability (highest for GFP pseudotypes) when the signal is low for each assay (Supplemental Figure 1). However, this bias decreases when signal becomes higher, resulting in the pseudotype assays becoming more concordant with authentic virus neutralization. The sigmoidal relationship between the amount of analyte detected and the readout in SEAP and luminescence assays could be a reason for the difference in bias between the pseudotyped virus assays. Variance at the lower end of the curve is less likely to be detected above background, compared to authentic virus neutralization and GFP pseudotyped virus neutralization where each infectious unit is counted, and variance has the same magnitude in both negative and positive samples. Furthermore, SEAP and luminescence detection kits often provide controls and stringent parameters for keeping background and noise to minimal levels.

To determine which neutralization platforms best correlated with authentic virus, linear regression analyses were performed. All pseudotyped virus neutralization platforms were significantly correlated to authentic virus neutralization regardless of the reporter with Luci. (Pearson's $r = 0.765$) and SEAP (Pearson's $r = 0.775$) having the highest correlations (Figure 2A). The pseudotyped virus neutralization assays were significantly correlated with each other with Pearson's r values as high as 0.971 between the Luci. and SEAP assays. Linear regression analyses demonstrated that ELISA titers to the RBD (Pearson's $r = 0.691$) and spike (Pearson's $r = 0.648$) are also significantly correlated with authentic virus neutralization titers (Figure 2B). Nucleocapsid ELISA was significantly correlated with authentic virus neutralization, but has the worst correlation with authentic virus neutralization (Pearson's $r = 0.514$), which has been shown previously for pseudotyped virus neutralization¹⁶. This is also congruent with the observation that antibodies targeting the RBD domain of spike are highly neutralizing¹⁷. A principle component analysis (PCA) was performed using all three ELISAs and all three

pseudotyped virus platforms as variables (Supplemental Figure 2). The resulting PCA plots shows distinct clustering of the samples with the highest authentic virus neutralization titers and a gradient from poorly neutralizing samples (in the bottom left) to highly neutralizing samples (in the top right). Finally, the average difference between the log(ND₅₀) for authentic virus neutralization and each pseudotyped virus neutralization is -0.487 for the GFP pseudotype, 0.191 for the Luci. pseudotype, and 0.069 for the SEAP pseudotype.

To assess granularity in the different ELISA results, cut-off values were used to categorize responses as high positive, low positive, or negative. Determination of cut-off values (RBD ratio: 15, nucleocapsid ratio:10, and spike: 6) was done by finding internal nadir present in histograms for the different ELISAs. The stratification of RBD ELISA responses into high and low groups did not result in significantly different responses in any of the neutralization assays (Figure 3A), suggesting that high RBD values do not necessarily correlate to higher neutralization titers, despite RBD ELISA positivity being associated with neutralization (Figure 2A). Similar results were obtained for the spike ELISA (Figure 3B) and nucleocapsid ELISA (Figure 3C). There was, however, a trend for increased neutralization in the high positive group versus the low positive group for each neutralization assay, regardless of ELISA assay, justifying future studies specifically designed to test the granularity of these assays. Congruent with the findings in Figure 2, highly positive ELISA results were significantly better at neutralizing than the negative samples for each ELISA.

While all the serological assays were significantly correlated with authentic virus neutralization, some assays performed better than others at predicting authentic virus neutralization (Supplemental Table 2). Based on correlation with authentic virus neutralization, the most accurate assays were the Luci. and SEAP pseudotyped virus neutralization assays. GFP pseudotyped virus neutralization, spike ELISA, and RBD ELISA form a second tier of assays which are still quite accurate at predicting authentic virus neutralization. Furthermore, the GFP pseudotyped virus neutralization was able to detect antibodies at significantly higher

dilutions than the other assays, making it the most sensitive assay tested. Despite nucleocapsid antigen being the basis for several common commercial antibody tests, nucleocapsid was the least predictive of authentic virus neutralization.

Collectively, these data demonstrate that VSV-ΔG pseudotyped virus neutralization platforms, especially Luci. and SEAP based platforms, are better at predicting authentic virus neutralization than ELISA regardless of the viral antigen tested. Not only are the Luci. and SEAP based pseudotype platforms most strongly correlated to authentic virus neutralization, they also have the lowest average difference in log(ND₅₀) compared to authentic virus neutralization. Previous reports have only compared ELISA titers to pseudotyped virus neutralization¹⁶, ELISA to authentic virus neutralization¹⁸, or only one type of ELISA and one pseudotyped virus platform against authentic virus neutralization¹⁰. Our studies provide one of the most comprehensive comparisons amongst multiple ELISA antigens, pseudotyped virus neutralization platforms, and authentic virus neutralization.

Of note, several spike and RBD positive samples showed very little authentic virus neutralization despite having moderate to high neutralization on the pseudotyped virus platforms. Furthermore, one sample appeared to show antibody dependent enhancement (ADE) in the authentic virus neutralization assay (1.8-fold increased plaque forming units (PFU)), but still showed low but detectable neutralization in all the pseudotyped virus platforms. While there is no definitive role for ADE during human SARS-CoV-2 infection, ADE has been demonstrated *in vitro* with other human coronaviruses¹⁹. Further characterization of this sample and screening for and characterizing similar samples will lead to a better understanding of the risk of ADE during SARS-CoV-2 infection. Recent evidence suggests that several SARS-CoV-2 variants, including B.1.351 and P.1, have decreased neutralization when treated with monoclonal antibodies or polyclonal sera derived from patients infected with early strains of SARS-CoV-2²⁰⁻²². Future studies need to assess how the mutations present in the variants differentially affect ELISA, pseudotyped virus neutralization, and authentic virus neutralization.

In addition to accuracy, the serological assays differ in several key features (Supplemental Table 2), and the assay of choice may have to be determined by the settings. The requirement for a BSL3 laboratory makes authentic virus assays technically challenging and unfeasible for many clinical and research applications. This can be overcome by pseudotyped viruses. However, creation and validation of the different pseudotyped viruses is not trivial and read-outs may require specialized equipment (e.g. luminometer for the Luci platform). Most laboratories have ready access to the equipment needed for performing ELISAs, making the technical requirements for these assays low. ELISAs can also be completed within several hours while the pseudotyped virus neutralization platforms require 12-24-hour incubations and authentic virus neutralization requires 48-72 hours. If all technical requirements have been met and are available, the assays are all relatively inexpensive, except for the Luci platform which requires expensive reagents for reading the results. If turnaround time is a priority, the RBD and spike ELISAs would provide the fastest results with minor decreases in predicting authentic virus neutralization response. Alternatively, in resource-limited settings like field hospitals, the GFP based pseudotyped virus neutralization assay requires only a basic fluorescence microscope for readout and is more predictive of authentic virus neutralization than any of the ELISAs. Overall, this study shows that all six serological assays, to varying degrees, correlated with authentic virus neutralization, and the optimal serological assay for assessing a protective antibody response is going to be institution and question specific.

206 Table 1: SARS-CoV-2 Protein ELISA Values

PCR	Sample	ELISA		
		RBD ratio (sample/negative)	N ratio (sample/negative)	Spike (AUC x 100)
Positive	1	17.92	13.75	9.26
Positive	2	16.48	14.00	12.90
Positive	3	9.62	7.20	3.36
Positive	4	8.79	4.45	1.97
Positive	5	18.41	8.74	14.19
Positive	6	10.56	6.46	8.21
Positive	7	12.01	6.20	3.43
Positive	8	19.10	16.61	11.38
Positive	9	15.54	5.85	2.87
Positive	10	16.73	10.95	3.08
Positive	11	21.74	9.42	5.30
Positive	12	24.08	14.14	7.24
Positive	13	20.13	15.73	3.97
Positive	14	19.59	13.09	5.39
Positive	15	8.12	5.32	2.43
Positive	16	13.76	10.37	3.89
Positive	17	6.94	5.59	3.51
Positive	18	21.43	5.11	12.55
Positive	19	21.20	1.76	5.04
Positive	20	18.76	8.33	8.03
Positive	21	20.48	17.47	6.39
Positive	22	18.39	9.25	7.29
Positive	23	17.04	9.90	6.26
Positive	24	17.90	5.82	6.74
Negative	25	1.04	0.71	
Negative	26	2.15	1.14	
Negative	27	1.80	1.17	
Negative	28	1.72	2.81	
Negative	29	2.27	3.57	
Negative	30	0.99	0.94	
Negative	31	1.99	1.01	
Negative	32	1.45	0.68	
Negative	33	1.75	1.25	
Negative	34	1.07	0.75	

207

208

209 Table 2: Authentic Virus and Pseudotyped Virus Neutralization Summary Statistics

PCR	Sample	Authentic Virus		GFP Pseudotype		Luci. Pseudotype		SEAP Pseudotype	
		AUC	ND ₅₀	AUC	ND ₅₀	AUC	ND ₅₀	AUC	ND ₅₀
Positive	1	0.565	380	0.739	908	0.520	276	0.582	382
Positive	2	0.630	671	0.804	2305	0.692	662	0.645	535
Positive	3	0.166	1	0.367	153	0.137	33	0.164	60
Positive	4	0.321	1	0.309	116	0.237	78	0.287	109
Positive	5	0.831	2527	0.860	5744	0.821	2162	0.836	2834
Positive	6	0.662	745	0.723	1505	0.668	692	0.706	722
Positive	7	0.206	51	0.661	672	0.467	253	0.412	181
Positive	8	0.794	2612	0.810	2997	0.764	1341	0.814	1922
Positive	9	0.095	8	0.613	665	0.314	112	0.322	121
Positive	10	0.284	86	0.633	586	0.412	193	0.368	154
Positive	11	0.649	782	0.822	3747	0.531	307	0.518	296
Positive	12	0.702	682	0.792	1995	0.597	447	0.645	527
Positive	13	0.646	606	0.837	3229	0.638	551	0.606	392
Positive	14	0.468	275	0.701	716	0.444	208	0.413	186
Positive	15	0.232	93	0.599	462	0.301	116	0.269	105
Positive	16	0.825	2769	0.833	2317	0.820	2606	0.848	2910
Positive	17	0.429	281	0.770	3222	0.470	248	0.493	259
Positive	18	0.886	35373	0.887	50378	0.886	37272	0.890	64851
Positive	19	0.808	2008	0.842	3409	0.694	662	0.700	635
Positive	20	0.593	495	0.733	983	0.565	369	0.634	589
Positive	21	0.414	175	0.744	1373	0.603	417	0.707	850
Positive	22	0.462	266	0.724	1001	0.610	444	0.719	972
Positive	23	0.192	63	0.636	390	0.450	215	0.423	197
Positive	24	0.414	190	0.617	441	0.370	163	0.474	266
Negative	25	0.031	9	0.151	40	0.000	1	0.000	1
Negative	26	0.035	3	0.191	57	0.000	1	0.000	1
Negative	27	0.141	33	0.035	10	0.000	1	0.000	1
Negative	28	0.212	54	0.007	2	0.000	1	0.000	1
Negative	29	0.176	43	0.051	15	0.000	1	0.060	18
Negative	30	0.000	1	0.231	82	0.000	1	0.000	1
Negative	31	0.192	37	0.000	1	0.000	1	0.060	17
Negative	32	0.110	18	0.209	68	0.000	1	0.000	1
Negative	33	0.051	4	0.000	1	0.000	1	0.000	1
Negative	34	0.148	38	0.059	18	0.000	1	0.000	1

210

211

Figure 1

Figure 1: Comparison of neutralization assays by sample groups. Area under the curve (AUC) (top) and neutralization dilution – 50% (ND₅₀) (bottom) calculations by neutralization assay type. AUC and ND₅₀ values were calculated and used to compare authentic virus neutralization (black), GFP pseudotype neutralization (pink), luciferase pseudotype (teal), and SEAP pseudotype (purple). * p < 0.05, ** p < 0.01, ***p < 0.001 (mixed-effects model with the Geisser-Greenhouse correction and Tukey multiple comparisons post-test and p-value adjustment). n = 34 samples run on each assay.

223 Figure 2

224

225

Figure 2: Correlation of SARS-CoV-2 serologic assays. A) SARS-CoV-2 specific ELISA assays and B) VSV pseudotyped virus neutralization assays were compared by simple linear regression. The Pearson's r values (a metric of correlation) and p-values corresponding to each graph are to the lower left of each set of graphs. The shade of background corresponds to the degree of correlation between the two assays. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$.

233 Figure 3

236

237

Figure 3: Comparison of high positive, low positive, and negative ELISA groups across neutralization assays. A) RBD B) spike (S) and C) nucleocapsid (N) positive samples were divided into high and low positives by finding cut off values using histograms (RBD ratio: 15, N ratio: 10, and Spike: 6). Log(ND₅₀) values for the corresponding samples were then graphed and compared by Kruskal-Wallis test with Dunn's multiple comparisons tests. Significance thresholds: * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, p **** $p < 0.0001$.

References

- 1 Corman, V. M. *et al.* Detection of 2019 novel coronavirus (2019-nCoV) by real-time RT-PCR. *Euro Surveill* **25**, 2000045, doi:10.2807/1560-7917.ES.2020.25.3.2000045 (2020).
- 2 Wang, W., Tang, J. & Wei, F. Updated understanding of the outbreak of 2019 novel coronavirus (2019-nCoV) in Wuhan, China. *Journal of medical virology* **92**, 441-447, doi:10.1002/jmv.25689 (2020).
- 3 Wu, F. *et al.* A new coronavirus associated with human respiratory disease in China. *Nature* **579**, 265-269, doi:10.1038/s41586-020-2008-3 (2020).
- 4 Zhou, P. *et al.* A pneumonia outbreak associated with a new coronavirus of probable bat origin. *Nature* **579**, 270-273, doi:10.1038/s41586-020-2012-7 (2020).
- 5 Grzelak, L. *et al.* A comparison of four serological assays for detecting anti-SARS-CoV-2 antibodies in human serum samples from different populations. *Science translational medicine* **12**, eabc3103, doi:10.1126/scitranslmed.abc3103 (2020).
- 6 Pérez-García, F. *et al.* Comparative evaluation of six immunoassays for the detection of antibodies against SARS-CoV-2. *Journal of virological methods*, 114047, doi:10.1016/j.jviromet.2020.114047 (2020).
- 7 Whitman, J. D. *et al.* Evaluation of SARS-CoV-2 serology assays reveals a range of test performance. *Nature Biotechnology* **38**, 1174-1183, doi:10.1038/s41587-020-0659-0 (2020).
- 8 Ju, B. *et al.* Human neutralizing antibodies elicited by SARS-CoV-2 infection. *Nature* **584**, 115-119, doi:10.1038/s41586-020-2380-z (2020).
- 9 van der Heide, V. Neutralizing antibody response in mild COVID-19. *Nature Reviews Immunology* **20**, 352-352, doi:10.1038/s41577-020-0325-2 (2020).

- 10 von Rhein, C. *et al.* Comparison of potency assays to assess SARS-CoV-2 neutralizing antibody capacity in COVID-19 convalescent plasma. *Journal of virological methods* **288**, 114031-114031, doi:10.1016/j.jviromet.2020.114031 (2020).
- 11 Yu, J. *et al.* DNA vaccine protection against SARS-CoV-2 in rhesus macaques. *Science* **369**, 806, doi:10.1126/science.abc6284 (2020).
- 12 Polack, F. P. *et al.* Safety and Efficacy of the BNT162b2 mRNA Covid-19 Vaccine. *New England Journal of Medicine*, doi:10.1056/NEJMoa2034577 (2020).
- 13 Deshpande, G. *et al.* Neutralizing antibody responses to SARS-CoV-2 in COVID-19 patients. *Indian Journal of Medical Research* **152**, 82-87, doi:10.4103/ijmr.IJMR_2382_20 (2020).
- 14 Oguntuyo, K. Y. *et al.* Quantifying absolute neutralization titers against SARS-CoV-2 by a standardized virus neutralization assay allows for cross-cohort comparisons of COVID-19 sera. *medRxiv*, doi:10.1101/2020.08.13.20157222 (2020).
- 15 Amanat, F. *et al.* A serological assay to detect SARS-CoV-2 seroconversion in humans. *Nature Medicine* **26**, 1033-1036, doi:10.1038/s41591-020-0913-5 (2020).
- 16 McAndrews, K. M. *et al.* Heterogeneous antibodies against SARS-CoV-2 spike receptor binding domain and nucleocapsid with implications for COVID-19 immunity. *JCI Insight* **5**, doi:10.1172/jci.insight.142386 (2020).
- 17 Piccoli, L. *et al.* Mapping Neutralizing and Immunodominant Sites on the SARS-CoV-2 Spike Receptor-Binding Domain by Structure-Guided High-Resolution Serology. *Cell* **183**, 1024-1042.e1021, doi:10.1016/j.cell.2020.09.037 (2020).
- 18 Heaney, C. D. *et al.* Comparative performance of multiplex salivary and commercially available serologic assays to detect SARS-CoV-2 IgG and neutralization titers. *medRxiv*, doi:10.1101/2021.01.28.21250717 (2021).

293 19 Lee, W. S., Wheatley, A. K., Kent, S. J. & DeKosky, B. J. Antibody-dependent
294 enhancement and SARS-CoV-2 vaccines and therapies. *Nature Microbiology* **5**, 1185-
295 1191, doi:10.1038/s41564-020-00789-5 (2020).

296 20 Chen, R. E. *et al.* Resistance of SARS-CoV-2 variants to neutralization by monoclonal
297 and serum-derived polyclonal antibodies. *Nature Medicine*, doi:10.1038/s41591-021-
298 01294-w (2021).

299 21 Hoffmann, M. *et al.* SARS-CoV-2 variants B.1.351 and B.1.1.248: Escape from
300 therapeutic antibodies and antibodies induced by infection and vaccination. *bioRxiv*,
301 2021.2002.2011.430787, doi:10.1101/2021.02.11.430787 (2021).

302 22 Wang, P. *et al.* Antibody Resistance of SARS-CoV-2 Variants B.1.351 and B.1.1.7.
303 *Nature*, doi:10.1038/s41586-021-03398-2 (2021).
304
305

Methods

Data Availability

Authors can confirm that all relevant data are included in the paper and/or its supplementary information files.

RBD/N ELISA

SARS-CoV-2 RBD protein was diluted to a concentration of 1.5 µg/ml in PBS and added at 50 µl per well to a 96-well ELISA plate. The ELISA plates were sealed and allowed to incubate at 4°C overnight. The next day the coating solution was removed, and the plates were blocked at room temperature (RT) using 3% milk (200 µl per well) for a minimum of 1 hour but not exceeding 4 hours. While the plates were being blocked, the samples were prepared by diluting the plasma 1:50 in 1% milk. Following the blocking period, the milk was removed, and the plates were washed 3x with 0.1% phosphate buffered saline containing 0.1% Tween-20 (PBS-T) using 200 µl per well. The diluted plasma was added to the blocked plate at 50 µl per well along with 2 positive controls (α SARS-CoV-2 RBD antibody at 1:5000, 1:25000, 1:125000, and 1:625000 dilutions and plasma from a naturally infected donor at a 1:50 dilution) and a known negative, pre-pandemic plasma sample (1:50). The samples were incubated for 1.5 hours at RT and then removed and washed 3x with 200 µl 0.1% PBS-T. Goat α human IgG horseradish peroxidase (HRP) conjugated secondary antibody was diluted 1:2500 in 1% milk and 50 µl was added to each well of the washed plate and incubated at RT for 30 minutes. Following the incubation period, the secondary was removed, and the plate was washed 3x with 0.1% PBS-T. OPD substrate was prepared directly before use and added at 50 µl per well for exactly 8 minutes. The O-phenylenediamine dihydrochloride (OPD) substrate was stopped by adding 50 µl of 3M HCl and then the plate was read using a spectrophotometer at 490nm.

Spike ELISA

SARS-CoV-2 spike protein was diluted to a concentration of 2 µg/ml in PBS and added at 50 µl per well to a 96-well ELISA plate. The ELISA plates were sealed and allowed to incubate at 4°C overnight. The next day the coating solution was removed, and the plates were blocked using 3% milk (200 µl per well) for a minimum of 1 hour but not exceeding 4 hours. While the plates were being blocked, the samples were prepared by creating a 3-fold serial dilution starting at 1:100 and ending at 1:8100 (1% milk as diluent). Following the blocking period, the milk was removed, and the plates were washed 3x with 0.1% PBS-T using 200 µl per well. The diluted plasma was added to the blocked plate at 50 µl per well along with 2 positive controls (α SARS-CoV-2 RBD antibody at 1:5000, 1:25000, 1:125000, and 1:625000 dilutions and plasma from a naturally infected donor at 1:100, 1:300, 1:900, 1:2700, and 1:8100 dilutions) and a known negative, pre-pandemic plasma sample (1:100). The samples were incubated for 1.5 hours at RT and then removed and washed 3x with 200 µl 0.1% PBS-T. Goat α human IgG HRP conjugated secondary antibody was diluted 1:2500 in 1% milk and 50 µl was added to each well of the washed plate and incubated at RT for 30 minutes. Following the incubation period, the secondary was removed, and the plate was washed 3x with 0.1% PBS-T. OPD substrate was prepared directly before use and added at 50 µl per well for exactly 8 minutes. The OPD substrate was stopped by adding 50 µl of HCL acid and then the plate was read using a spectrophotometer at 490nm. Spike data is presented as either AUC or AUC × 100 in order to plot it on the same scale as the other ELISAs.

Tissue culture

VeroE6 cells stably expressing TMPRSS2 (Vero-TMPRSS2) (XenoTech) were cultured in Eagle's Minimal Essential Medium (EMEM) supplemented with 10% fetal bovine serum (FBS), 100 U/mL penicillin, 100 µg/mL streptomycin, and 2 mM GlutaMax (Gibco). Media was

supplemented with 1 mg/mL G418 every other passage. All tissue culture was performed in a humidified incubator set to 37° C and 5% CO₂.

SARS-CoV-2 neutralizing antibody assay

Serially diluted plasma samples were mixed with diluted (approximately 6 PFU/cm²) SARS-CoV-2 (2019n-CoV/USA_WA1/2020) in EMEM supplemented with 5% FBS, 100 U/mL penicillin, 100 µg/mL streptomycin, and 2 mM GlutaMax. Mixtures were incubated for 1 h in a humidified incubator at 37° C and 5% CO₂. After 1 h, culture media was removed from approximately 90% confluent Vero-TMPRSS2 cells grown in 6-well plates and replaced with virus/plasma mixtures. Plates were returned to the incubator for 1 h at 37° C and 5% CO₂. Plates were rocked manually every 15 minutes. After incubation, an agarose overlay containing Minimal Essential Media (MEM) supplemented with 5% FBS 100 U/mL penicillin, 100 µg/mL streptomycin, 2 mM GlutaMax, 0.075% sodium bicarbonate, 0.01 M 4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid (HEPES), and 1% low melting temperature agarose (SeaPlaque; Lonza) was added to each well. Once agarose hardened at RT, plates were returned to the incubator at 37° C and 5% CO₂. After 48 h, cells were fixed with 10% neutral buffered formalin for 1 h, the agar plugs were removed, and then cells were stained with crystal violet for 5 – 10 minutes. Upon rinsing with H₂O, plaques were visualized and counted. All samples were run in duplicate

VSV-ΔG-GFP-SARS-CoV-2-S Neutralizing antibody assay

Serially diluted plasma samples were mixed with diluted and mixed with Spike/VSV-ΔG-GFP pseudotypes in EMEM supplemented with 5% FBS, 100 U/mL penicillin, 100 µg/mL streptomycin, and 2 mM GlutaMax. Mixtures were incubated for 1 h in a humidified incubator at 37° C and 5% CO₂. After 1 h, culture media was removed from approximately 90% confluent Vero-TMPRSS2 cells grown in 96-well plates and replaced with virus/plasma mixtures. Plates

were returned to the incubator at 37° C and 5% CO₂. After 24 h, IU were quantified manually using an EVOS fluorescence microscope. All samples run in duplicate.

Luciferase Assay

20 hours prior to assay set up, Vero-TMRSS2 were plated in a 96-well plate at 20,000 cells per well in DMEM supplemented with 5% FBS and 1 mg/mL G418. For assay set up, plasma samples were initially diluted 1:100 and serially diluted 1:3 in DMEM supplemented with 5% FBS. Diluted samples were mixed 1:1 with Spike/VSV-ΔG-Luciferase pseudotyped virus diluted to final 250 IU per well in serum free DMEM. Mixtures were incubated for 1 hour in a humidified incubator at 37° C and 5% CO₂. After the incubation period, culture medium was removed from Vero-TMPRSS2 cells and virus/plasma mixture was added to the cells in triplicate. Plates were incubated for approximately 18 hours in a humidified incubator at 37° C and 5% CO₂. After the incubation period, Luc-Screen Extended-Glow (ThermoFisher) buffers were added to the wells according to manufacturer's instructions and incubated for a minimum of 10 minutes at room temperature protected from light. Luminescence was measured with a luminometer using one second integration time.

SEAP Assay

20 hours prior to assay set up, Vero-TMRSS2 were plated in a 96-well plate at 20,000 cells per well in DMEM supplemented with 5% FBS and 1 mg/mL G418. For assay set up, plasma samples were initially diluted 1:100 and serially diluted 1:3 in DMEM supplemented with 5% FBS. Diluted samples were mixed 1:1 with purified Spike/VSV-ΔG--SEAP pseudotyped virus diluted to final 250 IU per well in serum free DMEM. Mixtures were incubated for 1 hour in a humidified incubator at 37° C and 5% CO₂. After the incubation period, culture medium was removed from Vero-TMPRSS2 cells and virus/plasma mixture was added to the cells in triplicate. Plates were incubated for approximately 28 hours in a humidified incubator at 37° C

and 5% CO₂. After the incubation period, Quanti-Blue (InvivoGen) solution was combined with 20 µL supernatant according to manufacturer's instructions and incubated for a minimum of 15 minutes at 37° C protected from light. Optical density was measured at 620-655 nm.

SARS-CoV-2/VSV pseudotype production

VSV-ΔG pseudotypes displaying the full-length SARS-CoV-2 spike (Wuhan-Hu-1 strain) were generated essentially as described²³ with the following modifications. Baby hamster kidney (BHK-21) cells in 10 cm dishes were transfected using Lipofectamine 2000 according to the manufacturer's instructions with 24 µg of a plasmid encoding a codon-optimized cDNA for the SARS-CoV-2 spike¹⁵, which was generously provided by Florian Krammer. Approximately 20-24 hours later the transfected cells were infected at a multiplicity of 5 with VSV-G pseudotyped ΔG-GFP, luciferase, or SEAP. Virus was adsorbed for 1 hr, the inoculum was removed, cells were rinsed once with serum-free DMEM and then 4 ml of hybridoma supernatant containing the I1 monoclonal antibody²⁴ was added for 30 minutes to neutralize residual VSV-ΔG pseudotyped virus from the inoculum and then replaced with DMEM containing 20% fetal bovine serum. The supernatant containing the spike-ΔG pseudotypes was collected 22-24 hours later, cell debris was removed by centrifugation at 450 x g for 10 minutes. For the ΔG-GFP and luciferase pseudotypes, the supernatant was aliquoted and stored at -80° C. For the ΔG-SEAP pseudotypes, the supernatant was transferred to a Beckman SW41 tube, underlayered with sterile 20% sucrose in PBS, and virus was pelleted at 35,000 rpm for 45 minutes in a SW41 swinging bucket rotor. Pelleting virus was required to separate it from SEAP released from the infected cells. The pellets were resuspended in DMEM containing 20% FBS and stored at -80° C.

Statistics

Area under the curve (AUC) and neutralization dilution – 50% (ND₅₀) analyses were performed in GraphPad Prism (version 9.0.0): non-linear regression (dose-response). Pearson's r values for comparing assays by percent maximum AUC were calculating using simple linear regression analysis in GraphPad Prism. AUC and ND₅₀ values for the different assays were compared by mixed-effects model with the Geisser-Greenhouse correction and Tukey multiple comparisons post-test and p-value adjustment in GraphPad Prism (version 9.0.0). Kruskal-Wallis tests with Dunn's multiple comparisons tests. were performed to compare neutralizing antibody responses between highly positive ELISA samples, low positive ELISA samples, and negative samples. Principle component analysis (PCA) was performed in GraphPad Prism (version 9.0.0) with principle components selected based on parallel analysis. A 95% percentile level was used, and 1000 simulations were performed for the PCA. The Bland-Altman analyses were performed in GraphPad Prism (version 9.0.0).

447 Methods References

- 448 15 Amanat, F. *et al.* A serological assay to detect SARS-CoV-2 seroconversion in humans.
449 *Nature Medicine* **26**, 1033-1036, doi:10.1038/s41591-020-0913-5 (2020).
450 23 Whitt, M. A. Generation of VSV pseudotypes using recombinant Δ G-VSV for studies on
451 virus entry, identification of entry inhibitors, and immune responses to vaccines. *Journal*
452 *of virological methods* **169**, 365-374, doi:10.1016/j.jviromet.2010.08.006 (2010).
453 24 Lefrancois, L. & Lyles, D. S. The interaction of antibody with the major surface
454 glycoprotein of vesicular stomatitis virus. I. Analysis of neutralizing epitopes with
455 monoclonal antibodies. *Virology* **121**, 157-167 (1982).

456

457

Acknowledgements

The authors wish to thank Amy E. Davis, Virginia Hargest, Rebekah Honce, Brandi Livingston, Victoria Meliopoulos, Bridgett Sharp, Maria Smith, and Kristin Wiggins for aiding the Schultz-Cherry lab's COVID-19 response; the members of the Thomas, McGargill, and Schultz-Cherry labs for technical assistance and feedback on the work; Dr. Gang Wu and the Center for Applied Bioinformatics at St. Jude along with Michael Meagher, Timothy Lockey, and the St. Jude Good Manufacturing Practice (GMP) facility; Tamanna Shamrin and Rishi Kodela for creation and management of the clinical database. This work was funded by ALSAC, the NIAID for the St. Jude Center of Excellence for Influenza Research and Surveillance (CEIRS contract HHSN27220140006C) and the NIAID Collaborative Influenza Vaccine Innovation Centers (CIVIC) contract 75N93019C00052, and by the University of Tennessee Research Foundation. JHE is supported by the American Society of Hematology Scholar Award. Work in the Krammer laboratory is partially funded by the NIAID Collaborative Influenza Vaccine Innovation Centers (CIVIC) contract 75N93019C00051, NIAID Center of Excellence for Influenza Research and Surveillance (CEIRS, contract # HHSN272201400008C), by the generous support of the Cohen Foundation, the JPB Foundation and the Open Philanthropy Project (research grant 2020-215611 (5384), and by anonymous donors.

Author contributions

Conceptualization: N.W., K.W., S.C., E.K.R., S.S.-C.
Conception, design, and oversight of parent study; SJTRC: J.H., H.H., K.J.A., R.D., A.H.G., J.H.E., J.W., and P.G.T.
Formal Analysis: NW., K.W., S.C., E.K.R., T.M., L.T.
Investigation: N.W., K.W., S.C., E.K.R.
Methodology: NW., K.W., S.C., E.K.R., C.Y.L. E.K.A., P.F., M.A.M.
Resources: F.K., M.A.W.

484 Sample acquisition and curation: K.J.A., A.G., A.H.G., E.K.A., J.H.E., J.W., and P.G.T.
 485 Writing, original draft: N.W., E.K.R.
 486 Writing, review, and editing: N.W., K.W., S.C., E.K.R., M.A.W., H.H., J.H.E., A.H.G., T.M., T.L.,
 487 D.R.H., M.A.M., F.K., J.W., P.G.T., S.S.-C.
 488 Visualization: N.W.
 489 Supervision: S.S.-C.

490

491 Competing interest's declaration

492 The Icahn School of Medicine at Mount Sinai has filed patent applications relating to SARS-
 493 CoV-2 serological assays and NDV-based SARS-CoV-2 vaccines which name FK as inventor.
 494 FK would also like to note the following, which could be perceived as a conflict of interest: He
 495 has previously published work on influenza virus vaccines with S. Gilbert (University of Oxford),
 496 has consulted for Curevac, Merck and Pfizer (before 2020), is currently consulting for Pfizer,
 497 Seqirus and Avimex, his laboratory is collaborating with Pfizer on animal models of SARS-CoV-
 498 2, his laboratory is collaborating with N. Pardi at the University of Pennsylvania on mRNA
 499 vaccines against SARS-CoV-2, his laboratory was working in the past with GlaxoSmithKline on
 500 the development of influenza virus vaccines and two of his mentees have recently joined
 501 Moderna. No other others have conflicts of interest to report.