medRxiv preprint doi: https://doi.org/10.1101/2021.04.15.21255554; this version posted April 20, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

Frequency of variants of hereditary cancer risk syndrome in 1682 Brazilian patients genotyped by next-generation sequencing

Short Title: Variants of hereditary cancer risk in 1682 Brazilian patients

Jarbas Maciel de Oliveira[1,2*], Nuria Bengala Zurro[3], Antonio Victor Campos Coelho[3], Marcel Pinheiro Caraciolo[3], Rodrigo Bertollo de Alexandre[3], George de Vasconcelos Carvalho Neto[3], Ivana Grivicich[2] and João Bosco de Oliveira Filho[3*]

[1] Hospital Memorial Arcoverde, Avenida José Bonifácio 1121, Arcoverde, Pernambuco 56503-740, Brazil

[2] Programa de Pós Graduação em Biologia Celular e Molecular Aplicada à Saúde, Universidade Luterana do Brasil, Av. Farroupilha 8001, Canoas - Rio Grande do Sul 92425-020, Brazil

[3] Hospital Israelita Albert Einstein, Avenida Albert Einstein 627, São Paulo 05652-900, Brazil

* Corresponding author

jarbasmaciel@yahoo.com.br (JMO) or joao.bosco@einstein.br (JBOF)

Funding: This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

medRxiv preprint doi: https://doi.org/10.1101/2021.04.15.21255554; this version posted April 20, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

Abstract

Hereditary cancer risk syndromes are a group of disorders caused by germline variants in a growing number of genes. Most studies on hereditary cancer have been conducted in white populations. Here we report the largest study in Brazilian individuals with multiple, self-reported, ethnicities. We genotyped 1682 individuals from all regions of the country who were referred to genetic testing for hereditary cancer risk with multigene Next-generation sequencing (NGS) panels. Most were women, and had a personal and/or family history of cancer. The majority of cancer cases were breast and ovarian. We identified a total of 321 pathogenic/likely pathogenic (P/LP) variants in 305 people (18.1%), corresponding to 166 unique variants. These variants were distributed among 32 genes, and most were detected on *BRCA1* and *BRCA2* (129 patients, 26.2% and 14.3% of all P/LP hits, respectively). The prevalence of any genes transheterozygosity in our sample was 0.89% (15/1682). The *BRCA1/BRCA2* double heterozygosity rate was 0.78% (1/129) for *BRCA* variants carriers and 0.06% (1/1682) overall. We classified patients according to the NCCN and Brazilian National Health Agency (ANS) genetic testing recommendation criteria. We found that the criteria had false negative rates of 17.3% and 44.2%, meaning that both failed to detect a substantial part of P/LP positive patients. Therefore, our results show that NGS adds to knowledge on the Brazilian spectrum of germline variants associated with cancer risk and indicate that Brazilian testing guidelines should be improved.

medRxiv preprint doi: https://doi.org/10.1101/2021.04.15.21255554; this version posted April 20, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

Keywords: breast cancer, cancer genomics, familial cancer history, population genetics, colorectal cancer.

Introduction

Hereditary cancer risk syndromes are a group of disorders caused by germline pathogenic variants in a growing number of genes. They are the main predisposing factor in about 5-10% of all diagnosed cancers. Accurate genetic diagnosis in patients with hereditary cancer risk syndromes may reduce morbidity and mortality by allowing the adoption of specific preventative and therapeutic measures. Additionally, family members at risk can be tested and counseled, extending the clinical benefits to many individuals [1].

Currently, there are over 30 hereditary cancer syndromes already described, the most prevalent being hereditary breast and ovarian cancer syndrome (HBOC). Other syndromes include Li-Fraumeni, Cowden, Lynch, Familial Adenomatous Polyposis (APC), multiple Endocrine Neoplasia Type 1 or Wermer's Syndrome (MEN1), multiple Endocrine Neoplasia Syndrome - Type 2 (RET), and Von Hippel-Lindau Syndrome (VHL), to cite the most common. Most syndromes are inherited in an autosomal-dominant manner, with variable penetrance, and are caused by mutations in oncogenes or tumor suppressor genes [2]. Although initially hard to diagnose genetically, the advent of next-generation sequencing (NGS) has allowed for testing using panels with dozens to hundreds of genes simultaneously and inexpensively [3]. This has caused a great increase in the clinical use of these NGS panels, while a growing body of evidence show their clinical utility as compared to single-gene testing [4, 5].

medRxiv preprint doi: https://doi.org/10.1101/2021.04.15.21255554; this version posted April 20, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

Although large studies have been published on the diagnostic utility of such panels, these are mostly restricted to white populations. The aim of this study was to describe the frequency and type of germline pathogenic variants in cancer susceptibility genes in patients with clinical criteria of hereditary cancer syndrome by NGS in the genetically admixed Brazilian population.

Materials and methods

Inclusion and exclusion criteria

The cohort to the retrospective study included 1682 Brazilian patients who received a multi-gene NGS panel for risk of hereditary cancer in the Genomika diagnostics unit, a genetics laboratory of the Albert Einstein Israeli Hospital (Genomika-Einstein hereafter). As this study took place in a private diagnostic laboratory this cohort was not specifically selected for any sex, age, ethnicity, or history of cancer. Both patients with or without criteria for hereditary risk cancer syndrome were considered. After genetic assessment we classified patients according to the USA National Comprehensive Cancer Network (NCCN, www.nccn.org) guidelines: genetic/familial high-risk assessment versions 3.2019 and 1.2020. Non-Brazilian patients were excluded.

Research ethics statement

This project was approved by the Institutional review board from the Research Ethics Committee of the Institute of Integral Medicine Professor Fernando Figueira (CEP-IMIP) in Recife, Pernambuco and all individuals

medRxiv preprint doi: https://doi.org/10.1101/2021.04.15.21255554; this version posted April 20, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

provided written consent for multi-gene testing (protocol number CAAE 29567220.4.1001.0071).

Multigene genotyping panel

Genotyping was performed through panels that covered the entire coding region including 20 bp flanking each exon and noncanonical splice region. Patients were sequenced with panels that ranged from 27 to 78 genes, including genes mostly associated with hereditary cancer risk, such as *APC, ATM, BARD1, BMPR1A, BRCA1, BRCA2, CDH1, CDK4, CHEK2, EGFR, EPCAM, GREM1 HOXB13, MEN1, MITF, MLH1, MRE11A, MSH2, MSH6, MUTYH, NBN, PALB2, PMS2, POLD1, POLE, PTEN, RAD50, RAD51C, RAD51D, RB1, RET, SMAD4, STK11, TP53, VHL.* The complete list of genes is available in the Table A.1).

Library preparation for NGS

Laboratory procedures were performed at Genomika-Einstein units (Recife - PE and São Paulo - SP). Genomic DNA was extracted from peripheral blood or saliva by automation (QIASymphony platform, QIAGEN, Hilden, Germany) using the DNA Mini Kit extraction kit (QIAGEN). The quality and quantity of the extracted DNA were assessed by fluorometry (Qubit, Thermo Fisher, Waltham, Massachusetts). The genomic DNA was enzymatically fragmented and enriched. The fragments were barcoded via multiplexed PCR technology by QIAseq Targeted DNA Panels (CDHS 174272-2274 QIAGEN). The genes were sequenced on MiSeq or Next-Seq 550 instruments (Illumina,

medRxiv preprint doi: https://doi.org/10.1101/2021.04.15.21255554; this version posted April 20, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

San Diego, CA) using MiSeq Reagent Kit v2 (300-cycles) or Next-Seq 550 High-throughput kit (300 cycles) with 100% coverage and 50X minimum depth.

Sanger sequencing

In cases where variant confirmation was deemed necessary (low read depth, complex indels), the genotype was determined by conventional PCR followed by Sanger sequencing on an ABI 3500 (Thermo Fisher, Waltham, Massachusetts) automatic sequencer. The variant call was made via the reference transcript from the ATG initiation codon aligned against the GRCh37 / hg19 reference genome.

Alu insertion identification

Relevant genetic screenings of breast/ovarian cancer families of Portuguese ancestry demonstrate that the c.156_157ins*Alu* variant is the most frequent *BRCA2* rearrangement. Detection of this variant was performed as published elsewhere [6] . Briefly, PCR of *BRCA2* exon 3 was performed and the *Alu* insertion is detected by differential agarose gel electrophoresis.

Variant classification and bioinformatic analysis

Bioinformatic analysis was performed using GATK 3.0 best practices. VCFs were annotated using Annovar and in-house databases. The clinical significance of the variants was determined according to the criteria of the American College of Genomics and Genetics (ACMG) 2015 guidelines for sequence variant interpretation and 2018 guideline update [7, 8]. Each variant

medRxiv preprint doi: https://doi.org/10.1101/2021.04.15.21255554; this version posted April 20, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

classification were performed independently by two specialists. Allelic frequency was examined through access to population databases as Genome Aggregation Database (gnomAD) [9], Exome Aggregation Consortium (ExAC) [10], and The 1000 Genomes Project [11]. Disease information were upload form ClinVar [12] and the Leinden Open Variation Database (LOVD) [13]. All variant information used for classification were updated and upgraded in an in-house local knowledgebase.

Results

Study sample characteristics and cancer histories

Between July 2016 and July 2019, we collected samples and clinical/demographic characteristics from 1682 Brazilian individuals referred to a private laboratory to undergo genotyping with a cancer hereditary risk multigenic panel (Table 1).

Among the individuals, 1557 (92.6%) were women with mean age of 47.4 ± 12.3 (range 11-88) years and 125 (7.4%) were men with mean age of 51.4 ± 15.2 (range 14-91) years. Regarding ethnic ancestry, 1007 (59.9%) individuals reported unknown background, 330 (19.6%) reported having multiple ethnicities (admixed), 259 (15.4%) white, 26 (1.5%) black, 22 (1.3%) Ashkenazi Jewish, 21 (1.2%) native indigenous and 17 Asian (1.0%) (Table 1).

Personal cancer history was reported by 1126 (66.9%) individuals, with breast or ovary cancers identified in 971 individuals (57.7%); colorectal cancer in 71 (4.2%); thyroid/parathyroid in 36 (2.1%); prostate cancer in 15 (0.9%); stomach cancer in 14 (0.8%); uterus, kidney, and sarcomas in 13 each (0.8%); leukemia in 11 (0.6%); non-melanoma skin and pancreas cancers in 9 each

medRxiv preprint doi: https://doi.org/10.1101/2021.04.15.21255554; this version posted April 20, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

(0.5%); and melanoma in 6 (0.4%). Other types of cancer (lung, liver, bladder, larynx etc.) were identified in 17 individuals (1.0%). Overall, 268 individuals (15.9%) did not present malignant neoplasms and for 293 individuals (17.4%) the history information was not available.

Family cancer history (considering first-, second- or third-degree relatives) was reported by 1182 (70.3%) individuals: 806 (47.9%) reported breast cancer family history; 310 (23.0%) colorectal cancers; 339 (20.2%) prostate cancer; 219 (13.0%) head and neck tumors; 172 (10.2%) lung cancer; 155 (9.2%) stomach cancer; 119 (7.1%) breast and ovarian and 48 (2.9%) ovarian cancer exclusively. No family history was reported by 163 (9.7%) individuals and 337 (20.0%) did not provide information regarding family history (Table 1, Table A.2).

Genetic findings

Pathogenic or likely pathogenic variants were found in 305 (18.1%) of the 1682 individuals. Additionally, 1252 variants of uncertain significance (VUS) were found in 753 (44.8%) individuals (Fig. 1A, Table A.3). The remaining 624 (37.1%) patients did not present any variants of clinical interest whatsoever (negatives).

The 305 individuals collectively had 321 pathogenic/likely pathogenic variant hits (corresponding to 166 unique variants) in 32 genes: *APC, ATM, BARD1, BRCA1, BRCA2, BRIP1, CDH1, CDK4, CHEK2, EPCAM, FANCC, MEN1, MITF, MLH1, MRE11A, MSH2, MSH6, MUTYH, NBN, PALB2, PMS2, POLD1, PTCH2, PTEN, RAD50, RAD51, RAD51C, RAD51D, RB1, RECQL, RET* and *TP53*.

medRxiv preprint doi: https://doi.org/10.1101/2021.04.15.21255554; this version posted April 20, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

The genes that most commonly presented pathogenic or likely pathogenic variants were *BRCA1* (84/321 = 26.2%), *BRCA2* (46, 14.3%) and *PALB2* (25, 7.8%). Lynch syndrome genes (*MLH1*, *MSH2*, *MSH6* and *PMS2*) came next with 17 (5.3%) and were tied with *TP53*, also with 17 (5.3%) hits. Next came *ATM* (15, 4.7%), *CHEK2* (14, 4.4%), *RAD51C* (9, 2.8%), *RAD51* (8, 2.5%), *RAD50* (7, 2.2%), *BRIP1* (6, 1.9%), *BARD1* (4, 1.2%), *PTEN* (4, 1.2%), *RAD51D* (4, 1.2%) and *APC* (3, 0,9%). Other genes amounted 17 (3.0%) variants (*NBN, RET,* biallelic *MUTYH, MITF, CDH1, EPCAM, FANCC, MRE11, POLD1, RB1, RECQL, MEN1, CDK4* and *PTCH2*). Moreover, monoallelic *MUTYH* variants, considered to have low penetrance, were detected in 42 (13.1%) cases (Fig. 1B).

In terms of unique variants, we detected 24 in *BRCA1*, 32 in *BRCA2*, 16 in *PALB2*, 16 in the Lynch syndrome genes, eight in *TP53*, 14 in *ATM,* eight in *CHEK2*, six in *RAD51C*, nine in *RAD51*, two in *RAD50,* six in *BRIP1*, four in *PTEN*, two in *RAD51*, three in *APC*, two in *NBN*, eight in *MUTYH*, two in *CDH1* and one each in *BARD1*, *RET*, *MITF*, *EPCAM*, *FANCC*, *MRE11*, *POLD1*, *RB1, RECQL, MEN1, CDK4* and *PTCH2*.

Among the 321 pathogenic/likely pathogenic variant hits, 84 (26.2%) were point mutations in conserved splice sites, 73 (22.7%) were frameshift deletions, 65 (20.2%) were missense point mutations, 58 (18.1%) were nonsense point mutations, 22 (6.9%) were frameshift duplications, 10 (3.1%) were frameshift insertions, four (1.3%) were frameshift indels, four inframe deletions (1.3%), and a single inframe insertion (0.3%) (Fig. 1C).

Most of the 1252 VUS hits (corresponding to 886 unique variants) were identified in *ATM* (198 hits, 15.8%), followed by *BRCA2* (86, 6.9%), *MHS6* (63,

medRxiv preprint doi: https://doi.org/10.1101/2021.04.15.21255554; this version posted April 20, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

5.0%), *BRIP1* (55, 4.4%), *RAD50* (52, 4.2%), *CHEK2* (38, 3.0%), *PALB2* (34, 2.7%), *MHS2* (31, 2.5%), *MLH1* (17, 1.4%), *TP53* (17, 1.4%), *BRCA1* (17, 1.4%). Other genes amounted to 644 (51.4%) VUS (Fig. 1D).

Individuals with multiple variants (transheterozygotes)

Among the 305 individuals with pathogenic/likely pathogenic variants, 290 (95.1%) were single heterozygotes, 14 (4.6%) presented two variants in different genes (any two of these: *APC, ATM, BRCA1, BRCA2, CDK4, CHEK2, MEN1, MUTYH, PALB2, PMS2, RAD51, RAD51C* or *TP53*) and a single patient (0.3%) presented three pathogenic variants in different genes (*BRCA1*, *MSH6* and *MUTYH*). The most common combination in the patients with two variants was a variant in a high-penetrance gene, such as *BRCA1* and *BRCA2*, alongside a variant with lower penetrance, such as monoallelic *MUTYH* or in *CHEK2*. No patients presented more than one pathogenic/likely pathogenic variants in the same gene (Fig. 2, Table A.4). We observed a single individual *BRCA1*/*BRCA2* double heterozygote, making the double heterozygotes prevalence among *BRCA* variants carriers about 0.78% (1/129) and 0.06% (1/1682) overall. Thus, the overall transheterozygosity prevalence was about 0.89% (15/1682) in our sample.

Positivity profiles according to testing criteria of the USA National Comprehensive Cancer Network and Brazilian National Health Agency

The sampled individuals were assessed for hereditary cancer testing indication according to two sets of clinical criteria: the NCCN for hereditary

medRxiv preprint doi: https://doi.org/10.1101/2021.04.15.21255554; this version posted April 20, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

breast/ovarian, colorectal, stomach or pancreas cancer syndromes, version 3.2019 and 1.2020 and the criteria of the Brazilian National Health Agency (ANS).

Among the 1682 individuals in study, 1008 (59.9%) met NCCN 3.2019 and 1.2020 criteria for testing, 368 (21.9%) did not meet the criteria and 306 did not have sufficient information for classification. Forty-five patients among the 305 that presented pathogenic/likely pathogenic (positive patients) were in the last group, leaving 260 classifiable patients. The true positive rate of NCCN criteria was 215/260 (82.7%) and the false negative rate was 45/260 (17.3%). The F1-measure was 33.9%. Regarding ANS criteria, 660 (39.2%) met the criteria for testing, 691 did not meet and 331 did not have sufficient information for classification. The true positive rate of ANS was 145/260 (55.7%) and the false negative rate was 115/260 (44.2%), and F1-measure was 31.5% (Table 2).

The group of 215 patients fulfilling testing criteria according to the NCCN had the following genetic profiles: 72 (36.4%) individuals having variants in *BRCA1*, 29 (14.6%) in *BRCA2*, 16 (8.1%) in *PALB2*, 14 (7.1%) in *TP53* , 11 (5.6%) in *ATM*, 9 (4.5%) in *CHEK2*, 7 (3.5%) in *RAD50* and *RAD51*, 6 (3.5%) in *RAD51C*, 4 (2.0%) in *MSH6*, 3 (1.5%) in *BRIP1*, *MSH2* and *RAD51D*, 2 (1.0%) in *APC*, *BARD1* and *CDH1* e 8 (4.0%) in other genes (*MEN1, MITF, MLH1, NBN, PMS2, POLD1, PTEN* and *RECQL*).

The group of 45 patients who did not meet the NCCN criteria had the following profiles: 9 (20.0%) individuals with variants in *MUTYH,* 7 (16.7%) in *BRCA2*, 6 (14.3%) in *BRCA1*, 5 (11.9%) in *PALB2*, 2 (4.8%) in *ATM, CHEK2, MLH1, MSH6, PTEN, RET* and *TP53*, besides 10 (23.8%) in other genes

medRxiv preprint doi: https://doi.org/10.1101/2021.04.15.21255554; this version posted April 20, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

(*BARD1, BRIP1, CDK4, FANCC, MRE11A, MSH2, PMS2, PTCH2, RAD51* and *RAD51C*).

In summary, both NCCN 1.2020/3.2019 and ANS criteria failed to detect a substantial part of positive patients. The Brazilian criteria fared even worse, missing about 44% of positive patients, versus approximately 17% with NCCN 1.2020/3.2019.

Geographic distribution of variants

We stratified our data according to each Brazilian state. The results are available in the Table A.5.

Discussion

Cancers were the second most common cause of death worldwide in 2018, with 9.6 million deaths (1 every 6 deaths). Lung, prostate, colorectal, stomach and liver are the most common in men, whereas women are more affected by breast, colorectal, lung, cervix and thyroid [14]. The Brazilian National Cancer Institute (INCA) estimates 450,000 new cancer cases per year (excluding non-melanoma skin cancers) for the 2020-2022 triennium [15].

Most tumors are sporadic, caused by somatic genetic events. However, about 10% of the patients carry germline pathogenic variants in cancer predisposition genes such as *BRCA1* or *BRCA2* [1]. The identification of these patients is of utmost clinical importance because early diagnosis of hereditary cancer risk syndromes may improve vigilance and treatment. Therefore, genetic investigation is already a tool of modern oncology [16].

medRxiv preprint doi: https://doi.org/10.1101/2021.04.15.21255554; this version posted April 20, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

Next generation sequencing (NGS) technologies reduced sequencing costs allowing rapid, precise, and simultaneous analysis of a great quantity of genetic material. NGS has been used to identify genetic causes of rare diseases and cancers, in clinical and research settings. Previously, patients in investigation of HBOC or Lynch syndrome risk would be genotyped by limited, specific panels. Nowadays, as NGS is being largely utilized by clinical laboratories, the patients may undergo genotyping by multigenic panels that allow a better stratification as well as personalized oncologic management [17].

Here, we report the results of our study, the largest conducted in Brazil, with 1682 individuals that underwent genotyping with 27- to 78-genes panels intended to detect germinative variants in cancer predisposition genes. Women are the majority in our sample (92.6%). Consequently, the most prevalent cancers were breast and ovarian (57.7%), colorectal (4.2%) and thyroid/parathyroid (2.1%).

The overall positivity rate was 18.1% (305/1682 individuals); 16.0% (269/1682) if individuals with monoallelic *MUTYH* variants are excluded. Considering just *BRCA1/BRCA2* pathogenic/likely pathogenic variants, the positivity rate was only 7.7% (129/1682 individuals). This demonstrates the importance of NGS multigene panels: it reduces the rate of false negatives, providing more information for oncologic management and prognosis.

The *BRCA* pathogenic variants positivity rate from our study seems low, but it is not unexpected since our cohort was not specifically selected for any sex, age, ethnicity, or history of cancer. Indeed, since there is significant allelic heterogeneity in *BRCA* genes, there is great variability in positivity rates in other studies [18]. In Brazil, studies of HBOC patients observed positivity rates between

medRxiv preprint doi: https://doi.org/10.1101/2021.04.15.21255554; this version posted April 20, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

1.3% [19] and 27.3% [20]. In other countries, the positivity rate among sporadic and high-risk hereditary cancer cases varied between 2.6% in the USA [21] and 27.9% in Japan [22] (reviewed in [23]). Therefore, our positivity rate is indeed within these ranges.

We compared the mutational profile of *BRCA1* and *BRCA2* genes we observed with other 20 Brazilian studies. Among the 56 variants we detected, 33 (19 in *BRCA1* and 14 in *BRCA2*) were recurrent in other studies. The five most recurrent variants were: *BRCA1*:c.5266dupC, found by other 18 studies [18, 20, 24-39], *BRCA1*:c.3331_3334delCAAG, found in other 12 studies [18, 20, 24, 27, 29, 31, 32, 34, 35, 37, 39, 40], *BRCA2*:c.2808_2811delACAA found in other seven studies [18, 29-31, 34, 35, 39], *BRCA1*: c.1687C>T found in six studies [26, 31, 32, 34, 35] and *BRCA1*:c.211A>G found in other five studies [35-38, 40].

The five most prevalent *BRCA1* mutations among the individuals sampled by our study were c.5266dupC, c.5074+2T>C, c.3331_3334delCAAG, c.1687C>T and c.211A>G. These results partially agree with a worldwide survey that included Brazilian families of women carriers of *BRCA* pathogenic variants. Rebbeck et al. (2018) [41] surveyed 101 Brazilian families and observed c.5266dupC, c.3331_3334delCAAG and c.1687C>T among the five most prevalent *BRCA1* mutations. Therefore, our results agree with three variants.

The five most prevalent *BRCA2* variants in our patients were c.2808_2811delACAA, c.9382C>T, c.6024dupG, c.8488-1G>A and c.5216dupA. Only c.2808_2811delACAA was in common with the Rebbeck et al. results [41]. Interestingly, they observed c.156_157insAlu among the five most prevalent variation among 49 Brazilian families, but our results place it in sixth place, tied with other variants.

medRxiv preprint doi: https://doi.org/10.1101/2021.04.15.21255554; this version posted April 20, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

The recurrence in independent samples from Brazil indicate the variants are representative of the *BRCA1* and *BRCA2* mutational spectrum in Brazil. Indeed, the *BRCA1*:c.5266dupC and *BRCA2*:c.156_157insAlu (found in other four studies [31, 34, 35, 42]) variants are considered as founder mutations coming from European populations and may be found in the individuals from all regions of Brazil [35]. Indeed, we detected *BRCA1:* c.5266dupC in individuals from Pernambuco and Alagoas, Northeast Brazil states, as well as in São Paulo (Southeast) and Paraná (South) and *BRCA2*:c.156_157insAlu was detected in an individual from Piauí, also a Northeast state.

We also observed some recurrent pathogenic variants in other genes: *TP53*:c.1010G>A, found in four other studies [20, 38-40], *CHEK2*:c.349A>G, found in two other studies [38, 43]; *MUTYH*:c.1147delC [43]; *MUTYH*:c.1187G>A [38], *MSH2*:c.2152C>T [44]; *MSH6*:c.3848_3862del [45], *MLH1*:c.677G>A [44] and *PALB2*:c.2257C>T [46]. These observations could mean that these variants are important co-players alongside the recurrent *BRCA* variants in the risk of hereditary cancers in Brazilian populations.

Transheterozygosity, i.e. heterozygosity at two different loci [47], is rare among patients at risk of hereditary cancers. The prevalence of double heterozygosity among *BRCA1/BRCA2* carriers ranges between 1.8% and 1.85% [48, 49] in the Jewish Ashkenazi populations. Among non-Ashkenazi Europeans, the prevalence is lower, between 0.22% and 0.87% [48]. An Italian study reported 0.62% [50] and a Korean study reported 1.2% [51]. Therefore, our estimate of 0.78% in the Brazilian sample seems plausible.

There has been controversy regarding the consequences over the phenotype of *BRCA1*/*BRCA2* double heterozygote. Earlier reports suggest that

medRxiv preprint doi: https://doi.org/10.1101/2021.04.15.21255554; this version posted April 20, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

they do not have worse phenotypes when comparing to single-variant carriers [52, 53]. Bell et al. (2002) [54] analyzed three independent samples of tumor tissue from the same *BRCA1/BRCA2* double heterozygote patient and observed that either one or other gene had suffered loss of heterozygosity (LOH) genetic events, suggesting that both genes are functionally equivalent in initiating tumorigenesis in breast tissue. Therefore, double heterozygote status would be equivalent to single-carrier status. Other case reports, case series and review of case reports also suggested that double heterozygote status was not associated with more severe phenotype [48, 55, 56]. In contrast, Randall et al. (1998) [57] observed LOH in both genes in ovarian tumors, perhaps signaling that the pathogenic events in each tissue are different and not always *BRCA1* mutations are equivalent to *BRCA2* counterparts.

However, more recent large surveys have been suggesting that double heterozygote have different phenotypes when compared to single heterozygote individuals. A large survey of 8162 German families with breast/ovarian cancer cases found eight double heterozygote individuals. They observed that they were younger at the onset of their first cancer episode when compared with their single heterozygote relatives and had more severe disease [58]. A worldwide survey of 32,295 women compared the clinical history of double heterozygote individuals with breast cancer to their single heterozygote counterparts. The same was done for ovarian cancer cases. They observed that clinical characteristics of double heterozygote individuals resemble phenotypes from *BRCA1*-only single heterozygotes, but tumor LOH microsatellite markers patterns are intermediate to the patterns of *BRCA1*-only single heterozygote and *BRCA2*-only single heterozygote. Thus, the authors suggest that the joint

medRxiv preprint doi: https://doi.org/10.1101/2021.04.15.21255554; this version posted April 20, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

effect of *BRCA1*/*BRCA2* variants are compatible with an additive model [47]. Another recent study also suggests that the double heterozygote phenotype more closely resembled that of *BRCA1* mutation carriers with poor prognosis factors and proposed a co-dominant model for the joint effects [59].

In summary, *BRCA1/BRCA2* double heterozygote may have early onset of disease and the phenotype is perhaps similar to the "severe end of spectrum of *BRCA1* mutation carriership" [48]. We found a single *BRCA1/BRCA2* double heterozygote woman in our sample. She presented personal history of breast cancer. Since she is in her late 40s, her first cancer case must have happened much earlier, however, detailed data on her phenotype is currently lacking.

Literature on *BRCA*/other genes double heterozygotes is sparser. Thus, comparison with our results was difficult due to the possible different combinations of variants. For example, Sokolenko et al. (2014) [60] found seven digenic combinations double heterozygotes among *BRCA1* and other DNA double-strand repair genes (*BRCA1/CHEK2, BRCA1/ATM, BRCA1/BLM, CHEK2/BLM, CHEK2/ATM, NBS1/ATM, and NBS1/BLM*) in Russian patients: none were observed in our study. They did not observe differences from single heterozygote individuals. In contrast, a case series including a German *BRCA1/PALB2* double heterozygous patient had no early onset but had severe disease (multifocal triple negative ductal carcinoma) [61]. Another case report presented a case of a double heterozygote *APC/MLH1* man with Kashmir/Egyptian ancestry, which had a history of six jejunal cancers [62]. Therefore, double heterozygosity may have unusual effects on cancer phenotypes. However, we did not find *APC/MLH1* double heterozygous, but

medRxiv preprint doi: https://doi.org/10.1101/2021.04.15.21255554; this version posted April 20, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

found an individual *APC/TP53* double heterozygote, so it is another case warranting further investigation.

Perhaps the joint effects of non-*BRCA* variants can be explained in an additive manner as well. Thus, some combinations will result in severe phenotypes while others will not, due to the specific penetrance of each variant. It seems that non-*BRCA* double heterozygotes will need to be analyzed on a case-by-case basis.

We found that NCCN 1.2020 criteria missed a substantial proportion of patients that had pathogenic/likely pathogenic variants. Other authors observed the same. We found that NCCN criteria missed a substantial proportion of patients that had pathogenic/likely pathogenic variants. Other authors observed the same. For example, Grindedal et al. (2017) [63] investigated *BRCA* mutations in a Norwegian breast cancer cohort and assessed some testing criteria, including NCCN, and found a false negative rate of 15.8%. Yang et al. (2018) [64] investigated 4196 patients genotyped with 40- to 80- genes panels and showed a false negative rate of 13.5% with NCCN criteria. Both were not much far from our estimate of 17% with NCCN 1.2020/3.2019.

Other authors proposed changes to NCCN criteria. Alemar et al. (2017) [34] found that adding criteria that are not included in the NCCN and ANS criteria (e.g. some ASCO criteria [65, 66]) achieved a higher predictive value, while other authors compared other four algorithms (BOADICEA [67, 68], BRCAPRO, Myriad [69] and Manchester score [70]) and observed that the pedigree-based BOADICEA most accurately predicted *BRCA1/BRCA2* variant carrier status in a Southeastern Brazilian population [71]. Although NCCN criteria were imperfect, ANS criteria fared worse and a reformulation is warranted.

medRxiv preprint doi: https://doi.org/10.1101/2021.04.15.21255554; this version posted April 20, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

Recently, a panel of Brazilian experts proposed recommendations for improving testing criteria for HBOC risk patients in Brazil. Besides modifying testing criteria, the expert panel also recommended offering risk-reducing surgeries for positive patients. For negative patients, investigating both maternal and paternal lineages is warranted, so the result of models estimating cancer risk can be communicated to the patient. They also suggested that VUS should always be reported and periodically reassessed, but no urgent clinical action is justified since most VUS are constantly reclassified to benign/likely benign categories. Furthermore, patients should be contacted whenever any update in testing protocols or management options should appear [72].

Our study had some limitations. The NGS panel detects small deletions and duplications up to 17 base-pairs, but large deletions and duplications are not detected by this methodology. Other structural chromosomal changes, such as inversions and translocations are not detected either. If some of these changes are suspected, we recommend using methodologies like array CGH, MLPA, qPCR or FISH to confirm the variant found. Expansion variants of trinucleotide repeats, deep intronic variants or regulatory regions such as promoters are not detectable in the present test. Epigenetic changes are also not detectable by this test. At least for *BRCA* genes though, large rearrangements seem to be uncommon in Brazilian populations [35, 42]. Moreover, further phenotype information was missing for several patients, precluding further analyses.

medRxiv preprint doi: https://doi.org/10.1101/2021.04.15.21255554; this version posted April 20, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

Conclusion

In conclusion, we genotyped 1682 Brazilian individuals with clinical criteria of hereditary cancer syndrome from all regions of the country with NGS multigenic panels suited for the detection of germline pathogenic variants associated with cancer susceptibility, making it the largest Brazilian study of this nature to date. We observed several *BRCA1* and *BRCA2* recurrent mutations, confirming their presence in the Brazilian mutational spectrum and generated data for other 30 genes and 110 variants with modest penetrance. We also estimated the prevalence of *BRCA* as well as non-*BRCA* double heterozygotes in the Brazilian population. More studies are necessary to discover the implication of transheterozygosity over the phenotype of affected individuals.

Appendix

Table A.1. Genotyped genes.
Table A.2. Ancestry, genetic findings summary, testing criteria, personal and family history.
Table A.3. Genetic Findings.
Table A.4. Positivity rate per gene or combination of genes, alongside number of unique pathogenic/likely pathogenic variants detected in each gene. The variants are detailed for transheterozygotes.
Table A.5. Patients per Brazilian state.

References

1. Samadder NJ, Giridhar KV, Baffy N, Riegert-Johnson D, Couch FJ. Hereditary Cancer Syndromes-A Primer on Diagnosis and Management: Part 1: Breast-Ovarian Cancer Syndromes. *Mayo Clin Proc*. Jun 2019;94(6):1084-1098. doi:10.1016/j.mayocp.2019.02.017
2. Nagy R, Sweet K, Eng C. Highly penetrant hereditary cancer syndromes. *Oncogene*. 2004;23(38):6445-6470. doi:10.1038/sj.onc.1207714

medRxiv preprint doi: https://doi.org/10.1101/2021.04.15.21255554; this version posted April 20, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

3. Zelli V, Compagnoni C, Cannita K, et al. Applications of Next Generation Sequencing to the Analysis of Familial Breast/Ovarian Cancer. *High Throughput*. Jan 10 2020;9(1)doi:10.3390/ht9010001

4. Xuan J, Yu Y, Qing T, Guo L, Shi L. Next-generation sequencing in the clinic: Promises and challenges. *Cancer Letters*. 2013;340(2):284-295. doi:10.1016/j.canlet.2012.11.025

5. Yohe S, Thyagarajan B. Review of Clinical Next-Generation Sequencing. *Archives of Pathology & Laboratory Medicine*. 2017;141(11):1544-1557. doi:10.5858/arpa.2016-0501-ra

6. Machado PM, Brandao RD, Cavaco BM, et al. Screening for a BRCA2 rearrangement in high-risk breast/ovarian cancer families: evidence for a founder effect and analysis of the associated phenotypes. *J Clin Oncol*. May 20 2007;25(15):2027-34. doi:10.1200/JCO.2006.06.9443

7. Richards S, Aziz N, Bale S, et al. Standards and guidelines for the interpretation of sequence variants: a joint consensus recommendation of the American College of Medical Genetics and Genomics and the Association for Molecular Pathology. *Genet Med*. May 2015;17(5):405-24. doi:10.1038/gim.2015.30

8. Biesecker LG, Harrison SM, ClinGen Sequence Variant Interpretation Working G. The ACMG/AMP reputable source criteria for the interpretation of sequence variants. *Genet Med*. Dec 2018;20(12):1687-1688. doi:10.1038/gim.2018.42

9. Karczewski KJ, Francioli LC, Tiao G, et al. The mutational constraint spectrum quantified from variation in 141,456 humans. *Nature*. 2020/05/01 2020;581(7809):434-443. doi:10.1038/s41586-020-2308-7

10. Karczewski KJ, Weisburd B, Thomas B, et al. The ExAC browser: displaying reference data information from over 60 000 exomes. *Nucleic Acids Research*. 2017;45(D1):D840-D845. doi:10.1093/nar/gkw971

11. A global reference for human genetic variation. *Nature*. 2015;526(7571):68-74. doi:10.1038/nature15393

12. Landrum MJ, Lee JM, Benson M, et al. ClinVar: improving access to variant interpretations and supporting evidence. *Nucleic Acids Research*. 2018;46(D1):D1062-D1067. doi:10.1093/nar/gkx1153

13. Fokkema IFAC, Taschner PEM, Schaafsma GCP, Celli J, Laros JFJ, Den Dunnen JT. LOVD v.2.0: the next generation in gene variant databases. *Human Mutation*. 2011;32(5):557-563. doi:10.1002/humu.21438

14. Organization WH. WHO report on cancer: setting priorities, investing wisely and providing care for all. 2020.

15. Instituto Nacional de Câncer José Alencar Gomes da Silva. Ministério da Saúde. Estimativa 2020: incidência de câncer no Brasil. Rio de Janeiro: Instituto Nacional de Câncer José Alencar Gomes da Silva; 2019. p. 120.

16. Yoshida R. Hereditary breast and ovarian cancer (HBOC): review of its molecular characteristics, screening, treatment, and prognosis. *Breast Cancer*. Aug 29 2020;doi:10.1007/s12282-020-01148-2

17. Suwinski P, Ong C, Ling MHT, Poh YM, Khan AM, Ong HS. Advancing Personalized Medicine Through the Application of Whole Exome Sequencing and Big Data Analytics. *Front Genet*. 2019;10:49. doi:10.3389/fgene.2019.00049

18. Alemar B, Herzog J, Brinckmann Oliveira Netto C, et al. Prevalence of Hispanic BRCA1 and BRCA2 mutations among hereditary breast and ovarian cancer patients from Brazil reveals differences among Latin American

medRxiv preprint doi: https://doi.org/10.1101/2021.04.15.21255554; this version posted April 20, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

populations. *Cancer Genet*. Sep 2016;209(9):417-422. doi:10.1016/j.cancergen.2016.06.008

19. Dillenburg CV, Bandeira IC, Tubino TV, et al. Prevalence of 185delAG and 5382insC mutations in BRCA1, and 6174delT in BRCA2 in women of Ashkenazi Jewish origin in southern Brazil. *Genet Mol Biol*. Jul 2012;35(3):599-602. doi:10.1590/S1415-47572012000400009

20. Cipriano NM, Jr., de Brito AM, de Oliveira ES, et al. Mutation screening of TP53, CHEK2 and BRCA genes in patients at high risk for hereditary breast and ovarian cancer (HBOC) in Brazil. *Breast Cancer*. May 2019;26(3):397-405. doi:10.1007/s12282-018-00938-z

21. Susswein LR, Marshall ML, Nusbaum R, et al. Pathogenic and likely pathogenic variant prevalence among the first 10,000 patients referred for next-generation cancer panel testing. *Genet Med*. Aug 2016;18(8):823-32. doi:10.1038/gim.2015.166

22. Sugano K, Nakamura S, Ando J, et al. Cross-sectional analysis of germline BRCA1 and BRCA2 mutations in Japanese patients suspected to have hereditary breast/ovarian cancer. *Cancer Sci*. Oct 2008;99(10):1967-76. doi:10.1111/j.1349-7006.2008.00944.x

23. Okano M, Nomizu T, Tachibana K, et al. The relationship between BRCA-associated breast cancer and age factors: an analysis of the Japanese HBOC consortium database. *J Hum Genet*. Mar 2021;66(3):307-314. doi:10.1038/s10038-020-00849-y

24. Lourenço JJ, Vargas FR, Bines J, et al. BRCA1 mutations in Brazilian patients. *Genetics and Molecular Biology*. 2004;27:500-504.

25. Dufloth RM, Carvalho S, Heinrich JK, et al. Analysis of BRCA1 and BRCA2 mutations in Brazilian breast cancer patients with positive family history. *Sao Paulo Med J*. Jul 7 2005;123(4):192-7. doi:10.1590/s1516-31802005000400007

26. Gomes MC, Costa MM, Borojevic R, et al. Prevalence of BRCA1 and BRCA2 mutations in breast cancer patients from Brazil. *Breast Cancer Res Treat*. Jul 2007;103(3):349-53. doi:10.1007/s10549-006-9378-6

27. Esteves VF, Thuler LC, Amendola LC, et al. Prevalence of BRCA1 and BRCA2 gene mutations in families with medium and high risk of breast and ovarian cancer in Brazil. *Braz J Med Biol Res*. May 2009;42(5):453-7. doi:10.1590/s0100-879x2009000500009

28. Ewald IP, Izetti P, Vargas FR, et al. Prevalence of the BRCA1 founder mutation c.5266dupin Brazilian individuals at-risk for the hereditary breast and ovarian cancer syndrome. *Hered Cancer Clin Pract*. Dec 20 2011;9(1):12. doi:10.1186/1897-4287-9-12

29. Carraro DM, Koike Folgueira MA, Garcia Lisboa BC, et al. Comprehensive analysis of BRCA1, BRCA2 and TP53 germline mutation and tumor characterization: a portrait of early-onset breast cancer in Brazil. *PLoS One*. 2013;8(3):e57581. doi:10.1371/journal.pone.0057581

30. Silva FC, Lisboa BC, Figueiredo MC, et al. Hereditary breast and ovarian cancer: assessment of point mutations and copy number variations in Brazilian patients. *BMC Med Genet*. May 15 2014;15:55. doi:10.1186/1471-2350-15-55

31. Fernandes GC, Michelli RA, Galvão HC, et al. Prevalence of BRCA1/BRCA2 mutations in a Brazilian population sample at-risk for hereditary breast cancer and characterization of its genetic ancestry. *Oncotarget*. Dec 6 2016;7(49):80465-80481. doi:10.18632/oncotarget.12610

medRxiv preprint doi: https://doi.org/10.1101/2021.04.15.21255554; this version posted April 20, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

32. Maistro S, Teixeira N, Encinas G, et al. Germline mutations in BRCA1 and BRCA2 in epithelial ovarian cancer patients in Brazil. *BMC cancer*. Dec 3 2016;16(1):934. doi:10.1186/s12885-016-2966-x

33. de Oliveira ES, Soares BL, Lemos S, et al. Screening of the BRCA1 gene in Brazilian patients with breast and/or ovarian cancer via high-resolution melting reaction analysis. Article. *Fam Cancer*. Apr 2016;15(2):173-181. doi:10.1007/s10689-015-9858-0

34. Alemar B, Gregório C, Herzog J, et al. BRCA1 and BRCA2 mutational profile and prevalence in hereditary breast and ovarian cancer (HBOC) probands from Southern Brazil: Are international testing criteria appropriate for this specific population? *PLOS ONE*. 2017;12(11):e0187630. doi:10.1371/journal.pone.0187630

35. Palmero EI, Carraro DM, Alemar B, et al. The germline mutational landscape of BRCA1 and BRCA2 in Brazil. *Sci Rep*. Jun 15 2018;8(1):9188. doi:10.1038/s41598-018-27315-2

36. Brianese RC, Nakamura KDM, Almeida F, et al. BRCA1 deficiency is a recurrent event in early-onset triple-negative breast cancer: a comprehensive analysis of germline mutations and somatic promoter methylation. *Breast Cancer Res Treat*. Feb 2018;167(3):803-814. doi:10.1007/s10549-017-4552-6

37. Cotrim DP, Ribeiro ARG, Paixão D, et al. Prevalence of BRCA1 and BRCA2 pathogenic and likely pathogenic variants in non-selected ovarian carcinoma patients in Brazil. *BMC cancer*. Jan 3 2019;19(1):4. doi:10.1186/s12885-018-5235-3

38. Bandeira G, Rocha K, Lazar M, et al. Germline variants of Brazilian women with breast cancer and detection of a novel pathogenic ATM deletion in early-onset breast cancer. *Breast Cancer*. Sep 28 2020;doi:10.1007/s12282-020-01165-1

39. Da Costa E Silva Carvalho S, Cury NM, Brotto DB, et al. Germline variants in DNA repair genes associated with hereditary breast and ovarian cancer syndrome: analysis of a 21 gene panel in the Brazilian population. *BMC Medical Genomics*. 2020;13(1)doi:10.1186/s12920-019-0652-y

40. Felix GE, Abe-Sandes C, Machado-Lopes TM, et al. Germline mutations in BRCA1, BRCA2, CHEK2 and TP53 in patients at high-risk for HBOC: characterizing a Northeast Brazilian Population. *Hum Genome Var*. 2014;1:14012. doi:10.1038/hgv.2014.12

41. Rebbeck TR, Friebel TM, Friedman E, et al. Mutational spectrum in a worldwide study of 29,700 families with BRCA1 or BRCA2 mutations. *Hum Mutat*. May 2018;39(5):593-620. doi:10.1002/humu.23406

42. Ewald IP, Cossio SL, Palmero EI, et al. BRCA1 and BRCA2 rearrangements in Brazilian individuals with Hereditary Breast and Ovarian Cancer Syndrome. *Genet Mol Biol*. Apr-Jun 2016;39(2):223-31. doi:10.1590/1678-4685-gmb-2014-0350

43. Felicio PS, Grasel RS, Campacci N, et al. Whole-exome sequencing of non-BRCA1/BRCA2 mutation carrier cases at high-risk for hereditary breast/ovarian cancer. *Hum Mutat*. Dec 16 2020;doi:10.1002/humu.24158

44. Soares BL, Brant AC, Gomes R, et al. Screening for germline mutations in mismatch repair genes in patients with Lynch syndrome by next generation sequencing. *Fam Cancer*. Jul 2018;17(3):387-394. doi:10.1007/s10689-017-0043-5

medRxiv preprint doi: https://doi.org/10.1101/2021.04.15.21255554; this version posted April 20, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

45. Rosa RCA, Santis JO, Teixeira LA, et al. Lynch syndrome identification in a Brazilian cohort of endometrial cancer screened by a universal approach. *Gynecol Oncol*. Oct 2020;159(1):229-238. doi:10.1016/j.ygyno.2020.07.013

46. Gomes R, Spinola PDS, Brant AC, et al. Prevalence of germline variants in consensus moderate-to-high-risk predisposition genes to hereditary breast and ovarian cancer in BRCA1/2-negative Brazilian patients. *Breast Cancer Res Treat*. Oct 30 2020;doi:10.1007/s10549-020-05985-9

47. Rebbeck TR, Friebel TM, Mitra N, et al. Inheritance of deleterious mutations at both BRCA1 and BRCA2 in an international sample of 32,295 women. *Breast Cancer Research*. Nov 11 2016;18112. doi:10.1186/s13058-016-0768-3

48. Leegte B, van der Hout AH, Deffenbaugh AM, et al. Phenotypic expression of double heterozygosity for BRCA1 and BRCA2 germline mutations. *J Med Genet*. Mar 2005;42(3):e20. doi:10.1136/jmg.2004.027243

49. Lavie O, Narod S, Lejbkowicz F, et al. Double heterozygosity in the BRCA1 and BRCA2 genes in the Jewish population. *Ann Oncol*. Apr 2011;22(4):964-966. doi:10.1093/annonc/mdq460

50. Zuradelli M, Peissel B, Manoukian S, et al. Four new cases of double heterozygosity for BRCA1 and BRCA2 gene mutations: clinical, pathological, and family characteristics. *Breast Cancer Res Treat*. Nov 2010;124(1):251-8. doi:10.1007/s10549-010-0853-8

51. Noh JM, Choi DH, Nam SJ, et al. Characteristics of double heterozygosity for BRCA1 and BRCA2 germline mutations in Korean breast cancer patients. *Breast Cancer Res Treat*. Jan 2012;131(1):217-22. doi:10.1007/s10549-011-1718-5

52. Friedman E, Bar-Sade Bruchim R, Kruglikova A, et al. Double heterozygotes for the Ashkenazi founder mutations in BRCA1 and BRCA2 genes. *Am J Hum Genet*. Oct 1998;63(4):1224-7. doi:10.1086/302040

53. Moslehi R, Russo D, Phelan C, Jack E, Antman K, Narod S. An unaffected individual from a breast/ovarian cancer family with germline mutations in both BRCA1 and BRCA2. *Clin Genet*. Jan 2000;57(1):70-3. doi:10.1034/j.1399-0004.2000.570111.x

54. Bell DW, Erban J, Sgroi DC, Haber DA. Selective loss of heterozygosity in multiple breast cancers from a carrier of mutations in both BRCA1 and BRCA2. *Cancer Res*. May 15 2002;62(10):2741-3.

55. Thiffault I, Hamel N, Pal T, et al. Germline truncating mutations in both MSH2 and BRCA2 in a single kindred. *Br J Cancer*. Jan 26 2004;90(2):483-91. doi:10.1038/sj.bjc.6601424

56. Musolino A, Naldi N, Michiara M, et al. A breast cancer patient from Italy with germline mutations in both the BRCA1 and BRCA2 genes. *Breast Cancer Res Treat*. May 2005;91(2):203-5. doi:10.1007/s10549-004-7704-4

57. Randall TC, Bell KA, Rebane BA, Rubin SC, Boyd J. Germline mutations of the BRCA1 and BRCA2 genes in a breast and ovarian cancer patient. *Gynecol Oncol*. Sep 1998;70(3):432-4. doi:10.1006/gyno.1998.5081

58. Heidemann S, Fischer C, Engel C, et al. Double heterozygosity for mutations in BRCA1 and BRCA2 in German breast cancer patients: implications on test strategies and clinical management. *Breast Cancer Research and Treatment*. Aug 2012;134(3):1229-1239. doi:10.1007/s10549-012-2050-4

59. Le Page C, Rahimi K, Rodrigues M, et al. Clinicopathological features of women with epithelial ovarian cancer and double heterozygosity for BRCA1 and

medRxiv preprint doi: https://doi.org/10.1101/2021.04.15.21255554; this version posted April 20, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

BRCA2: A systematic review and case report analysis. *Gynecol Oncol*. Feb 2020;156(2):377-386. doi:10.1016/j.ygyno.2019.11.019

60. Sokolenko AP, Bogdanova N, Kluzniak W, et al. Double heterozygotes among breast cancer patients analyzed for BRCA1, CHEK2, ATM, NBN/NBS1, and BLM germ-line mutations. *Breast Cancer Res Treat*. Jun 2014;145(2):553-62. doi:10.1007/s10549-014-2971-1

61. Pern F, Bogdanova N, Schürmann P, et al. Mutation analysis of BRCA1, BRCA2, PALB2 and BRD7 in a hospital-based series of German patients with triple-negative breast cancer. *PLoS One*. 2012;7(10):e47993. doi:10.1371/journal.pone.0047993

62. Lindor NM, Smyrk TC, Buehler S, et al. Multiple jejunal cancers resulting from combination of germline APC and MLH1 mutations. *Fam Cancer*. Dec 2012;11(4):667-9. doi:10.1007/s10689-012-9561-3

63. Grindedal EM, Heramb C, Karsrud I, et al. Current guidelines for BRCA testing of breast cancer patients are insufficient to detect all mutation carriers. *BMC cancer*. 2017;17(1):438-438. doi:10.1186/s12885-017-3422-2

64. Yang S, Axilbund JE, O'Leary E, et al. Underdiagnosis of Hereditary Breast and Ovarian Cancer in Medicare Patients: Genetic Testing Criteria Miss the Mark. *Ann Surg Oncol*. Oct 2018;25(10):2925-2931. doi:10.1245/s10434-018-6621-4

65. Statement of the American Society of Clinical Oncology: genetic testing for cancer susceptibility, Adopted on February 20, 1996. *J Clin Oncol*. May 1996;14(5):1730-6; discussion 1737-40. doi:10.1200/JCO.1996.14.5.1730

66. Lu KH, Wood ME, Daniels M, et al. American Society of Clinical Oncology Expert Statement: collection and use of a cancer family history for oncology providers. *J Clin Oncol*. Mar 10 2014;32(8):833-40. doi:10.1200/JCO.2013.50.9257

67. Antoniou AC, Pharoah PD, McMullan G, et al. A comprehensive model for familial breast cancer incorporating BRCA1, BRCA2 and other genes. *Br J Cancer*. Jan 7 2002;86(1):76-83. doi:10.1038/sj.bjc.6600008

68. Antoniou AC, Pharoah PP, Smith P, Easton DF. The BOADICEA model of genetic susceptibility to breast and ovarian cancer. *Br J Cancer*. Oct 18 2004;91(8):1580-90. doi:10.1038/sj.bjc.6602175

69. Parmigiani G, Berry D, Aguilar O. Determining carrier probabilities for breast cancer-susceptibility genes BRCA1 and BRCA2. *Am J Hum Genet*. Jan 1998;62(1):145-58. doi:10.1086/301670

70. Evans DG, Eccles DM, Rahman N, et al. A new scoring system for the chances of identifying a BRCA1/2 mutation outperforms existing models including BRCAPRO. *J Med Genet*. Jun 2004;41(6):474-80. doi:10.1136/jmg.2003.017996

71. Teixeira N, Maistro S, Del Pilar Estevez Diz M, et al. Predictability of BRCA1/2 mutation status in patients with ovarian cancer: How to select women for genetic testing in middle-income countries. *Maturitas*. Nov 2017;105:113-118. doi:10.1016/j.maturitas.2017.06.002

72. Achatz MI, Caleffi M, Guindalini R, Marques RM, Nogueira-Rodrigues A, Ashton-Prolla P. Recommendations for Advancing the Diagnosis and Management of Hereditary Breast and Ovarian Cancer in Brazil. *JCO Glob Oncol*. 2020;6:439-452. doi:10.1200/JGO.19.00170

medRxiv preprint doi: https://doi.org/10.1101/2021.04.15.21255554; this version posted April 20, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

Table 1. Demographic and clinical characteristics of Brazilian individuals tested genotyped with NGS hereditary cancer panel.

Demographic	**N**	**%**
Sex	1682	100
Female	1557	92.6
Male	125	7.4
Age at diagnosis (years)		
<30	79	4.7
30-40	452	26.9
41-50	526	31.3
51-65	465	27.6
>65	160	9.5
Ethnicity		
Multiple ethnicities (admixed)	330	19.6
White	259	15.4
Black	26	1.5
Jewish Ashkenazi	22	1.3
Native indigenous	21	1.2
Asian	17	1.0
No information	1007	59.9
Personal cancer history [a]		
Overall	1126	66.9
Breast or ovarian	971	57.7
Colorectal	71	4.2
Thyroid/parathyroid	36	2.1
Prostate	15	0.9
Stomach	14	0.8
Uterus	13	0.8
Kidney	13	0.8
Sarcomas	13	0.8
Leukemia	11	0.6
Skin (non-melanoma)	9	0.5
Pancreas	9	0.5
Other	17	1.0
No cancer history	268	15.9
No information	293	17.4

medRxiv preprint doi: https://doi.org/10.1101/2021.04.15.21255554; this version posted April 20, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

Family cancer history (first-, second- or third-degree relatives) [a]		
Overall	1182	70.3
Breast or ovarian	973	57.8
Colorectal	310	18.4
Thyroid/parathyroid	108	6.4
Prostate	339	20.2
Stomach	155	9.2
Uterus	122	7.3
Kidney	45	2.7
Sarcomas	22	1.3
Leukemia	139	8.3
Skin (non-melanoma)	72	4.3
Pancreas	83	4.9
Other	235	14.0
No cancer history	163	9.7
No information	337	20.0

[a] Number of individuals with personal history exceeds 1682 because some individuals reported multiple cancer types.

medRxiv preprint doi: https://doi.org/10.1101/2021.04.15.21255554; this version posted April 20, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

Table 2. NCCN 3.2019/1.2020 and Brazilian ANS testing criteria performance.

Decision	NGS testing		***Row totals***
	Positive patients	VUS/Negative patients	
NCCN 3.2019/1.2020			
At risk	215	793	***1008***
No risk	45	323	***368***
Column totals	***260***	***1116***	***1376***
Insufficient information: 306			
Brazilian ANS			
At risk	145	515	***660***
No risk	115	576	***691***
Column totals	***260***	***1091***	***1351***
Insufficient information: 331			

NCCN: sensitivity = 82.7%, specificity = 28.9%, false negative rate = 17.3%, F1-measure = 33.9%.

ANS: sensitivity = 55.8%, specificity = 52.8%, false negative rate = 44.2% F1-measure = 31.5%.

medRxiv preprint doi: https://doi.org/10.1101/2021.04.15.21255554; this version posted April 20, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

Figure captions

Figure 1. Variants profiles. (A) Distribution of patients according to genetic findings. P/LP = patients with pathogenic/likely pathogenic, VUS = patients with variants of uncertain significance (N = 1682). (B) Distribution of P/LP variants per gene. (C) P/LP variants functional annotation. (D) Distribution of VUS per gene. Lynch syndrome genes have been grouped together in (C) and (D).

Figure 2. Heatmap representing the distribution of pathogenic/likely pathogenic (P/LP) transheterozygotes in the sample. The top line represents the number of patients in each column. The single heterozygotes (n=290) are distributed in the main diagonal. Transheterozygotes (n=15) are distributed in the inferior half (14 double-heterozygotes and 1 triple-heterozygote). The total number of variants are 321 ($290 \times 1 + 14 \times 2 + 1 \times 3 = 321$).


A
P/LP
(18.1%)
Negative
(37.1%)
VUS
(44.8%)
B
BRCA1 26.2%
BRCA2 14.3%
MUTYH 13.1%
PALB2 7.8%
MLH1/MSH2/MSH6/PMS2 5.3%
TP53 5.3%
ATM 4.7%
CHEK2 4.4%
RAD51C 2.8%
RAD51 2.5%
RAD50 2.2%
BRIP1 1.9%
BARD1 1.2%
PTEN 1.2%
RAD51D 1.2%
Other genes 5.9%
0 25 50 75 100
N=321 P/LP variants
C
Frameshift indel
(1.25%)
Inframe deletion
(1.25%)
Frameshift insertion
(3.12%)
Inframe insertion
(0.31%)
Frameshift duplication
(6.85%)
Splice site SNV
(26.17%)
Nonsense
(18.07%)
Frameshift deletion
(22.74%)
Nonsynonymous SNV
(20.25%)
D
ATM 15.8%
MLH1/MSH2/MSH6/PMS2 9.7%
APC 7.2%
BRCA2 6.9%
POLE 5.9%
POLD1 4.8%
BRIP1 4.4%
RAD50 4.2%
RET 3.8%
CHEK2 3.0%
MUTYH 3.0%
PALB2 2.7%
NBN 2.5%
CDH1 2.2%
BARD1 1.8%
Other genes 22.3%
0 100 200 300 400
N=1252 VUS


Total 305 patients, 321 P/LP patients
85 45 40 25 16 14 11 8 7 7 6 6 5 4 4 4 2 3 2 1 2 2 0 1 1 1 1 1 1 1 0 1
BRCA1 78
BRCA2 1 40
MUTYH 3 37
PALB2 24
RAD51 1 16
TP53 13
ATM 1 1 11
CHEK2 3 8
RAD51C 1 7
RAD50 7
BRIP1 6
MSH2 6
MSH6 1 1 5
BARD1 4
PTEN 4
RAD51D 4
APC 1 2
MLH1 3
CDH1 2
PMS2 1 1
NBN 2
RET 2
CDK4 1
EPCAM 1
POLD1 1
PTCH2 1
RB1 1
RECQL 1
FANCC 1
MITF 1
MEN1 1
MRE11A 1
BRCA1 BRCA2 MUTYH PALB2 RAD51 TP53 ATM CHEK2 RAD51C RAD50 BRIP1 MSH2 MSH6 BARD1 PTEN RAD51D APC MLH1 CDH1 PMS2 NBN RET CDK4 EPCAM POLD1 PTCH2 RB1 RECQL FANCC MITF MEN1 MRE11A
Number of Patients
>40
20-40
10-20
1-10
0