Health effects of European colonization: An investigation of skeletal remains from 19th to early 20th century migrant settlers in South Australia

Angela Gurr^{1,2*}, Jaliya Kumaratilake^{1,2}, Alan Henry Brook^{3,4}, Stella Ioannou¹, F. Donald Pate⁵, Maciej Henneberg^{1,2,5,6}

1. Biological Anthropology and Comparative Anatomy Research Unit, Adelaide Medical School, University of Adelaide, Adelaide, South Australia
2. Discipline of Anatomy and Pathology, Adelaide Medical School, University of Adelaide, Adelaide, South Australia
3. School of Dentistry, University of Adelaide, Adelaide, South Australia
4. Institute of Dentistry, Queen Mary, University of London, London, United Kingdom
5. Archaeology, Flinders University, Adelaide, South Australia
6. Institute of Evolutionary Medicine, University of Zurich, Zurich, Switzerland

Key words: Colonial Migration, Health, Economic Status, Metabolic Deficiencies, Adelaide

Running title: Skeletal manifestations of health deficiencies in Colonial South Australia

*Corresponding author: Email: angela.gurr@adelaide.edu.au (AG)

Abstract

Nineteenth century medical understanding of human metabolism was limited, therefore, the incidence of metabolic deficiencies was not fully recorded. In addition, the transition from agricultural based mode of life to the industrial one significantly changed the pattern of these metabolic deficiencies. They were further altered by colonisation of distant continents. Palaeopathological study of skeletal remains from the early industrialised colonial era allowed light to be shed on the metabolic stresses produced by this new mode of life. Aims of this study were to investigate manifestations of disease in skeletal remains from 65 (20 adults, 45 sub-adults) migrant settlers buried in the “free ground” of St Mary’s Anglican Church Cemetery (1847 to 1927). An area allocated for burials paid for by the South Australian Government. Skeletal manifestations were determined and interpreted in terms of their multiple aetiologies. Findings were compared with those published for two 19th century British samples. Skeletal manifestations, commonly related to metabolic deficiencies, were observed. Areas of abnormal porosity of bone cortices were seen in 9 adults and 12 sub-adults, flaring of metaphyses was seen in one sub-adult, flaring of costochondral junctions of the ribs was seen in one sub-adult. Porous lesions of orbital roof bones (Types 3 to 5) were seen on three sub-adults. Micro-CT scans of tooth samples located interglobular dentine in three individuals. Comparison of St Mary’s findings with St Martins, Birmingham, and St Peter’s, Wolverhampton, UK, showed more individuals from St Mary’s had areas of abnormal porosity of bone cortices possibly related to vitamin C deficiency. However, St Mary’s sample displayed fewer changes attributable to vitamin D deficiency as expected in a country with greater UV irradiation. This indicates that, although the early industrialisation produced metabolic stresses, change of the environment through colonisation of new continents altered the distribution of metabolic deficiencies.

Abstract

Nineteenth century medical understanding of human metabolism was limited, therefore, the incidence of metabolic deficiencies was not fully recorded. In addition, the transition from agricultural based mode of life to the industrial one significantly changed the pattern of these metabolic deficiencies. They were further altered by colonisation of distant continents. Palaeopathological study of skeletal remains from the early industrialised colonial era allowed light to be shed on the metabolic stresses produced by this new mode of life. Aims of this study were to investigate manifestations of disease in skeletal remains from 65 (20 adults, 45 sub-adults) migrant settlers buried in the “free ground” of St Mary’s Anglican Church Cemetery (1847 to 1927). An area allocated for burials paid for by the South Australian Government. Skeletal manifestations were determined and interpreted in terms of their multiple aetiologies. Findings were compared with those published for two 19th century British samples. Skeletal manifestations, commonly related to metabolic deficiencies, were observed. Areas of abnormal porosity of bone cortices were seen in 9 adults and 12 sub-adults, flaring of metaphyses was seen in one sub-adult, flaring of costochondral junctions of the ribs was seen in one sub-adult. Porous lesions of orbital roof bones (Types 3 to 5) were seen on three sub-adults. Micro-CT scans of tooth samples located interglobular dentine in three individuals. Comparison of St Mary’s findings with St Martins, Birmingham, and St Peter’s, Wolverhampton, UK, showed more individuals from St Mary’s had areas of abnormal porosity of bone cortices possibly related to vitamin C deficiency. However, St Mary’s sample displayed fewer changes attributable to vitamin D deficiency as expected in a country with greater UV irradiation. This indicates that, although the early industrialisation produced metabolic stresses, change of the environment through colonisation of new continents altered the distribution of metabolic deficiencies.

34 Introduction

35
36 Although medical knowledge and medical literature had been well developed in the 19th
37 century, they did not include the modern understanding of human metabolism and thus could
38 not record the occurrence of metabolic deficiencies. Pattern of these deficiencies had been
39 significantly changed by the transition from agricultural based mode of life to the industrial
40 one and further altered by colonisation of far away continents. Palaeopathological study of
41 skeletal remains from the early industrialisation colonial era allows to shed light on the
42 metabolic stresses produced by the new mode of life.

43
44 Industrialisation of Britain in the 19th century altered the landscape, the economy, and the
45 lives of many people. This had health consequences. The expansion of urban centres of
46 manufacture offered opportunities for employment, however, the majority of cities lacked the
47 infrastructure to cope with the increase in the population size due to a migration of workers.
48 Many people had to live under overcrowded and unsanitary conditions, where the water
49 supplies could be contaminated [1-3]. These conditions, long hours of working inside
50 factories powered by coal fired engines, and diets lacking in essential nutrients contributed to
51 the poor health of the working classes in a way different from traditional agricultural
52 communities [2, 4-7].

53
54 Rapid development of urban centres, also affected the traditional small industries in rural
55 regions of Britain. The inability to compete with the low production costs of commercial
56 items in large factories caused an increase in rural unemployment [8-10]. In addition,
57 importation of large quantities of raw materials to feed the urban industries at cheaper prices,
58 such as iron, copper and tin ores, caused the down turn of regional mining industries [9, 10].

Poor harvests and the rapid spread of potato blight, further affected farmers [8, 11]. The net outcome was economic hardship in regional rural areas of the United Kingdom.

The British Government planned to reduce over-crowding in industrial cities, unemployment and economic hardships in regional areas by resettling people into the new colony of South Australia [12, 13]. The South Australian Act (1834) allowed the sale of land in the proposed settlement to individuals who would establish primary industries such as farming, mining and manufacturing [13-15]. Thus, emigration to South Australia was extensively advertised in Britain [16]. The vast size and natural environment of South Australia for farming, its Mediterranean climate and rich minerals could have attracted people to migrate to the new colony who wished to create and develop their own opportunities, rather than going to an established colony such as in North America [16]. The development of the new Australian colony required builders, mechanics, agricultural labourers, and miners [17]. A high number of migrants to South Australia came from the counties of Cornwall, Devon, Dorset and Somerset, followed by Lancashire, Middlesex, Staffordshire and Warwickshire [16, 18, 19].

Migrants who could not afford the cost of the voyage to South Australia could apply for an “assisted passage”. Migration via this route was regulated by The Colonial Land and Emigration Commission (CLEC), whose agents selected equal numbers of male and female healthy young people of “good character” [16, 17, 20]. The policy was designed to maintain a regular supply of skilled workers to landowners, new industries and to advance the development of new settlements [16, 17, 20]. The assisted passage to the new colony was funded by the British Government and The South Australian Company. Additional financial help, for individuals with an assisted passage, could be applied for from local charitable organisations. This money covered the cost of transport to the port of departure, the

84 compulsory deposit required for bedding, utensils and a set of “all weather” clothing for the
85 long voyage [16, 20].

86

87 In total, 186,054 individuals migrated to South Australia between 1836 and 1900 from the
88 United Kingdom. The government “assisted passage” was received by 123,039 (66%)
89 migrants [17]. The above mentioned selection criteria were not applied to individuals who
90 paid the full costs of their passage and had adequate funds to support themselves in the
91 colony.

92

93 The health of migrants on arrival in South Australia was of a high priority for the British
94 Government. Therefore, the Colonial Land and Emigration Commission monitored
95 conditions on board ships. Each ship had a Surgeon Superintendent, who was accountable for
96 the health and wellbeing of all passengers [16, 20, 21].

97

98 Political disagreements within the new South Australian Government delayed the surveying
99 of land for migrant settlements and the development of supporting infrastructure [12]. This
100 affected the initial establishment of farms, food production, industrial enterprises, and
101 permanent housing for migrants. These issues also delayed the development of the economy,
102 caused unemployment and led to the removal of the first Governor, Captain John Hindmarsh.
103 The second Governor, Lieutenant Colonel George Gawler, was appointed by the British
104 Government in 1838 [19, 22]. Governor Gawler had a proactive approach to leadership and
105 commissioned multiple infrastructure projects to rapidly develop the colony [22, 23].

106 Payment of the expenditure associated with these developments was rejected by the British
107 Government, who were unwilling to cover the high costs, and recalled Governor Gawler to
108 London in 1841 [19, 24].

The third South Australian Governor, Captain George Grey, was appointed the same year, and the colony faced its first economic depression [19, 25]. The lack of funds to continue public works meant Governor Grey faced retrenchment between 1841 and 1845. He redirected the majority of the unemployed workforce to agricultural industries [12, 19]. The eventual outcome was an increase of agricultural goods, particularly wheat and animal products and the establishment of export industries to other Australian colonies and Britain [12, 15].

Climatic conditions of South Australia, high summer temperatures and limited rainfall, often resulted in periods of drought and poor harvests. These conditions could have contributed to the poor economic growth and the unemployment experienced during the establishment of the colony. Throughout this period, the life of many settlers was harsh, particularly those who were unemployed and had to depend on charitable organisations and the government for their survival [23]. Government of South Australia State Records [26], indicate that 446 sick and destitute people received help in the form of food rations from the Emigration Department in 1839 - 1840. This number increased to 904 persons for the period of 1840 -1841 [23]. In 1843, an act was passed relating to the care of the needy and this was followed by the establishment of the Destitute Board in 1849. This board offered support to the elderly, chronically infirm, and some widows, and initiated the construction of a ‘Destitute Asylum’ in the city of Adelaide during the 1850s [23, 27, 28].

The Destitute Asylum was modelled on the British workhouse system [27], with strict regulations, such as compulsory wearing of uniforms for “inmates”, and severe penalties if regulations were not followed [23, 29]. This “indoor” relief at the Destitute Asylum was the last resort for individuals who had to prove that they had no relatives able to support them

[29]. Expectant mothers with no other support were also admitted to the asylum for up to six months and received “Lying-In-Home” relief [23, 29]. Furthermore, “deserted” women with children, who had no means of support were allowed admission to the asylum [29].

Individuals, who still had their own accommodation but no other support, were given weekly food rations and fire wood as “outdoor” relief from the Asylum [29]. People who lived in rural areas either had to travel long distances into the city of Adelaide to receive “outdoor” relief or cope as best they could within their own community.

Other burdens that early migrant settlers had to face were the spread of infectious diseases such as diphtheria, typhoid, typhus fever, and tuberculosis [30-34]. In addition, individuals who could not afford the cost of local medical services would have been badly affected, as the only public hospital was located at the city of Adelaide [35, 36]. People, who lived in rural areas may also have had limited access to health services [36]. Chronic conditions such as metabolic deficiencies, together with some of the above-mentioned diseases, can cause changes in the bone morphology. A number of skeletal abnormalities, for example, abnormal porosity in the cortex of bones, bending of long bones, enlargement and flaring of costochondral junctions of ribs, and porous lesions of the bones of the orbital roof, have previously been interpreted as signs of metabolic deficiencies, [37-41]. However, these skeletal lesions can be produced by a range of aetiologies [42-44].

St Marys-on-the-Sturt was an early rural migrant settlement, which was established from the late 1830’s [45]. It was a small village located eight kilometres south of the city of Adelaide (Fig.1A to C). The majority of the settlers were of British origin, who wished to establish an Anglican Church (St Mary’s) in the village. They also allocated the land surrounding the church for a cemetery (Fig.1D), with the first burial in 1847 [46]. The cemetery had a section

referred to as the “free ground”, where individuals or their families, who had no funds to support the cost of a burial were interred. The burials in the “free ground” were unmarked and the costs were paid for by the Government of South Australia [23, 47, 48]. This section of the cemetery was located at the rear of the church building (Fig. 1D) and was in use from 1847 to 1927. [45]

Fig 1. Location maps. A: Location of South Australia in relation to other Australian States and Territories. **B:** Location of the city of Adelaide, capital of South Australia. **C:** Location of St Mary’s Anglican Church Cemetery in relation to the city of Adelaide. **D:** St Mary’s Church and associated cemetery. The hatched rectangle shows the “free ground” area which is located at the rear of the church building.

The focus of this study was to investigate abnormalities seen in the skeletal remains excavated from the “free ground” area of St Mary’s Anglican Church Cemetery and understand the lifeways of these people. This study cannot ascertain the socioeconomic status of each individual buried in the “free ground” of St Mary’s Cemetery from the observed skeletal manifestations. However, the location of their burial suggests they had faced economic hardship around the time of death. Parish records of the St Mary’s Church Cemetery, list occupations of individuals who were buried in the “free ground” area, as, among others, cabinet maker, blacksmith, schoolteacher, and accountant. Some of these people had skilled trades and were considered to be more than just labourers. The probability is that this was a mixed group, mainly people who were from a working class.

Aims and Objectives:

The aims of this study were to

- i) Investigate manifestations of disease in the skeletal remains of a group of settlers buried in the “free ground” area of St Mary’s Anglican Church Cemetery, South Australia, from 1847 to 1927.
- ii) Compare the findings of the St Mary’s samples with those published for skeletal samples from two 19th century British communities.
- iii) Compare the number of individuals buried during the study period (1847 to 1927) in other sections of St Mary’s Cemetery to the number of people and the percentage of burials at the “free ground” area to understand the economic context of life in the new colony.

Material and Methods

Skeletal remains of 65 individuals (20 adults and 45 sub-adults), excavated from the “free ground” section of St Mary’s Anglican Church Cemetery, South Road, Adelaide, South Australia (Site code: SMB - St Mary’s Burial) in 2000, were available for examination. No other sections of the cemetery were excavated. These individuals were buried in the “free ground” between 1847 and 1927 in unmarked graves, which prevented identification of individuals. Sex and age at death were estimated immediately post excavation using the standard physical anthropological methods [47].

The skeletal remains are part of an archaeological collection; therefore, no destructive analysis was permitted.

Scoring of skeletal material:

Each skeleton was arranged in the anatomical position, and an inventory of the bones was prepared as described by Buikstra et al.[49] and Mitchell and Brickley [50]. Skeletal

manifestations related to metabolic deficiencies were recorded as described by Brickley[51], Brickley and Ives [37], Heron [52], Mays et al., [39], Ortner et al., [41, 53], and Ortner & Ericksen,[40] (Table 1 and Table 2). Results that are presented in Table 1 are shown as the number of individuals with the observed manifestation (-n-) over the total number (-N-) of individuals with the bone available for observation (n/N).

Porous lesions on the bones of the orbital roof were quantitated using the scoring system described by Stuart-Macadam [54], i.e., Type 1 - “capillary-like impression on bone”, Type 2- “scattered fine foramina”, Type 3 to 5 -ranged from “large and small isolated foramina to outgrowths from trabecular bone that extended to the surface of the outer table. Investigation of the internal structure of tooth samples for interglobular dentine (IGD) was part of the study. The traditional methods used for such investigations were histological techniques and these require sectioning of the tooth sample. Consequently, a non-destructive method introduced to detect IGD, using X-Ray Computed Tomography (micro-CT) was used. Cost of this method allowed only a selection of tooth samples from St Mary’s skeletal collection to be investigated.

One tooth each from 19 individuals was selected. The individuals were selected from a broad age range (~2 years to 60+ years of age). Collected teeth included two permanent incisors, three permanent canines, three permanent premolars, nine permanent first molars, and two primary molars, as the same tooth type was not available from each individual. Each tooth was scanned using the Bruker SkyScan 1276 Micro-CT scanner at Adelaide Microscopy, The University of Adelaide [55]. The scanner was set at source voltage: 100 kV, source current 200 μ A, camera binning: 4032 x 2688, filter: aluminium and copper, pixel size: 9.0 μ m. Tooth sample from SMB 63, was additionally scanned using the pixel size was 5.21 μ m.

Micro-CT scan datasets were reconstructed into a visual image using NRecon, a volumetric reconstruction software. These reconstructed scan data sets were viewed as either two-dimensional (2D) or three-dimensional (3D) images using Dataviewer, a ‘volume rendering’ software and Avizo 9 software [56]. The 2D and 3D images were analysed to identify mineralisation defects in the teeth. Dentine defects seen on the micro-CT scans were scored, as described by Colombo et al. [57] and Veselka et al. [58].

St Mary’s Parish Records and Cemetery Survey:

Parish records from St Mary’s Anglican Church, South Australia, in relation to potential burials in the “free ground” area of the cemetery from 1847 to 1927 were used. These documents recorded the burial location of some individuals, but not all [47]. The parish records also provided an approximate date of burials and the occupation for some individuals. Calculations for Figure 3 used the data from these records. The date of death and approximate date of burial for individuals who paid for their burial in provenanced graves, i.e., in the “leased ground” of the main section of the cemetery with a memorial marker/gravestone (Fig. 1D), was recorded during a survey of the St Mary’s Cemetery.

Comparison of observed metabolic deficiencies from St Mary’s samples with those of British samples:

Findings of the St Mary’s samples were compared with those published for two 19th century to early 20th century British skeletal samples. One sample was from St Martin’s-in-the-Bullring Church, Birmingham, England (N= 406) [37, 39, 59], where the majority of burials were between 1810 and 1864, with declining numbers of individuals interred until 1915 [59]. St Martin’s was located in an industrial city where many individuals from the working classes might have been buried, this sample was considered appropriate as many individuals

interred in the “free ground” of St Mary’s, or their families if they were infants, were likely from a similar socioeconomic working-class background at the time of death. The published results for a second British skeletal sample were from a section of St Peter’s Collegiate Church, Wolverhampton, England, referred to as the “overflow burial ground” (1819 to approximately 1900) [60, 61]. Wolverhampton was originally a market town with a mix of agriculture and small industries. The industrial development of local mining activities, during the 19th century, contributed to the increase in population size [60]. Published findings from St Peter’s skeletal samples were considered an appropriate comparison sample to the St Mary’s sample as some individuals in St Peter’s sample could have been miners, and it is known that many migrants to the new colony were from mining communities [16, 17, 62, 63].

Results

Demography:

Estimated age at death range and estimation of sex for the skeletal samples from St Mary’s “free ground” are shown in Figure 2. Among the 45 sub-adults in this sample, 36 were under two years of age.

Fig 2. Demography. *Distribution of the excavated St Mary’s “free ground” skeletal samples by age at death and sex [47].*

Parish Records and Cemetery Survey Data:

In the main cemetery area, surrounding St Mary’s Church, a total of 227 people were buried in “provenanced” graves between 1847 and 1927. The comparison of the number of individuals recorded in parish records as *possibly* buried in the “free ground” area of St

Mary's Cemetery (n=195), with the number of provenanced burials (n=227), by the decade of burial are presented in Figure 3.

Fig 3. St Mary's Parish Records: A Comparison of the number of people possibly buried in the "free ground" area (n=195) with those buried in provenanced graves (n= 227) of St Mary's cemetery, according to parish records [47], by decades of burial. Red dashed line indicates the percentage of burials in the "free ground" area of the cemetery.

Summary of observed skeletal manifestations:

Pathological manifestations relating to health deficiencies, possibly metabolic, were identified on the skeletal remains of some individuals buried at St Mary's "free ground" area. They are presented below:

Abnormal porosity in the bone cortex:

Nine adults and 12 sub-adults were observed with at least one area of abnormal porosity in the cortical parts of following bones: maxillae – the posterior surface, alveolar process, and palatine processes (Fig. 4); medial surfaces of the coronoid process of the mandible; and the greater wing of the sphenoid. One infant, SMB 56, (approximately 6-9 months of age), in addition to the above bones, showed areas of abnormal porosity in cortices of the lateral and basilar portion of the occipital bone, scapulae, ribs, vertebral arches, ilia, and the extremities of long bones. The prevalences of abnormal porosity are presented in Table 1.

Fig 4. SMB 56. Palate, inferior view, showing areas of abnormal porosity that extended throughout the alveolar process of the maxillae.

Table 1. Prevalence of observed skeletal manifestations associated with health deficiencies observed in different bones of the skeletons of individuals from St Mary's sample.

Age (years)	0-1	1-4	5-9	10-14	15-19	20-29	30-39	40-49	50-59	60+	Total	Total Prev. %
Abnormal porosity in the cortex of bones listed below:												
Maxillae:												
posterior surface	1/5	2/11	0/2	0/2	0/1	0/1	0/2	0/3	0/2	0/1	3/30	10
Palatine processes	1/4	0/13	0/2	0/3	0/1	0/1	0/3	1/5	1/6	0/1	3/39	8
Alveolar process	2/3	4/13	1/2	1/2	0/1	0/1	1/3	0/5	0/6	0/1	9/37	24
Mandible:												
Coronoid process	1/9	0/13	0/2	0/3	0/1	0/1	0/3	1/5	0/6	0/1	2/44	5
Alveolar process	1/9	0/15	0/2	1/3	0/1	0/1	0/3	0/5	0/7	0/2	3/48	6
Greater wing of the sphenoid bones	0/3	1/6	1/3	1/2	0/1	0/1	0/3	0/5	0/6	0/2	3/32	9
Other skeletal changes to bones listed below:												
Ribs:												
Enlargement costochondral junctions	1/4	0/5	0/1	0/2	0/1	0/1	0/3	0/3	0/2	0/0	1/22	5
Long bones:												
Flaring-distal metaphysis	0/13	1/14	0/3	0/1	0/1	0/1	0/3	0/5	0/7	0/1	1/49	2
Orbital roofs porous lesions (any Type)	0/7	2/14	1/3	3/3	1/1	0/1	0/3	0/5	2/8	0/2	9/46	20
Orbital roofs porous lesions (Types 3 or 4)	0/7	1/14	1/3	1/3	0/1	0/1	0/3	0/5	0/6	0/2	3/46	7

Note. Results presented as the number of individuals with the observed sign (-n-) over the total number (-N-) of individuals with the bone available for observation (n/N), with age groups of the individuals.

Enlargement and flaring of the costochondral junctions of ribs:

One infant, SMB 56, had enlargement and flaring of the costochondral junctions of ribs (Table 1).

Enlargement and flaring of the metaphyses:

One sub-adult, SMB 8, showed flaring and enlargement of the distal metaphyses of the femora (Table 1).

Porous lesions on the bones of the orbital roof:

Three sub-adults, SMB 4A, SMB 19 and SMB 28, age range from 4 years to 13 years, showed porous lesions on the bones of the orbital roof of Types 3 and 4 [64], which are often referred to as ‘cribra orbitalia’. The lesions of SMB 28 were composed of small and large pores on the right and left orbital roof respectively (Fig. 5). The pores on the right orbital roof seemed to penetrate the cortical bone, while those on the left appeared to be exposed trabecular bone (Fig. 5). This sub-adult also displayed areas of abnormal porosity in the cortex of the greater wing of the sphenoid bones bilaterally (Table 1). The prevalence rate of porous lesions seen on orbital roof bones in St Mary’s sample are presented in Table 1.

Fig 5. SMB 28. Lesions on the orbital roof bones. Red arrow indicates small pores in the right cortical bone. Blue arrow indicates possible exposed trabecular bone on the left.

Dentine - Interglobular dentine (IGD)

Lesions in the internal structure of the tooth seen in micro-CT scans were areas of unmineralised collagen matrix in the dentine. Such lesions have been described as interglobular dentine [65, 66]. Tooth samples from three individuals, SMB 6 (~ 40 to 45 years of age), SMB 63 (~55 to 60 + years of age), and SMB 70 (~ 7 to 12 years of age) (Fig. 6) were seen with areas of IGD. Individual SMB 70 had suffered from congenital syphilis and had been treated with mercury [67]. Therefore, this sub-adult was excluded from this part of the study as the toxic treatment could have resulted in IGD. The tooth sample from SMB

63 (a permanent lower lateral incisor) was selected for further investigation using a higher resolution of the micro-CT scanner (pixel size: 5.21 μm). This revealed areas of IGD in three separate incremental linear arrangements, one of the areas of IGD was observed opposite an external enamel hypoplastic defect (Fig. 6). The crown of this tooth type commences mineralisation at approximately 30 weeks of intrauterine life (± 1 month) and completes at approximately 3.5 years (± 1 year).

Fig 6. Micro-CT images. SMB 63. (a) Lower lateral permanent incisor with a fracture in crown. (b) Transverse slice: at the level of IGD in the crown. (c) Transverse slice: arrows show four areas of IGD. (d) Transverse slice: arrows show three areas of IGD, image was angled to show location of IGD in three concentric layers. (e) White arrow shows IGD (internal) opposite an enamel hypoplastic defect *red arrow* (external).

Sub-adults from the St Mary's sample who showed two or more macroscopic pathological manifestations of a health deficiency are listed in Table 2.

Table 2. St Mary's sub-adult samples, with their age range , showing two or more macroscopic pathological manifestation of a health deficiency.

Burial Code	SMB 27B	SMB 28	SMB 51	SMB 56	SMB 58	SMB 70
Age range at death (years)	0 to 5	10 to 15	10 to 15	0 to 5	0 to 5	7 to 12
Greater wing of the sphenoid bones †	--	P	--	--	A	P
Zygomatic bones: internal surface †	--	--	--	--	P	--
Maxillae: posterior surface †	A	A	--	P	P	A
Maxillae: palatine process †	A	A	A	P	A	A
Maxillae: alveolar process †	P	P	--	P	P	P
Mandible: coronoid process medial surface †	A	A	A	P	A	--
Mandible: alveolar process †	A	A	P	P	P	--
Orbital roofs: † Type: (number given)	P 1	P 4	P 1	--	A	A
Scapulae: † supraspinous or infraspinous areas	--	A	A	P	A	P
Pelvic bones †	A	A	A	P	A	P
Ribs: enlargement of costochondral junctions	--	A	A	P	A	--
Long bones: Flaring of distal metaphyses	--	A	--	A	A	A

Key: † = Areas of abnormal porosity (lesions) in the bone cortices observed

P = condition present

A = condition absent

-- = condition unobservable due to skeletal part missing

Findings of St Mary's samples and demographic profiles were compared with published results for 19th century skeletal samples buried at St Martin's-in-the-Bullring Church, Birmingham, England, and St Peter's Collegiate Church, (overflow burial ground), Wolverhampton, England. A comparison of the number and percentage of adults and sub-adults from each sample is presented in Table 3.

Table 3. Demographic profiles of St Mary's, St Martins and St Peter's cemeteries

Cemetery	Total Sample Size N =	Adults		Sub- adults	
		number	%	number	%
St Mary's (SA)	65	20	31	45	69
St Martin's (UK)	406	242	60	164	40
St Peter's (UK)	150	92	61	58	39

A comparison of the macroscopic observed skeletal manifestations among the sub-adult samples from the three cemeteries is presented in Figure 7.

Fig 7. Comparison of the three cemeteries. *Skeletal manifestations observed in the sub-adults samples from St Mary's, St Martin's and St Peter's Cemeteries and the number of individuals affected.*

Abnormal porosity in the bone cortex:

Sub-adults from St Martin's and St Peter's Cemeteries were observed with areas of abnormal porosity in the cortices of the following bones: infra orbital area, maxillae, hard palate, coronoid processes of the mandible, orbital, parietal and occipital bones, and scapulae [37, 59, 61]. St Mary's had more sub-adults with this manifestation (abnormal porosity in the bone cortex) compared to St Martin's and St Peter's samples (Fig. 7). Nine adults from the St Mary's sample were also seen with one area of abnormal porosity in the cortices of bones.

However, this skeletal manifestation was not observed in any adults from St Martin's or St Peter's samples [37, 59, 61].

Bending /bowing distortion of long bones:

This skeletal abnormality was seen only among sub-adults of St Martins and St Peter's cemeteries (Fig. 7) [39, 59, 61]. Furthermore, no adults from either of the three cemeteries were seen with this skeletal abnormality.

Enlargement and flaring of metaphyses:

St Martin's had a greater number of sub-adults with flaring of metaphyses than St Mary's and St Peter's sub-adult samples (Fig. 7) [39, 59, 61]. No adults from the three cemeteries were seen with this skeletal manifestation.

Porous lesions on the bones of the orbital roof:

The scores for this specific skeletal manifestation from St Martin's samples cannot be compared between St Mary's and St Peter's samples as they were not available separately for the Types 3 and 4. However, published results for this cemetery do state that 18 sub-adults and 17 adults were observed with "varying degrees" of porous lesions on the bones of the orbital roof (Fig. 7). [59].

Discussion

The "free ground" area of St Mary's Anglican Church Cemetery was allocated for the interment of people from the community of St Marys-on-the-Sturt, South Australia, who did not have the funds to cover the cost of a burial. According to the parish records from St

Mary's Church, for the decades: 1840s, 1850s, and the 1860s; 80%, 80%, and 83% respectively, of the total number of people buried at the St Mary's Cemetery were interred at the "free ground" area (Fig. 3). This indicates that for approximately 30 years, over 80% of the individuals who were buried at the cemetery or their families, could not pay for the burials. This was the period of establishment of the new colony, during which time, the colony experienced economic recession and had a high unemployment rate [12, 19]. Therefore, a good percentage of the early settlers at St Marys-on-the-Sturt region may not have had regular employment to earn an adequate amount of money to support themselves and save for the latter part of their life. They may have had to depend on charitable organisations or the Government for their survival. This early economic hardship in the colony is reflected in the need for relief from the Destitute Asylum for many living in or near the city of Adelaide. Introduction of an old age pension, after the turn of the 20th century, reduced the need for the Destitute Asylum [29].

Thirty six out of the 45 subadults (80%) buried at the "free ground" area were less than two years of age (Fig. 2). St Mary's Church parish records list some infants as "unbaptised - no service" [47]. It is possible that some of these infants were stillborn or died a short time after birth. Such infants could have contributed to the high percentage of sub-adults in the St Mary's sample. There were no official regulations regarding the location of burial of a still born infant in the early colony or the requirement to register the birth of a still born child until 1936 [48, 68]. Burial practices of perinates, excavated at the 19th century Parramatta Convict Hospital in NSW, Australia, showed they were afforded very little respect or care [69].

The percentage of burials at the St Mary's "free ground" area began to reduce from the 4th decade after the establishment of cemetery and reached 8% of the total burials in the decade of the 1920's (Fig. 3). This demonstrates that there was a gradual improvement of the economic status of some individuals who lived at St Marys-on-the-Sturt region. This local improvement could have followed the national trend of economic development of the colony of South Australia. This also shows that the majority of the skeletons studied here come from the earlier period in the life of the colony and practically characterise its health status during the 19th century, rather than reaching into the 20th century.

Bones of the skeleton are a dynamic tissue and undergo remodelling during life, possibly according to the variations in forces acting on them during life. Therefore, disease manifestations seen in skeletal remains, particularly in bone cortices, occurred during the last remodelling before death [70, 71].

Abnormal porous lesions on bone cortices result from defective calcification of the bone matrix [40, 41, 51, 72]. Vitamin C and vitamin D deficiencies affect collagen synthesis [73, 74], and affect mineralisation [65, 75, 76], respectively. Abnormal porosity of the bone cortices seen in archaeological skeletal samples could be produced by both processes. In addition, porous lesions seen on the bones of the orbit roof could be caused by an anaemia. Anaemia has often been associated with a deficiency of iron. [54, 77, 78]. When haemoglobin becomes inadequate, due to the lack of iron in the body, the red marrow compensates by the overproduction of red blood cells and thus produces a proliferation of red bone marrow. This can result in an expansion of the trabecular bone, amongst other pathological processes, between the tables of the skull [78]. In extreme conditions, the surface bone cortex of the

orbital roof may thin out and expose the underlying trabecular bone, which could appear as porous lesions.

Abnormal porous lesions that were seen in the cortical bone of 9 adults and 12 sub-adults of St Mary's skeletal collection could have resulted from either vitamin C or D deficiencies (Table 1). Presence of abnormal porosities in the bone cortices of the greater wing of the sphenoid, bilaterally, has been considered in the past as pathognomonic of vitamin C deficiency [40, 41]. Three of St Mary's sub-adults would fit into this category (Table 1). However, Brickley [38] has suggested that abnormal porous lesions should be present on more than one bone/location of the skeleton to be considered as a diagnosis of vitamin C deficiency. Accordingly, six of St Mary's sub-adults including two of the three sub-adults mentioned above with lesions on the sphenoid bones, would fit into the "multiple" lesions category (Table 2). Furthermore, one sub-adult in this category had flaring of the costochondral junctions of the ribs. This skeletal manifestation could be attributed to either vitamin C or vitamin D deficiencies (Brickley 2008). The remaining nine adults and six sub-adults only had one area of abnormal porosity on one location of the skeleton. Therefore, could not be considered as individuals with a deficiency of vitamin C. The areas of abnormal porosity seen on these individuals could represent a manifestations of vitamin D deficiency, [38, 39]. However, none of them had any other of the common signs of vitamin D deficiency, such as bowing and bending distortions of the long bones [38, 39, 79]. One sub-adult did have flaring of the distal metaphyses of the femora but without any other pathological manifestation observed elsewhere in the skeleton, this sub-adult could not be included in either of the above vitamin C or D categories (Table 1).

Three of St Mary's sub-adults had porous lesions on the bones of the orbital roof of Types 3 and 4 [54]. These "types" of lesion have been considered as an indicator of anaemia. This could result from a dietary deficiency of iron, and/or vitamin B12, malabsorption of iron from the gut due to chronic diseases or gastrointestinal and blood parasites, chronic blood loss and/or genetic conditions or a combination of these conditions [42, 54, 78, 80-83]. One sub-adult, SMB 28, also had areas of abnormal porosity on the greater wings of the sphenoid bones, (bilaterally). Therefore, it is likely that this individual could have had a co-occurrence of anaemia and vitamin C deficiency (Brickley, 2016; Fain, 2005; Ortner et al., 2001; Ortner & Ericksen, 1997). Vitamin C enhances the absorption of iron from the gut, so a deficiency of vitamin C in SMB 28 could have intensified any anaemia present.

Tooth development commences in utero and continues until approximately 21 to 23 years of age [84, 85]. In dentine, after the development of the tooth, the remodelling process is minimal compared to bone [65, 66, 86], therefore, any pathological changes (lesions) would remain for the rest of the life. The areas of IGD, seen in the two adults from St Mary's sample, would have resulted from health insults that caused an interruption of the mineralisation process in dentine during the development of the tooth. It is unknown if these individuals spent their younger lives in the UK or South Australia, indicating that the IGD could have occurred before migration. The three areas of IGD seen in SMB 63 could have resulted from three separate episodes of health insults during their childhood. Furthermore, one of the health insults that caused an area of IGD may have also resulted in the enamel hypoplastic defect (Fig. 6.e) [87-89]. The formation of IGD has been widely related to vitamin D deficiency [57, 58, 90-93]. However, neither of the two St Mary's adults showed any other skeletal pathological manifestations, including porous lesions or bending abnormalities of long bones.

557

558 **Comparison of St Mary's "free ground" skeletal samples with those of St Martin's and**
 559 **St Peter's samples in Britain:**
 560

561 The working-class background of some of the people buried at the "free ground" area of St
 562 Mary's Cemetery could be similar to those published for St Martin's and St Peter's
 563 cemeteries in Britain [59, 61]. However, the demographic profile of the St Mary's sample is
 564 somewhat different from those of the British samples, as the majority of St Mary's sample is
 565 made up of sub-adults (Fig. 2). Most of these sub-adults were under two years of age and
 566 according to St Mary's Church parish records, one of the common causes of death was
 567 gastrointestinal conditions [47]. Most of the sub-adult deaths could have resulted from
 568 dehydration resulting from diarrhoea or vomiting, as emergency access to medical help
 569 would have been difficult due to the location of the public hospital in the city of Adelaide,
 570 some 8 km away from St Marys-on-the-Sturt.

571 Areas of abnormal porosity in the cortices of bones, that could be considered as a deficiency
 572 of vitamin C, were seen in sub-adults from St Mary's, St Martin's and St Peter's skeletal
 573 samples [37, 61]. The prevalence of such lesions was higher among the subadults from the St
 574 Mary's sample (13 %) compared to that from St Martin's (4 %) and St Peter's samples (2 %).
 575 The lower prevalence of this manifestation (probable vitamin C deficiency) among the sub-
 576 adults of the British samples could be due to the availability of fresh fruits and vegetables in
 577 Birmingham and Wolverhampton. The initial difficulties and delays in establishing farms and
 578 the production of foods, during the first decades of the development of South Australia,
 579 indicates that foods particularly those rich in vitamin C, may not have been readily available.
 580 This was especially true if there was a scarcity of water [13, 94]. Some of the sub-adults
 581 affected by this deficiency, from all three of the cemeteries, could have been breastfed infants
 582 [37, 61]. Therefore, their deficiency could have resulted from a dietary deficiency of the

mother [51, 53, 95]. Infants could also have been affected by the feeding and weaning practices of the period [96-98]. Elevated nitrogen isotope values (+ 1.7‰) observed in skeletal remains of some infants from the St Mary's sample [99], relative to those of adult females in the same sample, suggested that breastmilk was a principal source of diet for infants. However, these findings do not indicate whether the child received adequate amounts of milk during breast feeding.

The number of individuals with the manifestation of enlargement and flaring of the costochondral junctions of ribs and flaring of the metaphyses of long bones was higher among St Martin's and St Peter's sub-adult samples compared to those of St Mary's samples (Fig. 7) [39, 61]. The manifestation of bending distortions of long bones was only seen among sub-adults of St Martin's and St Peter's samples, and not observed in any of sub-adult samples from St Mary's (Fig. 7). Skeletal remains of adults from the three cemeteries were free of the above-mentioned manifestations. These skeletal abnormalities have been linked to a chronic deficiency of vitamin D [39, 61]. Among the St Mary's subadults the lower incidence of the manifestations linked to vitamin D deficiency (Fig. 7) could be due to the abundance of sunlight (UV rays) in South Australia.

St Martin's had the higher number of subadults with porous lesions in the bones of the orbital roof (Fig. 7) [59]. However, the scores for these porous lesions, as described by Stuart-Macadam [64], were not available (Brickley, personal communication, 2018). This lack of data made it difficult to compare St Martin's to St Mary's and St Peter's samples in this aspect (Fig 7). Nevertheless, it is clear that many of St Martin's sub-adults had experienced chronic systemic health insults before death [78].

A comparison between the South Australian and British samples assisted in the evaluation of the level of manifestations related to health deficiencies observed in the migrant settlers buried in the “free ground” area of St Mary’s Anglican Church Cemetery. It also contributes to a wider picture of life and death in the mid-19th to early 20th century community of St Marys-on-the-Sturt.

Conclusion

Skeletal manifestations observed on individuals excavated from the “free ground” area of St Mary’s Anglican Church Cemetery, South Australia, show that St Mary’s sub-adults had a higher incidence of abnormal porosity on the bone cortex compared to the British samples. Furthermore, skeletal abnormalities such as bending distortion of long bones were more common among the British samples compared to St Mary’s samples. This indicates that, although the early industrialisation produced metabolic stresses, change of the environment through colonisation of new continents altered the distribution of metabolic deficiencies.

The new colony of South Australia, with its natural resources and Mediterranean climate could have offered many early British migrant the hope of a better life. However, unpreparedness of the Government of the new colony for settlement of the early migrants, and the refusal of the British Government to bear the cost of the establishment of the new infrastructure, led to an economic recession and mass unemployment. These caused marked hardship to the migrants. Subsequently, the timely action of the third governor of the colony helped the South Australian economy to recover. The pattern of burials at the “free ground” area of St Mary’s Cemetery from 1847 to 1927 showed the evolution of the economic status of some migrants who had settled at St Marys-on-the-Sturt. Furthermore, this pattern of

633 burials in the St Mary's Cemetery was a good reflection of the wider economic development
634 of the colony.

635

636

637

638

639

640

641

642

643

644

645

646

647

648

649

650

651

652

653

654

655

656

657

658

659

660

661

662

663

664 *The authors have no conflicts of interest to declare.*

665

666

667 *This research received no specific grant from any funding agency, commercial entity or not-*
668 *for-profit organization.*

669

670

671 *The data that support the findings of this study are available in the supporting information or*
672 *from the corresponding author upon request.*

673

674

675

676

677 **Professional and Funding Acknowledgements**

678 The authors acknowledge The University of Adelaide including the Faculty of Health and
679 Medical Sciences Honours and PhD Scholarships for AG. Fr. William Deng, St Mary's
680 Anglican Church and the Department of Archaeology, Flinders University, for allowed
681 access to the St Mary's skeletal sample. The authors acknowledge the facilities, and the
682 scientific and technical assistance, of the Australian Microscopy & Microanalysis Research
683 Facility at the Adelaide Microscopy, University of Adelaide, especially Ruth Williams for
684 her assistance with micro-CT scanning and post processing software.

685

686

687

688

689

690

691

692

693

694

695

696

697

698

699

700

701

References

1. Betsinger TK, DeWitte SN. The Bioarchaeology of Urbanization The Biological, Demographic, and Social Consequences of Living in Cities. 1st ed. 2020. ed. Switerland: Springer International Publishing; 2020.
2. Szreter S. Economic Growth, Disruption, Deprivation, Disease, and Death: On the Importance of the Politics of Public Health for Development. *Population and Development Review*. 1997;23(4):693-728. doi: 10.2307/2137377.
3. Szreter S. Urbanization, Mortality, and the Standard of Living Debate. New Estimates of the Expectation of Life at Birth in Nineteenth-Century British Cities. In: *Health and Wealth*. Boydell & Brewer; 2005. p. 165.
4. Ashworth WJ. The industrial revolution the State, knowledge and global trade. London : Bloomsbury Academic; 2017.
5. Brimblecombe P. London air pollution, 1500–1900. *Atmospheric Environment* (1967). 1977;11(12):1157-62. doi: 10.1016/0004-6981(77)90091-9.
6. Dewitte SN, Hughes-Morey G, Bekvalac J, Karsten J. Wealth, health and frailty in industrial-era. London. *Annals of Human Biology*. 2016;43(3):241-54. doi: 10.3109/03014460.2015.1020873.
7. Newman SL, Gowland RL. Dedicated Followers of Fashion? Bioarchaeological Perspectives on Socio-Economic Status, Inequality, and Health in Urban Children from the Industrial Revolution (18th–19th C), England. *International Journal of Osteoarchaeology*. 2017;27(2):217-29. doi: 10.1002/oa.2531.
8. Lokan J. From Cornish Miner to Farmer in Nineteenth-Century South Australia: A Case Study. *Cornish studies*. 2008;16(1):48-77. doi: 10.1386/corn.16.1.48_1.
9. Payton P. Cornish Emigration in Response to Changes in the International Copper Market in 1860s. Exeter, UK: University of Exeter; 1995.
10. Rowe J. Cornwall in the age of the Industrial Revolution. Liverpool: Liverpool University Press; 1953.
11. Lidwell-Durnin J. Cultivating famine: data, experimentation and food security, 1795–1848. *The British journal for the history of science*. 2020;53(2):159-81. doi: 10.1017/S0007087420000199.
12. Price AG. The foundation and settlement of South Australia, 1829-1945 : a study of the colonization movement, based on the records of the South Australian Government and on other authoritative documents. Adelaide: Libraries Board of South Australia; 1973.
13. Wilkinson GB. South Australia, its advantages and its resources : being a description of that colony, and a manual of information for emigrants / by George Blakiston Wilkinson. London : J. Murray; 1848.
14. Nance C. The South Australian social experiment 1836-71 : a study of some of the aspects of South Australia's early social development. Adelaide Flinders University of South Australia 1977.
15. Wilkinson GB. The working man's handbook to South Australia, with advice to the farmer, and detailed information for the several class of labourers and artisans. London: Murray; 1849.
16. Haines R. Emigration and the labouring poor : Australian recruitment in Britain and Ireland, 1831-60. New York : Macmillan; St. Martin's Press; 1997.
17. Haines R. Nineteenth century government assisted immigrants from the United Kingdom to Australia : schemes, regulations and arrivals, 1831-1900 and some vital statistics 1834-1860. South. Australia: Flinders University of South Australia; 1995.

- 751 18. Lawrence S, Davies P. Cornish tin-streamers and the Australian gold rush: technology
752 transfer in alluvial mining. *Post-medieval archaeology*. 2015;49(1):99-113. doi:
753 10.1179/0079423615Z.00000000073.
- 754 19. Pike D. *Paradise of dissent : South Australia 1829-1857* / Douglas Pike. 2nd ed.
755 Melbourne : Melbourne University Press; 1967.
- 756 20. Haines R. *Life and death in the age of sail : the passage to Australia*. Sydney: UNSW
757 Press; 2003.
- 758 21. Haines R. Medical Superintendence and Child Health on Government-Assisted
759 Voyages to South Australia in the Nineteenth Century. *Health and History*.
760 2001;3(2):1-29. doi: 10.2307/40111403.
- 761 22. Gibbs RM. *Under the Burning Sun: A history of colonial South Australia, 1836-1900*.
762 Adelaide: Southern Heritage; 2013.
- 763 23. Dickey B. Rations, residence, resources : a history of social welfare in South
764 Australia since 1836 , with contributions from Elaine Martin and Rod Oxenberry.
765 Martin E, Oxenberry R, editors. Netley, S. Aust. : Wakefield Press; 1986.
- 766 24. Dare R. Paupers' rights: Governor Grey and the poor law in South Australia.
767 *Australian historical studies*. 1992;25(99):220-43. doi: 10.1080/10314619208595908.
- 768 25. Jaensch D. *The Flinders history of South Australia. Political history*. Netley, S. Aust:
769 Wakefield Press; 1986.
- 770 26. Government of South Australia SR. Alphabetical index to return of sick and destitute
771 emigrants who have received relief from the emigration department 1993 [cited 2021
772 25.01]. Available from: [https://archives.sa.gov.au/finding-information/discover-our-](https://archives.sa.gov.au/finding-information/discover-our-collection/health-and-welfare/destitution-south-australia#navbar)
773 [collection/health-and-welfare/destitution-south-australia#navbar](https://archives.sa.gov.au/finding-information/discover-our-collection/health-and-welfare/destitution-south-australia#navbar).
- 774 27. Piddock S. "An Irregular and Inconvenient Pile of Buildings": The Destitute Asylum
775 of Adelaide, South Australia and the English Workhouse. *International Journal of*
776 *Historical Archaeology*. 2001;5(1):73-95. doi: 10.1023/A:1009597226492.
- 777 28. Piddock S. *A Space of Their Own: The Archaeology of Nineteenth Century Lunatic*
778 *Asylums in Britain, South Australia and Tasmania*. Orser CE, editor. New York, NY:
779 Springer 2007.
- 780 29. Geyer M. *Behind the wall : the women of the destitute asylum Adelaide, 1852-1918*.
781 Adelaide, S. Aust: Axiom, in association with the Migration Museum; 1994.
- 782 30. The Adelaide Observer. Prevention of Infection from Typhus Fever. The Adelaide
783 Observer. 1845 18.01.1845 (col. 2).
- 784 31. The South Australia Register. Pollution of the Torrens. The South Australian
785 Register. 1873 16.12.1874.
- 786 32. The South Australia Register. Drinking Typhoid. The South Australian Register. 1898
787 17.02.1898;Sect. Register.
- 788 33. The South Australian Register. Summer Cholera-Its Causes. The South Australian
789 Register,. 1847 10.02.1847;Sect. Medicine and Surgery.
- 790 34. Woodruff P. *Two million South Australians*. Kent Town, S. Aust: Peacock
791 Publications; 1984.
- 792 35. Forbes ILD. *From Colonial Surgeon to Health Commission : the government*
793 *provision of health services in South Australia, 1836-1995*. Myrtle Bank, S. Aust:
794 I.L.D. Forbes 1996.
- 795 36. Manning GH. *A colonial experience, 1838-1910 : a woman's story of life in Adelaide,*
796 *the District of Kensington and Norwood together with reminiscences of colonial life /*
797 *Underdale, S. Aust. : Gillingham Printers; 2001.*
- 798 37. Brickley M, Ives R. Skeletal manifestations of infantile scurvy. *American Journal of*
799 *Physical Anthropology*. 2006;129(2):163-72. doi: 10.1002/ajpa.20265.

38. Brickley M. The bioarchaeology of metabolic bone disease. 1st ed. Ives R, editor. Amsterdam ; Boston: Amsterdam ; Boston : Elsevier/Academic Press; 2008.
39. Mays S, Brickley M, Ives R. Skeletal manifestations of rickets in infants and young children in a historic population from England. *American Journal of Physical Anthropology*. 2006;129(3):362-74. doi: 10.1002/ajpa.20292.
40. Ortner DJ, Ericksen MF. Bone changes in the human skull probably resulting from scurvy in infancy and childhood. *International Journal of Osteoarchaeology*. 1997;7(3):212-20. doi: 10.1002/(SICI)1099-1212(199705)7:3<212::AID-OA346>3.0.CO;2-5.
41. Ortner DJ, Kimmerle EH, Diez M. Probable evidence of scurvy in subadults from archeological sites in Peru. *American Journal of Physical Anthropology*. 1999;108(3):321. doi: 10.1002/(SICI)1096-8644(199903)108:3<321::AID-AJPA7>3.0.CO;2-7.
42. Brickley M. Cribra orbitalia and porotic hyperostosis: A biological approach to diagnosis. *American journal of physical anthropology*. 2018;167(4):896. doi: 10.1002/ajpa.23701.
43. Rivera F, Lahr. M, Beatriz. M. New evidence suggesting a dissociated etiology for cribra orbitalia and porotic hyperostosis. *American Journal of Anthropology*. 2017;164:79-96. doi: 10.17863/CAM.10366.
44. Wapler U, Crubézy E, Schultz M. Is cribra orbitalia synonymous with anemia? Analysis and interpretation of cranial pathology in Sudan. *American Journal of Physical Anthropology*. 2004;123(4):333-9. doi: 10.1002/ajpa.10321.
45. Davies J, St Mary's. A Pioneer Walk Through the Churchyard of St Mary's South Road. St Mary: St Mary's Anglican Church; 1991.
46. Somerset M, Ragless R. St Mary's Walk: A Historical and Social Perspective of a Significant but Little-known Suburb of Adelaide. Mitcham, South Australia: City of Mitcham; 2009.
47. Anson TJ. The Bioarchaeology of the St. Mary's Free Ground Burials: Reconstruction of colonial South Australian lifeways. Adelaide: University of Adelaide; 2004.
48. Nicol R. At the end of the road : government, society and the disposal of human remains in the nineteenth and twentieth centuries. St Leonards, N.S.W. : Allen & Unwin; 1994.
49. Buikstra JE, Ubelaker DH, Haas J, Aftandilian D, Field Museum of Natural H. Standards for data collection from human skeletal remains : proceedings of a seminar at the Field Museum of Natural History, organized by Jonathan Haas / volume editors, Jane E. Buikstra and Douglas H. Ubelaker, assistant editor, David Aftandilian ; contributions by D. Aftandilian ... [et al.]. Fayetteville, Ark.: Fayetteville, Ark. : Arkansas Archeological Survey; 1994.
50. Mitchell PD, Brickley M. Updated Guidelines for the Standards for Recording Human Remains. Southampton: British Association for Biological Anthropology and Osteology & Institute of Field Archaeologists; 2017.
51. Brickley M. The Bioarchaeology of Metabolic Bone Disease. Boston: Elsevier Academic Press; 2008.
52. Heron C. Anne L. Grauer (Ed.): A Companion to Paleopathology. *Chromatographia*. 2013;76(23-24):1795-6. doi: 10.1007/s10337-013-2546-z.
53. Ortner DJ, Butler W, Cafarella J, Milligan L. Evidence of probable scurvy in subadults from archeological sites in North America. *American Journal of Physical Anthropology*. 2001;114(4):343-51. doi: 10.1002/ajpa.1046.
54. Stuart-Macadam P. Anemia in roman Britain: Poundbury camp. In: Bush H, Zvelebil M, editors. *Health in Past Societies: Biocultural Interpretations of Human Skeletal*

850 Remains in Archaeological Contexts. Oxford: British Archaeological Reserach Series;
851 1991. p. 101-13.

852 55. Bruker. Products: <http://bruker-microct.com/products/downloads.htm>. 2018.
853 Available from: <http://bruker-microct.com/products/downloads.htm>.

854 56. ThermoFisher Scientific. 3D Visulaization & Analysis Software: Amiro-Avizo 2019.
855 Available from: [https://www.thermofisher.com/au/en/home/industrial/electron-](https://www.thermofisher.com/au/en/home/industrial/electron-microscopy/electron-microscopy-instruments-workflow-solutions/3d-visualization-analysis-software.html)
856 [microscopy/electron-microscopy-instruments-workflow-solutions/3d-visualization-](https://www.thermofisher.com/au/en/home/industrial/electron-microscopy/electron-microscopy-instruments-workflow-solutions/3d-visualization-analysis-software.html)
857 [analysis-software.html](https://www.thermofisher.com/au/en/home/industrial/electron-microscopy/electron-microscopy-instruments-workflow-solutions/3d-visualization-analysis-software.html).

858 57. Colombo A, Ortenzio L, Bertrand B, Coqueugniot H, Knüsel CJ, Kahlon B, et al.
859 Micro-computed tomography of teeth as an alternative way to detect and analyse
860 vitamin D deficiency. Journal of Archaeological Science: Reports. 2019;23:390-5.
861 doi: 10.1016/j.jasrep.2018.11.006.

862 58. Veselka B, Brickley MB, D'Ortenzio L, Kahlon B, Hoogland MLP, Waters-Rist AL.
863 Micro-CT assessment of dental mineralization defects indicative of vitamin D
864 deficiency in two 17th-19th century Dutch communities. Am J Phys Anthropol.
865 2019;1-10. Epub 2019/03/19. doi: 10.1002/ajpa.23819. PubMed PMID: 30882907.

866 59. Brickley M, Buteux S, Adams J, Cherrington R. St. Martin's Uncovered:
867 Investigations in the churchyard of St. Martin's-in- the- Bull Ring, Birmingham,
868 2001. Oxford: Oxbow Books; 2006.

869 60. Adams J, Colls K. "Out of the darkness, cometh light' Life and Death in Nineteenth-
870 Century Wolverhampton:Excavation of the overflow burial ground of St Peter's
871 Collegiate Church, Wolverhampton 2001-2002. Oxford: BAR Publishing- British
872 Archaeological Reports; 2007.

873 61. Arabaolaza I, Ponce P, Boylston A. Skeletal Analysis. . In: Adams J, Colls K, editors.
874 '*Out of darkness cometh light*' Life and death in Nineteenth-Century Wolverhampton
875 Excavation of the overflow buril ground of St Peter's Collegiate Church,
876 Wolverhampton 2001-2002. Oxford: British Archaeolohical Reports (BAR)
877 Publishing; 2007. p. 39-70.

878 62. Haines R, Shlomowitz R. Immigration from the United Kingdom to colonial
879 Australia: A statistical analysis. Journal of Australian studies. 1992;16(34):43-52. doi:
880 10.1080/14443059209387107.

881 63. Shlomowitz R. Nominated and selected government-assisted immigration from the
882 United Kingdom to Australia, 1848-1900 : annual estimates, by colony. Bedford Park,
883 S. Aust: Flinders University of South Australia, Discipline of Economic History;
884 1991.

885 64. Stuart Macadam P. Anaemia in Roman Britain. M. BHZ, editor. Oxford: Tempvs
886 Reparatvm; 1991.

887 65. Atkinson ME, White FH. Principles of Anatomy and Oral Anatomy for Dental
888 Students. Edinburgh: Churchill Livingstone; 1992.

889 66. Nanci A. Ten Cate's Oral Histology : Development, Structure, and Function. Saint
890 Louis: Saint Louis: Elsevier; 2012.

891 67. Ioannou S, Henneberg M, Henneberg RJ, Anson T. Diagnosis of Mercurial Teeth in a
892 Possible Case of Congenital Syphilis and Tuberculosis in a 19th Century Child
893 Skeleton. Journal of Anthropology (2010). 2015;2015:1-11. doi:
894 10.1155/2015/103842.

895 68. South Australian Government. State Records: Registration of Life Events. 2021 [cited
896 2021 31.03]. Available from: [https://archives.sa.gov.au/finding-information/discover-](https://archives.sa.gov.au/finding-information/discover-our-collection/registration-life-events/registration-life-events)
897 [our-collection/registration-life-events/registration-life-events](https://archives.sa.gov.au/finding-information/discover-our-collection/registration-life-events/registration-life-events).

- 898 69. Donlon D, Casey M, Haak W, Adler C, Hack W. Early colonial burial practices for
899 perinates at the Parramatta convict hospital, NSW. *Australasian Historical*
900 *Archaeology*. 2008;26:71-83.
- 901 70. Enlow DH. principles of bone remodeling : an account of post-natal growth and
902 remodeling processes in long bones and the manabile. Springfield, Illinois.: Charles C
903 Thomas; 1963.
- 904 71. Kenkre JS, Bassett JHD. The bone remodelling cycle. *Annals of Clinical*
905 *Biochemistry*. 2018;55(3):308-27. doi: 10.1177/0004563218759371.
- 906 72. Snoddy AME, Buckley HR, Elliott GE, Standen VG, Arriaza BT, Halcrow SE.
907 Macroscopic features of scurvy in human skeletal remains: A literature synthesis and
908 diagnostic guide. *American journal of physical anthropology*. 2018;167(4):876. doi:
909 10.1002/ajpa.23699.
- 910 73. Stone N, Meister A. Function of ascorbic acid in the conversion of proline to collagen
911 hydroxyproline. *Nature*. 1962;194:555.
- 912 74. Stark RJ. A proposed framework for the study of paleopathological cases of subadult
913 scurvy. *International Journal of Paleopathology*. 2014;5:18-26. doi:
914 10.1016/j.ijpp.2014.01.005.
- 915 75. Christakos S, Ajibade DV, Dhawan P, Fechner AJ, Mady LJ. Vitamin D: metabolism.
916 *Endocrinology and metabolism clinics of North America*. 2010;39(2):243. doi:
917 10.1016/j.ecl.2010.02.002.
- 918 76. St-Arnaud R, Arabian A, Travers R, Barletta F, Raval-Pandya M, Chapin K, et al.
919 Deficient Mineralization of Intramembranous Bone in Vitamin D-24-Hydroxylase-
920 Ablated Mice Is Due to Elevated 1,25-Dihydroxyvitamin D and Not to the Absence of
921 24,25-Dihydroxyvitamin D1. *Endocrinology (Philadelphia)*. 2000;141(7):2658-66.
922 doi: 10.1210/endo.141.7.7579.
- 923 77. Camaschella C. Iron-Deficiency Anemia. *N Engl J Med*2015. p. 1832-43.
- 924 78. Godde K, Hens SM. An epidemiological approach to the analysis of cribra orbitalia as
925 an indicator of health status and mortality in medieval and post-medieval London
926 under a model of parasitic infection. *American journal of physical anthropology*.
927 2021. doi: 10.1002/ajpa.24244.
- 928 79. Buikstra JE. Ortner's Identification of Pathological Conditions in Human Skeletal
929 Remains. 3 ed. Buikstra JE, editor: Elsevier Science; 2019.
- 930 80. Blackwell S, Hendrix PC. Less Common Anemias BEYOND IRON DEFICIENCY.
931 *Clinician Reviews*. 2001;11(4):56.
- 932 81. Ferguson A, Brydon WG, Brian H, Williams A, Mackie MJ. Use of whole gut
933 perfusion to investigate gastrointestinal blood loss in patients with iron deficiency
934 anaemia. *Gut*. 1996;38(1):120. doi: 10.1136/gut.38.1.120.
- 935 82. Jelliffe DB, Blackman V. Bahima disease. Possible "milk anemia" in late childhood.
936 *The Journal of pediatrics*. 1962;61:774.
- 937 83. Weiss G, Goodnough LT. Anemia of Chronic Disease. *The New England Journal of*
938 *Medicine*. 2005;352(10):1011-23. doi: 10.1056/NEJMra041809.
- 939 84. AlQahtani S. The london atlas: developing an atlas of tooth development and testing
940 its quality and performance measures. ProQuest Dissertations Publishing; 2012.
- 941 85. AlQahtani SJ, Hector MP, Liversidge HM. Brief communication: The London atlas of
942 human tooth development and eruption. *American Journal of Physical Anthropology*.
943 2010;142(3):481-90. doi: 10.1002/ajpa.21258.
- 944 86. Hillson S. Tooth Development in Human Evolution and Bioarchaeology. Cambridge:
945 Cambridge : Cambridge University Press; 2014.
- 946 87. Brook AH, Smith JM. Hypoplastic enamel defects and environmental stress in a
947 homogeneous Romano-British population. *Eur J Oral Sci*. 2006;114:370-4.

- 948 88. Nikiforuk G, Fraser D. Etiology of Enamel Hypoplasia and Interglobular Dentin: The
949 Roles of Hypocalcemia and Hypophosphatemia. *Metabolic Bone Disease and Related*
950 *Research*. 1979;2(1):17-23. doi: 10.1016/0221-8747(79)90014-6.
- 951 89. Seow WK. Developmental defects of enamel and dentine: challenges for basic
952 science research and clinical management.(Report). *Australian Dental Journal*.
953 2014;59(s1):143. doi: 10.1111/adj.12104.
- 954 90. Brickley MB, D'ortenzio L, Kahlon B, Schattmann A, Ribot I, Raguin E, et al.
955 Ancient Vitamin D Deficiency: Long-Term Trends. *Current Anthropology*.
956 2017;58(3):420-7. doi: 10.1086/691683.
- 957 91. Brickley MB, Kahlon B, D'Ortenzio L. Using teeth as tools: Investigating the mother-
958 infant dyad and developmental origins of health and disease hypothesis using vitamin
959 D deficiency. *American journal of physical anthropology*. 2019. doi:
960 10.1002/ajpa.23947.
- 961 92. D'Ortenzio L, Kahlon B, Peacock T, Salahuddin H, Brickley M. The rachitic tooth:
962 Refining the use of interglobular dentine in diagnosing vitamin D deficiency.
963 *International Journal of Paleopathology*. 2018;22:101-8. doi:
964 10.1016/j.ijpp.2018.07.001.
- 965 93. Mellanby M. Diet and the Teeth: An Experimental Study. Part III. The Effect of Diet
966 on Dental Structure and Disease in Man. *Journal of the American Medical*
967 *Association*. 1934;102(18):1523-. doi: 10.1001/jama.1934.02750180075026.
- 968 94. Smith PA, Pate FD, Piddock S. These Enchanted Hills: Transforming cultural
969 landscapes in the Adelaide Hills Face Zone, South Australia. *Landscape Research*.
970 2020.
- 971 95. Jaffe HL. *Metabolic, degenerative, and inflammatory diseases of bones and joints*.
972 Philadelphia: Philadelphia : Lea & Febiger; 1972.
- 973 96. Fildes VA. *Breasts, bottles, and babies : a history of infant feeding*. Edinburgh:
974 Edinburgh : Edinburgh University Press; 1986.
- 975 97. Lewis M. The Problem of Infant Feeding: The Australian Experience from the Mid-
976 Nineteenth Century to the 1920s. *Journal of the History of Medicine and Allied*
977 *Sciences*. 1980;35(2):174. doi: 10.1093/jhmas/XXXV.2.174.
- 978 98. Molleson T, Cox M. *The Spitalfields Project, Vol. 2: The Anthropology, The Middling*
979 *Sort*. York, England.: Council of British Archaeology; 1993.
- 980 99. Pate DF, Anson TJ. Stable Isotope And Dietary Composition In The Mid-Late 19th
981 Century Anglican Population, Adelaide, South Australia. *Journal Of The*
982 *Anthropological Society of South Australia*. 2012;35(July 2012):1- 16.

St Mary's Cemetery

Alveolar process

Palantine process

Horizontal plate
of palatine bone

0

1

2

3

4

5

6

7

8cm