Research letter

Preliminary study on the pathogenesis of anal fistula

Jingyi Zhu^{1,2}, MD; Qingming Wang^{1,2}, MD; Zubing Mei^{1,2}, MD. PhD

Author affiliations

1. Department of Anorectal Surgery, Shuguang Hospital Affiliated to Shanghai University of Traditional Chinese Medicine, Shanghai, China
2. Anorectal Disease Institute of Shuguang Hospital, Shanghai, China

Running tittle

The pathogenesis of anal fistula (28 characters)

Corresponding Author

Dr. Qingming Wang, MD

Department of Anorectal Surgery, Shuguang Hospital, Shanghai University of

Traditional Chinese Medicine,

Anorectal Disease Institute of Shuguang Hospital,

528 Zhangheng Road, Shanghai 201203, China

Tel: +86 21 20256187

Fax: +86 21 20256187

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

E-mail: davidwqm@163.com or davidwqm@gmail.com

Authors' contributions

Study concept and design: Jingyi Zhu, Qingming Wang

Acquisition of data: Jingyi Zhu

Analysis and interpretation of data: Jingyi Zhu, Qingming Wang

Drafting of the manuscript: all authors.

Critical revision of the manuscript for important intellectual content: all authors.

Obtained funding: Qingming Wang

Corresponding author: Qingming Wang

Funding

This work was supported by the Science and Technology Commission of Shanghai

Municipality (grant no. 19401933900) (to Q.W.). The sponsor did not have any

involvement in the study design; collection, analysis, or interpretation of data; writing

of the report; or the decision to submit the article for publication.

Conflicts of interest/Competing interests

The authors declare that they have no conflicts of interest.

This work was supported by the Science and Technology Commission of Shanghai

Municipality (grant no. 19401933900).

ABSTRACT

BACKGROUND

Anal gland infection is one of the main pathogenic factors of anal fistula. The anal gland is mainly consist of columnar epithelial cells and goblet cells. Goblet cells could secrete mucins which maintain the surface mucous gel layer of intestine to prevent columnar epithelial cells from the direct acting of bacteria. The absence or diminished secretion of goblet cells could lead to the exposure of columnar epithelial cells to bacteria. In recent years, studies have found that most of the microorganisms in perianal abscess and anal fistula are intestinal bacteria. So, it can be considered that the occurrence and development of anal fistula is closely related to gut microbiota. However, the molecular mechanism of gut microbiota acting on the epithelial cells of anal gland leading to the occurrence and development of anal fistula has not been explored.

METHODS

Anal fistula tissues were collected from 30 patients, and HE staining was employed to observe pathological changes of anal gland. Stool specimens were collected from normal group (14 normal subjects), preoperative group (21 subjects before surgery) and postoperative group (16 subjects after surgery). High-throughput 16S rRNA gene sequencing was performed to explore differences among different groups.

RESULTS

It was found by HE staining that normal anal gland is composed of a large number of columnar epithelial cells as well as goblet cells, while the anal gland of patients with anal fistula is seriously lack of goblet cells. In the normal ones, a large number of vacuoles formed by the dye dissolution of mucin particles can be seen at the top of the goblet cells, and a large amount of mucin secretion can be seen in the center of the anal gland. The results of high-throughput 16S rRNA gene sequencing showed that there were significant statistical differences in species richness, species evenness and species composition between the preoperation group and the normal group ($P_1=0.00004$), and between the preoperative group and the postoperative group ($P_2=0.00003$). It also showed that PWY-6471 metabolic pathway was the most significant difference between the preoperative group and the postoperative group ($P=0.0033$).

CONCLUSIONS

The occurrence of anal fistula may be related to the abnormal permeability of mucus layer and the change of intestinal flora composition. The absence or diminished secretion of goblet cells may cause exception of the permeability of mucus layer which makes the columnar epithelial cells of anal gland exposed to the direct contact of peptidoglycan. The metabolite

of pathogenic microorganisms could activate the signal pathway (i.g. NLR/NF- κ B) which leading to the overexpression of inflammatory factors and cause anal fistula.

INTRODUCTION

Anal gland infection is one of the main pathogenic factors of anal fistula (AF). The anal gland is mainly consisted of columnar epithelial cells and goblet cells. Goblet cells could secrete mucins which maintain the surface mucous gel layer of intestine to prevent columnar epithelial cells from the direct acting of bacteria. The absence or diminished secretion of goblet cells could lead to the exposure of columnar epithelial cells to bacteria. In recent years, studies have found that most of the microorganisms in perianal abscess and AF are intestinal bacteria. So, it can be considered that the occurrence and development of AF is closely related to gut microbiota (GM). However, the molecular mechanism of GM acting on the epithelial cells of anal gland leading to the occurrence and development of AF has not been explored.

METHODS

Stool specimens were collected from normal group (14 normal subjects), preoperative group (21 subjects before surgery) and postoperative group (16 subjects after surgery). QIAamp DNA Stool Mini Kit was used for DNA extraction and quality control. Subsequently, 16S rRNA genes were amplified using a specific primer with the barcode (338F-518R). Then Ion Plus Fragment Library Kit (Life Technologies Corporation) was used in library construction. High-throughput sequencing analysis of the 16S

rRNA was performed by Ion GeneStudio S5 series gene sequencer (Life Technologies Corporation). Finally, the differences of GM among the groups were explored by analyzing the diversity of sequencing results, species classification annotation, functional annotation and pathway enrichment prediction.

AF tissues were collected from 30 patients and fixed in Bouin solution for 12 hours and washed in 70% ethanol to remove excess Bouin solution. Fixed AF were then processed by washing in a graded, increasing ethanol concentration (70%, 80%, 95%, and 100%) for 45 min each, and then permeabilized in CitriSolv (Fisher Scientific) for 30 min. The processed AF were then embedded in paraffin, sectioned, and the 5- μ m-thick sections were transferred to slides. Then HE staining was performed using an HE staining kit (Solarbio, Beijing, China) according to the manufacturers instructions and observe pathological changes of anal gland under ordinary light microscope.

RESULTS

It was found by HE staining that normal anal gland is composed of a large number of columnar epithelial cells as well as goblet cells, while the anal gland of patients with AF is seriously lack of goblet cells. In the normal ones, a large number of vacuoles formed by the dye dissolution of mucin particles can be seen at the top of the goblet cells, and a large amount of

mucin secretion can be seen in the center of the anal gland. The results of high-throughput 16S rRNA gene sequencing showed that there were significant statistical differences in species richness, species evenness and species composition between the preoperation group and the normal group ($P_1=0.00004$), and preoperative group and the postoperative group ($P_2=0.00003$). It also showed that PWY-6471 metabolic pathway was the most significant difference between the preoperative group and the postoperative group ($P=0.0033$).

DISCUSSIONS

Persistent or intermittent discharge of purulent, bloody or mucinous secretions as well as local redness, swelling, heat and pain are the main clinical features of AF.

Epidemiological studies show that the incidence rate of AF is 0.17%². The number of new AF cases is about 41.3 thousand with the 24.2814 million resident population of Shanghai. However, there is still a lack of in-depth studies on the pathogenesis of AF to date. The existing theories of anal gland infection, central space infection, and immunity can't accurately explain the pathogenesis of AF.

(1) The abnormal cellular composition of anal gland could be a causative factor in the pathogenesis of AF.

Studies by Swidsinski A and Johansson ME found that the mucus layer secreted by colonic goblet cells can prevent intestinal bacteria from colonic epithelial cells effectively³⁻⁴. Johansson ME also proposed that colonic goblet cells can secrete mucoprotein 2 (MUC2) continuously to maintain and renew the inner mucus layer in another study⁵. The inner mucus layer is an adhesive mucus layer in normal colon that bacteria cannot penetrate, while the outer mucus layer is a non-adhesive mucus layer that bacteria can penetrate⁶. The change of core structure in mucus layer is associated with attenuated secretory responses in goblet cell⁷. The further studies by Grootjans J found that mucus secreted by colonic goblet cells can remove bacteria in crypt and restore mucus layer⁸⁻⁹. Therefore, the integrity of intestinal barrier function is closely associated with goblet cell secretion. (Fig 1.)

Studies by Mitalas LE found that the epithelial cells in the internal opening of AF were derived from the epithelial lining of Morgagni crypt and anal gland¹⁰. However, the specific mechanisms between the occurrence of AF and the abnormalities of goblet cells in anal gland has not been reported yet in the literature. Our studies on HE staining showed that the occurrence of AF could be associated with intestinal barrier dysfunction caused by the abnormalities of goblet cells in anal gland. So, it is suggested that the abnormalities of goblet cells in anal gland could cause excessive inflammatory factors release in columnar epithelial cells.

The abnormalities of goblet cells could cause the abnormal penetrability of mucus layer, and then the columnar epithelial cells are exposed to the direct contact of microorganisms. Finally, columnar epithelial cells release inflammatory factors excessively could cause AF.

(2) The abnormal composition of GM could be a causative factor in the pathogenesis of AF.

The large density of microbial community in the distal intestine is even one order of magnitude higher than the total number of human cells¹¹. The results of intestinal meta-genomic sequencing showed that more than 99% of the genes were bacterial genes¹²⁻¹³. The distal GM system could be regarded as a special "micro organ". It is composed of special "cells" like other human organs and has a symbiotic relationship with the host. It is responsible for a variety of physiological functions which including energy metabolism and the development and regulation of the immune system. However, not all GM populations could build a harmonious relationship with their host¹⁴⁻¹⁵. To date, an increasing number of studies has indicated that the occurrence and development of many diseases are related to the changes of GM system such as constipation, irritable bowel syndrome, nervous system diseases, cardiovascular diseases, obesity, metabolic syndrome, autoimmune diseases, brain diseases, asthma and allergic diseases¹⁶⁻¹⁹.

Microbial detection of distal AF tissues has been carried out in some studies, and it is found that the microbial species is mainly intestinal origin²⁰. However, there is almost completely absent from the literature about the composition of faecal microbial community in AF and the difference between AF and normal population so far. Therefore, the main research objectives in the next years should be the composition of GM, the effect of pathogenic GM on epithelial cells and the pathogenesis in AF.

This may help to determine the GM based AF treatment strategy²¹.

Our studies on HE staining and high-throughput sequencing of 16S rRNA suggested that the abnormalities of goblet cells in anal gland and the changes of GM composition may together result the direct contact between columnar epithelial cells and pathogenic microorganisms which could release inflammatory factors excessively and then cause AF.

(3) Peptidoglycan (PGN), a metabolite of GM, promotes the release of inflammatory factors in columnar epithelial cells may be the pathogenesis of AF.

Inflammation is an important part of the pathogenesis in AF²². GM and their residues may potentially promote persistent inflammation procedure of AF, and PGN is a powerful inflammatory stimulator among them²³⁻²⁴.

PGN is a major component of Gram-positive and Gram-negative bacteria cell wall. It provides structural strength and enables bacteria to resist osmotic pressure.

PGN can stimulate monocyte phagocytes and endothelial cells to release immune regulatory substances such as tumor necrosis factor- α (TNF- α), interleukin (IL-1, IL-6, IL-8, IL-12, etc.), interferon α , etc. A study found that 90% of patients could be detected the presence of PGN and 60% of patients could be detected the response of host cells to PGN through sequence and analyse AF tissues by 16S rRNA high-throughput and immunohistochemical analysis²⁵. Studies by Van Onkelen RS found that IL-1 β , IL-8, IL-12p40 and TNF- α were specifically expressed in AF tissues²⁶, and the response of columnar epithelial cells which contain PGN and the response of host cells to PGN was the pathogeny of AF. The results above suggest that the overexpression of PGN may be an important pathogenic mechanism of AF.

(4) PGN-mediated NLR/NF- κ B signaling pathway release inflammatory factors.

Host cells can internalize PGN through different mechanisms and trigger NLR/NF- κ B signaling pathway: ① Cells wrap bacteria into phagolysosome through phagocytosis and release PGN into cytoplasm after the degradation of bacteria. Some pathogens can also replicate avoid the phagolysosomes in host cells and release PGN into cytoplasm. ② Extracellular PGN fragments can enter host cells through endocytosis and be transported to cytosol through lysosomal membrane transporters (SLC15A3/4). Alternatively, the oligopeptide translocon carries the PGN

fragments into the host cells through the oligopeptide transporter (hPepT1) expressed in the intestine. ③ Some bacteria can also directly transport PGN to the cytoplasm of host cells through their secretion system. ④ It also contributes to PGN internalization that outer membrane vesicles (OMV) released by Gram-negative bacteria²⁷. (Fig 2.)

PGN is one of the components in the cell wall of bacteria, which covers bacterial cells with grid-like structure. The main degradation products of PGN are γ -d-glutamyl-meso-diaminopimelic acid (iE-DAP) and muramyl dipeptide (MDP). IE-DAP is common presented in Gram-negative bacteria and MDP is widely presented in both Gram-negative and Gram-positive bacteria.

NOD1 and NOD2 proteins can sense iE-DAP and MDP, respectively. Both NOD1 and NOD2 belong to NOD-like-receptor (NLR). They are pattern recognition receptors located in the cytoplasmic matrix. They are mainly expressed in antigen presenting cells and epithelial cells. It is mainly composed of three domains: the leucine-rich repeats (LRRs) domain involved in ligand recognition at the carboxyl-terminal, the central NOD domain (also known as NACHT domain) which promotes oligomerization and has ATPase activity and the caspase-recruitment domain (CARD) which is composed of the recruitment structure of apoptotic protease²⁸. The oligomerization of NLR can promote the recruitment of downstream receptor interacting protein 2 (RIP2) after it is activated. Then it recruit

receptor-interacting serine/threonine-protein kinase 2 (RIPK2) through CARD-CARD domain interaction. After RIPK2 is activated, it can recruit and activate TGF β -activated kinase 1 (TAK1) which can activate NF- κ B signaling pathway.

NF- κ B is one of the important proteins in nuclear transcription factors and a kind of transcription activator which exists in various eukaryotic cells. In human body, it mainly exists in the form of NF- κ B-I κ Bs complex which depends on the degradation of I κ BS by IKK and ubiquitinase to activate. IKK regulates the phosphorylation of I κ B α , an inhibitor of NF- κ B, which leads to the degradation of I κ B α after polyubiquitin and cause the nuclear translocation of NF- κ B. NF- κ B signaling pathway can induce the expression of TNF- α , IL-1, IL-6, IL-8, IL-12 and other inflammatory factors. (Fig 3.)

Studies have found that NLR is ubiquitous in epithelial cells²⁹⁻³⁰ which suggest that PGN could up-regulate NLR/NF- κ B signaling pathway and promote the release of inflammatory factors in columnar epithelial cells of anal gland which may be an important pathogenic mechanism of AF.

In conclusion, in view of the abnormal microbial species and the abnormal expression of inflammatory factors in AF patients. And studies confirm that PGN can affect the release of inflammatory factors in epithelial cells by mediating NLR/NF- κ B signaling pathway. We propose a scientific hypothesis that the NLR / NF- κ B signaling pathway up-regulated PGN

promotes the release of inflammatory factors in anal gland columnar epithelial cells may be an important pathogenic mechanism of AF.

REFERENCES

- [1] Jiang W; Zhang HX; Sui N; et al., Epidemiological survey of common anorectal diseases among Chinese urban residents, *China public health*, 2016, 32(10): 1293-1296.
- [2] Swidsinski A; Loening-Baucke V; Theissig F; et al., Comparative study of the intestinal mucus barrier in normal and inflamed colon, *Gut*, 2007, 56(3): 343-350.
- [3] Johansson ME; Gustafsson JK; Holmén-Larsson J; et al., Bacteria penetrate the normally impenetrable inner colon mucus layer in both murine colitis models and patients with ulcerative colitis, *Gut*, 2014, 63(2): 281-291.
- [4] Birchenough GM; Nyström EE; Johansson ME; et al., A sentinel goblet cell guards the colonic crypt by triggering Nlrp6-dependent Muc2 secretion, *Science*, 2016, 352(6293): 1535-1542.
- [5] Meyer-Hoffert U; Hornef MW; Henriques-Normark B; et al., Andersson M; Secreted enteric antimicrobial activity localises to the mucus surface layer, *Gut*, 2008, 57(6): 764-771.
- [6] Johansson ME; Hansson GC; The goblet cell: a key player in ischaemia-reperfusion injury, *Gut*, 2013, 62(2): 188-189.
- [7] van der Post S; Jabbar KS; Birchenough G; et al., Structural weakening of the colonic mucus barrier is an early event in ulcerative colitis pathogenesis, *Gut*, 2019, 68(12): 2142-2151.

- [8] Grootjans J; Hundscheid IH; Lenaerts K; et al., Ischaemia-induced mucus barrier loss and bacterial penetration are rapidly counteracted by increased goblet cell secretory activity in human and rat colon, *Gut*, 2013, 62(2): 250-258.
- [9] Mitalas LE; van Onkelen RS; Monkhurst K; et al., Identification of epithelialization in high transsphincteric fistulas, *Techniques in Coloproctology*, 2012, 16(2): 113-117.
- [10] Backhed F; Host-Bacterial Mutualism in the Human Intestine, *Science*, 2005, 307(5717): 1915-1920.
- [11] Lepage P; Leclerc MC; Joossens M; et al., A metagenomic insight into our gut's microbiome, *Gut*, 2013, 62(1): 146-158.
- [12] Qin J; Li R; Raes J; et al., A human gut microbial gene catalogue established by metagenomic sequencing, *Nature*, 2010, 464(7285): 59-65.
- [13] Dethlefsen L; McFall-Ngai M; Relman DA; An ecological and evolutionary perspective on human-microbe mutualism and disease, *Nature*, 2007, 449(7164): 811-818.
- [14] Borody TJ; Khoruts A; Fecal microbiota transplantation and emerging applications, *Nature Reviews Gastroenterology & Hepatology*, 2012, 9(2): 88-96.
- [15] Gomes AC; Hoffmann C; Mota JF; The human gut microbiota: Metabolism and perspective in obesity, *Gut Microbes*, 2018, 9(4): 308-325.
- [16] Parker A; Fonseca S; Carding SR; Gut microbes and metabolites as modulators of blood-brain barrier integrity and brain health, *Gut Microbes*, 2020, 11(2): 135-157.
- [17] Yazdanbakhsh M; Kremsner PG; van Ree R; Allergy, parasites, and the hygiene hypothesis, *Science*, 2002, 296(5567): 490-494.

- [18]Round JL; Mazmanian SK; The gut microbiota shapes intestinal immune responses during health and disease, *Nature Reviews Immunology*, 2009, 9(5): 313-323.
- [19]Van Onkelen RS; Mitalas LE; Gosselink MP; et al., Assessment of microbiota and peptidoglycan in perianal fistulas, *Diagnostic Microbiology and Infectious Disease*, 2013, 75(1): 50-54.
- [20]Siegmund B; Feakins RM; Bamias G; et al., Results of the Fifth Scientific Workshop of the ECCO (II): Pathophysiology of Perianal Fistulizing Disease, *Journal of Crohn's and Colitis*, 2016, 10(4): 377-386.
- [21]Laman JD; Schoneveld AH; Moll FL; et al., Significance of peptidoglycan, a proinflammatory bacterial antigen in atherosclerotic arteries and its association with vulnerable plaques, *The American Journal of Cardiology*, 2002, 90(2): 119-123.
- [22]Inohara N; Ogura Y; Fontalba A; et al., Host Recognition of Bacterial Muramyl Dipeptide Mediated through NOD2, *Journal of Biological Chemistry*, 2003, 278(8): 5509-5512.
- [23]Van Onkelen RS; Gosselink MP; van Meurs M; et al., Pro-inflammatory cytokines in cryptoglandular anal fistulas, *Techniques in Coloproctology*, 2016, 20(9): 619-625.
- [24]Wolf AJ, Reyes CN; Liang W; et al., Hexokinase Is an Innate Immune Receptor for the Detection of Bacterial Peptidoglycan, *Cell*, 2016, 166(3): 624-636.
- [25]Strober W; Murray PJ; Kitani A; et al., Signalling pathways and molecular interactions of NOD1 and NOD2, *Nat Rev Immunol*, 2006, 6(1): 9-20.

- [26] Kelly P; Meade KG; O'Farrelly C; Non-canonical Inflammasome-Mediated IL-1 β Production by Primary Endometrial Epithelial and Stromal Fibroblast Cells Is NLRP3 and Caspase-4 Dependent, *Front Immunol*, 2019, 10: 102.
- [27] Zhu S; Ding S; Wang P; et al., Flavell RA; Nlrp9b inflammasome restricts rotavirus infection in intestinal epithelial cells, *Nature*, 2017, 546(7660): 667-670.
- [28] Zhang ZJ; Zheng D; Wang QM; et al., "Anal carbuncle prescription" for internal and external use to promote wound healing after high perianal abscess surgery: a clinical study of 50 cases, *Jiangsu traditional Chinese medicine*, 2018, 50(8): 41-43.

Figure legends

Fig 1. Microorganisms penetrate the mucus layer and act on the columnar epithelial cells of anal gland.

Fig 2. Host cells can internalize PGN through different mechanisms and trigger NLR/NF- κ B signaling pathway

Fig 3. The NLR / NF- κ B signaling pathway mediated by PGN promotes the release of inflammatory factors.