

Supplementary Information

Article title

Potential of machine learning to predict early ischemic events after carotid endarterectomy or stenting: A comparison with surgeon predictions

Journal name

Neurosurgical Review

Author names

Kazuya Matsuo, Atsushi Fujita, Kohkichi Hosoda, Jun Tanaka, Taichiro Imahori, Taiji Ishii, Masaaki Kohta, Kazuhiro Tanaka, Yoichi Uozumi, Hidehito Kimura, Takashi Sasayama, Eiji Kohmura.

Affiliation and e-mail address of the corresponding author

Kazuya Matsuo, MD, PhD, Department of Neurosurgery, Kobe University Graduate School of Medicine, Kobe, Japan. E-mail: kkmatsuo@outlook.jp

Supplemental Table 1. Patient characteristics of CEA group and CAS group.

Variable	CEA group (n=95)	CAS group (n=70)	<i>p</i> value
Age, year, mean (SD)	74.0 (8.22)	74.5 (6.74)	0.70
pre-treatment mRS, median [IQR]	0 [0-1]	0 [0-2]	0.41
LDL-cholesterol, mg/dL, mean (SD)	89.7 (32.7)	90.7 (23.9)	0.83
Prior medical histories, n (%)			
Hypertension	84 (88)	52 (74)	0.02
Diabetes mellitus	30 (32)	28 (40)	0.32
Arteriosclerotic disease	36 (38)	35 (50)	0.15
Anatomical and pathophysiological features			
Contralateral occlusion, n (%)	9 (9.5)	5 (7.1)	0.78
Stenosis at a high position, n (%)	8 (8.4)	5 (7.1)	1
Type III Aorta, n (%)	33 (35)	31 (44)	0.26
ICA-PSV, cm/sec, mean (SD)	259 (125)	294 (132)	0.09
Mobile plaque, n (%)	14 (15)	5 (7.1)	0.15
Plaque ulceration, n (%)	33 (35)	6 (8.6)	<0.0001
Plaque with hyperintense signal on TOF, n (%)	43 (45)	18 (26)	0.01
Previous neck irradiation, n (%)	7 (7.4)	8 (11)	0.42
Symptomatic, n (%)	45 (47)	19 (27)	0.01
Crescendo TIA or stroke in evolution, n (%)	3 (3.2)	7 (10)	0.10
Outcome, n (%)			
Ischemic stroke within 30 days	20 (21)	26 (37)	0.03
Major ischemic stroke	3 (3.2)	0 (0)	0.27
Minor ischemic stroke	0 (0)	3 (4.3)	0.07
Asymptomatic DWI hyperintense lesions	17 (18)	23 (33)	0.04

CAS = carotid artery stenting; CEA = carotid endarterectomy; DWI = diffusion weighted imaging; IQR = interquartile range; LDL = low density lipoprotein; mRS = modified Rankin scale; ICA-PSV = internal carotid artery-peak systolic velocity; TIA = transient ischemic attack; TOF = time-of-flight.

Supplemental table 2. Optimized hyperparameters of five machine learning models

Model	Hyperparameters
XGBoost	learning_rate = 0.3 n_estimators = 360 max_depth = 1 min_child_weight = 6.5 gamma = 0.95 subsample = 0.65 colsample_bytree = 0.65 objective = 'binary:logistic' reg_alpha = 0.001 reg_lambda = 0.1 max_delta_step = 1 early_stopping_rounds = 20
Random forest	criterion = 'entropy' max_depth = 4 max_features = 4 min_samples_leaf = 0.0001 min_samples_split = 0.0001 n_estimators = 40
Logistic regression	penalty = 'l2' C = 0.05 solver = 'saga'
SVM	kernel = 'linear' C = 2.5 coef0 = 0 degree = 1 gamma = 0.0001 probability = True

Neural Network	<pre>models.Sequential() model.add(layers.Dense(16, activation='relu', input_shape=(17,))) model.add(layers.Dropout(0.2)) model.add(layers.BatchNormalization()) model.add(layers.Dense(16, activation='relu')) model.add(layers.Dropout(0.2)) model.add(layers.BatchNormalization()) model.add(layers.Dense(1, activation='sigmoid')) EarlyStopping (patience=20, restore_best_weights=True) optimizer=Adam (lr=0.01) batch_size=8 epochs=41</pre>
----------------	--

SVM = support vector machine

* A subset of the program code generated for this study is available at GitHub and can be accessed at [BLINDED FOR REVIEW].