

SUPPLEMENTAL TABLES

Supplemental Table 1: Characteristics of the cohort. ApoB indicates Apolipoprotein B; ASCVD, Atherosclerotic Cardiovascular Disease; BMI, Body Mass Index; HDL-C, High Density Lipoprotein Cholesterol; HES, Hospital Episode Statistics; IQR, Interquartile Range; LDL-C, Low Density Lipoprotein Cholesterol; Lp(a), Lipoprotein (a); MI, Myocardial Infarction; PCE, Pooled Cohort Equation; SD, Standard Deviation; TC, Total Cholesterol

Categories	Sub-categories	Whole Cohort	Training Subset	Validation Subset
n (%)	All	408,422	5,000	403,422
	Women	220,802 (54.06)	2,916 (58.32)	217,886 (54.01)
	Men	187,620 (45.94)	2,084 (41.68)	185,536 (45.99)
Age at baseline, y (\pm SD)	All	56.91 (8.00)	55.97 (7.95)	56.93 (8.00)
	Women	56.72 (7.92)	55.95 (7.90)	56.73 (7.92)
	Men	57.15 (8.09)	55.99 (8.03)	57.16 (8.09)
BMI, kg/m ² (\pm SD)		27.42 (4.75)	26.97 (4.52)	27.42 (4.76)
Blood Pressure, mmHg (\pm SD)	Diastolic	82.27 (10.11)	82.12 (10.09)	82.27 (10.11)
	Systolic	138.29 (18.64)	137.22 (18.66)	138.30 (18.64)
Blood lipids, mmol/L (\pm SD)	LDL-C	3.57 (0.87)	3.72 (0.81)	3.57 (0.87)
	HDL-C	1.45 (0.38)	1.49 (0.38)	1.45 (0.38)
	TC	5.71 (1.14)	5.92 (1.06)	5.71 (1.15)
	APO-B	1.03 (0.24)	1.07 (0.23)	1.07 (0.24)
nmol/L (\pm SD)	Lp(a)	44.09 (49.45)	44.63 (49.97)	44.09 (49.45)
Hypertension, n (%)		91,201 (22.33)	659 (13.18)	90,542 (22.44)
Diabetes, n (%)		20,620 (5.05)	98 (1.96)	20,522 (5.09)
Current Smoker at baseline, n (%)		41,253 (10.10)	490 (9.80)	40,763 (10.10)
10-year risk of first ASCVD event (PCE)	n (%)	360,407 (88.24)	4,720 (94.40)	355,687 (88.17)
	Risk % (\pm SD)	8.38 (7.66)	7.36 (7.15)	8.39 (7.66)
HES CAD Outcome, n (%)	Prevalent (all cases)	32,694 (8.00)	219 (4.38)	32,475 (8.05)
HES MI Outcome, n (%)	Prevalent (all cases)	14,924 (3.65)	97 (1.94)	14,827 (3.67)
HES MI-Revascularisation Outcome, n (%)	Prevalent (all cases)	20,769 (5.08)	131 (2.62)	20,638 (5.12)
Mortality, n (%) Before March 2020	All-cause	24,229 (5.93)	247 (4.94)	23,982 (5.94)
Follow-up, y (\pm IQR)		11.04 (1.40)	11.10 (1.35)	11.04 (1.40)

Supplemental Table 2: Association between PRS_{CAD} and prevalent CAD in the training dataset. Non-adjusted and adjusted models are used in logistic regression with prevalent cardiovascular outcomes. Data are presented as logistic-regression odd ratios (OR) per SD increase of PRS_{CAD} with their 95% confidence intervals (95% CI), logistic regression P-value significance and corresponding area under the receiver operating characteristic curve (AUC) values. MI, Myocardial Infarction; MI-REVASC: MI or Revascularisation Procedure; PRS_{CAD}, Polygenic Risk Score for Coronary Artery Disease; PCE, Pooled Cohort Equation; PCs, First Ten Genetic Principal Components; CI: Confidence Interval

LDPred Rho Threshold	Covariates	CAD (n = 219)			MI (n = 97)			MI-REVASC. (n = 131)		
		OR (95% CI)	P-value	AUC	OR (95% CI)	P-value	AUC	OR (95% CI)	P-value	AUC
0.1	PRS _{CAD}	1.36 (1.19 to 1.56)	9.15E-06	0.588	1.27 (1.03 to 1.55)	2.21E-02	0.570	1.45 (1.21 to 1.72)	3.59E-05	0.603
	PRS _{CAD} + PCs	1.36 (1.19 to 1.56)	1.12E-05	0.599	1.27 (1.03 to 1.55)	2.26E-02	0.603	1.45 (1.21 to 1.73)	3.81E-05	0.615
	PRS _{CAD} + AGE	1.38 (1.20 to 1.58)	6.90E-06	0.691	1.27 (1.04 to 1.56)	2.06E-02	0.663	1.46 (1.22 to 1.74)	2.91E-05	0.691
	PRS _{CAD} + AGE + SEX + PCs	1.38 (1.20 to 1.59)	4.47E-06	0.737	1.28 (1.04 to 1.57)	1.71E-02	0.736	1.47 (1.23 to 1.75)	2.16E-05	0.754
0.03	PRS _{CAD}	1.42 (1.24 to 1.63)	5.72E-07	0.599	1.36 (1.11 to 1.67)	2.57E-03	0.591	1.53 (1.28 to 1.82)	2.06E-06	0.619
	PRS _{CAD} + PCs	1.42 (1.23 to 1.63)	6.81E-07	0.609	1.36 (1.11 to 1.67)	2.73E-03	0.620	1.53 (1.28 to 1.83)	2.21E-06	0.631
	PRS _{CAD} + AGE	1.43 (1.25 to 1.65)	3.78E-07	0.695	1.37 (1.12 to 1.68)	2.37E-03	0.671	1.54 (1.29 to 1.84)	1.58E-06	0.698
	PRS _{CAD} + AGE + SEX + PCs	1.44 (1.26 to 1.66)	2.14E-07	0.740	1.38 (1.13 to 1.69)	1.86E-03	0.741	1.55 (1.30 to 1.85)	1.06E-06	0.760
0.01	PRS _{CAD}	1.44 (1.26 to 1.65)	1.78E-07	0.600	1.43 (1.17 to 1.76)	5.10E-04	0.597	1.57 (1.32 to 1.87)	5.26E-07	0.621
	PRS _{CAD} + PCs	1.44 (1.26 to 1.66)	1.94E-07	0.611	1.43 (1.17 to 1.75)	5.76E-04	0.627	1.57 (1.32 to 1.88)	5.68E-07	0.634
	PRS _{CAD} + AGE	1.46 (1.27 to 1.68)	7.73E-08	0.694	1.44 (1.18 to 1.77)	4.34E-04	0.672	1.59 (1.33 to 1.90)	3.14E-07	0.698
	PRS _{CAD} + AGE + SEX + PCs	1.48 (1.29 to 1.70)	3.45E-08	0.742	1.46 (1.19 to 1.78)	3.10E-04	0.744	1.61 (1.35 to 1.92)	1.75E-07	0.762
0.003	PRS _{CAD}	1.43 (1.25 to 1.64)	3.14E-07	0.594	1.45 (1.18 to 1.77)	3.69E-04	0.586	1.54 (1.29 to 1.84)	1.50E-06	0.606
	PRS _{CAD} + PCs	1.44 (1.25 to 1.65)	3.15E-07	0.609	1.44 (1.18 to 1.77)	4.37E-04	0.621	1.54 (1.29 to 1.84)	1.60E-06	0.623
	PRS _{CAD} + AGE	1.46 (1.27 to 1.68)	9.70E-08	0.692	1.46 (1.19 to 1.79)	2.89E-04	0.669	1.57 (1.31 to 1.87)	7.36E-07	0.692
	PRS _{CAD} + AGE + SEX + PCs	1.48 (1.29 to 1.71)	3.67E-08	0.739	1.47 (1.20 to 1.81)	1.99E-04	0.744	1.59 (1.33 to 1.90)	3.65E-07	0.759
0.001	PRS _{CAD}	1.37 (1.19 to 1.58)	7.59E-06	0.583	1.39 (1.13 to 1.71)	1.67E-03	0.576	1.43 (1.20 to 1.71)	8.20E-05	0.588
	PRS _{CAD} + PCs	1.38 (1.20 to 1.58)	7.54E-06	0.601	1.38 (1.13 to 1.70)	1.98E-03	0.613	1.43 (1.20 to 1.71)	8.69E-05	0.608
	PRS _{CAD} + AGE	1.40 (1.22 to 1.61)	2.51E-06	0.688	1.40 (1.14 to 1.72)	1.31E-03	0.665	1.45 (1.21 to 1.73)	4.28E-05	0.684
	PRS _{CAD} + AGE + SEX + PCs	1.42 (1.23 to 1.63)	1.13E-06	0.734	1.41 (1.15 to 1.73)	1.01E-03	0.740	1.47 (1.23 to 1.75)	2.48E-05	0.751
0.0003	PRS _{CAD}	1.12 (0.98 to 1.29)	8.83E-02	0.533	0.97 (0.79 to 1.19)	7.60E-01	0.513	1.05 (0.89 to 1.25)	5.61E-01	0.510
	PRS _{CAD} + PCs	1.13 (0.99 to 1.30)	7.58E-02	0.564	0.98 (0.80 to 1.20)	8.42E-01	0.573	1.06 (0.89 to 1.26)	5.07E-01	0.548
	PRS _{CAD} + AGE	1.13 (0.99 to 1.30)	7.18E-02	0.674	0.98 (0.81 to 1.20)	8.82E-01	0.651	1.07 (0.90 to 1.27)	4.76E-01	0.664
	PRS _{CAD} + AGE + SEX + PCs	1.13 (0.98 to 1.29)	8.50E-02	0.725	0.98 (0.80 to 1.20)	8.11E-01	0.727	1.06 (0.89 to 1.26)	5.29E-01	0.735

Supplemental Table 3: Association between PRS_{CAD} and prevalent cardiovascular outcomes in multivariable models. Non-adjusted and adjusted models are used in logistic regression with CAD, MI and MI-REVASC. outcomes. Data are presented as logistic-regression odd ratios (OR) per SD increase of PRS_{CAD} and their 95% confidence intervals (95% CI), Nagelkerke's pseudo-R² corresponding area under the receiver operating characteristic curve (AUC) values.

BMI, Body Mass Index; HDL-C, High Density Lipoprotein Cholesterol; LDL-C, Low Density Lipoprotein Cholesterol; Lp(a), Lipoprotein (a); MI, Myocardial Infarction; MI-REVASC: MI or Revascularisation Procedure; PCE, Pooled Cohort Equation; PCs, First Ten Genetic Principal Components; PRS_{CAD}, Polygenic Risk Score for Coronary Artery Disease; TC, Total Cholesterol

Outcome	Logistic Regression Models	Event (n)	Control (n)	Total (n)	OR (95% CI)	Pseudo-R ²	AUC
CAD	Model 1: AGE + SEX	32,475	370,947	403,422	NA	0.127	0.742
	Model 2: PRS _{CAD} + PCs	32,475	370,947	403,422	1.50 (1.49 to 1.52)	0.0280	0.613
	Model 3: PRS _{CAD} + AGE + SEX + PCs	32,475	370,947	403,422	1.56 (1.54 to 1.58)	0.158	0.766
	Model 4: PCE	27,568	328,119	355,687	NA	0.107	0.749
	Model 5: PRS _{CAD} + PCE + PCs	27,568	328,119	355,687	1.51 (1.49 to 1.53)	0.132	0.758
	Model 6: PRS _{CAD} + AGE + SEX + PCs + LDL + HDL + TC + BMI + SBP + DIABETES + SMOKING	27,516	327,628	355,144	1.55 (1.53 to 1.57)	0.205	0.797
MI	Model 1: AGE + SEX	14,827	388,595	403,422	NA	0.0990	0.744
	Model 2: PRS _{CAD} + PCs	14,827	388,595	403,422	1.58 (1.56 to 1.61)	0.0280	0.629
	Model 3: PRS _{CAD} + AGE + SEX + PCs	14,827	388,595	403,422	1.63 (1.60 to 1.65)	0.129	0.772
	Model 4: PCE	12,517	343,170	355,687	NA	0.0800	0.748
	Model 5: PRS _{CAD} + PCE + PCs	12,517	343,170	355,687	1.58 (1.55 to 1.61)	0.105	0.761
	Model 6: PRS _{CAD} + AGE + SEX + PCs + LDL + HDL + TC + BMI + SBP + DIABETES + SMOKING	12,495	342,649	355,144	1.60 (1.57 to 1.63)	0.172	0.807
MI-REVASC.	Model 1: AGE + SEX	20,638	382,784	403,422	NA	0.123	0.756
	Model 2: PRS _{CAD} + PCs	20,638	382,784	403,422	1.66 (1.63 to 1.68)	0.0370	0.640
	Model 3: PRS _{CAD} + AGE + SEX + PCs	20,638	382,784	403,422	1.73 (1.70 to 1.75)	0.162	0.789
	Model 4: PCE	17,402	338,285	355,687	NA	0.0960	0.758
	Model 5: PRS _{CAD} + PCE + PCs	17,402	338,285	355,687	1.67 (1.65 to 1.70)	0.130	0.774
	Model 6: PRS _{CAD} + AGE + SEX + PCs + LDL + HDL + TC + BMI + SBP + DIABETES + SMOKING	17,373	337,771	355,144	1.71 (1.68 to 1.74)	0.205	0.817

Supplemental Table 4: Association between PRS_{CAD} and MI incidence in adjusted Cox regression models. Adjusted Cox regression results from PRS_{CAD} models including age and the first ten genetic principal components (PCs) as covariates were obtained in women and men, by quartiles of age (Q1 to Q4) and for the whole group (all age). Data are presented as estimated hazard ratios (HR) for incident MI outcome per SD increase of PRS_{CAD}, with their 95% confidence intervals (95% CI), and Cox regression P-value significance.

Outcome	Sex Group	Age Group	Age Window	Max-Min Difference in Age Group	N Event	N Total	Ratio N Event / 100% of Event	HR (95% CI)	P-value
MI Incident	All	All Ages	from 40 to 73	33	7,746	393,725	100.00%	1.53 (1.49 to 1.56)	2.69E-296
		Quartile 1	from 40 to 51	11	889	100,284	11.44%	1.89 (1.77 to 2.02)	1.15E-79
		Quartile 2	from 51 to 58	7	1,602	99,321	20.68%	1.59 (1.51 to 1.67)	2.43E-74
		Quartile 3	from 58 to 63	5	2,224	98,008	28.71%	1.42 (1.36 to 1.48)	4.22E-61
		Quartile 4	from 63 to 73	10	3,031	96,112	39.13%	1.48 (1.42 to 1.53)	2.14E-98
	Women	All Ages	from 40 to 71	31	2,295	216,020	100.00%	1.45 (1.39 to 1.51)	2.02E-70
		Quartile 1	from 40 to 51	11	208	55,176	9.06%	1.57 (1.37 to 1.79)	9.04E-11
		Quartile 2	from 51 to 58	7	427	56,422	18.61%	1.43 (1.30 to 1.57)	9.17E-14
		Quartile 3	from 58 to 63	5	662	54,227	28.85%	1.40 (1.30 to 1.51)	7.90E-18
		Quartile 4	from 63 to 71	8	998	50,195	43.49%	1.47 (1.38 to 1.57)	1.68E-33
	Men	All Ages	from 40 to 73	33	5,451	177,705	100.00%	1.56 (1.52 to 1.60)	1.02E-228
		Quartile 1	from 40 to 51	11	681	45,108	12.49%	2.00 (1.86 to 2.16)	1.93E-72
		Quartile 2	from 51 to 58	7	1,175	42,899	21.56%	1.65 (1.56 to 1.75)	1.71E-63
		Quartile 3	from 58 to 63	5	1,562	43,781	28.66%	1.43 (1.36 to 1.50)	7.19E-45
		Quartile 4	from 63 to 73	10	2,033	45,917	37.30%	1.48 (1.41 to 1.54)	1.04E-66

Supplemental Table 5: Comparison between adjusted Cox regression models for MI incidence including PCE and PRS_{CAD}. Data are presented as estimated hazard ratios (HR) for incident MI outcome per SD increase of PRS_{CAD}, with their 95% confidence intervals (95% CI), and Cox regression p-value significance. *Sex adjustment was only performed in “all” group.

MI: Myocardial Infarction; PRS_{CAD}: Polygenic Risk Score for Coronary Artery Disease; PCE: Pooled Cohort Equation; PCs: First Ten Genetic Principal Components

Outcome	Sex Group	Age Group	PRS _{CAD} + AGE + SEX* + PCs				PRS _{CAD} + PCE + PCs			
			N Event	N Total	HR (95% CI)	P-value	N Event	N Total	HR (95% CI)	P-value
MI Incident	All	All Ages	7,746	393,725	1.53 (1.49 to 1.56)	2.69E-296	6,735	347,654	1.51 (1.48 to 1.55)	5.81E-245
		Quartile 1	889	100,284	1.89 (1.77 to 2.02)	1.15E-79	797	89,246	1.88 (1.75 to 2.01)	1.61E-69
		Quartile 2	1,602	99,321	1.59 (1.51 to 1.67)	2.43E-74	1,391	87,841	1.55 (1.47 to 1.64)	8.64E-59
		Quartile 3	2,224	98,008	1.42 (1.36 to 1.48)	4.22E-61	1,938	86,339	1.40 (1.34 to 1.46)	8.17E-49
		Quartile 4	3,031	96,112	1.48 (1.42 to 1.53)	2.14E-98	2,609	84,228	1.47 (1.41 to 1.53)	7.01E-82
	Women	All Ages	2,295	216,020	1.45 (1.39 to 1.51)	2.02E-70	1,976	191,065	1.43 (1.37 to 1.5)	4.34E-57
		Quartile 1	208	55,176	1.57 (1.37 to 1.79)	9.04E-11	190	49,124	1.55 (1.34 to 1.79)	2.52E-09
		Quartile 2	427	56,422	1.43 (1.30 to 1.57)	9.17E-14	373	49,965	1.36 (1.23 to 1.51)	1.72E-09
		Quartile 3	662	54,227	1.40 (1.30 to 1.51)	7.90E-18	572	47,885	1.37 (1.26 to 1.49)	5.42E-14
		Quartile 4	998	50,195	1.47 (1.38 to 1.57)	1.68E-33	841	44,091	1.46 (1.37 to 1.57)	1.88E-27
	Men	All Ages	5,451	177,705	1.56 (1.52 to 1.60)	1.02E-228	4,759	156,589	1.54 (1.50 to 1.59)	4.50E-189
		Quartile 1	681	45,108	2.00 (1.86 to 2.16)	1.93E-72	607	40,122	1.99 (1.84 to 2.16)	1.72E-63
		Quartile 2	1,175	42,899	1.65 (1.56 to 1.75)	1.71E-63	1,018	37,876	1.62 (1.52 to 1.73)	8.73E-51
		Quartile 3	1,562	43,781	1.43 (1.36 to 1.50)	7.19E-45	1,366	38,454	1.41 (1.34 to 1.49)	1.71E-36
		Quartile 4	2,033	45,917	1.48 (1.41 to 1.55)	1.04E-66	1,768	40,137	1.47 (1.40 to 1.54)	1.74E-55

Supplemental Table 6: MI incidence in individuals at high genetic risk. Adjusted Cox regression results were obtained for individuals above percentile thresholds (ranging between the 80th and the 99th percentile). Hazard ratios (HR) for MI incidence and their 95% confidence intervals (95% CI) were compared in all individuals, and separately in women and in men, for all ages and for each quartile of age (Q1 to Q4).

Outcome	Sex Group	Age Group	Age Distribution	Percentile Thres.	N event	N total	HR (95% CI)	P-value
MI Incident	All	All Ages	from 40 to 73	80%	2,528	76,217	2.02 (1.93 to 2.12)	2.03E-185
				90%	1,454	37,919	2.19 (2.07 to 2.32)	1.80E-159
				95%	806	18,881	2.36 (2.19 to 2.54)	2.39E-117
				97.50%	435	9,409	2.51 (2.28 to 2.76)	1.95E-77
				99%	211	3,727	3.04 (2.65 to 3.49)	4.52E-57
		Quartile 1	from 40 to 51	80%	374	19,683	2.92 (2.55 to 3.33)	9.30E-56
				90%	236	9,793	3.29 (2.83 to 3.82)	2.68E-55
				95%	141	4,874	3.63 (3.03 to 4.34)	1.24E-44
				97.50%	71	2,437	3.44 (2.70 to 4.39)	1.86E-23
				99%	39	964	4.48 (3.25 to 6.17)	6.13E-20
		Quartile 2	from 51 to 58	80%	543	19,322	2.13 (1.92 to 2.36)	2.26E-46
				90%	317	9,616	2.33 (2.06 to 2.63)	2.52E-41
				95%	173	4,794	2.44 (2.08 to 2.86)	1.99E-28
				97.50%	102	2,382	2.89 (2.36 to 3.53)	4.20E-25
				99%	48	946	3.45 (2.59 to 4.60)	3.28E-17
		Quartile 3	from 58 to 63	80%	669	18,933	1.77 (1.61 to 1.93)	1.02E-34
				90%	389	9,412	1.96 (1.75 to 2.18)	2.99E-33
				95%	206	4,695	2.01 (1.74 to 2.32)	1.34E-21
				97.50%	110	2,341	2.09 (1.73 to 2.53)	4.61E-14
				99%	54	927	2.63 (2.00 to 3.44)	2.54E-12
		Quartile 4	from 63 to 73	80%	953	18,270	1.90 (1.76 to 2.05)	2.20E-60
				90%	528	9,084	1.98 (1.80 to 2.18)	3.29E-46
				95%	289	4,517	2.10 (1.86 to 2.37)	3.57E-33
				97.50%	155	2,248	2.22 (1.89 to 2.61)	3.73E-22
				99%	75	887	2.64 (2.10 to 3.33)	9.71E-17

Outcome	Sex Group	Age Group	Age Distribution	Percentile Thres.	N event	N total	HR (95% CI)	P-value
MI Incident	Women	All Ages	from 40 to 71	80%	704	42,500	1.80 (1.65 to 1.97)	2.25E-38
				90%	436	21,166	2.16 (1.95 to 2.40)	2.21E-47
				95%	252	10,549	2.41 (2.11 to 2.75)	1.58E-39
				97.50%	148	5,253	2.80 (2.37 to 3.30)	1.18E-33
				99%	75	2,086	3.51 (2.79 to 4.42)	1.24E-26
		Quartile 1	from 40 to 51	80%	73	10,963	2.18 (1.64 to 2.89)	9.07E-08
				90%	49	5,469	2.78 (2.01 to 3.82)	4.32E-10
				95%	36	2,723	3.98 (2.78 to 5.70)	5.17E-14
				97.50%	21	1,359	4.38 (2.79 to 6.88)	1.40E-10
				99%	14	538	7.20 (4.18 to 12.4)	1.02E-12
		Quartile 2	from 51 to 58	80%	130	11,155	1.76 (1.44 to 2.17)	6.84E-08
				90%	82	5,561	2.16 (1.70 to 2.75)	4.03E-10
				95%	48	2,774	2.42 (1.79 to 3.27)	8.15E-09
				97.50%	30	1,381	2.98 (2.05 to 4.32)	8.74E-09
				99%	16	549	3.93 (2.38 to 6.47)	8.25E-08
		Quartile 3	from 58 to 63	80%	191	10,655	1.63 (1.38 to 1.93)	1.08E-08
				90%	128	5,295	2.18 (1.80 to 2.65)	2.24E-15
				95%	66	2,646	2.14 (1.66 to 2.75)	5.01E-09
				97.50%	36	1,320	2.29 (1.64 to 3.21)	1.28E-06
				99%	18	525	2.84 (1.78 to 4.54)	1.25E-05
		Quartile 4	from 63 to 71	80%	311	9,728	1.83 (1.60 to 2.10)	8.73E-19
				90%	182	4,838	2.04 (1.74 to 2.40)	3.15E-18
				95%	108	2,402	2.35 (1.92 to 2.87)	5.99E-17
				97.50%	63	1,192	2.68 (2.08 to 3.46)	3.65E-14
				99%	29	473	3.00 (2.07 to 4.35)	5.60E-09

Outcome	Sex Group	Age Group	Age Distribution	Percentile Thres.	N event	N total	HR (95% CI)	P-value
MI Incident	Men	All Ages	from 40 to 73	80%	1,843	33,698	2.12 (2.00 to 2.24)	1.83E-150
				90%	1,041	16,730	2.21 (2.07 to 2.37)	4.09E-117
				95%	570	8,316	2.34 (2.14 to 2.55)	1.16E-81
				97.50%	296	4,147	2.37 (2.11 to 2.66)	3.67E-47
				99%	139	1,639	2.77 (2.34 to 3.27)	2.65E-32
		Quartile 1	from 40 to 51	80%	300	8,722	3.17 (2.73 to 3.69)	1.80E-50
				90%	187	4,324	3.45 (2.92 to 4.09)	4.34E-47
				95%	106	2,150	3.56 (2.90 to 4.39)	3.24E-33
				97.50%	51	1,077	3.20 (2.41 to 4.26)	1.43E-15
				99%	25	427	3.74 (2.51 to 5.58)	1.01E-10
		Quartile 2	from 51 to 58	80%	420	8,160	2.28 (2.02 to 2.57)	1.35E-41
				90%	242	4,048	2.40 (2.08 to 2.77)	7.72E-34
				95%	137	2,008	2.60 (2.17 to 3.11)	9.26E-26
				97.50%	74	999	2.76 (2.18 to 3.49)	3.17E-17
				99%	34	395	3.14 (2.23 to 4.42)	4.88E-11
		Quartile 3	from 58 to 63	80%	484	8,273	1.82 (1.63 to 2.02)	1.25E-27
				90%	267	4,112	1.88 (1.65 to 2.15)	6.41E-21
				95%	139	2,051	1.88 (1.58 to 2.24)	1.18E-12
				97.50%	77	1,018	2.05 (1.63 to 2.58)	8.29E-10
				99%	36	402	2.38 (1.71 to 3.32)	2.69E-07
		Quartile 4	from 63 to 73	80%	658	8,526	1.95 (1.78 to 2.14)	3.40E-45
				90%	361	4,231	1.99 (1.77 to 2.23)	2.96E-32
				95%	197	2,099	2.08 (1.80 to 2.41)	1.46E-22
				97.50%	96	1,052	1.96 (1.60 to 2.41)	1.28E-10
				99%	48	412	2.46 (1.85 to 3.28)	7.41E-10

Supplemental Table 7: Net Reclassification Improvement of PRS_{CAD} for MI incidence, sex and age stratified. Net reclassification improvement for events (NRI Events), for non-events (NRI Non-Events) and overall net reclassification improvement (NRI) were determined with their 95% confidence intervals (95% CI), for a risk threshold of 2% ($\text{NRI}^{0.02}$) and for continuous NRI ($\text{NRI}^{>0}$) for all individuals, but also in women and in men. 95% CIs were calculated using bootstrapping method.

Categorized Net Reclassification Improvement ($\text{NRI}^{0.02}$)				
Sex Group	Age Group	$\text{NRI}^{0.02}$ Events	$\text{NRI}^{0.02}$ Non-Events	Overall $\text{NRI}^{0.02}$
All	All Ages	0.0669 (0.0535 to 0.0792)	-0.0217 (-0.0245 to -0.0180)	0.0452 (0.0333 to 0.0573)
	Quartile 1	0.190 (0.133 to 0.242)	-0.0326 (-0.0451 to -0.0211)	0.158 (0.111 to 0.199)
	Quartile 2	0.134 (0.0950 to 0.165)	-0.0544 (-0.0702 to -0.0327)	0.0794 (0.0527 to 0.104)
	Quartile 3	0.0245 (0.000711 to 0.0425)	-0.00615 (-0.0129 to 0.000821)	0.0183 (-0.00557 to 0.0365)
	Quartile 4	-0.00488 (-0.0288 to 0.0101)	0.0431 (0.0295 to 0.0637)	0.0383 (0.0200 to 0.0536)
Women	All Ages	0.0775 (0.0519 to 0.108)	-0.0159 (-0.0235 to -0.00828)	0.0616 (0.0416 to 0.0864)
	Quartile 1	0.0198 (0.000 to 0.0673)	-0.0000562 (-0.00138 to 0.0000817)	0.0197 (-0.0000817 to 0.0668)
	Quartile 2	0.0306 (0.00249 to 0.0686)	-0.00308 (-0.00702 to -0.00116)	0.0275 (0.000479 to 0.0625)
	Quartile 3	0.0772 (0.0199 to 0.132)	-0.0218 (-0.0427 to -0.00767)	0.0554 (0.00856 to 0.0923)
	Quartile 4	0.138 (0.086 to 0.184)	-0.0550 (-0.0700 to -0.0400)	0.0828 (0.0420 to 0.123)
Men	All Ages	-0.0187 (-0.0345 to -0.00366)	0.0689 (0.0548 to 0.0845)	0.0502 (0.0385 to 0.0634)
	Quartile 1	0.331 (0.277 to 0.385)	-0.131 (-0.150 to -0.107)	0.199 (0.157 to 0.248)
	Quartile 2	-0.0917 (-0.155 to -0.00357)	0.161 (0.0572 to 0.258)	0.0693 (0.0263 to 0.121)
	Quartile 3	-0.0522 (-0.0827 to -0.0313)	0.190 (0.136 to 0.237)	0.138 (0.101 to 0.162)
	Quartile 4	-0.0287 (-0.0435 to -0.0150)	0.103 (0.0723 to 0.141)	0.0744 (0.0540 to 0.104)
Continuous Net Reclassification Improvement ($\text{NRI}^{>0}$)				
All	All Ages	0.158 (0.140 to 0.177)	0.165 (0.152 to 0.178)	0.324 (0.297 to 0.349)
	Quartile 1	0.287 (0.223 to 0.346)	0.258 (0.230 to 0.290)	0.545 (0.470 to 0.627)
	Quartile 2	0.168 (0.126 to 0.211)	0.182 (0.149 to 0.212)	0.349 (0.288 to 0.413)
	Quartile 3	0.124 (0.0850 to 0.157)	0.143 (0.121 to 0.166)	0.267 (0.216 to 0.315)
	Quartile 4	0.155 (0.123 to 0.187)	0.147 (0.132 to 0.163)	0.302 (0.260 to 0.345)
Women	All Ages	0.137 (0.0989 to 0.176)	0.159 (0.127 to 0.191)	0.296 (0.235 to 0.356)
	Quartile 1	0.130 (0.00676 to 0.263)	0.182 (0.119 to 0.248)	0.313 (0.152 to 0.489)
	Quartile 2	0.118 (0.0317 to 0.205)	0.140 (0.0953 to 0.189)	0.258 (0.144 to 0.368)
	Quartile 3	0.130 (0.0520 to 0.198)	0.129 (0.0892 to 0.162)	0.258 (0.158 to 0.346)
	Quartile 4	0.147 (0.0937 to 0.199)	0.152 (0.123 to 0.182)	0.299 (0.230 to 0.360)
Men	All Ages	0.172 (0.148 to 0.191)	0.174 (0.162 to 0.187)	0.346 (0.314 to 0.372)
	Quartile 1	0.321 (0.262 to 0.388)	0.281 (0.251 to 0.313)	0.602 (0.525 to 0.683)
	Quartile 2	0.188 (0.139 to 0.243)	0.196 (0.168 to 0.224)	0.384 (0.315 to 0.458)
	Quartile 3	0.118 (0.0807 to 0.160)	0.152 (0.130 to 0.175)	0.271 (0.221 to 0.326)
	Quartile 4	0.150 (0.114 to 0.187)	0.155 (0.135 to 0.174)	0.305 (0.257 to 0.355)

Supplemental Table 8: Association between PRS_{CAD} and all-cause mortality in adjusted Cox regression models. Adjusted Cox regression models including age and the first ten genetic principal components (PCs) as covariates were obtained all individuals, but also in women and in men, by quartiles of age (Q1 to Q4) and for the whole group (all age). Data are presented as estimated hazard ratios (HR) for mortality per SD increase of PRS_{CAD} with their 95% confidence intervals (95% CI), and Cox regression P-value significance.

Outcome	Sex Group	Age Group	Age Window	Max-Min Difference in Age Group	N Event	N Total	Ratio N Event / 100% of Event	HR (95% CI)	P-value
Mortality	All	All Ages	from 40 to 73	33	23,982	403,422	100.00%	1.08 (1.06 to 1.09)	5.46E-30
		Quartile 1	from 40 to 51	11	1,841	100,856	7.68%	1.11 (1.06 to 1.16)	6.40E-06
		Quartile 2	from 51 to 58	7	3,831	100,855	15.97%	1.05 (1.01 to 1.08)	4.22E-03
		Quartile 3	from 58 to 63	5	6,557	100,855	27.34%	1.08 (1.05 to 1.10)	2.33E-09
		Quartile 4	from 63 to 73	10	11,753	100,856	49.01%	1.08 (1.06 to 1.10)	3.78E-17
	Women	All Ages	from 40 to 71	31	9,604	217,886	100.00%	1.04 (1.02 to 1.06)	3.17E-04
		Quartile 1	from 40 to 51	11	830	55,289	8.64%	1.07 (1.00 to 1.15)	4.99E-02
		Quartile 2	from 51 to 58	7	1,649	56,704	17.17%	0.95 (0.91 to 1.00)	5.41E-02
		Quartile 3	from 58 to 63	5	2,651	54,737	27.60%	1.04 (1.00 to 1.08)	5.92E-02
		Quartile 4	from 63 to 71	8	4,474	51,156	46.58%	1.07 (1.03 to 1.10)	2.67E-05
	Men	All Ages	from 40 to 73	33	14,378	185,536	100.00%	1.10 (1.09 to 1.12)	6.26E-32
		Quartile 1	from 40 to 51	11	1,011	45,567	7.03%	1.15 (1.08 to 1.22)	1.64E-05
		Quartile 2	from 51 to 58	7	2,182	44,151	15.18%	1.13 (1.08 to 1.18)	3.36E-08
		Quartile 3	from 58 to 63	5	3,906	46,118	27.17%	1.10 (1.07 to 1.14)	6.73E-10
		Quartile 4	from 63 to 73	10	7,279	49,700	50.63%	1.09 (1.07 to 1.12)	1.33E-13

Supplemental Table 9: Mortality in individuals at high genetic risk. Adjusted Cox regression results were obtained for individuals belonging above percentile thresholds (ranging between the 80th and the 99th percentile). Hazard ratios (HR) for mortality and their 95% confidence intervals (95% CI) were compared in all individuals, but also separately in women and in men, for all ages and for each quartile of age (Q1 to Q4).

Outcome	Sex Group	Age Group	Age Distribution	Percentile Thres.	N event	N total	HR (95% CI)	P-value
Mortality	All	All Ages	from 40 to 73	80%	5,251	75,434	1.14 (1.11 to 1.18)	1.22E-17
				90%	2,717	37,626	1.18 (1.13 to 1.23)	6.86E-16
				95%	1,392	18,780	1.21 (1.15 to 1.28)	3.49E-12
				97.50%	737	9,349	1.29 (1.20 to 1.39)	8.27E-12
				99%	302	3,733	1.33 (1.18 to 1.49)	1.07E-06
		Quartile 1	from 40 to 51	80%	444	19,728	1.28 (1.15 to 1.42)	7.83E-06
				90%	246	9,840	1.39 (1.22 to 1.59)	1.23E-06
				95%	137	4,906	1.54 (1.29 to 1.83)	1.36E-06
				97.50%	75	2,447	1.67 (1.33 to 2.11)	1.27E-05
				99%	28	981	1.52 (1.04 to 2.20)	2.85E-02
		Quartile 2	from 51 to 58	80%	850	19,321	1.15 (1.06 to 1.24)	3.81E-04
				90%	448	9,638	1.20 (1.09 to 1.33)	2.18E-04
				95%	219	4,824	1.17 (1.02 to 1.34)	2.73E-02
				97.50%	123	2,399	1.33 (1.11 to 1.59)	2.02E-03
				99%	51	958	1.39 (1.06 to 1.84)	1.89E-02
		Quartile 3	from 58 to 63	80%	1,400	18,771	1.09 (1.03 to 1.16)	3.43E-03
				90%	709	9,377	1.09 (1.01 to 1.18)	2.26E-02
				95%	376	4,667	1.17 (1.05 to 1.30)	3.18E-03
				97.50%	196	2,326	1.21 (1.05 to 1.40)	7.58E-03
				99%	81	928	1.27 (1.02 to 1.58)	3.32E-02
		Quartile 4	from 63 to 73	80%	2,589	17,583	1.15 (1.10 to 1.20)	4.87E-10
				90%	1,331	8,755	1.18 (1.11 to 1.25)	2.33E-08
				95%	681	4,362	1.20 (1.11 to 1.29)	5.37E-06
				97.50%	357	2,165	1.26 (1.13 to 1.40)	2.01E-05
				99%	153	856	1.33 (1.14 to 1.56)	4.13E-04

Outcome	Sex Group	Age Group	Age Distribution	Percentile Thres.	N event	N total	HR (95% CI)	P-value
Mortality	Women	All Ages	from 40 to 71	80%	2,044	41,534	1.09 (1.04 to 1.15)	3.46E-04
				90%	1,081	20,708	1.16 (1.09 to 1.24)	3.76E-06
				95%	559	10,336	1.20 (1.10 to 1.30)	3.78E-05
				97.50%	300	5,148	1.30 (1.16 to 1.45)	9.82E-06
				99%	119	2,060	1.28 (1.07 to 1.54)	6.83E-03
		Quartile 1	from 40 to 51	80%	198	10,860	1.27 (1.08 to 1.49)	3.72E-03
				90%	108	5,421	1.36 (1.11 to 1.66)	3.19E-03
				95%	59	2,706	1.46 (1.12 to 1.90)	5.03E-03
				97.50%	35	1,348	1.73 (1.23 to 2.43)	1.48E-03
				99%	15	538	1.84 (1.10 to 3.07)	1.92E-02
		Quartile 2	from 51 to 58	80%	335	11,006	1.02 (0.906 to 1.15)	7.25E-01
				90%	177	5,494	1.09 (0.930 to 1.27)	2.96E-01
				95%	87	2,749	1.05 (0.849 to 1.31)	6.35E-01
				97.50%	49	1,369	1.20 (0.900 to 1.59)	2.16E-01
				99%	18	550	1.10 (0.693 to 1.75)	6.81E-01
		Quartile 3	from 58 to 63	80%	545	10,403	1.04 (0.946 to 1.14)	4.24E-01
				90%	278	5,196	1.06 (0.936 to 1.20)	3.59E-01
				95%	148	2,589	1.14 (0.963 to 1.34)	1.29E-01
				97.50%	77	1,292	1.19 (0.949 to 1.49)	1.32E-01
				99%	31	517	1.19 (0.838 to 1.70)	3.27E-01
		Quartile 4	from 63 to 71	80%	969	9,263	1.12 (1.04 to 1.20)	1.75E-03
				90%	519	4,597	1.21 (1.10 to 1.32)	5.46E-05
				95%	268	2,290	1.23 (1.09 to 1.40)	8.19E-04
				97.50%	146	1,133	1.34 (1.14 to 1.58)	4.97E-04
				99%	58	454	1.31 (1.01 to 1.69)	4.30E-02

Outcome	Sex Group	Age Group	Age Distribution	Percentile Thres.	N event	N total	HR (95% CI)	P-value
Mortality	Men	All Ages	from 40 to 73	80%	3,216	33,892	1.18 (1.13 to 1.22)	3.44E-16
				90%	1,642	16,912	1.19 (1.13 to 1.26)	2.05E-11
				95%	830	8,447	1.21 (1.13 to 1.30)	9.32E-08
				97.50%	442	4,197	1.30 (1.18 to 1.43)	7.26E-08
				99%	184	1,672	1.35 (1.17 to 1.56)	4.62E-05
		Quartile 1	from 40 to 51	80%	247	8,867	1.30 (1.12 to 1.50)	4.06E-04
				90%	137	4,420	1.42 (1.19 to 1.70)	1.35E-04
				95%	77	2,202	1.58 (1.25 to 2.00)	1.13E-04
				97.50%	40	1,100	1.62 (1.18 to 2.23)	2.66E-03
				99%	15	441	1.48 (0.89 to 2.47)	1.30E-01
		Quartile 2	from 51 to 58	80%	512	8,319	1.23 (1.12 to 1.36)	3.44E-05
				90%	272	4,144	1.29 (1.14 to 1.47)	8.27E-05
				95%	131	2,077	1.23 (1.03 to 1.47)	2.19E-02
				97.50%	77	1,027	1.47 (1.17 to 1.85)	8.50E-04
				99%	33	409	1.59 (1.13 to 2.24)	8.33E-03
		Quartile 3	from 58 to 63	80%	857	8,367	1.13 (1.05 to 1.22)	1.72E-03
				90%	427	4,185	1.11 (1.00 to 1.23)	4.26E-02
				95%	228	2,078	1.19 (1.04 to 1.36)	1.03E-02
				97.50%	117	1,036	1.22 (1.02 to 1.47)	3.34E-02
				99%	51	411	1.34 (1.01 to 1.76)	3.96E-02
		Quartile 4	from 63 to 73	80%	1,633	8,307	1.17 (1.11 to 1.24)	2.41E-08
				90%	823	4,147	1.16 (1.08 to 1.25)	4.55E-05
				95%	428	2,057	1.20 (1.09 to 1.32)	3.09E-04
				97.50%	217	1,026	1.22 (1.06 to 1.39)	4.30E-03
				99%	95	402	1.34 (1.10 to 1.64)	4.35E-03