Previously infected vaccinees broadly neutralize SARS-CoV-2 variants

Hans C. Leier^{1,4}, Timothy A. Bates^{1,4}, Zoe L. Lyski¹, Savannah K. McBride¹, David X. Lee¹,
Felicity J. Coulter¹, James R. Goodman², Zhengchun Lu¹, Marcel E. Curlin^{3,*}, William B.
Messer^{1,3,*}, Fikadu G. Tefesse^{1,*}

Affiliations

¹ Department of Molecular Microbiology & Immunology, Oregon Health & Science University; Portland, OR 97239, United States.

² Medical Scientist Training Program, Oregon Health & Science University; Portland, OR 97239, United States.

³ Division of Infectious Diseases, Oregon Health & Science University; Portland, OR 97239, United States.

⁴ These authors contributed equally to this work.

***Correspondence to:** Fikadu G. Tefesse tefesse@ohsu.edu, William B. Messer messer@ohsu.edu, Marcel E. Curlin, curlin@ohsu.edu

Abstract

We compared the serum neutralizing antibody titers before and after two doses of the BNT162b2 COVID-19 vaccine in ten individuals who recovered from SARS-CoV-2 infection prior to vaccination to 20 individuals with no history of infection, against clinical isolates of B.1.1.7, B.1.351, P.1, and the original SARS-CoV-2 virus. Vaccination boosted pre-existing levels of anti-SARS-CoV-2 spike antibodies 10-fold in previously infected individuals, but not to levels significantly higher than those of uninfected vaccinees. However, neutralizing antibody titers increased in previously infected vaccinees relative to uninfected vaccinees against every variant tested: 5.2-fold against B.1.1.7, 6.5-fold against B.1.351, 4.3-fold against P.1, and 3.4-fold against original SARS-CoV-2. Our study indicates that a first-generation COVID-19 vaccine provides broad protection from SARS-CoV-2 variants in individuals with previous infection.

42

43

44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62

63
64

Main

In recent months, multiple coronavirus disease 2019 (COVID-19) vaccine candidates have successfully concluded phase 3 trials¹, with the three candidates authorized for emergency use by the U.S. Food and Drug Administration reporting efficacies of 95% (BNT162b2 [Pfizer-BioNTech]), 94% (mRNA-1273 [Moderna]), and 66% (Ad26.COV2.S [Janssen])²⁻⁴. When combined with the substantial portion of many communities estimated to have gained natural immunity through infection with severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2)⁵, the rollout of safe and effective vaccines has raised the possibility that high levels of population immunity could soon be reached. Clouding this prospect is the emergence and global spread of SARS-CoV-2 variants of concern (VOCs), such as those first identified in the United Kingdom (lineage B.1.1.7)⁶, South Africa (B.1.351)⁷, Brazil (P.1)⁸, and California (B.1.429)⁹. Most VOCs possess partially overlapping combinations of spike mutations that enhance binding to the SARS-CoV-2 cellular receptor angiotensin-converting enzyme 2 (ACE2), increasing transmissibility¹⁰ (Supplementary Table 1). More concerning has been the emergence of spike mutations with the potential to escape neutralizing antibodies raised against earlier lineages of SARS-CoV-2 through infection with an original lineage or by first-generation COVID-19 vaccines¹¹⁻¹³. Recent population studies have validated these findings, showing surges of reinfections in regions with extensive transmission of B.1.35^{17,14} and P.1⁸, and large declines in vaccine efficacy against B.1.351¹⁴⁻¹⁶.

To investigate whether vaccination of individuals previously infected by SARS-CoV-2 confers greater protection from VOCs than vaccination of individuals with no evidence of

previous infection, in a cohort of BNT162b2 vaccinees (Supplementary Table 2) we identified a group of 10 study participants who had received a positive COVID-19 PCR test result prior to vaccination, along with an age- and sex-balanced group of 20 participants who had not. While vaccination of previously infected individuals boosted the 50% maximal effective concentration (EC₅₀) of antibodies against the immunodominant SARS-CoV-2 spike receptor-binding domain (RBD) ten-fold (pre-vaccination geometric mean titer [GMT], 82.15; post-vaccination GMT, 823.3), the vaccine-elicited antibody titers of uninfected individuals (GMT, 699.5) were not significantly lower (Fig. 1A). Similarly, post-vaccination levels of RBD-binding IgG (Fig. 1B) and IgA (Fig. 1C) did not differ significantly between the two groups.

We then measured the pre- and post-vaccination neutralizing activity of the two groups of sera against an early SARS-CoV-2 isolate (USA-WA1/2020) and isolates of B.1.1.7, B.1.351, and P.1 (Fig. 2). Pre-vaccination sera from previously infected participants provided higher levels of neutralization against USA-WA1/2020 (GMT, 39.0) than against the three VOCs (GMT, 25.7 for B.1.1.7; GMT<20 for B.1.351; GMT, 31.2 for P.1), consistent with previous reports of convalescent sera^{12,17}. Similarly, post-vaccination sera from uninfected participants showed greater neutralization of USA-WA1/2020 than of the VOCs (GMT, 578.6 for USA-WA1/2020; 223.0 for B.1.1.7; 47.5 for B.1.351; 171.9 for P.1). However, post-vaccination serum from previously infected individuals possessed significantly higher neutralizing activity against every SARS-CoV-2 lineage relative to post-vaccination serum from uninfected participants: neutralizing antibody titers increased by a factor of 3.5 against USA-WA1/2020 (95% confidence interval [CI], 2.8 to 4.0); by a factor of 5.2 against B.1.1.7 (95% CI, 2.37 to 9.8); by a factor of 6.5 against B.1.351 (95% CI, 3.4 to 12.3); and by a factor of 4.3 against P.1 (95% CI, 2.8 to 6.5). Notably, there was no

significant difference ($P=0.2736$, Wilcoxon rank-sum test) between the post-vaccination neutralizing antibody titers of previously infected participants against B.1.351 (GMT, 307.3; 95% CI, 91.0 to 1038) and those of uninfected participants against USA-WA1/2020 (GMT, 578.6; GMT, 332.5 to 1007), suggesting that first-generation COVID-19 vaccines could retain near-complete efficacy against even the most resistant VOCs when administered following natural infection.

Overall, our findings provide important evidence for broad and potent neutralizing antibody responses against emerging SARS-CoV-2 variants, even with exposure to only wildtype SARS-CoV-2 antigen. This reinforces a recent report that natural infection with B.1.351 elicits a similar cross-reactive neutralizing antibody response against B.1.351, P.1, and original SARS-CoV-2¹⁸. While these and other laboratory results must be validated by ongoing population-level studies, they indicate a novel role for COVID-19 vaccines in protecting hard-hit populations from future waves of the pandemic.

References

1. G. Forni, A. Mantovani, COVID-19 vaccines: where we stand and challenges ahead. *Cell Death Differ.* **28**, 626-639 (2021).
2. L. R. Baden *et al.*, Efficacy and Safety of the mRNA-1273 SARS-CoV-2 Vaccine. *N. Eng. J. Med.* **384**, 403-416 (2021).
3. F. P. Polack *et al.*, Safety and Efficacy of the BNT162b2 mRNA Covid-19 Vaccine. *N. Eng. J. Med.* **383**, 2603-2615 (2020).
4. J. Sadoff *et al.*, Safety and Efficacy of Single-Dose Ad26.COV2.S Vaccine against Covid-19. *N. Eng. J. Med.*, <https://doi.org/10.1056/NEJMoa2101544> (2021).
5. X. Chen *et al.*, Serological evidence of human infection with SARS-CoV-2: a systematic review and meta-analysis. *Lancet Glob. Health* **9**, E598-E609 (2021).
6. N. G. Davies *et al.*, Estimated transmissibility and impact of SARS-CoV-2 lineage B.1.1.7 in England. *Science*, <https://doi.org/10.1126/science.abg3055> (2021).
7. H. Tegally *et al.*, Emergence and rapid spread of a new severe acute respiratory syndrome-related coronavirus 2 (SARS-CoV-2) lineage with multiple spike mutations in South Africa. medRxiv, 2020.12.21.20248640 (2020).
8. E. C. Sabino *et al.*, Resurgence of COVID-19 in Manaus, Brazil, despite high seroprevalence. *Lancet* **397**, 452-455 (2021).
9. W. Zhang *et al.*, Emergence of a Novel SARS-CoV-2 Variant in Southern California. *JAMA* **325**, 1324-1326 (2021).

- 130 10. J. A. Plante *et al.*, The Variant Gambit: COVID's Next Move. *Cell Host Microbe* **29**,
131 508-515 (2021).
- 132 11. R. E. Chen *et al.*, Resistance of SARS-CoV-2 variants to neutralization by monoclonal
133 and serum-derived polyclonal antibodies. *Nat. Med.* **27**, 717-726 (2021).
- 134 12. M. Hoffmann *et al.*, SARS-CoV-2 variants B.1.351 and P.1 escape from neutralizing
135 antibodies. *Cell*, <https://doi.org/10.1016/j.cell.2021.03.036> (2021).
- 136 13. P. Wang *et al.*, Antibody Resistance of SARS-CoV-2 Variants B.1.351 and B.1.1.7.
137 *Nature*, <https://doi.org/10.1038/s41586-021-03398-2> (2021).
- 138 14. E. Mahase, Covid-19: Novavax vaccine efficacy is 86% against UK variant and 60%
139 against South African variant. *BMJ*, n296 (2021).
- 140 15. T. Kustin *et al.*, Evidence for increased breakthrough rates of SARS-CoV-2 variants of
141 concern in BNT162b2 mRNA vaccinated individuals. medRxiv, 2021.04.06.21254882
142 (2021).
- 143 16. S. A. Madhi *et al.*, Efficacy of the ChAdOx1 nCoV-19 Covid-19 Vaccine against the
144 B.1.351 Variant. *N. Eng. J. Med.*, <https://doi.org/10.1056/NEJMoa2102214> (2021).
- 145 17. T. A. Bates *et al.*, Neutralization of SARS-CoV-2 variants by convalescent and
146 vaccinated serum. medRxiv, 2021.04.04.21254881 (2021).
- 147 18. T. Moyo-Gwete *et al.*, Cross-Reactive Neutralizing Antibody Responses Elicited by
148 SARS-CoV-2 501Y.V2 (B.1.351). *N. Eng. J. Med.*,
149 <https://doi.org/10.1056/NEJMc2104192> (2021).
- 150 19. T.A. Bates *et al.*, Cross-reactivity of SARS-CoV structural protein antibodies against

151 SARS-CoV-2. *Cell Rep.* **34**, 108737 (2021).

152 20. J.B. Case *et al.*, Growth, detection, quantification, and inactivation of SARS-CoV-2.

153 *Virology* **548**, 39-48 (2020)

154

155

156

157

158

159

160

161

162

163

164

165

166

167

168

169 **Acknowledgments**

170 The authors thank the clinical staff at Oregon Health & Science University who assisted in
171 enrolling study participants and collecting the serum samples used in this report.

172 **Funding**

- 173 M.J. Murdock Charitable Trust
- 174 National Institutes of Health training grant T32AI747225
- 175 Oregon Health & Science University Innovative IDEA grant 1018784
- 176 National Institutes of Health grant R01AI145835

177 **Competing interests**

178 Authors declare that they have no competing interests.

179 **Data and materials availability**

180 All data are available in the main text or the supplementary materials.

Figure 1. Anti-SARS-CoV-2 antibody levels elicited by vaccination and natural infection. (A) Half maximal effective concentration (EC₅₀) of total pan-Ig antibodies specific to the spike RBD were measured by enzyme-linked immunosorbent assay (ELISA) in serum collected from donors previously infected with SARS-CoV-2 pre- and post-vaccination with the BNT162b2 vaccine. (B) EC₅₀ of RBD-binding IgA. (C) EC₅₀ of RBD-binding IgG. Statistical comparisons were made using the Wilcoxon rank-sum test. LOD denotes limit of detection.

● Individuals with no history of infection (n=20) ■ Previously infected individuals (n=10)

Figure 2. Immunity from natural SARS-CoV-2 infection boosted by subsequent

vaccination broadens protection against variants of concern. Neutralization of

SARS-CoV-2 variants by pre-and post-vaccination sera collected from previously

infected and naïve individuals. Shown are the results of a 50% focus reduction

neutralization testing (FRNT₅₀) for an early SARS- CoV-2 isolate USA-WA1/2020 (A)

and VOC clinical isolates of B.1.1.7 (B), B.1.351 (C), and P.1 (D). Horizontal bars

represent geometric mean titer and I bars represent 95% confidence intervals.

Statistical comparisons were made using the Wilcoxon rank-sum test. LOD denotes limit

of detection.

Supplementary Materials for

Previously infected vaccinees broadly neutralize SARS-CoV-2 variants

Hans C. Leier, Timothy A. Bates, Zoe L. Lyski, Savannah K. McBride, David X. Lee,
Felicity J. Coulter, James R. Goodman, Zhengchun Lu, Marcel E. Curlin, William B.
Messer, Fikadu G. Tafesse

Contains:

Materials and methods

Supplementary Tables 1-2

Materials and Methods

Cohort

This study was conducted in accordance with the Oregon Health & Science University (OHSU) Institutional Review Board (IRB#00022511). Study participants were enrolled at OHSU immediately after receiving the first dose of the BNT162b2 (Pfizer-BioNTech) COVID-19 vaccine. After receiving informed consent, 4-6 mL whole blood was drawn and centrifuged 10 minutes at 1000 x g. A second blood sample was obtained at least 14 days after participants received their second vaccine dose. Serum was stored at -20°C until heat inactivation in preparation for FRNTs and ELISAs. n=10 participants were identified with a positive COVID-19 PCR test prior to vaccination and n=20 age- and sex-balanced participants who did not have a history of positive PCR test or RBD-binding antibodies in an ELISA. See Supplementary Table 1 for more details of the cohort.

ELISA

The 96-well ELISA plates (Corning, Cat# 3590) were coated with recombinant RBD protein prepared as previously described¹⁹ at a concentration of 1 ug/mL in PBS and incubated overnight at 4°C. Coating antigen was removed, plates were washed once with PBST containing 0.05% Tween (wash buffer) and blocked for 1 hour at RT with 5% milk prepared in PBST containing 0.05% Tween (dilution buffer). 100 mL of 1:20 (pre-vaccine), or 1:200 (post- vaccine) dilution of serum in dilution buffer was added to each starting well. Three-fold serial dilutions were performed in dilution buffer. Plates were incubated at room temperature for 1 hour. The plates were washed 3 times with wash

buffer and 100 mL of 1:10,000 dilution of anti-human GOXHU IGG/A/M HRP (Invitrogen, Cat# A18847) detection antibody was added and incubated at room temperature for 1 hour. After washing the plates 3 times with wash buffer, 100 mL of colorimetric detection reagent containing 0.4 mg/ml o-phenylenediamine and 0.01% hydrogen 45 peroxide in 0.05 M citrate buffer (pH 5) were added and the reaction was stopped after 20 minutes by the addition of 100 mL 1 M HCl. Optical density (OD) at 492 nm was measured using a CLARIOstar plate reader. Plates were normalized, background removed, and endpoint antibody titers determined by the highest dilution with a positive signal, defined as 4-fold above background. Final endpoint titer values below the detection limit of 20 were set to 19.

Viruses

SARS-CoV-2 isolates were obtained from BEI Resources (Supplementary Table 2): Isolate USA/CA_CDC_5574/2020 [B.1.1.7] (BEI Resources NR-54011); Isolate hCoV-54 19/South Africa/KRISP-K005325/2020 [B.1.351] (BEI Resources NR-54009); Isolate hCoV-19/Japan/TY7-503/2021 [P.1] (BEI Resources NR-54982); and Isolate USA-WA1/2020 [early isolate] (BEI Resources NR-52281). Isolates were propagated and titrated in Vero E6 cells as previously described²⁰.

FRNT

Serial dilutions of patient sera and virus neutralization were carried out in biological duplicate in a 96-well plate format. Briefly, each sample was added in duplicate 1:10 to dilution media (Opti-MEM, 2% FBS), and four-fold serial dilutions were made for a range of 1:10 – 1:2560. An equal volume of dilution media containing 100 FFU of SARS-CoV-

2 was added to each well (final dilutions of sera, 1:20–1:5120) and incubated 1 h at 37°C. The virus-serum mixtures were then added to monolayers of Vero E6 cells, incubated with occasional rocking 1 h at 37°C, and covered with overlay media (Opti-MEM, 2% FBS, 1% methylcellulose). Overlay media was then removed, and plates were fixed for 1 h in 4% paraformaldehyde in PBS. Foci were developed as described. Plates were imaged with a CTL Immunospot Analyzer, then foci were counted using CTL ImmunoSpot (7.0.26.0) Percent neutralization values for FRNT50 were compiled and analyzed using python (v3.7.6) with numpy (v1.18.1), scipy (v1.4.1), and pandas (v1.0.1) data analysis libraries. Replicate data were fit with a three-parameter logistic model and final FRNT50 values below the lowest dilution of 20 were set to 19 while values above the maximum dilution of 5120 were set to 5121. FRNT50 curves were plotted using python with the Matplotlib (v3.1.3) data visualization library.

Statistical analysis

Aggregated ELISA endpoint titers and FRNT50 values were analyzed in GraphPad Prism (v9.0.2). Comparisons were performed using the Wilcoxon rank-sum test. Reported group averages are geometric means, with error bars representing 95% confidence intervals.

Supplementary Table 1. List of mutations in B.1.1.7, B.1.351 and P.1 SARS-CoV-89 2
clinical isolates.*

Lineage	GISAID Clade	GISAID ID	Spike mutations (RBD highlighted)	Non-Spike mutations
B.1.1.7	GR	EPI_ISL_683466	H69del, V70del, Y145del, N501Y , A570D, D614G, P681H, T716I, S982A, D1118H	N D3L, N G204R, N R203K, N S235F, NS8 Q27stop, NS8 R52I, NS8 Y73C, NSP3 A890D, NSP3 A1305V, NSP3 I1412T, NSP3 T183I, NSP6 F108del, NSP6 G107del, NSP6 S106del, NSP12 P323L, NSP13 K460R, NSP14 E347G
B.1.351	GH	EPI_ISL_678570	D80A, D215G, L242del, A243del, L244del, K417N , E484K , N501Y , D614G, A701V	E P71L, N T205I, NS3 Q57H, NS3 S171L, NSP2 T85I, NSP3 K837N, NSP5 K90R, NSP6 F108del, NSP6 G107del, NSP6 S106del, NSP12 P323L
P.1			D138Y, D614G, K417T , E484K , N501Y , G181V, H655Y, L18F, P26S, R190S, T20N, T1027I, V1176F	N G204R, N P80R, N R203K, NS3 S253P, NS8 E92K, NSP3 K977Q, NSP3 S370L, NSP3 T186A, NSP6 F108del, NSP6 G107del, NSP6 S106del, NSP12 P323L, NSP13 E341D

*Obtained from BEI Resources

Supplementary Table 2. Description of cohort.

	Previously infected (n=10)	No history of infection (n=20)
Sex – no. (%)		
Female	6 (60)	15 (75)
Male	4 (40)	5 (25)
Median age – yrs (range)	36.5 (23-61)	44 (23-60)
Median time between vaccine doses – days (range)	22 (20-25)	21 (21-32)
Median time from second vaccine dose to second serum sample – days (range)	18 (14-28)	17 (14-28)
Median time from positive COVID-19 PCR test to first vaccine dose – days (range)	52 (13-302)	NA