EVIDENCE FOR BIOLOGICAL AGE ACCELERATION AND TELOMERE SHORTENING IN COVID19 SURVIVORS

Alessia Mongelli¹, Veronica Barbi¹, Michela Gottardi Zamperla¹, Sandra Atlante¹, Luana Forleo¹, Marialisa Nesta², Massimo Massetti², Alfredo Pontecorvi², Simona Nanni², Antonella Farsetti³, Oronzo Catalano⁴, Maurizio Bussotti⁵, Laura Dalla Vecchia⁵, Tiziana Bachetti⁶, Fabio Martelli⁷, Maria Teresa La Rovere^{6,8}, Carlo Gaetano^{1,8,*}.

1. Laboratory of Epigenetics, Istituti Clinici Scientifici Maugeri IRCCS, Via Maugeri 10, 27100, Pavia, Italy
2. Università Cattolica del Sacro Cuore; Foundation "Policlinico Universitario A. Gemelli IRCCS", Rome, Italy.
3. Institute for Systems Analysis and Computer Science "A. Ruberti" (IASI), National Research Council (CNR) 00185, Rome, Italy.
4. Cardiac rehabilitation Unit, Istituti Clinici Scientifici Maugeri IRCCS, Via Maugeri 10, 27100 Pavia, Italy
5. Cardiorespiratory Rehabilitation Department, IRCCS Maugeri Clinical Scientific Institutes, Milan.
6. Scientific Direction, Istituti Clinici Scientifici Maugeri IRCCS, Via Maugeri 4, 27100 Pavia, Italy.
7. Laboratory of Molecular Cardiology, Policlinico San Donato IRCCS, San Donato Milanese, 20097 Milan, Italy.
8. Department of Cardiology, Istituti Clinici Scientifici Maugeri IRCCS, Montescano (Pavia), 27040 Montescano (Pavia), Italy.

*Corresponding Carlo Gaetano, Laboratory of Epigenetics, Istituti Clinici Scientifici Maugeri IRCCS, Via Maugeri 10, 27100, Pavia, Italy ; carlo.gaetano@icsmaugeri.it; ph.00390382592262.

ABSTRACT

Introduction & Background: the SARS-CoV-2 infection determines the COVID19 syndrome characterized, in the worst cases, by severe respiratory distress, pulmonary and cardiac fibrosis, inflammatory cytokines release, and immunosuppression. This condition has led to the death of about 2.15% of the total infected world population so far. Among survivors, the presence of the so-called persistent post-COVID19 syndrome (PPCS) is a common finding. In patients who survived the SARS-CoV-2 infection, overt PPCS presents one or more symptoms such as fatigue, dyspnea, memory loss, sleep disorders, and difficulty concentrating. The pathophysiology of PPCS is currently poorly understood, and whether epigenetic mechanisms are involved in this process is unexplored.

Methods & Results: In this study, a cohort of 117 COVID19 survivors (post-COVID19) and 144 non-infected volunteers (COVID19-free) were analyzed using pyrosequencing of defined CpG islands previously identified as suitable for biological age determination. Besides, telomere length (TL) and ACE2 and DPP-4 receptor expression were determined. The results show a consistent biological age increase in the post-COVID19 population ($58,44 \pm 14,66$ ChronoAge Vs. $67,18 \pm 10,86$ BioAge, $P < 0,0001$), determining a DeltaAge acceleration of $10,45 \pm 7,29$ years ($+5.25$ years above range of normality) compared to $3,68 \pm 8,17$ years for the COVID19-free population ($P < 0,0001$). A significant telomere shortening parallels this finding in the post-COVID19 cohort

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

compared to COVID19-free subjects (post-COVID19 TL: $3,03 \pm 2,39$ Kb vs. COVID19-free: $10,67 \pm 11,69$ Kb; $P < 0,0001$). Additionally, ACE2 expression was decreased in post-COVID19 patients compared to COVID19-free, while DPP-4 did not change.

Conclusion: In light of these observations, we hypothesize that some epigenetic alterations are associated with the post-COVID19 condition, particularly in the younger (< 60 years). Although the consequences of such modifications on the long-term clinical outcome remain unclear, this finding might help indicating a direction to investigate the pathophysiology at the onset of the persistent post-COVID19 syndrome.

Keywords: biological age, COVID19, post-COVID19, telomeres, epigenetics, DNA methylation, ACE2, DPP-4, DeltaAge.

Running title: Biological clock acceleration in younger post-COVID19 survivors.

BACKGROUND

In the last trimester of 2019, a new severe acute respiratory syndrome (SARS) outbreak emerged due to a new Coronavirus (SARS-CoV-2) diffusion determining the COVID19 syndrome. At the time of this writing, the virus infected about 143 million worldwide, causing so far > 3 million deaths (source: <https://coronavirus.jhu.edu/map.html>). More than 122 million survived the COVID19 syndrome (source: <https://www.statista.com/statistics/1087466/covid19-cases-recoveries-deaths-worldwide/>). Not surprisingly, nowadays, more and more frequently, survivors are coming to clinicians' attention for the presence of post-COVID19 symptoms. Among others, the most commonly reported are fatigue, dyspnoea, memory loss, sleep disorders, and difficulty concentrating (1), which recently contributed to defining the so-called persistent post-COVID19 syndrome (PPCS) (1).

It has been observed that SARS-CoV-2 infected people who developed the adult respiratory distress syndrome (ARDS) also accumulated an excessive deposition of extracellular matrix in the alveolar compartment, causing pulmonary function's worsening (2). Fibrosis accumulation has also been observed in the cardiovascular and nervous systems. The infection frequently caused increased circulating Troponin T and Brain Natriuretic Peptides, suggesting the presence of myocardium damage with possible activation of a remodeling process (3). Besides, a compromised heart function with reduced contractility has often been observed (4). Other effects were on the vascular system and included altering the coagulation cascade and clotting (5)(6). In this case, the fibrinogen pathway's fast activation enhanced the risk of thrombotic events, leading to acute stroke and pulmonary embolism (7)(6). Interestingly, coagulation problems have been seen in post-COVID survivors and in PPCS patients, in which anticoagulants are routinely prescribed.

Similar to what was observed in 2003 SARS and the Middle East respiratory syndrome (MERS) epidemics, in COVID19 survivors, an increase of psychological disorders that include depression, posttraumatic stress, and anxiety has been observed (8). Similar disorders are often present in the general population exposed to stress determined by sociological factors often linked to economic damage or the sudden loss of some beloved ones (8). Remarkably, during the SARS or MERS epidemics, neuropsychiatric symptoms have been reported in hospital workers and caregivers (9). A similar situation has been reported in hospital workers assisting COVID19 patients (10).

Despite all the variety and significance of the symptoms reported by numerous COVID19 survivors with or without PPCS, valuable biomolecular markers to monitor this condition are still lacking.

Upon SARS-CoV-2 infection, the angiotensin-converting enzyme 2 (ACE2) expression level in the vascular system tends to decrease (11). This enzyme is involved in regulating the renine-angiotensin system (RAS); ACE2 contrasts the activity of the related angiotensin-converting enzyme (ACE) by converting the Angiotensin-II in Angiotensin (1-7) (12)(13). A low expression of ACE2 causes an accumulation of Angiotensin II, which may associate with exacerbating conditions leading to respiratory distress, hypertension, arrhythmia, cardiac hypertrophy, left ventricular function failure, atherosclerosis, and aortic aneurysms (13)(14). Moreover, ACE2 is negatively correlated with aging; it is relatively abundant in young and healthy people with significantly less risk of CVDs, while a lower quantity has been observed in the elderly (15).

The dipeptidyl-peptidase IV (DPP-4) is the receptor of MERS coronavirus (MERS-CoV) and has been reported in some cases to function as a co-receptor of SARS-CoV-2 (16). DPP4 expression increases on the surface of senescent cells (17), and its transmembrane form can cleave many molecules such as chemokines, neuropeptides, and incretin hormones. Because DPP-4 inactivates incretin hormones, high levels of DPP-4 have been observed in patients with type 2 diabetes mellitus and cardiovascular diseases. DPP-4 inhibitors have been used to treat T2DM, cardiac ischemia, and systolic dysfunction (18) (19). Some evidence suggests that DPP-4 inhibitors might reduce the coronavirus's entrance into the airway, envisaging an additional therapeutic tool for COVID19 treatment (20). Whether the level of ACE2 and DPP4 in peripheral blood may represent valuable biomarkers to monitor recovery from COVID19 or the onset of PPCS is unclear.

In humans, telomere shortening is associated *in vivo* with the aging process, and, *in vitro*, it characterizes the cellular replicative senescence (21). Telomeres possess properties that make them suitable as biomarkers in several diseases or conditions, including cancer, CVDs, or aging (22) (23). The inverse correlation between telomere length (TL) and chronological age has been used for age prediction (24). Interestingly, among individuals infected by Sars-CoV-2, a reduced TL has been associated with the risk of developing more severe symptoms suggesting that TL at the moment of the infection might influence the clinical outcome (25). At present, little is known about the telomere dynamics during Sars-CoV-2 infection and in COVID19 survivors and whether this parameter might help predict the risk of developing PPCS.

In recent years, several studies aimed to identify biological or molecular markers of aging that correlate with chronological age and could be helpful to estimate the biological vs. chronological age (26). Some of these parameters have been defined based on modifications of the DNA methylome that correlate with the chronological age and might be used in age prediction models to define the biological age molecularly: the so-called DNAmAge (28). Many of these studies focalized on healthy or diseased individuals for forensic or public health problems (27) (28). Since the seminal Horvath's work (29), several methods have been developed to estimate variation in methylation levels in selected DNA CpGs. These approaches apply to determining DNAmAge and have been used to emphasize the difference with chronological age: the so-called DeltaAge. Some methods are based on evaluating many CpGs, explored using a genome-wide array or next-generation sequencing technologies. However, others have been developed, taking into account a reduced number of CpGs analyzed by pyrosequencing (30)(31)(32). All systems are based on DNA methylation values

obtained from whole blood samples. Because of their practicality, several pyrosequencing blood-based age-prediction models became recently popular, especially for forensic applications. These simplified methods have the additional advantage of being rapid and suitable to most laboratory settings without requiring bioinformatics (24)(33). A recent comparative study (34) analyzed the performance of some of these "reductionist" methods and found that the algorithm proposed by Bekaert B. et al., among others, worked well for biological age prediction in young and old subjects. This algorithm considers a prediction result correct for individuals aged 60 or higher when the predicted age matches the chronological age within a ± 5.2 years range (24)(35). Considering that most post-COVID19 subjects fall within age 50 to 60 or higher, this method has been deemed suitable for the present study (24).

All the procedures to determine the epigenetic biological clock estimate DNAmAge and DeltaAge. A positive DeltaAge is considered an acceleration of the biological clock, while a negative DeltaAge indicates a bioage younger than the chronological one. In some studies, this parameter turned helpful to evaluate the risk of the onset of cardiovascular and neurodegenerative diseases, cancer, and the occurrence of death by all-cause (36).

In infectious diseases, the application of these methods is still limited. However, a DeltaAge acceleration has been observed in people infected by the Human Immunodeficiency Virus (HIV), the Cytomegalovirus, or bacteria such as the *Helicobacter pylori* (36). Besides, in post-mortem brain tissue, the DNAmAge of chronically HIV-positive individuals was higher than negative controls. Interestingly, a partial reversion of the accelerated DNAmAge has been recently observed following antiretroviral therapy (37)(38). Hence, HIV infection might change the epigenome landscape, potentially enhancing the risk of developing age-related diseases such as neurocognitive disorders (39). Similarly, in people infected by Cytomegalovirus, DNA methylation analyses performed in circulating leucocytes revealed an increased DeltaAge (40). The long-term consequences of these epigenomic alterations remain to be ascertained.

The present study investigates whether in COVID19 survivors there is a DNAmAge alteration and a DeltaAge acceleration, which, in association with other molecular parameters such as the telomere length and ACE2 expression in peripheral blood, might typify a set of biomarkers valuable in other and future studies exploring the risk of PPCS-associated pathophysiological manifestations.

RESULTS.

Evaluation of DNAmAge and DeltaAge in COVID19 survivors.

A cohort of 117 COVID19 survivors came to the attention of our physicians. Upon approval by the Ethical Committee and informed consent signing, peripheral blood was collected in EDTA vacutainers. A group of 144 age- and sex-matched COVID19-free volunteers with some risk factors partially overlapping with the post-COVID19 patients were recruited among the hospital workers and non-COVID19 patients (see Table I). Genomic DNA was extracted from the whole blood by a robotized station as described in methods. After bisulfite conversion and PCR amplification, pyrosequencing was performed. DNAmAge calculations were completed according to Bekaert et al. (24). Results indicate that the y-axis intercept differs significantly between the COVID19-free (Fig. 1A) and the post-COVID19 (Fig. 1B) population. The post-COVID19 group intercepted the y-axis at

value 35.22, while the COVID19-free at 17.76. This difference determined an increment of DNAmAge of about 9 years in the post-COVID19 compared to the same group's chronological age. No difference was appreciable in the controls (Table II). Accordingly, the vast majority (76.6%) of post-COVID19 had an average DeltaAge acceleration of 10.45 years (Figure 2, red dots). Considering that this method has a tolerance of about ± 5.2 years (24)(35), the corrected average accelerated DeltaAge for this group was 5.25. On the contrary, all the COVID19-free volunteers had a DeltaAge of 3.68, falling well into the range of normality (24) (Figure 2, blue squares). The post-COVID19/COVID19-free DeltaAge ratio was 2.84 (Table II). Interestingly, the DeltaAge distribution within the two groups showed that the COVID19-free samples were evenly distributed between the normal (39,9%) and the accelerated range (48.9%) while the 12.8% had a decelerated biological clock. On the contrary, 76.6 % of post-COVID19 had an accelerated DeltaAge, with only the 23.4% falling in the normal or decelerated range (Figure 3A). Interestingly, while the COVID19-free DeltaAge was distributed evenly among the different age groups, the increase of DeltaAge in the post-COVID19 population was well represented among the younger people (age 56 ± 12.8 years; Figure 3A, B). The older individuals, in both COVID19-free and COVID19-survivors groups, did not show signs of DeltaAge acceleration. This result indicates that the younger survivors might be more sensitive to the SARS-CoV-2-dependent remodeling of the epigenome landscape (Figure 3B).

Table I: Clinical information and risk factors in 117 COVID19 survivors and 144 COVID19-free volunteers.

CLINICAL DATA	COVID19-free	Post-COVID19
Samples (n)	144 (Male 66,0 %; Female 34,0%)	117 (Male 60,7%; Female 39,3%)
BMI $\geq$ 30	9,0%	15,3%
Smokers	37,5%	16,9%
diabetics	11,1%	12,1%
Hypertension	40,3%	36,3%
Clinical history of CVDs	33,3%	27,4%
Antecedent lung involvement	1,6%	20,2%
COVID19-related complications		
Pneumonia	/	57,3%

Oxygen therapy	/	52,4%
Artificial ventilation	/	35,5%
Length of viral positivity (average) in weeks	/	4,84

188

189

190

191 **Figure 1.** Biological age determination in COVID19-free (blue squares) vs. post-COVID19 (red
192 dots). A) Linear regression of COVID19-free volunteers DNAmAge. B) Linear regression of
193 DNAmAGE in the post-COVID19 subjects. In both graphs, the black dashed line is the bisector and
194 represents the perfect correlation between chronological and biological age. The post-COVID19
195 group (right panel) showed a statistically significant DNAmAge acceleration.; $P < 0,0001$ (two-sided
196 T-test).

197

Figure 2. DeltaAge distribution between COVID-free (left; blue squares) and post-COVID survivors (right; red dots). The black dashed lines indicate the ± 5 years limit of the normal range according to the method. *P*-value of $<0,0001$ (Two sided T-test).

Figure 3. A) DeltaAge range of distribution within each age group. Specifically, in the COVID19-free cohort (blue bars), 12.8% of the participants were decelerated (mean: -8.7 ± 5)*; 39.0% were falling within normality, while 48.2 % presented an accelerated DeltaAge (mean: $+5.15 \pm 4.34$). Interestingly, only a negligible portion (0.9%) of post-COVID19 patients (red bars) are in the decelerated range. While 22.5 % were within the normal range, the vast majority (76.6 %) bore an accelerated bioclock (mean: $+8.7 \pm 5.79$). The average chronological age is reported above each bin. * DeltaAge mean values are considered after subtraction of the ± 5.2 normality range distribution. B) The graph shows DeltaAge distribution according to the different chronological age groups. A significance of $P < 0.0001$ between the younger COVID19-free group (< 60) and the corresponding post-COVID19 patients is shown (Two-sided T-test)

Telomere length quantification.

TL shortening has been reported as a risk factor for developing more severe COVID19 syndrome (25). We investigated this parameter, which is also associated with the progression of the aging process (23). Comparing COVID19-free vs. post-COVID19 individuals revealed the presence of a significant chromosome ends shortness in the COVID19 survivors' group (Figure 4A). Specifically, in the COVID19-free (Fig.4A; blue squares) volunteers, TL was 3.5 fold longer than in the post-COVID19 group (red dots). The correlation between DeltaAge distribution and TL indicates that post-COVID19 (Fig. 4C; red dots) compared with the COVID-free (Fig. 4B; blue squares) have shorter telomeres independently by an accelerated DeltaAge, suggesting that these two parameters might be regulated independently.

Figure 4. A) Telomere length analysis in COVID19 survivors (red dots) and COVID-free subjects. The graph shows that the COVID19-free group has longer chromosome ends compared to the post-COVID19. P-value < 0.0001 (Two-sided T-test). B & C) Correlation between DeltaAge and TL in COVID19-free volunteers (B; blue squares) and post-COVID19 (C; red dots) patients.

Peripheral blood expression of ACE2 and DPP-4

In a cell infected by SARS-Cov-2, the ACE2 expression decreases, but little is known about the intensity of this biomarker in post-COVID19 survivors. We evaluated the mRNA level of ACE2 (SARS-CoV and SARS-CoV-2 receptor) and DPP-4 (MERS-CoV receptor). The results are shown in Figures 5A and B. In the post-COVID19 population, at the time point in which the blood samples were taken, longer than four weeks from the end of the viral infection (see Table I), ACE2 expression was significantly reduced (Fig. 5A). The expression level of DPP4 was unchanged (Fig. 5B).

Figure 5. A) qPCR determination of ACE2 expression level in COVID19-free (blue squares) and post-COVID19 (Two-sided T-test: $P < 0.0001$). B) mRNA level determination of DPP4 in COVID19-free vs. post-COVID19 showing no difference between the two groups. C) Correlation between DeltaAge and the relative expression levels of ACE2 mRNA in the peripheral blood of COVID19-free (blue squares) vs. post-COVID individuals (red dots). D) Correlation between DeltaAge and the relative expression levels of DPP4 mRNA in the peripheral blood of COVID19-free (blue squares) vs. post-COVID individuals (red dots).

The increased DeltaAge of the post-COVID19 group correlated well with the lowest ACE2 expression level (Fig. 5C; $P < 0.01$). No differences were observed in the distribution of DPP4 by DeltaAge (Fig. 5D).

Discussion.

Nowadays, the global vaccination program against the SARS-CoV-2 is actively ongoing, and the incidence of COVID19 will soon decrease sensibly worldwide. Nevertheless, among the millions of COVID19 survivors, many will require long-term assistance due to increased post-COVID19 clinical sequelae defined as PPCS (1)(41). Despite the many manifestations associated with PPCS, there is a lack of molecular biomarkers potentially valuable monitoring PPCS onset and evolution. In this study, we took advantage of prior indication that biological age, defined as DNAmAge, could

be altered in the presence of viral or bacterial infections (37)(40) (42) and from the fact that shorter telomeres have been associated with the risk of developing a worse COVID19 (25). In this light, we found that a consistently accelerated DeltaAge (+5.22 years above normal range) characterized the post-COVID19 population, and particularly those chronologically under 60 years (Figure 3A and B). This observation was paralleled by a significant telomere shortening (Figure 4). Although the two parameters seem independent (Fig. 4B and C), both alterations coexisted in the post-COVID population. The pathophysiology at the basis of these findings remains unclear; however, they may reflect a modified epigenetic environment, particularly evident among the younger COVID19 survivors (Fig.3). It is well known that the progression of aging is associated with critical metabolic changes. Some of these changes occur in the level of metabolites regulating the function of essential epigenetic enzymes, such as the decrease in NAD⁺ levels, the cofactor of Sirtuins (43), and the reduction in alpha-ketoglutaric acid (44), the cofactor for all dioxygenases, including those deputed to proteins and DNA demethylation (45). Although very speculative, it may be that older adults are relatively less sensitive to SARS-CoV-2-dependent epigenetic changes due to changes in their metabolic landscape. Additional experiments are necessary to elucidate this relevant aspect. In light of this consideration, a further question could be whether epigenetic changes might exist antecedently the first viral contact, persisting perhaps or worsening progressively up to the post-COVID19 period.

Several epigenetic phenomena have been associated with the SARS-CoV-2 infection (46), including the epigenetic regulation of ACE2 and IL-6, which has been reported to interest the development of a worse COVID19 due to excessive inflammation (47). Besides, the SARS-CoV-2 has been found to induce changes in DNA methylation, which affect the expression of immune response inhibitory genes that could, in part, contribute to the unfavorable progression of COVID-19 (48). Finally, it is noteworthy that a recently identified signature made of 44 variably methylated CpGs has been predictive of subjects at risk of developing worse symptoms after SARS-CoV-2 infection (49). Interestingly, none of these newly identified CpGs overlap with those involved in the DNAmAge prediction used in this (24) or other studies (29). Hypothetically, it might be possible that distinct signals are regulating the structure of the epigenomic regions determining a higher risk of developing a worse COVID19 syndrome and those associated with DNAmAge prediction.

Even though epigenetics might provide clinically relevant information about COVID19 (33) progression, no data is so far available about the involvement of epigenetic processes in the onset of the post-COVID19 syndrome or PPCS. Although none of the post-COVID survivors included in our study declared symptoms of PPCS, our evidence indicates the presence of changes in the methylation level of some CpGs associated with biological age calculation. This observation might reflect a more extensive phenomenon underlining unprecedented changes in the epigenome associated with the SARS-CoV-2 infection. A long-term follow-up of patients with an accelerated DeltaAge might be envisaged.

Telomere length is a marker of aging: progressive telomere shortening is a well-characterized phenomenon observed in older adults compared to the younger and attributed to the so-called telomeric attrition. This condition is worsened by the absence of telomerase activity physiologically silenced early post-natally and along adulthood (23). An accelerated TL shortening is a parameter associated with an increased risk of developing cardiovascular diseases and other disorders (50). In COVID19, patients bearing shorter telomeres in their peripheral leukocytes have been proposed to risk a worse prognosis (51). In the post-COVID19 group analyzed here, the average TL was $3,03 \pm$

2,39 Kb compared to the $10,67 \pm 11,69$ Kb of the control group ($P < 0.0001$). As shown in Table 2, the chronological ages of the two cohorts were about comparable. Hence it is unlikely that the aging process could have been a determinant eliciting the difference. Accordingly, our results suggest that the observed TL shortening could be independent of DeltaAge (Figure 4B and C), indicating that the SARS-CoV-2 infection might directly contribute to telomere erosion in the blood cellular component.

ACE2 is a crucial component of the SARS-CoV-2 infection process. SARS-CoV-2 uses the ACE2 receptor to invade human alveolar epithelial cells and other cells, including cardiac fibroblasts (52). In infected individuals, ACE2 is often down-regulated due to the infection (11)(47). The enzyme is expressed in several tissues, including alveolar lung cells, gastrointestinal tissue, vascular cells, and the brain; however, it is relatively under-represented in circulating blood cells. In all cases studied, the total relative ACE2 mRNA level in the peripheral blood of non-COVID19 or post-COVID19 subjects was significantly lower than that of the MERS-CoV receptor DPP4. However, in the post-COVID19 group, the ACE2 mRNA expression was reduced significantly compared to controls, while DPP4 elicited similar expression levels in both groups. Interestingly, the accelerated DeltaAge, predominant in the younger Post-COVID19 survivors, significantly correlated with the lower ACE2 mRNA level, suggesting for an adverse effect of DNAmAge on ACE2 density in peripheral blood (Figure 5B and C).

The two groups considered in this study were not significantly different in terms of age, sex, and known clinical conditions before SARS-CoV-2 infection except for a relatively higher incidence of BMI > 30 (15.3% vs. 9%) in the post-COVID19 population compared to controls as well as for a record of more frequent lung diseases (20.2% vs. 1.6%; see Table I). The origin of the persistent reduction in ACE2 expression in the post-COVID19 group remains unsolved, and a longitudinal study should be performed monitoring this parameter.

Conclusions.

This study has many significant limitations, including the small number of subjects investigated and the low number of CpGs considered. Although we used a valid forensic method to establish the biological age in the examined groups (24)(35) adopting other methods evaluating a large set of CpGs might be recommended (29)(53). However, the application of the latter procedures is limited by the elevated cost and relative complexity and may not be feasible at the laboratory level in many hospitals.

Nevertheless, it is shown here that individuals belonging to a group of COVID19 survivors had a significant acceleration of their biological age, occurring mainly in the younger individuals. This type of information has been correlated with TL shortening and the expression of ACE2 mRNA. It is too early to extrapolate whether relevant clinical indications may arise from this and other studies assessing the role of epigenetic changes in COVID19 syndrome (49) (48). However, a warning might be raised that sequelae of SARS-CoV-2 infection might rely on persistent epigenomic modifications, possibly underlying the presence of a COVID19 epigenetic memory. The epigenomic landscape of actual post-COVID19 survivors and in future COVID19 survivors from SARS-CoV-2 variants should then be taken under consideration to gain predictive prognostic insights and to monitor the response to treatment.

Table II: Summary

	COVID19-free	Post-COVID19	P-VALUE
Samples (n)	144 (Male 66,0 %; Female 34,0%)	117 (Male 60,7%; Female 39,3%)	
Chronological age (years)	62,48 ± 9,04	58,44 ± 14,66	Ns
Biological age (years)	63,81 ± 13,66	67,18 ± 10,86	Ns
Chronological Vs Biological (p-value)	Ns	<0,0001	
DeltaAge (years)	3,68 ± 8,17	10,45 ± 7,29	<0,0001
Ratio	1	2.84	
DeltaAge distribution			
Decelerated (%)	12,8	0,9	
Normal (%)	39,0	22,5	
Accelerated (%)	48,2	76,6	
Telomere length (kb)	10,67 ± 11,69	3,03 ± 2,39	<0,0001
ACE2 expression (2 [^] (-dct))	0,001390 ± 0,002298	0,0003801 ± 0,0004463	<0,0001
DPP4 expression (2 [^] (-dct))	0,1038 ± 0,089	0,1152 ± 0,069	ns

351 **Materials and methods.**

352 The samples have been classified into two groups: COVID19-free (n= 144), a heterogeneous group
353 that includes healthies, cardiovascular diseases, and obstructive sleep apnea patients, while the post-
354 COVID19 group includes all of the previous patients who had SARS-CoV-2 infection (n= 117). The
355 clinical features of both populations are summarised in Table 2.

356 DNA extraction from whole blood.

357 Blood samples collected in EDTA and 200 ul have been used to perform the extraction using QIAmp
358 DNA Blood Mini Kit (Qiagen, cat. 55106) associated with automated system QIAcube (Qiagen, cat.
359 9002160) according to manufacturer instructions. Then 2ul of DNA has been quantified with
360 QIAxpert (Qiagen, cat. 9002340)

361 Bisulfite Conversion.

362 1ug of DNA has been used to convert with Epiect fast DNA bisulfite (Qiagen, cat. 59824) following
363 the manufacturer instructions associated with RotorGene 2plex HRM (Qiagen, cat.9001560) and
364 QIAcube automated system. Then 2ul of converted DNA has been quantified with QIAxpert.

365 Polymerase chain reactions for pyrosequencing.

366 PCR reaction mixes were performed using PyroMark PCR kit (Qiagen, cat. 978103) following the
367 manufacturer's instructions. The sequences of primer used are available on Supplementary methods.

368 Pyrosequencing.

369 The amplicons have been sequenced in order to check the level of methylation in each CpG site.
370 PyroMark Q24 Advanced Reagents (Qiagen, cat. 970902) have been loaded in the PyroMark Q24
371 Cartridge (Qiagen, cat.979202) following the manufacturer instructions, and 5ul of PCR product has
372 been added to the reaction mix: Pyromark Binding Buffer (supplied in PyroMark Q24 Advanced
373 Reagents kit), StreptavidinSepharose High performance (GE Healthcare, cat. GE17-5113-01) and
374 H2O DNase/RNase free. Samples have been shaken at room temperature for 15 minutes at 1400rpm.
375 Successively, the samples have been undergone to PyroMark Q24 Vacuum Station (Qiagen, cat.
376 9001515) procedure in which the target sequences have been purified and put into an annealing buffer
377 containing the sequencing primer [0,375 uM]. The sequences of oligos are available in supplementary
378 data. The plate containing the sequence to analyze and the primer has been heated at 80°C for
379 5minutes. Successively, the PyroMark Q24 Advanced (Qiagen, cat. 9001514) has been set to analyze
380 the target sequences (Supplementary methods).

381 DNAmAge estimation.

382 Bekaert's algorithm has been applied to estimate the biological age of the population (24) as reported
383 in Daunay et al. (35):

384 $26.44119 - 0.201902 * ASPA - 0.239205 * EDARADD + 0.0063745 * ELOVL22 + 0.6352654 * PDE4C$

385

386

387 Telomere length quantification.

388 The chromosome end has been quantified by PCR Real-Time of Absolute Human Telomere Length
389 Quantification qPCR Assay Kit (ScienCell, cat. 8918) following the manufacturer's instructions.
390

391 RNA extraction.

392 According to manufacturer instruction, the total RNA has been isolated from whole blood using
393 QIAmp RNA blood mini (Qiagen, cat. 52304) and automatized extractor QIAcube. The RNA has
394 been quantified with QIAxpert.

395 cDNA synthesis and qPCR Real-Time.

396 Omniscript RT kit (Qiagen, cat. 205113) has been used to convert total RNA into cDNA following
397 the manufacturer's instructions.

398 The qPCR RealTime has been performed on RotorGene 2plex HRM using the RT2 SYBR Green
399 ROX FAST Mastermix (Qiagen, cat 330620). The sequences of primers are available on
400 supplementary data. To perform the amplification; the machine has been set:

401 Initial denaturation: 95° C, 5 minutes;

402 Denaturation: 95° C, 15 seconds;

403 Annealing: 60° C, 30 seconds;

404 Elongation: 72° C, 30 seconds;

405 Final elongation: 72° C, 1 minute.

406 Denaturation, annealing, and elongation have been repeated for 45 times.

407

408 Data analysis.

409 All data has been analyzed with GraphPad Prism 8.4.3. Method to calculate the p-value: Two sided
410 T-test.

411

412

413

414

415

416

417

418

419

420 List of abbreviations:

421 adult respiratory distress syndrome (ARDS)
 422 angiotensin-converting enzyme 2 (ACE2)
 423 biological age based on DNA methylation (DNAmAge)
 424 cardiovascular diseases (CVDs)
 425 coronavirus disease 19 (COVID19)
 426 Cytosine-Guanosine dinucleotides (CpGs)
 427 dipeptidyl-peptidase IV (DPP-4)
 428 Middle East respiratory syndrome (MERS)
 429 MERS coronavirus (MERS-CoV)
 430 persistent post-COVID19 syndrome (PPCS)
 431 renine- angiotensine system (RAS)
 432 severe acute respiratory syndrome (SARS)
 433 severe acute respiratory syndrome CoronaVirus 2 (SARS-CoV-2)
 434 Telomere length (TL)
 435 type 2 diabetes mellitus (T2DM)

436 Declarations:

- 437 • This study have been approved by the Ethical committee of Istituti Clinici Scientifici
 438 Maugeri Spa SB di Pavia. Document: METH-COVID19 - Studio epigenetico sulle
 439 conseguenze dell'infezione da SarsCoV-2 (ID: 2543 CE)
- 440 • Author Contributions: Conceptualization,: AF, CG, TB and MTLR.; methodology: AM,
 441 VB, MGZ, SA; validation, CG AF, AP, FM, MM, MN.; formal analysis: AM, CG,
 442 MTLR, TB, SN; investigation: AM, VB, MGZ,SA, LF ; resources: MN, VB, SN; data
 443 curation: AM; writing—original draft preparation: AM,CG; writing—review and editing:
 444 AF, TB, MTLR; supervision: CG; project administration: CG; funding acquisition: CG,
 445 MTLR.
- 446 • Funding:This research was funded by ITALIAN MINISTRY OF EDUCATION,
 447 UNIVERSITY AND RESEARCH, grant number PRIN2017S55RXB to A.F. and grant
 448 number PRIN2015HPMLFY to A.P., the ITALIAN MINISTRY OF HEALTH, grant
 449 number RF 2010-2318330 to A.F.; “RicercaCorrente” and “Progetto di Rete Cardiovascolare
 450 IRCCS: CardioCovid” to MTLR and “progetto di rete Aging IRCCS: progetto IRMA e
 451 progetto SIRI” to C.G.; F.M.is supported by the Italian Ministry of Health, (Ricerca Corrente
 452 and 5 x 1000), Telethon Foundation (#446 GGP19035A), AFM-Telethon (# 23054) and EU
 453 Horizon 2020 project COVIRNA (Grant #101016072).; ASSOCIAZIONE ITALIANA PER
 454 LA RICERCA SUL CANCRO AIRC under IG 2019, grant number ID 22858 project, S.N.
 455 This article partly based upon work from EU-CardioRNA COST Action CA17129.

- 456 • Acknowledgements: The authors would like to thanks Marta Lovagnini, Elena Robbi,
457 Riccardo Sideri and Lorena Grano De Oro fpr valuable work and assistance providing patients
458 information tracking and blood sample collection.
- 459 • All authors have read and agreed to the published version of the manuscript.
- 460 • Conflicts of Interest:The authors declare no conflict of interest
- 461

REFERENCES

1. Oronsky B, Larson C, Hammond TC, Oronsky A, Kesari S, Lybeck M, et al. A Review of Persistent Post-COVID Syndrome (PPCS). *Clin Rev Allergy Immunol* [Internet]. 2021;(0123456789). Available from: <http://www.ncbi.nlm.nih.gov/pubmed/33609255>
2. Heshui Shi*, Xiaoyu Han*, Nanchuan Jiang*, Yukun Cao, Osamah Alwalid, Jin Gu, Yanqing Fan† CZ, Summary. Radiological findings from 81 patients with COVID-19 pneumonia in Wuhan, China: a descriptive study. 2020;(January):19–21.
3. Bansal M. Cardiovascular disease and COVID-19. *Diabetes Metab Syndr*. 2020;14(4)(January):337–9.
4. Tan W, Aboulhosn J. The cardiovascular burden of coronavirus disease 2019 (COVID-19) with a focus on congenital heart disease. 2020;(January).
5. Ranucci M, Sitzia C, Baryshnikova E, Di Dedda U, Cardani R, Martelli F, et al. Covid-19-Associated Coagulopathy: Biomarkers of Thrombin Generation and Fibrinolysis Leading the Outcome. *J Clin Med* [Internet]. 2020;9(11). Available from: <https://www.mdpi.com/2077-0383/9/11/3487>
6. Klok FA, Kruipb MJHA, Meerc NJM van der, Arbousd MS, Gommerse DAMPJ, Kantf KM, et al. Incidence of thrombotic complications in critically ill ICU patients with COVID-19. 2020;(January).
7. Walkey AJ, Summer R, Ho V, Alkana P. Acute respiratory distress syndrome: Epidemiology and management approaches. *Clin Epidemiol*. 2012;4(1):159–69.
8. Lai J, Ma S, Wang Y, Cai Z, Hu J, Wei N, et al. Factors Associated With Mental Health Outcomes Among Health Care Workers Exposed to Coronavirus Disease 2019. *JAMA Netw open*. 2020;3(3):e203976.
9. Liu X, Kakade M, Fuller CJ, Fan B, Fang Y, Kong J, et al. Depression after exposure to stressful events: Lessons learned from the severe acute respiratory syndrome epidemic. *Compr Psychiatry*. 2012;53(1):15–23.
10. Alexopoulos P, Soldatos R, Kontogianni E, Frouda M, Ioanna Aligianni S, Skondra M, et al. COVID-19 Crisis Effects on Caregiver Distress in Neurocognitive Disorder. *J Alzheimers Dis*. 2021;79(1):459–66.
11. Gue YX, Gorog DA. Reduction in ACE2 may mediate the prothrombotic phenotype in COVID-19. *Eur Heart J*. 2020;41(33):3198–9.
12. Xiao C. Li1,*, Jianfeng Zhang2,* and JLZ. The vasoprotective axes of the renin-angiotensin system: physiological relevance and therapeutic implications in cardiovascular, hypertensive and kidney diseases. *Physiol Behav*. 2016;176(1):139–48.
13. Wu CH, Mohammadmoradi S, Chen JZ, Sawada H, Daugherty A, Lu HS. Renin-angiotensin system and cardiovascular functions. *Arterioscler Thromb Vasc Biol*. 2018;38(7):E108–16.
14. Myall KJ, Mukherjee B, Castanheira AM, Lam JL, Benedetti G, Mak SM, et al. Persistent Post-COVID-19 Inflammatory Interstitial Lung Disease: An Observational Study of Corticosteroid Treatment. *Ann Am Thorac Soc*. 2021 Jan;
15. Sargiacomo C, Sotgia F, Lisanti MP. COVID-19 and chronological aging: Senolytics and other anti-aging drugs for the treatment or prevention of corona virus infection? *Aging (Albany NY)*. 2020;12(8):6511–7.

- 504 16. Wang N, Shi X, Jiang L, Zhang S, Wang D, Tong P, et al. Structure of MERS-CoV spike
505 receptor-binding domain complexed with human receptor DPP4. *Cell Res* [Internet].
506 2013;23(8):986–93. Available from: <http://dx.doi.org/10.1038/cr.2013.92>
- 507 17. Kim KM, Noh JH, Bodogai M, Martindale JL, Yang X, Indig FE, et al. Identification of
508 senescent cell surface targetable protein DPP4. *Genes Dev.* 2017;31(15):1529–34.
- 509 18. Enz N, Vliegen G, De Meester I, Jungraithmayr W. CD26/DPP4 - a potential biomarker and
510 target for cancer therapy. *Pharmacol Ther* [Internet]. 2019;198:135–59. Available from:
511 <https://doi.org/10.1016/j.pharmthera.2019.02.015>
- 512 19. Suda M, Shimizu I, Yoshida Y, Hayashi Y, Ikegami R, Katsuumi G, et al. Inhibition of
513 dipeptidyl peptidase-4 ameliorates cardiac ischemia and systolic dysfunction by up-
514 regulating the FGF-2/EGR-1 pathway. *PLoS One.* 2017;12(8):1–18.
- 515 20. Solerte SB, Di Sabatino A, Galli M, Fiorina P. Dipeptidyl peptidase-4 (DPP4) inhibition in
516 COVID-19. *Acta Diabetol* [Internet]. 2020;57(7):779–83. Available from:
517 <https://doi.org/10.1007/s00592-020-01539-z>
- 518 21. HAYFLICK L, MOORHEAD PS. The serial cultivation of human diploid cell strains. *Exp*
519 *Cell Res.* 1961 Dec;25:585–621.
- 520 22. Zhan Y, Hägg S. Telomere length and cardiovascular disease risk. *Curr Opin Cardiol.* 2019
521 May;34(3):270–4.
- 522 23. Bekaert S, De Meyer T, Van Oostveldt P. Telomere attrition as ageing biomarker. *Anticancer*
523 *Res.* 2005;25(4):3011–22.
- 524 24. Bekaert B, Kamalandua A, Zapico SC, Van De Voorde W, Decorte R. Improved age
525 determination of blood and teeth samples using a selected set of DNA methylation markers.
526 *Epigenetics.* 2015;10(10):922–30.
- 527 25. Froidure A, Mahieu M, Hoton D, Laterre PF, Yombi JC, Koenig S, et al. Short telomeres
528 increase the risk of severe COVID-19. *Aging (Albany NY).* 2020;12(20):19911–22.
- 529 26. Horvath S, Raj K. DNA methylation-based biomarkers and the epigenetic clock theory of
530 ageing. *Nat Rev Genet.* 2018 Jun;19(6):371–84.
- 531 27. C Zapico S, Ubelaker DH. Applications of physiological bases of ageing to forensic sciences.
532 Estimation of age-at-death. *Ageing Res Rev.* 2013 Mar;12(2):605–17.
- 533 28. Jylhävä J, Pedersen NL, Hägg S. Biological Age Predictors. *EBioMedicine.* 2017 Jul;21:29–
534 36.
- 535 29. Horvath S. DNA methylation age of human tissues and cell types. *Genome Biol.*
536 2013;14(10).
- 537 30. Jung S-E, Shin K-J, Lee HY. DNA methylation-based age prediction from various tissues
538 and body fluids. *BMB Rep.* 2017 Nov;50(11):546–53.
- 539 31. Vidaki A, Kayser M. Recent progress, methods and perspectives in forensic epigenetics.
540 *Forensic Sci Int Genet.* 2018 Nov;37:180–95.
- 541 32. Zubakov D, Liu F, Kokmeijer I, Choi Y, van Meurs JBJ, van IJcken WFJ, et al. Human age
542 estimation from blood using mRNA, DNA methylation, DNA rearrangement, and telomere
543 length. *Forensic Sci Int Genet.* 2016 Sep;24:33–43.
- 544 33. Weidner CI, Lin Q, Koch CM, Eisele L, Beier F, Ziegler P, et al. aging of blood can be

tracked by DNA methylation changes at just three CpG sites. *Genome Biol.* 2014 Feb;15(2):R24.

34. Daunay A, Baudrin LG, Deleuze J-F, How-Kit A. Evaluation of six blood-based age prediction models using DNA methylation analysis by pyrosequencing. *Sci Rep.* 2019 Jun;9(1):8862.

35. Daunay A, Baudrin LG, Deleuze JF, How-Kit A. Evaluation of six blood-based age prediction models using DNA methylation analysis by pyrosequencing. *Sci Rep.* 2019;9(1):1–10.

36. Declerck K, Vanden Berghe W. Back to the future: Epigenetic clock plasticity towards healthy aging. *Mech Ageing Dev* [Internet]. 2018;174(January):18–29. Available from: <https://doi.org/10.1016/j.mad.2018.01.002>

37. Horvath S, Levine AJ. HIV-1 infection accelerates age according to the epigenetic clock. *J Infect Dis.* 2015;212(10):1563–73.

38. Esteban-cantos A, Rodríguez-centeno J, Barruz P, Alejos B, Saiz-medrano G, Nevado J, et al. Epigenetic age acceleration changes 2 years after antiretroviral therapy initiation in adults with HIV : a substudy of the NEAT001 / ANRS143 randomised trial. 2021;197–205.

39. Levine AJ, Quach A, Moore DJ, Achim CL, Soontornniyomkij V, Masliah E, et al. Accelerated epigenetic aging in brain is associated with pre-mortem HIV-associated neurocognitive disorders. *J Neurovirol* [Internet]. 2016;22(3):366–75. Available from: <http://dx.doi.org/10.1007/s13365-015-0406-3>

40. Kananen L, Nevalainen T, Jylhävä J, Marttila S, Hervonen A, Jylhä M, et al. Cytomegalovirus infection accelerates epigenetic aging. *Exp Gerontol* [Internet]. 2015;72:227–9. Available from: <http://dx.doi.org/10.1016/j.exger.2015.10.008>

41. Walitt B, Bartrum E. A clinical primer for the expected and potential. 2021;6:1–7.

42. Sabry D, Ahmed R, Abdalla S, Fathy W, Eldemery A, Elamir A. Braf, Kras and Helicobacter pylori epigenetic changes-associated chronic gastritis in Egyptian patients with and without gastric cancer. *World J Microbiol Biotechnol.* 2016;32(6):1–8.

43. Bonkowski MS, Sinclair DA. Slowing ageing by design: the rise of NAD(+) and sirtuin-activating compounds. *Nat Rev Mol Cell Biol.* 2016 Nov;17(11):679–90.

44. Asadi Shahmirzadi A, Edgar D, Liao C-Y, Hsu Y-M, Lucanic M, Asadi Shahmirzadi A, et al. Alpha-Ketoglutarate, an Endogenous Metabolite, Extends Lifespan and Compresses Morbidity in Aging Mice. *Cell Metab.* 2020 Sep;32(3):447-456.e6.

45. Ito S, D'Alessio AC, Taranova O V, Hong K, Sowers LC, Zhang Y. Role of Tet proteins in 5mC to 5hmC conversion, ES-cell self-renewal and inner cell mass specification. *Nature.* 2010 Aug;466(7310):1129–33.

46. Atlante S, Mongelli A, Barbi V, Martelli F, Farsetti A, Gaetano C. The epigenetic implication in coronavirus infection and therapy. *Clin Epigenetics.* 2020;12(1).

47. Sang ER, Tian Y, Miller LC. Epigenetic Evolution of ACE2 and IL-6 Genes: Non-Canonical Interferon-Stimulated Genes Correlate to COVID-19 Susceptibility in Vertebrates. 2021;

48. Li S, Ma F, Yokota T, Garcia G, Palermo A, Wang Y, et al. Metabolic reprogramming and epigenetic changes of vital organs in SARS-CoV-2-induced systemic toxicity. *JCI Insight.* 2021;6(2):1–18.

- 587 49. Castro de Moura M, Davalos V, Planas-Serra L, Alvarez-Errico D, Arribas C, Ruiz M,
588 Aguilera-Albesa S, Troya J, Valencia-Ramos J, Vélez-Santamaria V, Rodríguez-Palmero A,
589 Villar-Garcia J, Horcajada JP, Albu S, Casasnovas C, Rull A, Reverte L, Dietl B, Dalm EM.
590 Epigenome-wide association study of COVID-19 severity with respiratory failure.
591 EBioMedicine. 2021;
- 592 50. Cheng F, Carroll L, Joglekar M V, Januszewski AS, Wong KK, Hardikar AA, et al. Diabetes,
593 metabolic disease, and telomere length. *lancet Diabetes Endocrinol*. 2021 Feb;9(2):117–26.
- 594 51. Sanchez-Vazquez R, Guío-Carrión A, Zapatero-Gaviria A, Martínez P, Blasco MA. Shorter
595 telomere lengths in patients with severe COVID-19 disease. *Aging (Albany NY)*. 2021
596 Jan;13(1):1–15.
- 597 52. Amendola A, Garoffolo G, Songia P, Nardacci R, Ferrari S, Bernava G, et al. Human
598 cardiosphere-derived stromal cells exposed to SARS-CoV-2 evolve into hyper-
599 inflammatory/pro-fibrotic phenotype and produce infective viral particles depending on the
600 levels of ACE2 receptor expression. *Cardiovasc Res*. 2021 Mar;
- 601 53. Hannum G, Guinney J, Zhao L, Zhang L, Hughes G, Sadda S, et al. Genome-wide
602 Methylation Profiles Reveal Quantitative Views of Human Aging Rates. *Mol Cell* [Internet].
603 2013;49(2):359–67. Available from:
604 <https://www.sciencedirect.com/science/article/pii/S1097276512008933>

605