1 Exposure to landscape fire smoke extremely reduced birthweight in low- and middle- 2 income countries

3 Jiajianghui Li^{1, #}, M.S.; Tianjia Guan^{2, #}, Ph.D; Qian Guo^{3, #}, M.S.; Guannan Geng⁴, Ph.D;
4 Huiyu Wang¹, M.S.; Fuyu Guo¹, Ph.D; Jiwei Li⁵, Ph.D; Tao Xue^{1, *}, Ph.D

- 5 1. Institute of Reproductive and Child Health / Ministry of Health Key Laboratory of
6 Reproductive Health and Department of Epidemiology and Biostatistics, School of Public
7 Health, Peking University Health Science Centre, Beijing, China, 100191
- 8 2. Department of Health Policy, School of Health Policy and Management, Chinese Academy of
9 Medical Sciences and Peking Union Medical College, Beijing, China, 100730
- 10 3. School of Energy and Environmental Engineering, University of Science and Technology
11 Beijing, Beijing 100083, China
- 12 4. School of Environment, Tsinghua University, Beijing, China, 100085
- 13 5. College of Computer Science and Technology, Zhejiang University, Hangzhou, China

14
15 * Corresponding to Tao Xue (txue@hsc.pku.edu.cn);
16 # Those authors equally contribute to the paper.

17 This work was supported by PKU-Baidu Fund, National Natural Science Foundation of China,
18 Peking University Health Science Centre (early career funding for Tao Xue), and CAMS
19 Innovation Fund for Medical Sciences.

20
21 Jiajianghui Li: lijiajianghuizi@stu.pku.edu.cn
22 Tianjia Guan: guan_tianjia@163.com
23 Qian Guo: guo_guo0825@163.com
24 Huiyu Wang: 193234751@qq.com
25 Fuyu Guo: fyguo@pku.edu.cn
26 Jiwei Li: jiwei_li@zju.edu.cn
27 Guannan Geng: guannangeng@tsinghua.edu.cn
28 Tao Xue: txue@hsc.pku.edu.cn

29

30 **Abstract**

31 Landscape fire smoke (LFS) has been associated with reduced birthweight, but evidence from
 32 low- and middle-income countries (LMICs) is rare. Here, we present a sibling-matched case–
 33 control study of 227,948 newborns to identify an association between fire-sourced fine
 34 particulate matter (PM_{2.5}) and birthweight in 54 LMICs from 2000 to 2014.

35 We selected mothers from the geocoded Demographic and Health Survey with at least two
 36 children and valid birthweight records. Newborns affiliated with the same mother were
 37 defined as a family group. Gestational exposure to LFS was assessed in each newborn using
 38 the concentration of fire-sourced PM_{2.5}. We determined the associations of the within-group
 39 variations in LFS exposure with birthweight differences between matched siblings using a
 40 fixed-effects regression model. Additionally, we analyzed the binary outcomes of low
 41 birthweight (LBW) or very low birthweight (VLBW).

42 According to fully adjusted models, a 1 µg/m³ increase in the concentration of fire-sourced
 43 PM_{2.5} was significantly associated with a 2.17 g (95% confidence interval [CI]: 0.56–3.77)
 44 reduction in birthweight, a 2.80% (95% CI: 0.97–4.66) increase in LBW risk, and an 11.68%
 45 (95% CI: 3.59–20.40) increase in VLBW risk.

46 Our findings indicate that gestational exposure to LFS harms maternal health.

47

48 **Introduction**

49 The natural cycle of landscape fires (e.g., wildfires, tropical deforestation fires, and
50 agricultural biomass burning) plays an important role in maintaining the terrestrial ecosystem.
51 Yet, landscape fire smoke (LFS) triggers a costly and growing global public health problem¹.
52 The frequency and intensity of landscape fire events have tended to increase^{2,3}, driven by
53 interactions between human activities (e.g., slash and slave agriculture) and climate change⁴.
54 Most emissions originate from fires located in tropical rainforests and savannas, where they
55 cause recurrent episodes of severe ambient pollution that affect mainly low- and middle-
56 income countries (LMICs)¹. Worldwide concern regarding large-scale natural forest burning
57 has sparked increasing interest in the harmful effects of LFS exposure on human health.

58 LFS is composed of hundreds of combustion products, such as carbon monoxide, nitrogen
59 oxides, and polycyclic aromatic hydrocarbons⁵. Particulate matter with an aerodynamic
60 diameter $\leq 2.5 \mu\text{m}$ (PM_{2.5}) is one of the most widely studied pollutants derived from LFS. The
61 24-h average PM_{2.5} concentrations on burning days can be many times higher than those on
62 normal days, and the peak level of ambient exposure can persist for several weeks⁶. Several
63 epidemiological studies have shown adverse effects of LFS exposure on human health. The
64 majority of the study outcomes involved morbidity and mortality caused by acute respiratory
65 and cardiovascular diseases, particularly among vulnerable individuals, such as children and
66 the elderly⁷⁻⁹. However, the health impacts of LFS on susceptible pregnant women, another
67 highly vulnerable group due to gestation-related physiological changes, such as an increased
68 breathing rate during pregnancy¹⁰, are rarely discussed. Previous studies have shown that
69 gestational exposure to air pollutants, including those derived from LPS and other emissions,
70 are associated with abnormal placental vascular function and, consequently, low birthweight
71 (LBW)¹¹, preterm birth¹², and small for gestational age¹³. Exposure to air pollution
72 induces subclinical disorders, such as inflammation, oxidative stress, increased blood
73 viscosity, mitochondrial methylation, and hypoxia^{14,15}, which renders the association between
74 LFS and adverse maternal outcomes biologically plausible.

Birth weight comprehensively reflects the intrauterine environment quality, fetal growth, and maternal nutritional status. Therefore, LBW is a global public health problem. LBW neonates are at higher risk for a range of diseases in later life (e.g., early growth retardation and adulthood cardiorespiratory diseases) compared with normal newborns^{16,17}. The estimated worldwide prevalence of LBW was 14.6% (95% confidence interval [CI] 12.4–17.1) in 2015, compared with 17.5% (95% CI 14.1–21.3) in 2000¹⁸. Although some progress has been made in reducing LBW globally, the average annual reduction rate is insufficient to achieve the 2025 WHO global target¹⁹. Additionally, the relevant burden of LBW is unequally distributed in the world, as 91% of LBW infants are from LMICs, particularly from southern Asia and sub-Saharan Africa¹⁸.

All risk factors for adverse birth defects should be screened thoroughly to protect maternal and infant health. In addition to conventional factors, such as nutrition and maternal condition, environmental factors, including ambient PM_{2.5} exposure, also contribute to a considerable global burden of adverse birth outcomes²⁰. Although a growing number of studies have suggested the deleterious effects of urban air pollution on LBW^{21,22}, little is known about fire-sourced PM_{2.5}. Furthermore, studies concerning the association between LFS and birthweight have focused primarily on high-income countries, and their conclusions have been inconsistent^{23–26}. As most landscape fires occur in LMICs (Figure 1), studies to assess the relevant health impacts of LFS among residents of such countries are warranted. Hence, we explored the association between birthweight and fire-sourced PM_{2.5} in 54 LMICs using a sibling-matched case–control study from 2000 to 2014, which was based on a series of global surveys.

Methods

Study population

The birthweight data and relevant individual variables were extracted from 113 individual Demographic and Health Surveys (DHSs) representing 54 LMICs from 2000 to 2014 (Figure 1). The DHSs were conducted by the US Agency for International Development to

characterize the demography and health of more than 90 developing countries. DHSs are usually conducted every 5 years in each country, using a two-stage sampling design. Information on demographic factors, socioeconomic factors, and health status was collected from a nationally representative probability sample of households. Well-trained fieldworkers conducted the surveys using uniformly designed questionnaires in the native language. The primary sampling units (e.g., villages or blocks) have begun to be geocoded in recent waves using global positioning systems (GPSs). The GPS-enhanced DHSs were matched with environmental exposure and thus included in our analyses. Please refer to the open-access website for detailed information on the DHS database: <https://www.dhsprogram.com/>.

All females aged 15–49 years within each selected household were eligible respondents of the survey. The health outcome analyzed in this study was birthweight. The DHSs asked the screened mothers to recall the birthweight of all children born during the 5 years leading up to the survey date. To characterize different levels of reduced birthweight, we examined two binary outcomes: LBW (birthweight < 2,500 g) and very LBW (VLBW, birthweight < 1,500 g). Other covariates, such as maternal age and birth time (in months), were extracted from the relevant questionnaires.

118 ***Exposure assessment***

We assessed ambient exposure to LFS as the PM_{2.5} concentration attributable to landscape fires (hereafter, defined as fire-sourced PM_{2.5}), referring to a previous study²⁷. The fire-sourced PM_{2.5} concentrations were estimated using the GEOS-Chem chemical transport model (CTM). The detailed settings of the GEOS-Chem are provided in the supplemental text.

The fire-sourced PM_{2.5} was derived from a comparison between two GEOS-Chem model runs. The two models were identical except for switching on and off fire emissions. We quantified the fraction (ρ_m) of fire-attributed PM_{2.5} using the following equation:

$$126 \quad \rho_{m,y} = (\text{PM}_{2.5,m,y}^{\text{on}} - \text{PM}_{2.5,m,y}^{\text{off}}) / \text{PM}_{2.5,m,y}^{\text{on}} \dots\dots (1),$$

127 where the superscripts *on* and *off* denote the two simulations switching on and off fire

emissions, respectively; and the subscripts m and y are month and year indicators, respectively.

Additionally, considering the potential errors in the CTM procedure²⁸, the bias correction approach has been widely applied to improve CTM performance in exposure assessments using ground-surface observational data²⁹. We utilized a well-developed and widely utilized product of satellite-based PM_{2.5} estimates³⁰ (PM_{2.5,y}^{satellite}) in the bias correction. This product has a spatial resolution of $0.05^\circ \times 0.05^\circ$ and was publicly available from 1998 to 2018 as the gridded annual mean (http://fizz.phys.dal.ca/~atmos/martin/?page_id=140#V4.GL.03). We prepared the GEOS-Chem simulations ($\rho_{m,y}$ and PM_{2.5,m,y}^{on}) in the same $0.05^\circ \times 0.05^\circ$ grid using an inversed distance-weighted downscaling approach before bias correction. The bias correction rate (η_y) was calculated as

$$\eta_y = \text{PM}_{2.5,y}^{\text{satellite}} \div (1/12 \times \sum_m \text{PM}_{2.5,m,y}^{\text{on}}) \dots\dots (2).$$

The monthly exposure indicator of fire-sourced or non-fire-sourced PM_{2.5} was finally derived as

$$[\text{Fire-sourced PM}_{2.5}]_{m,y} = \eta_y \times \rho_{m,y} \times \text{PM}_{2.5,m,y}^{\text{on}},$$

$$[\text{Non-fire-sourced PM}_{2.5}]_{m,y} = \eta_y \times (1 - \rho_{m,y}) \times \text{PM}_{2.5,m,y}^{\text{on}} \dots\dots (3).$$

We utilized the non-fire-sourced PM_{2.5} as a potential confounder in the subsequent epidemiological analysis. We also collected other environmental variables, including ambient temperature and humidity data, from an assimilated dataset called Modern-Era Retrospective analysis for Research and Applications, version 2 (please refer to the supplemental text for more details). The variables were downloaded as monthly gridded values and prepared in the $0.05^\circ \times 0.05^\circ$ grid in the same way as we processed the GEOS-Chem outputs.

To understand the associations between birthweight and environmental variables, we focused on exposure during gestation. As the DHSs did not record the specific duration of gestation, we utilized the 9-month average preceding the survey month as exposure time during

153 pregnancy.

154

155 *Study design*

156 To evaluate the association between fire-sourced PM_{2.5} and birthweight, we applied a sibling-
 157 matched case-control study, similar to our previous work³¹. We defined the descendants of
 158 the same mother as a family group of matched siblings and compared birthweights within
 159 each group. The family-level baseline birthweight (defined as the mean birthweight within
 160 each group) can be affected by many complex factors, including genetics, socioeconomic
 161 position, and quality of local medical services, which can be difficult to measure. We matched
 162 the mothers to control for those unmeasured confounders in the study design. A sibling-
 163 matched case-control design is a simple, robust, and cost-efficient method, particularly for
 164 large-population data with potential heterogeneity^{32,33}.

165 We extracted the females with at as least two valid records of newborns from all available
 166 individual DHSs. Each valid record was defined using the following inclusion criteria: (1) a
 167 valid birthweight, (2) a record of the GPS coordinates, (3) a valid birthdate, and (4) valid
 168 environmental exposure values (some of the satellite-based PM_{2.5} estimates were not available
 169 for some locations, such as small islands). Finally, our dataset assembled 113 individual
 170 surveys from 54 countries (Figure 1) and included 227,948 newborns classified into 109,137
 171 groups according to their mothers.

172 *Statistical analyses*

173 According to the sibling-matched method, we used a fixed-effects regression model to
 174 evaluate the associations between fire-sourced PM_{2.5} and LBW or VLBW. The model was
 175 specified as

$$176 \quad y_{i,j} \sim x_{i,j}\beta + z_{i,j}\gamma + \theta_i \dots \dots (4),$$

177 where the subscripts i and j denote the indices of the mother-affiliated group and child,

178 respectively; $x_{i,j}$ denotes the concentration of fire-sourced $PM_{2.5}$; $z_{i,j}$ denotes the adjusted
 179 covariates; $y_{i,j}$ denotes the outcome variable; and θ_i is a nuisance parameter for the fixed-effect
 180 to control for group-specific unmeasured confounders. The model was specified as Gaussian
 181 regression when setting $y_{i,j}$ as the birthweight and as logit regression when setting $y_{i,j}$ as the
 182 logit-transformed probability of LBW or VLBW. The adjusted covariates included maternal
 183 age; child sex; multiple births; non-fire-sourced $PM_{2.5}$; spline terms of birth order (with 5
 184 degrees of freedom [DF]), of temperature (3 DF), of humidity (3 DF), of the calendar year (5
 185 DF), and of the monthly index (4 DF); and a spatiotemporal effect parametrized by the
 186 interaction between country and calendar year. The nonlinear effects of month and year
 187 controlled for the seasonal periodic variation and long-term trend in child health, respectively.
 188 The spatiotemporal effect captured country-specific trends. The association between
 189 birthweight and LFS was evaluated by the regression coefficient β for 1 $\mu g/m^3$ increases in
 190 fire-sourced $PM_{2.5}$. The association with LBW or VLBW was evaluated by excess risk [= (1 –
 191 $\exp(\beta)) \times 100\%$] for a 1 $\mu g/m^3$ increase in fire-sourced $PM_{2.5}$. All environmental exposures
 192 were quantified using the same time window. The major time window was the 9 months
 193 before birth. We also calculated the alternative exposure indicator as an average during the 3
 194 or 6 months preceding birth.

195 In sensitivity analyses, we first derived the lag-distributed model to explore how the
 196 association varied during the exposure time window. Next, subpopulation-specified
 197 associations were estimated using interaction analyses to detect potential heterogeneity. Third,
 198 we developed nonlinear associations by replacing the linear term of fire-sourced $PM_{2.5}$ using a
 199 set of thin-plate spline functions. Fourth, to examine potential recall bias, we estimated the
 200 associations by strata of the durations from birth to survey time. Fifth, to test whether the
 201 estimated associations were attributable to LFS exposure or other direct damage related to
 202 landscape fires (e.g., destroying human habitats), we derived an indicator for the transported
 203 fire-sourced $PM_{2.5}$, according to a satellite image of the burned area (MCD64A1,
 204 <https://lpdaacsvc.cr.usgs.gov/appears/>). The fire-sourced $PM_{2.5}$ at the pixels with the zero-
 205 burned area during gestation were defined as smoke transported from other locations. The

206 associations with transported fire-sourced PM_{2.5} could be indicative of the effect of LFS
207 exposure via the inhalation pathway.

208 Family-level baseline birthweight and the correlated gestation length can be determined by
209 factors such as genetics and socioeconomic level, which may also affect the association
210 between the within-group change in birthweight and LFS exposure. The fixed-effects
211 regression models actually infer an association by examining the coherence between
212 derivation of individual-level birthweight from the family-level mean (i.e., the expected
213 birthweight given a specific family) and the corresponding exposure. To show that, we
214 defined the family-level baseline birthweight as y_i . Equation (4) was equivalently transformed
215 as $\Delta y_{ij} = (y_{ij} - y_i) \sim x_{ij}\beta + z_{ij}\gamma + (\theta_i - y_i)$, where $(\theta_i - y_i)$ is viewed as a new nuisance
216 parameter θ_i^* . The nature of fixed-effects model made it possible to explore how the effect of
217 fire-sourced PM_{2.5} varied according to family-level mean birthweight. Accordingly, we
218 derived a baseline-varying association model as follows:

$$219 \quad \Delta y_{ij} \sim f(y_i)x_{ij} + z_{ij}\gamma + \theta_i^* \dots\dots (5),$$

220 where $f(y_i)$ denotes the baseline-varying coefficient and is parametrized by a set of spline
221 functions.

222 All analyses were performed in R (version 4.0.2) (The R Foundation for Statistical
223 Computing, Vienna, Austria). Inference for the fixed-effects models was performed using the
224 *fixest* package. The lag-distributed effect was parametrized using the *dlnm* package, and the
225 thin-plate splines for the nonlinear association and baseline-varying effect were parametrized
226 using the *mgcv* package.

227 **Results**

228 ***Descriptive summary***

229 Among the 227,948 samples analyzed from 54 LMICs, there were 109,137 groups of siblings,
230 who were matched to their mothers. Each group had an average of 2.13 (standard deviation

[SD]: 0.36) samples. Approximately half of the samples were from Sub-Saharan Africa and one-quarter from South Asia. The mean birthweight was 3,082 (SD: 724) g. There were 31,854 LBW births and 2,912 VLBW births. More statistics on the analyzed samples are shown in Supplemental Table S1.

The spatial distribution of the long-term average fire-sourced $PM_{2.5}$ concentrations is displayed in Figure 1. Our modeling results indicate that the hotspots of LFS include regions of the Congo rainforest basin, Amazon rainforest basin, Indonesian forests, Siberia forests, and other forest areas. Our analyzed samples included newborns from the first three regions. Among our samples, the average concentration of full gestational exposure to fire-sourced $PM_{2.5}$ was 4.29 (SD: 5.53) $\mu g/m^3$. A detailed summary of the environmental exposures is presented in Table S1.

Gestational exposure to fire-sourced $PM_{2.5}$ varied widely between family groups; it was higher in those living closer to hotspot regions (e.g., forests). Approximately 92% of the variation in fire-sourced $PM_{2.5}$ exposure was between family groups, and 8% was within family groups. For the total variation in birthweight, 71.7% was between groups and 28.3% within groups (Table 1). The within- or between-group variations reflected temporal or spatial changes in those variables, respectively. Birthweight and fire-sourced $PM_{2.5}$ were negatively correlated in the temporal dimension but positively correlated in the spatial dimension. As the spatial patterns in birthweight and fire-sourced $PM_{2.5}$ were determined by geographic, socioeconomic, and cultural factors, the between-group correlations may have been confounded by their common determiners, such as urbanization level. The complex correlations between birthweight and LFS suggest that statistical inference of their associations should be cautiously determined based on well-designed epidemiological analysis.

254

255 Figure 1 Distribution of the fire-sourced PM_{2.5} concentrations (average for 2000–2014) and
 256 the locations of the analyzed samples (grey dots). The circles represent the rate of very-low-
 257 birthweight births among the analyzed samples by country.

Table 1 Between- or within-group variations in birthweight and gestational exposure to fire-sourced PM_{2.5} and their correlations.

	Standard deviation (% of total variance)		Correlation (R)
	Birthweight	Fire-sourced PM _{2.5}	
Total	724 (100%)	5.53 (100%)	0.1594
Between groups of matched siblings	613 (71.7%)	5.30 (91.7%)	0.1973
Within groups of matched siblings	386 (28.3%)	1.59 (8.3%)	-0.0035

Association between LFS and birthweight

Although the fully adjusted model indicated that a 1 µg/m³ increase in exposure to fire-sourced PM_{2.5} during the 9 months before birth was significantly associated with a 2.17 g (95% CI: 0.56–3.77) birthweight reduction, the association was sensitive to differences in the adjusted covariates and exposure time window (Figure 2). For example, the unadjusted model reported the estimated effect as a birthweight reduction of 0.84 g (95% CI: -0.53 to 2.22). The lag-specific associations did not display a clear time-varying pattern (Figure S1), which suggests a linear increase in strength in the association with increased length of the exposure time window (Figures S2 & 2). The subpopulation-specific results suggest that the association was potentially heterogeneous (Figure S3). The association was stronger for female infants, newborns of nulliparous mothers, or newborns of unemployed mothers, compared with the corresponding references (i.e., male infants, newborns of multiparous mothers, or newborns of employed mothers, respectively). The nonlinear analysis confirmed the adverse effect of LFS on birthweight (Figure 3). The sublinear association suggests a saturated effect of a ~60 g birthweight reduction for exposure to extreme fires, which contributed to a PM_{2.5} concentration > ~20 µg/m³.

276

277 Figure 2 Estimated associations between gestational exposure to fire-sourced PM_{2.5} and
 278 birthweight change, low birthweight, or very low birthweight. The dots represent the point-
 279 estimates, and bars represent the corresponding 95% confidence intervals.

280

281 Figure 3 The nonlinear association between gestational exposure to fire-sourced PM_{2.5} and
 282 change in birthweight. The solid line represents the point-estimates, the dashed line represents
 283 the 95% confidence intervals, the boxplots represent the distributions of different exposure
 284 levels by regions; and the red dots represent the mean exposure level.

285

286 *Association between LFS and VLBW*

287 Our fully adjusted models suggested that LFS exposure during the 9 months before birth was
 288 significantly associated with an increased risk of LBW or VLBW. According to the estimates,
 289 the risks of LBW and VLBW increased by 2.80% (95% CI: 0.97–4.66) and 11.68% (95% CI:
 290 3.59–20.40), respectively, for each 1 $\mu\text{g}/\text{m}^3$ increase in exposure to fire-sourced $\text{PM}_{2.5}$.
 291 Compared to LBW, VLBW was more strongly associated with fire-sourced $\text{PM}_{2.5}$. The
 292 estimated association for LBW was sensitive to different covariate adjustments (Figure 2) and
 293 was potentially heterogeneous among subpopulations (Figure S3). The association for VLBW
 294 was robust, given the adjustments for different covariates (Figure 2), and was less
 295 heterogeneous (Figure S3). Among all effect modifiers, only the type of cooking energy,
 296 which was indicated for indoor sources of particulate matter, significantly changed the
 297 association between VLBW and fire-sourced $\text{PM}_{2.5}$ (P-value = 0.02, Figure S3). The use of
 298 unclear cooking energy, as a competing risk factor for fire-sourced $\text{PM}_{2.5}$, significantly
 299 weakened the association between VLBW and fire-sourced $\text{PM}_{2.5}$.

300 To further explore why VLBW was more strongly associated with fire-sourced $\text{PM}_{2.5}$ than was
 301 LBW, we developed a model to estimate the baseline-varying association between birthweight
 302 and fire-sourced $\text{PM}_{2.5}$. As a result, the absolute change in birthweight for per unit exposure to
 303 fire-sourced $\text{PM}_{2.5}$ varied with the baseline of family-level mean birthweight (Figure S4).
 304 Samples with an extremely low or high baseline birthweight were more susceptible to fire-
 305 sourced $\text{PM}_{2.5}$ than were those with a moderate baseline birthweight, in terms of the absolute
 306 birthweight change (Figure S4). As a reduction in birthweight and a family-level baseline co-
 307 determined the increased incidence of LBW or VLBW, the relative change in birthweight
 308 reflected the relevant risk. Newborns from families with a lower baseline were more
 309 susceptible to fire-sourced $\text{PM}_{2.5}$, in terms of a relative birthweight reduction (Figure 4). This
 310 finding partially explains why gestational exposure to fire-sourced $\text{PM}_{2.5}$ was robustly and
 311 strongly associated with VLBW. According to our estimates, a 1 $\mu\text{g}/\text{m}^3$ increase in fire-
 312 sourced $\text{PM}_{2.5}$ was associated with a relative birthweight reduction of 1.47% (95% CI: 0.06–
 313 2.88), 0.54% (95% CI: 0.03–1.05), 0.16% (95% CI: 0.01–0.31), or 0.03% (95% CI: –0.04 to

0.10) for samples with a baseline birthweight of 1,500, 2,000, 2,500, or 3,000 g, respectively.

Sensitivity analyses

First, we explored potential recall bias using sensitivity analyses, and the estimated associations were not significantly different according to recall period (Figure S3). Second, we examined the spatial variations in the estimated associations. Although analysis of differences in region-specific associations reported non-significant results (Figure S3), we could not completely rule out the possibility of spatial heterogeneity in the estimated results. As most of our analyzed samples were from Sub-Saharan Africa, only those estimates were significant, and the remaining region-specific results had wide-ranging 95% CIs (Figure S3). Third, we explored whether the estimated adverse effects were attributable to ambient LFS exposure or other impacts (e.g., social stress) relevant to landscape fires. The estimated effects of exposure to fire-sourced PM_{2.5} were not significantly different between locations without burning and those with burning (Figure S3). These results confirm that ambient exposure was the major exposure pathway for our analyzed outcomes.

329

330 Figure 4 Baseline-varying association between gestational exposure to fire-sourced PM_{2.5} and
 331 relative change in birthweight. The solid line represents the point-estimates, the dashed line
 332 represents the 95% confidence intervals, and the boxplots represent the distributions of
 333 different birthweights by region.

334

335 Discussion

336 This study revealed an association between increased risk of VLBW and exposure to LFS
 337 during pregnancy in 54 LIMCs. We found that gestational exposure to fire-sourced PM_{2.5}
 338 potentially reduced birthweight, and the infants with a low family-level mean of birthweight
 339 were susceptible to the effect of LFS. We present the first sibling-matched study on the
 340 association between birthweight and fire-sourced PM_{2.5} from multiple LIMCs.

341 LFS contains numerous hazardous pollutants and exerts both short- and long-term effects on
 342 human health³⁴. An adverse birth outcome is one of the consequences. Holstius et al.²³
 343 estimated that the mean birthweight of fire-exposed infants are 7.0 g (95% CI: 11.8–2.2), 9.7
 344 g (95% CI: 14.5–4.8), and 3.3 g (95% CI: –0.6 to 7.2) lower than those of unexposed infants
 345 when wildfires occurred during the first, second, and third trimesters, respectively. In a
 346 retrospective cohort study in Brazil, fire-exposed mothers (defined as the highest PM_{2.5}
 347 concentration quartile) reported a higher risk of LBW than did unexposed mothers. The
 348 adjusted odds ratios for exposure during the second and third trimesters were 1.51 (95% CI:
 349 1.04–2.17) and 1.50 (95% CI: 1.06–2.15), respectively²⁴. Abdo et al.²⁵ showed that a 1
 350 µg/m³ increase in the wildfire-sourced PM_{2.5} concentration during the first trimester was
 351 associated with a 5.7 g (95% CI: 0.4–11.1) reduction in birthweight, in United States. In
 352 contrast, an Australian cross-sectional study reported that the average birthweight of male
 353 infants exposed to the 2003 Canberra wildfires was significantly higher than that of
 354 unexposed peers²⁶. That study attributed the positive association between wildfire smoke and
 355 birthweight change to the potentially increased levels of maternal blood glucose after
 356 exposure.²⁶ The mixed results might be due to differences among studies in terms of
 357 epidemiological design, population characterization, exposure time window, and competing
 358 risk factors. Our results are comparable with most previous findings. Additionally, although
 359 there are few studies on fire-sourced PM_{2.5} similar to ours, a large number of studies have
 360 focused on the effect of gestational exposure to urban ambient PM_{2.5} on birthweight. A recent
 361 meta-analysis, which included more than 2.7 million participants, reported that the pooled risk
 362 ratio of LBW for a 10 µg/m³ increase in PM_{2.5} exposure was 1.081 (95% CI 1.043–1.120)²¹.

363 Considering the abundance of toxic organic particles derived from biomass burning, the
364 association of birthweight with fire-sourced PM_{2.5} should be stronger than that with urban
365 PM_{2.5}.

366 The family-level mean birthweight varied widely between households (Table 2). These
367 variations may have been caused by many factors, including genetics, socioeconomic position
368 (e.g., nutrition), and indoor household pollution. As birthweight is positively correlated with
369 gestation length, a poor family-level baseline birthweight comprehensively indicates a
370 combined risk of short gestation and LBW. Because the cumulative effect of fire-sourced
371 PM_{2.5} was observed to increase with exposure duration (Figure S2), a larger birthweight at the
372 family level, as indicated for a longer gestation, increased the absolute effect of the per-unit
373 LFS exposure (Figure S4).

374 As our models did not control for gestation length due to the lack of relevant data, the
375 estimated associations reflected the joint outcome of LFS exposure by reducing gestation
376 length or birthweight. The fetus grows in a sublinear temporal pattern³⁵ during the late
377 gestational stage, which suggests its weight increases to an approximately stable level before
378 birth. Considering this sublinear relationship and given the different baseline birthweights, the
379 same reduction in gestation length led to different reductions in birthweight. The marginal
380 effect of a shorter gestation period on the reduction in birthweight is larger for an infant with a
381 lower baseline birthweight. For instance, if an infant is expected to reach the stable level of
382 fetal weight, a marginal reduction in its gestational age will not considerably change the
383 birthweight. As exposure to LFS has been reported to cause a short gestation²⁵, its indirect
384 effect on reduced birthweight can be enhanced by poor baseline maternal health.

385 Reducing the risk of LBW is one of the 2025 WHO global targets, and it contributes to
386 achieving the Sustainable Development Goals. Although major interventions to reduce LBW
387 have focused on improving maternal nutrition, environmental toxic pollutants, also identified
388 as risk factors, can additionally contribute to the relevant disease burden. For instance,
389 exposure to air pollution was linked to a global burden of 476,000 infant deaths in 2019

390 during the first month of life via increased risks of LBW and premature birth³⁶. LFS induces
 391 peak concentrations of PM_{2.5} lasting for a few weeks or months. Unlike other environmental
 392 exposures, such as urban PM_{2.5}, LFS exposure is unequally distributed spatially and
 393 temporally. The individual-level impacts on exposed mothers are not negligible. Particularly,
 394 LFS exposure frequently occurred in LMICs (Figure 1), where the fire-sourced PM_{2.5}
 395 adversely affected infants with other risk factors, such as malnutrition. However, due to the
 396 lower specificity of our model evaluating birthweight without adjusting for gestation length,
 397 we could not evaluate the clinical outcomes or disease burden of the LFS-attributed
 398 birthweight reductions. As there is an increasing probability of landscape fire under global
 399 climate change, assessing the adverse impacts of LFS exposure is of public health importance
 400 to protect maternal health in LMICs. Future studies should be conducted to answer this
 401 question.

402 This study was associated with the following limitations. First, the lack of data on specific
 403 gestation lengths in the DHSs limited our exploration. Without this information, we could not
 404 distinguish the exposure time window, which introduced potential exposure misclassifications
 405 into the epidemiological analyses. Also, we could not determine the clinical importance of the
 406 analyzed birthweight without normalizing it to gestation length. Second, characterizing LFS
 407 exposure is always challenging. Previous studies measured exposure using surrogates, such as
 408 the temporal duration of landscape fires or satellite remote sensing of fire points. To increase
 409 interpretability, we utilized fire-sourced PM_{2.5} as a direct indicator of ambient exposure to
 410 LFS. However, as the exposure assessments were based on a CTM, the uncertainties
 411 introduced by the modeling procedure could not be avoided completely. The potential errors
 412 enhanced the exposure misclassifications in this study and thus may have resulted in a biased
 413 association. Third, since the analyzed samples were living newborns, our estimates were
 414 potentially subjected to survival bias. For instance, fire-sourced PM_{2.5} was associated with
 415 extremely reduced birthweight, which increases the probability of stillbirth or neonatal death.
 416 In other words, our sample may have included fewer embryos more susceptible to fire-
 417 sourced PM_{2.5} than expected. Therefore, survival bias could have led to an underestimated

association by ignoring those susceptible embryos. Finally, a sibling-matched study requires a selected sample, which may have lowered the representativeness of this study. The inclusion of infants of the DHS in this study was determined by many factors, such as family size and planning, willingness to bear a child, socioeconomic position, and contraceptive behavior. However, some of those factors could not be easily measured, which impeded the development of a set of sampling weights to increase the representativeness of our estimates.

Conclusions

Using a global-scale sibling-matched case-control study, an association was identified between gestational exposure to fire-sourced $PM_{2.5}$ and reduced birthweight in LMICs. The association was sensitive to regression model settings and potential heterogeneity among different subpopulations. Newborns from families with a low baseline level of birthweight were susceptible to LFS-associated birthweight reduction, which was consistent with our finding of a strong and robust association between fire-sourced $PM_{2.5}$ and VLBW. This study adds to the epidemiological evidence on the adverse effects of LFS on birthweight. Relevant interventions regarding frequently occurring landscape fires, such as climate change mitigations, should be taken to protect maternal and infant health.

435 **References**

- 436 1. van der Werf GR, Randerson JT, Giglio L, et al. Global fire emissions and the
437 contribution of deforestation, savanna, forest, agricultural, and peat fires (1997-2009).
438 *Atmos Chem Phys* 2010; **10**(23): 11707-35.
- 439 2. Cascio WE. Wildland fire smoke and human health. *Sci Total Environ* 2018;
440 **624**: 586-95.
- 441 3. Turetsky MR, Kane ES, Harden JW, et al. Recent acceleration of biomass
442 burning and carbon losses in Alaskan forests and peatlands. *Nat Geosci* 2011; **4**(1):
443 27-31.
- 444 4. Hantson S, Pueyo S, Chuvieco E. Global fire size distribution is driven by
445 human impact and climate. *Global Ecol Biogeogr* 2015; **24**(1): 77-86.
- 446 5. Matz CJ, Egyed M, Xi GL, et al. Health impact analysis of PM2.5 from wildfire
447 smoke in Canada (2013-2015,2017-2018). *Science of the Total Environment* 2020;
448 **725**.
- 449 6. Barn PK, Elliott CT, Allen RW, Kosatsky T, Rideout K, Henderson SB.
450 Portable air cleaners should be at the forefront of the public health response to
451 landscape fire smoke. *Environ Health-Glob* 2016; **15**.
- 452 7. Johnston FH, Henderson SB, Chen Y, et al. Estimated global mortality
453 attributable to smoke from landscape fires. *Environ Health Perspect* 2012; **120**(5):
454 695-701.
- 455 8. Delfino RJ, Brummel S, Wu J, et al. The relationship of respiratory and
456 cardiovascular hospital admissions to the southern California wildfires of 2003.

- 457 *Occup Environ Med* 2009; **66**(3): 189-97.
- 458 9. Morgan G, Sheppard V, Khalaj B, et al. Effects of bushfire smoke on daily
459 mortality and hospital admissions in Sydney, Australia. *Epidemiology* 2010; **21**(1):
460 47-55.
- 461 10. Kolarzyk E, Szot WM, Lyszczarz J. Lung function and breathing regulation
462 parameters during pregnancy. *Arch Gynecol Obstet* 2005; **272**(1): 53-8.
- 463 11. Pedersen M, Giorgis-Allemand L, Bernard C, et al. Ambient air pollution and
464 low birthweight: a European cohort study (ESCAPE). *Lancet Respir Med* 2013; **1**(9):
465 695-704.
- 466 12. Padula AM, Yang W, Lurmann FW, Balmes J, Hammond SK, Shaw GM.
467 Prenatal exposure to air pollution, maternal diabetes and preterm birth. *Environ Res*
468 2019; **170**: 160-7.
- 469 13. Bijnens EM, Derom C, Gielen M, et al. Small for gestational age and
470 exposure to particulate air pollution in the early-life environment of twins. *Environ Res*
471 2016; **148**: 39-45.
- 472 14. Lee BE, Park HS, Kim YJ, Park EA, Hong YC, Ha EH. Air pollution exposure
473 and health effects in fetus. *J Prev Med Public Health* 2004; **37**(4): 291-9.
- 474 15. Janssen BG, Byun HM, Gyselaers W, Lefebvre W, Baccarelli AA, Nawrot TS.
475 Placental mitochondrial methylation and exposure to airborne particulate matter in
476 the early life environment: An ENVIRONAGE birth cohort study. *Epigenetics* 2015;
477 **10**(6): 536-44.
- 478 16. Samaras TT, Elrick H, Storms LH. Birthweight, rapid growth, cancer, and

- 479 longevity: a review. *J Natl Med Assoc* 2003; **95**(12): 1170-83.
- 480 17. Hviid A, Melbye M. The impact of birth weight on infectious disease
481 hospitalization in childhood. *Am J Epidemiol* 2007; **165**(7): 756-61.
- 482 18. Blencowe H, Krusevec J, de Onis M, et al. National, regional, and worldwide
483 estimates of low birthweight in 2015, with trends from 2000: a systematic analysis.
484 *Lancet Glob Health* 2019; **7**(7): e849-e60.
- 485 19. Organization WH. Global targets 2025 to improve maternal, infant and young
486 children nutrition. 2017. <https://www.who.int/nutrition/global-target-2025/en/>.
- 487 20. Malley CS, Kuylensstierna JC, Vallack HW, Henze DK, Blencowe H, Ashmore
488 MRJEI. Preterm birth associated with maternal fine particulate matter exposure: a
489 global, regional and national assessment. 2017; **101**: 173-82.
- 490 21. Li C, Yang M, Zhu Z, et al. Maternal exposure to air pollution and the risk of
491 low birth weight: A meta-analysis of cohort studies. *Environ Res* 2020; **190**: 109970.
- 492 22. Li XY, Huang SQ, Jiao AQ, et al. Association between ambient fine
493 particulate matter and preterm birth or term low birth weight: An updated systematic
494 review and meta-analysis. *Environ Pollut* 2017; **227**: 596-605.
- 495 23. Holstius DM, Reid CE, Jesdale BM, Morello-Frosch R. Birth Weight following
496 Pregnancy during the 2003 Southern California Wildfires. *Environ Health Persp* 2012;
497 **120**(9): 1340-5.
- 498 24. Cândido da Silva AM, Moi GP, Mattos IE, Hacon Sde S. Low birth weight at
499 term and the presence of fine particulate matter and carbon monoxide in the Brazilian
500 Amazon: a population-based retrospective cohort study. *BMC pregnancy and*

501 *childbirth* 2014; **14**: 309.

502 25. Abdo M, Ward I, O'Dell K, et al. Impact of Wildfire Smoke on Adverse
503 Pregnancy Outcomes in Colorado, 2007-2015. *Int J Environ Res Public Health* 2019;
504 **16**(19).

505 26. O'Donnell MH, Behie AM. Effects of wildfire disaster exposure on male birth
506 weight in an Australian population. *Evol Med Public Health* 2015; **2015**(1): 344-54.

507 27. Stowell JD, Geng G, Saikawa E, et al. Associations of wildfire smoke PM_{2.5}
508 exposure with cardiorespiratory events in Colorado 2011–2014. *Environment*
509 *international* 2019; **133**: 105151.

510 28. Solazzo E, Bianconi R, Pirovano G, et al. Operational model evaluation for
511 particulate matter in Europe and North America in the context of AQMEII. 2012; **53**:
512 75-92.

513 29. Zhang H, Wang J, García LC, et al. Improving Surface PM_{2.5} Forecasts in
514 the United States Using an Ensemble of Chemical Transport Model Outputs: 1. Bias
515 Correction With Surface Observations in Nonrural Areas. 2020; **125**(14):
516 e2019JD032293.

517 30. Van Donkelaar A, Martin RV, Brauer M, et al. Global Estimates of Fine
518 Particulate Matter Using a Combined Geophysical-Statistical Method with Information
519 from Satellites, Models, and Monitors. *Environmental Science & Technology* 2016;
520 **50**(7): 3762-72.

521 31. Xue T, Zhu T, Geng G, Zhang QJTLPH. Association between pregnancy loss
522 and ambient PM_{2.5} using survey data in Africa: a longitudinal case-control study,

1998–2016. 2019; **3**(5): e219-ee25.

32. Ha S, Mendola P. Are we ready to establish a causal relationship between air pollution and pregnancy loss? *The Lancet Planetary Health* 2019; **3**(5): e198-e9.

33. Curtis DJAohg. Use of siblings as controls in case-control association studies. 1997; **61**(4): 319-33.

34. Xu RB, Yu P, Abramson MJ, et al. Wildfires, Global Climate Change, and Human Health. *New Engl J Med* 2020; **383**(22): 2173-81.

35. Salomon L, Bernard J, Ville YJUiO, Obstetrics GTOJotlSoUi, Gynecology. Estimation of fetal weight: reference range at 20 – 36 weeks' gestation and comparison with actual birth-weight reference range. 2007; **29**(5): 550-5.

36. Impacts on Newborns. <https://www.stateofglobalair.org/health/newborns>.

37. Fountoukis C, Nenes A. ISORROPIA II: a computationally efficient thermodynamic equilibrium model for K⁺ Ca²⁺ Mg²⁺ NH₄⁺ Na⁺ SO₄²⁻ NO₃⁻ Cl⁻ H₂O aerosols. 2007.

38. Park RJ, Jacob DJ, Kumar N, Yantosca RM. Regional visibility statistics in the United States: Natural and transboundary pollution influences, and implications for the Regional Haze Rule. *Atmospheric Environment* 2006; **40**(28): 5405-23.

39. Fairlie TD, Jacob DJ, Park RJ. The impact of transpacific transport of mineral dust in the United States. *Atmospheric Environment* 2007; **41**(6): 1251-66.

40. Pye H, Liao H, Wu S, et al. Effect of changes in climate and emissions on future sulfate-nitrate-ammonium aerosol levels in the United States. *Journal of Geophysical Research: Atmospheres* 2009; **114**(D1).

545 41. Hoesly RM, Smith SJ, Feng L, et al. Historical (1750-2014) anthropogenic
546 emissions of reactive gases and aerosols from the Community Emission Data
547 System (CEDS). *Geoscientific Model Development* 2018; **11**: 369-408.

548