

1 **Women’s Awareness of Obstetric Fistula and its Associated Factors Among**
2 **Reproductive-Age Women in Ethiopia: A Multilevel Analysis Based on**
3 **National Survey Data.**

4 Walleign Alemnew (WA)¹, Bezawit Mulat (BM)^{2*}, Kegnie Shitu (KS)¹

5 ¹ Department of Health Education and Behavioral Sciences, Institute of Public Health College of
6 Medicine, University of Gondar, Gondar, Ethiopia

7 ² Department of Human Physiology, School of Medicine, College of Medicine and Health
8 Sciences, University of Gondar, Gondar, Ethiopia

9 Walleign Alemnew

10 E-mail: walleignaleminew@gmail.com

11 Bezawit Mulat

12 E-mail: mulatbezawit@gmail.com

13 Kegnie Shitu

14 Email: kegnsh@gmail.com

15

16 ***Corresponding Author**

17 Bezawit Mulat

18 E-mail: mulatbezawit@gmail.com

19 Cell Phone: +251963729635

20

21

22 **Abstract**

23 **Objective:** This study aimed to determine the magnitude and associated factors of women's
24 awareness of obstetric fistula (OF).

25 **Methods:** This community based crosssectional study was conducted among 15,683
26 reproductive-aged women. A multi-level multivariable logistic regression analysis employed. A
27 95% CI and p-value < 0.05 were used to declare statistical significance.

28 **Result:** The magnitude of women's awareness of OF was 38% [95%CI: 0.37, 0.39]. Individual
29 level variables such as being in the age group 20-25 [AOR=1.17; 95%CI:1.02, 1.35], 26-30
30 [AOR=1.50; 95%CI: 1.27, 1.76] and >30 [AOR=1.76; 95%CI: 1.50, 2.07], being Muslim
31 religion follower [AOR = 0.83; 95%CI: 0.73, 0.94], attending primary education [AOR=1.70;
32 95%CI: 1.53, 1.89], attending secondary education [AOR=3.43; 95%CI: 2.95, 3.99] and
33 attending higher education [AOR=5.88; 95%CI: 4.66, 7.42], history of pregnancy termination
34 [AOR=1.31; 95%CI: 1.13, 1.51], media exposure at least once a week [AOR= 1.33; 95%CI: 1.2,
35 1.49], internet use [AOR= 2.25; 95%CI: 1.84, 2.75], medium house hold wealth [AOR=1.17;
36 95%CI: 1.02, 1.34], rich house hold wealth [AOR=1.50; 95%CI:1.31, 1.72] and community level
37 factors including high community media exposure [AOR = 1.30; 95%CI: 1.05, 1.61], high
38 community ANC rate [AOR=1.66; 95%CI: 1.37, 2.02] and low health facility distance problem
39 [AOR=1.49; 95%CI: 1.23, 1.81] were significantly associated with awareness of OF.

40 **Conclusion:** The magnitude of women's awareness of OF was very low in Ethiopia. Maternal
41 age, religion, being educated, media exposure, being from a wealthier household, low health
42 facility distance problem, and high community ANC rate were significantly and positively
43 associated factors with awareness of OF.

44

45

46 **Background**

47 Obstetric fistula is a life-altering birth associated injury [1] that creates an abnormal connection
48 between the genital tract and the urinary tract (urogenital fistula) or the gastrointestinal tract
49 (most commonly, rectovaginal fistula) that occur as the result of obstetric trauma, typically from
50 prolonged obstructed labor[2, 3] . It is a public health concern for women and their communities
51 within developing nations, particularly in Africa and Southeast Asia [4] and it is found to be one
52 of the most visible indicators of maternal morbidity [5]. The World Health Organization
53 estimates there are 130,000 new cases of obstetric fistula each year, calculated from an
54 assumption that fistula is likely to occur in 2 percent of the 6.5 million cases of obstructed labor
55 that occur in developing countries [2, 6]. Of these, around 33,000 women are found in Sub-
56 Saharan Africa, and it affects about 1.62 women per 1000 reproductive age in Ethiopia [7, 8].
57 Currently around 36,000 and 39,000 women living with obstetric fistula and that every year there
58 are between 3,300 and 3,750 new cases of obstetric fistula in Ethiopia according to the problem
59 analysis report [9].

60 Pieces of evidence show that the risk of the vaginal fistula is common in settings with a lack of
61 inadequate emergency obstetric care, healthcare workforce shortages, and poor investment in
62 maternity services [6, 10]. Further, sociocultural issues such as early marriage, harmful cultural
63 practices like female genital mutilation, home-based childbirth, poor policy implementation of
64 female education, and misconceptions about childbirth practices are other contributing factors for
65 the high burden of obstetric fistula in Sub-Saharan Africa [11, 12]. Adequate information and
66 awareness of the risk factors, causes, and treatment options for vaginal fistula may help women
67 to take appropriate steps to prevent vaginal fistula. Generally, different researches reported that

68 still there is a low level of awareness about obstetric fistula in Africa, especially in Ethiopia. The
69 magnitude of awareness was found to be 40.8% and 39.5% in Bench sheka zone, south, Ethiopia,
70 and awi zone Amhara region, Ethiopia respectively [7, 9].

71 Despite the high burden of obstetric fistula in Sub-Saharan Africa, studies addressing the
72 awareness of obstetric fistula among women are limited, particularly in Ethiopia. Therefore, this
73 study aims to determine the magnitude of obstetric fistula awareness and factors that could
74 contribute to the awareness of obstetric fistula among women of reproductive age in Ethiopia.
75 Furthermore, findings from this analysis will assist policy-makers and public health
76 programmers to understand the magnitude of obstetric fistula awareness and the contributory
77 factors.

78 **Methods**

79 **Data source, population, and sampling procedure**

80 The present study was based on the most recent Ethiopian Demographic and Health Survey
81 (EDHS) data. The survey was conducted from January 18, 2016, to June 27, 2016. For this
82 survey, a complete list of 84, 915 Enumeration Areas (EAs) from the Ethiopian Population and
83 Housing Census was used as a sampling frame. A stratified two-stage cluster sampling technique
84 was employed. In the first stage, 645 EAs were selected. In the second stage, an average of 28
85 households was selected per cluster/EA. The data is freely available in public and we accessed
86 the dataset used for the present study after we registered and received an authentication letter
87 from the Demographic and Health Survey (DHS) program at The DHS Program - Ethiopia:
88 Standard DHS, 2016 Dataset. For this study, a total weighted sample of 15, 683 reproductive-
89 age women were included. Detailed information about the sampling strategy, questioner, or other
90 important information is available in the 2016 EDHS report.

91 **Variables of the Study**

92 The outcome variable of the present study was women's awareness of obstetric fistula. The
93 variable was dichotomized into **1** = "ever heard of fistula" and **0** = "never heard of fistula". The
94 independent variables were further classified into individual level (level 1) variables and
95 community level (level 2) variables. Individual-level variables included age, marital status,
96 religion, educational status, media exposure, internet access, family wealth index, sexual activity,
97 birth history, pregnancy termination, current working status, and current pregnancy status.
98 Whereas, community variables involved variables directly taken with no aggregation (residence
99 and contextual region), and variables obtained by aggregating individual values into their
100 respected community (community poverty, community female education, community media
101 exposure, community antenatal care service utilization rate, and community health facility
102 distance). Since the aggregate values of each variable didn't follow a normal distribution curve,
103 we categorized the aggregate values of a cluster into groups based on median values.

104 **Operational definitions**

105 **Household wealth quintile:** The wealth index classifications were done in quintiles: poorest;
106 poor; middle; rich; richest. These were computed using principal component analyses (PCA).
107 This variable was further categorized into three categories (Poor, Medium, and Rich) by merging
108 the lower two (poorest and poor) quantiles and the upper two (richest and rich) quantiles.

109 **Media Exposure:** This variable was computed from the frequency of exposure to the two
110 commonest mass media routes (radio and television). In this study exposure to
111 magazines/newspapers was excluded because little (<5%) women were exposed to this channel.
112 The variable was categorized into three categories: no exposure to either media route; exposure
113 to either one of the two routes; and exposure to both within a week.

114 **Contextual region:** For this survey regions were categorized into three categories (agrarian,
115 pastoral, and metropolitan) that may have a strong relationship to health information seeking and
116 awareness of obstetric fistula. The Tigray, Amhara, Oromia, SNNP, Gambella, and Beneshangul
117 Gumuz were recorded as agrarian. The Somali and Afar regions were merged to form the
118 pastoralist region and the city administrations- Addis Ababa, Dire Dawa, and Harar were
119 combined as metropolitan [1].

120 **Community female education:** This is the aggregate value of the educational levels of women
121 based on the average of proportions of educational levels in the community. It was defined as
122 low if the proportion of women with secondary education & above in the community was 0 –
123 12.4 % and high if the proportion was 12.5 –100 %.

124 **Community media exposure:** This variable was derived from the individual responses for
125 exposure to radio or television. It was defined as low if the proportion of women exposed to
126 media in the community was 0–18.7 % and high if the proportion was 18.8–100 %.

127 **Community ANC utilization rate:** This variable is also derived from the individual values for
128 ANC utilization. It was defined as low if the proportion of women who attended at least one
129 ANC visit in the community was 0 – 81.3 % and high if the proportion was between 81.4 –100
130 %.

131 **Community poverty:** With the same procedure, this variable is also derived from an individual
132 household's wealth index. It was defined as high if the proportion of women from the two lowest
133 wealth quintiles in a given community was 25.9–100 % and low if the proportion was 0–26 %.

134 **Community distance to the health facility:** The variable was aggregated from individual
135 perceived distance to a health facility is a big problem. It was as categorized as low if the

136 proportion of women who perceived health facility distance as a big problem in the community
137 was 0–42.2% and categorized as high if the proportion was between 42.2% and 100%.

138 **Data processing and Analysis**

139 Data were extracted from individual records (IR) files and further coding and transformations
140 were done using statistical software, STATA version 14. The weighted samples were used for
141 analysis to adjust for unequal probability of selection and non-response in the original survey.
142 Since the EDHS employed multi-stage stratified cluster sampling techniques, the data have a
143 hierarchical structure. Therefore, a multivariable multilevel binary logistic regression model was
144 used to estimate the fixed and random effects of the factors associated with awareness of fistula.
145 Four models were constructed. The first model also called an empty model which was fitted
146 without any explanatory variables. This model is important for understanding the community
147 variations, and we used it as a reference to estimate how much community factors were able to
148 explain the observed variations in the awareness of fistula. Moreover, this model was used to
149 justify the use of a multilevel statistical framework as it is a litmus paper on whether multilevel
150 or the traditional logistic regression should be used. It was assessed using the Likelihood Ratio
151 test (LR), Median Odds Ratio (MOR), Intra-class Correlation Coefficient (ICC), and
152 Proportional Change of Variance (PCV). The second model contained only individual-level
153 factors. The third contained only community-level factors. Whereas, the final (fourth) model
154 containing both individual and community-level factors. Moreover, the model comparison was
155 done using model deviance, a model with the lowest deviance selected for reporting and
156 interpretation results.

157

158

159

160 **Result**

161 **Individual-level factors**

162 A total of 15683 reproductive-age women were included in this study. The median age of the
163 participants was 27 with an inter-quartile range of 15. Nearly half (47.8%) of the participants
164 didn't attend any formal education and more than half (63.8%) of them were married. Only one-
165 third (33.3%) of them were employed during the survey. Regarding the reproductive history,
166 about 77% (12033), 67.5 (10587), and 7.2% (1135) of the participants were ever had sex, ever
167 had a child, and currently pregnant respectively. Moreover, only 18.5% (2,896), and 4.4% (693)
168 of women had been exposed to either television or radio at least once a week and had ever used
169 the internet respectively (Table 1).

170 **Table 1: Sociodemographic and sexual characteristics of the reproductive age**
171 **women in Ethiopia (n = 15, 683)**

Variables	Frequency	Proportion
Age		
15-19	3, 381	21.6
20-25	3, 655	23.3
26-30	2, 882	18.4
>=31	5, 765	36.7
Religion		
Cristian	10, 580	67.4
Muslim	4, 893	31.1
Others	205	1.3
Highest educational level		
No-formal education	7498	47.8
Primary	5490	35
Secondary	1818	11.6
Higher	877	5.6
Marital status		
Married	10014	63.8
Unmarried	5669	36.2
Ever had a child		
No	5096	32.5

Yes	10587	67.5
Sexual experience		
Never had sex	3650	23.3
Ever had sex	12033	76.7
Currently pregnant		
No or unsure	14548	92.8
Yes	1135	7.2
Ever had a terminated pregnancy		
No	14447	92.1
Yes	1236	7.9
Mass media exposure (TV/Radio)		
Both less than a week or not at all	11,702	74.6
Either at least once a week	2,896	18.5
Both at least once a week	1,085	6.9
Use of Internet		
Never	14,904	95.0
Ever	693	4.4
Currently working		
No	10,463	66.7
Yes	5,220	33.3
Household wealth Index		
Poor	8,420	53.7
Medium	3,100	19.8
Rich	4,164	26.5

172

173 **Community-level variables**

174 The majority (77.8%) of the participants were from rural residency. About 50.8% (7,714), 60.3%
 175 (9,452), 58.0% (9,089), and 62.8% (9,848) were from a community where there is high poverty
 176 low female education, low media exposure and low ANC utilization rate respectively (Table 2).

177 **Table 2: Summary result of community-level factors of women's awareness of obstetric**

178 **fistula in Ethiopia (n=15, 683)**

Variables	Frequency	Proportion
Residence		
Urban	3,476	22.2
Rural	12,207	77.8
Contextual Region		

Agrarian	13,870	88.5
Pastoral	792	5.0
Metropolitan	1,021	6.5
Community poverty		
Low	7,969	49.2
High	7,714	50.8
Community female education		
Low	9,452	60.3
High	6,231	39.7
Community media exposure		
Low	9,089	58.0
High	6,594	42.0
Community ANC utilization rate		
Low	9,848	62.8
High	5,835	37.2
Distance to Health facility is a big problem		
Low	6,656	42.4
High	9,027	57.6

179

180 **Awareness of obstetric fistula**

181 The overall prevalence of awareness of fistula among reproductive-age women was 38% (95%
182 CI: 37%-39%).

183 **Random effect and Model comparison**

184 As is depicted in table 4 the null model revealed that awareness of fistula was significantly
185 varied across communities/clusters. This was assessed by the ICC value in the null model by
186 which 36.0% of the total variation in awareness of fistula was due to the difference between
187 clusters. Moreover, the MOR in the null model, which was 3.63 (95% CI: (3.32, 3.97)),
188 indicated that there was a significant variation between clusters. This shows if a woman moves to
189 another area with a higher probability of awareness of fistula, her risk of having awareness of
190 fistula will increase 3.63 times. The final model that included both the individual and community
191 level factors explained the greatest variance (52 %) in fistula awareness as it was indicated by
192 PCV. All of the three parameters revealed that the effect of clustering was significant. Indeed,

193 based on model fit comparison indices, the final model was found to be the best-fitted model
 194 (had the lowest deviance) (Table 3).

195 **Table 3: Random effect model and model fitness for factors associated with women's**
 196 **awareness of obstetric fistula in Ethiopia (n=15, 683)**

Measure of variation	Model 1(null)	Model 2	Model 3	Model 4
Community level variations (SE)	1.84 (0.18)	0.85(0.07)	0.69 (.06)	0.68 (0.06)
Intra-class correlation coefficient	0.360	0.211	0.174	.171
Median odds ration	3.63 (3.32,3.97)	2.40 (2.24,2.58)	2.20 (2.07,2.35)	2.19 (2.06,2.34)
Proportional Change of Variance	Reference	0.414	0.516	0.52
Model fitness				
Deviance	18290.62	16908.39	17801.16	16775.54

197 **Individual and community level factors associated with women's awareness of**
 198 **obstetric fistula**

199 The multivariable multilevel logistic regression analysis showed that age, religion, education,
 200 ever pregnancy termination, media exposure, internet use, household wealth index, community
 201 media exposure, community health facility distance problem, and community ANC utilization
 202 rate were significantly associated with women's awareness of obstetric fistula at p-value <0.05.

203 The odds of having awareness of fistula was higher among women with the age of 20-25 years
 204 old (AOR =1.17, 95% CI: 1.02, 1.35)), 26-30 years old (AOR= 1.5, 95% CI: (1.27, 1.76)) and >
 205 30 years old (AOR= 1.76, 95% CI: 1.5, 2.07) as compared to women with the age of 15-19 years
 206 old. Women of Islamic religion followers had 17% lower odds of having fistula awareness
 207 (AOR= 0.83, 95% CI: (0.73, 0.94) than Christianity followers. Women with primary (AOR= 1.7,

208 95% CI: (1.53, 1.89)), secondary (AOR= 3.43; 95% CI: (2.95, 3.99)), and tertiary education
 209 (AOR= 5.88; 95% CI: (4.66, 7.42)) had higher odds of having awareness of fistula than women
 210 with no formal education respectively. Moreover, the odds of having fistula awareness were
 211 higher among women who had ever terminated pregnancy than their counterparts (AOR=1.31,
 212 95% CI: (1.13, 1.51)). Women who were exposed to radio and /or TV at least once a week were
 213 more likely to have awareness of fistula than those who were exposed less than a week/ not at all
 214 (AOR = 1.33 95% CI: (1.2, 1.49)). Moreover, the odds of having awareness of fistula were 2.3
 215 times higher among women who had ever used the internet (AOR= 2.25, 95% CI: (1.84, 2.75)).
 216 Regarding Woman's family wealth index, a woman from a family with the highest two quantiles
 217 (rich) and third quantile (medium) was more likely to have awareness of fistula as compared to a
 218 woman from a family with the lowest two quantiles (AOR=1.17, 95% CI: (1.02, 1.34)) and
 219 (AOR= 1.50, 95% CI: (1.31, 1.72)) respectively. The odds of having fistula awareness were 1.3
 220 (AOR=1.30 95%CI: (1.05, 1.61)), 1.66 (AOR=1.66, 95% CI: (1.37, 2.02)) and 1.49 (AOR=1.49,
 221 95% CI: (1.23, 1.81) times higher among a woman from a community with high media exposure,
 222 high ANC utilization, and high community health distance problem respectively (Table 4).

223 **Table 4: A multilevel logistic regression analysis of factors associated with women's**
 224 **awareness of obstetric fistula among reproductive age women in Ethiopia (n=15, 683)**

Variables	Model II AOR (95% CI)	Model III AOR (95% CI)	Model IV AOR (95% CI)
Age			
15-19	1		1
20-25	1.20 (1.04, 1.38) *		1.17 (1.02, 1.35) *
26-30	1.57 (1.33, 1.85) **		1.50 (1.27, 1.76) **
>30	1.87 (1.59, 2.20) **		1.76 (1.50, 2.07) **

Religion		
Christians	1	1
Muslim	0.77 (0.67, 0.87) **	0.83 (0.73, 0.94) **
Education		
No formal Education	1	1
Primary	1.79 (1.61, 1.99) **	1.70 (1.53, 1.89) **
Secondary	3.77 (3.25, 4.38) **	3.43 (2.95, 3.99) **
Higher	6.50 (5.16, 8.18) **	5.88 (4.66, 7.42) **
History of birth		
No	1	1
Yes	1.11 (0.96, 1.29)	1.14 (0.98, 1.33)
Sex history		
No	1	1
Yes	1.15 (0.98, 1.35)	1.16 (0.99, 1.35)
Ever terminated pregnancy		
No	1	1
Yes	1.31 (1.13, 1.52) **	1.31 (1.13, 1.51) **
Media exposure		
Less than a week/never	1	1
At least once a week	1.46 (1.31, 1.62) **	1.33 (1.2, 1.49) **
Internet use		
Never	1	1
Ever	2.43 (1.99, 2.96) **	2.25 (1.84, 2.75) **
Currently working		
No	1	1
Yes	1.12 (1.03, 1.23) *	1.10 (0.9, 1.19)
Household wealth index		
Poor	1	1
Medium	1.27 (1.11, 1.45) **	1.17 (1.02, 1.34) *
Rich	1.92 (1.69, 2.17) **	1.50 (1.31, 1.72) **

Residence		
Rural	1	1
Urban	1.62 (1.23, 2.15) **	1.11 (0.84, 1.48)
Contextual region		
Pastorals	1	1
Agrarians	1.28 (1.07, 1.53) *	1.09 (0.91, 1.31)
Metropolitans	1.29 (0.99, 1.70)	0.97 (0.74, 1.28)
Community-level poverty		
High	1	1
low poverty	1.17 (0.94, 1.46)	0.93(0.74, 1.16)
Community female education		
Low	1	1
High	1.57 (1.27, 1.93) **	1.09 (0.88, 1.34)
Community media exposure		
Low	1	1
High	1.49 (1.21, 1.84) **	1.30 (1.05, 1.61) *
Community ANC rate		
Low	1	1
High	1.81 (1.49, 2.20) **	1.66 (1.37,2.02) **
Health facility distance problem		
High	1	1
Low	1.55 (1.28, 1.88) **	1.49 (1.23, 1.81) **

225 *Note: * statistically significant at p value < 0.05, ** statistically significant at p value < 0.01,*
226 *Model II= a model contained only individual level factors, Model III= a model contained only*
227 *community level factors, Model IV= a model contained both the individual and community level*
228 *factors*

229

230 **Discussion**

231 The study aimed to assess the magnitude and associated factors of awareness of fistula in
232 Ethiopia using EDHS 2016 data. According to the findings of this study, the estimated
233 magnitude of fistula awareness in Ethiopia was 38% which is consistent with studies done in
234 Bench sheka zone, south, Ethiopia, and Awi zone Amhara region Ethiopia 40.8% and 39.5%
235 respectively [7, 9]. The possible justification for the similarity might be due to the same socio-

236 demographic characteristic of the respondents. Whereas the finding is lower than studies
237 conducted in different countries like Nigeria (57.8%), Ghana (45.8%), Nepal (72.99%), and
238 Dabat, northwest, Ethiopia (54.7%)[8, 13-16]. The difference might be due to the study period,
239 study population, study design, and socio-demographic characteristics. In addition to these, the
240 above studies were based on small sample size or they are mostly facility and community-based
241 cross-sectional studies.

242 In multivariable multilevel logistic regression analysis being in the age group 20-25, 26-30 years
243 and, being Muslim religion follower, attending primary education, attending secondary education
244 and attending higher education, history of pregnancy termination, media exposure at least once a
245 week, internet use, medium household wealth, rich households wealth, community ANC and low
246 health facility distance were significantly associated with fistula awareness in Ethiopia.

247 Older women had high odds of fistula awareness compared to young age women; this is might be
248 due to that older women have had different or multiple exposures to media, different health
249 education forums and they have had better opportunities to access/engage in health care services
250 including counseling services [11]. Attending formal education has had higher odds than not
251 attending formal education for fistula awareness. The finding agrees with studies done in the
252 Awi zone of North West Ethiopia, Bench sheka zone, south, Ethiopia Burkina Faso, and
253 Northern Ghana [7, 9, 11]. The possible explanation for these might be those attending formal
254 education have greater opportunities to get information, asking, and getting health services than
255 those who hadn't attended formal education.

256 Besides, our study reveals that previous experience or exposure to pregnancy termination had
257 statistical significance with the level of fistula awareness. Women who got pregnancy
258 termination history or experience were 1.31 times more likely to have a better awareness than
259 those who hadn't get pregnancy termination experiences. The possible justification may be

260 getting or seeking health care services may create an indirect opportunity for counseling and
261 health education; these may increase the level of awareness or their knowledge towards fistula.

262 In this study, community ANC rate has a significant association with awareness on fistula, those
263 who have high community ANC rate have high odds compared to low community ANC rate.
264 The finding is more supported with a study conducted in the Amhara region, Delanta woreda and
265 Asela, Oromia region [17, 18] this implies that more awareness regarding fistula might be linked
266 to ANC coverage increment where women can be counseled about maternal health issues.

267 Other important factors which have a significant association with fistula awareness were wealth
268 quin-tiles, and access to information dissemination platforms (mobile and internet utilization) in
269 this study, which had also been previously reported in studies [19-22] conducted in different
270 areas.

271 In this study, there was a significant association between women's awareness about fistula and
272 distance to the nearby health facility. For women who perceived health, facility distance as a low
273 problem in the community, their awareness is 1.55 more likely to be good than those women
274 who perceived health facility distance as a high problem. The finding is supported by the studies
275 in Ethiopia and China respectively [23, 24] that identify the distance to the health care facility
276 was linked with poor knowledge related to fistula and poor health-seeking behavior. This might
277 be due to as the health facility becomes near; women are willing to visit the health care facilities
278 and have got a chance to attend health-related information and education.

279 **Conclusion and recommendations**

280 The magnitude of awareness towards fistula was very low in Ethiopia. Maternal age, religion,
281 being educated, media exposure, from high wealth family, low distance of health facility

282 problem, and high community ANC rate are factors that were significantly and positively
283 associated with women's awareness towards fistula. Thus, community awareness creation
284 actions are mandatory to offer accurate information on the risk factors, causes, and treatment
285 options of obstetric fistula.

286 **List of abbreviations**

287 ANC Antenatal care, BCC Behavioral Change Communication, EDHS Ethiopian Demographic
288 and Health Survey, AOR Adjusted Odds Ratio, ICC Intra Class Correlation, MOR Median Odd
289 Ratio PCV Proportional Change in Variance

290 **Ethics approval**

291 We registered our research project at the Demographic and Health Survey by submitting a
292 concept note concerning the topic of interest. After evaluation of our concept note the
293 International Review Board of DHS program data archivists gave us a permission letter to
294 download use the dataset for this registered research project.

295 **Consent for publication**

296 Not applicable.

297 **Availability of data and materials**

298 The dataset used for the analysis the present study is freely available at: <https://dhsprogram.com>

299 **Competing interests**

300 The authors declare that they have no competing interests

301 **Funding**

302 There was no fund/grant received for this research project

303 **Authors' contributions**

304 WA, BM and KS were involved in inception, methods and acquisition of the data. KS did data
305 extraction and formal analysis. WA and BM did the interpretation of results and writing the draft
306 manuscript. WA, BM and KS were involved in project administration and revising the final
307 manuscript. All authors read and approved the final manuscript.

308 **Acknowledgments**

309 We would like to acknowledge the MEASURE DHS program for permitting us to obtain and use
310 dataset used for this study.

311

312

313

314

315 **References**

- 316 1. Swain D, Parida SP, Jena SK, Das M, Das H. Prevalence and risk factors of obstetric
317 fistula: implementation of a need-based preventive action plan in a South-eastern rural
318 community of India. *BMC women's health*. 2020;20(1):1-10.
- 319 2. UptoDate, Obstetric fistulas in resource-limited settings. Available from:
320 <https://www.uptodate.com/contents/obstetric-fistulas-in-resource-limited-settings/abstract/1>.
- 321 3. Animut M, Mamo A, Abebe L, Berhe MA, Asfaw S, Birhanu Z. "The sun keeps rising
322 but darkness surrounds us": a qualitative exploration of the lived experiences of women with
323 obstetric fistula in Ethiopia. *BMC women's health*. 2019;19(1):1-10.
- 324 4. Tunçalp Ö, Tripathi V, Landry E, Stanton CK, Ahmed S. Measuring the incidence and
325 prevalence of obstetric fistula: approaches, needs and recommendations. *Bulletin of the World
326 Health Organization*. 2014;93:60-2.
- 327 5. Yismaw L, Alemu K, Addis A, Alene M. Time to recovery from obstetric fistula and
328 determinants in Gondar university teaching and referral hospital, northwest Ethiopia. *BMC
329 women's health*. 2019;19(1):1-8.
- 330 6. Kasamba N, Kaye DK, Mbalinda SN. Community awareness about risk factors,
331 presentation and prevention and obstetric fistula in Nabitovu village, Iganga district, Uganda.
332 *BMC pregnancy and childbirth*. 2013;13(1):1-10.
- 333 7. Balcha WF, Nigussie AA, Beyene FY, Tesfu AA. Awareness and Its Associated Factors
334 of Obstetrics Fistula among Antenatal Care Attendees in Injibara Town Health Institutions, Awi
335 Zone, North West, Ethiopia, 2019. *Journal of Pregnancy*. 2020;2020.
- 336 8. Lucy I, Kingsley C, Japari M, Aaron M, Ayobami A, Steve T. Assessment of the
337 Knowledge and Obstetric Features of Women Affected By Obstetric Fistula at Obstetric Fistula
338 Centre in Bingham University Teaching Hospital, Jos. Nigeria. *Journal of Obstetrics and
339 Gynecology*. 2020;1(1):1-9.
- 340 9. Asefa Z, Amenu D, Berhe A. Awareness of Obstetric Fistula and its Associated Factors
341 among Reproductive-age Group Women in Bench Sheko Zone, Southwest, Ethiopia.
342 Community based Cross-Sectional Study. *J Women's Health Care*. 2020;9(509):2167-0420.20.
- 343 10. Adler A, Ronsmans C, Calvert C, Filippi V. Estimating the prevalence of obstetric
344 fistula: a systematic review and meta-analysis. *BMC pregnancy and childbirth*. 2013;13(1):1-14.
- 345 11. Morhason-Bello IO, Kareem YO, Abdus-Salam RA, Bello OO, Lawal OO, Akinlusi FM,
346 et al. Factors associated with the awareness of vaginal fistula among women of reproductive age:
347 findings from the 2018 Nigerian demographic health cross-sectional survey. *BMJ open*.
348 2020;10(11):e040078.
- 349 12. Deribe K, Fronterre C, Dejene T, Biadgilign S, Deribew A, Abdullah M, et al. Measuring
350 the spatial heterogeneity on the reduction of vaginal fistula burden in Ethiopia between 2005 and
351 2016. *Scientific reports*. 2020;10(1):1-10.
- 352 13. Ezeonu P, Ekwedigwe K, Isikhuemen M, Eliboh M, Onoh R, Lawani L, et al. Awareness
353 of obstetric vesicovaginal fistula among pregnant women in a rural hospital. *Advances in
354 Reproductive Sciences*. 2017;5(3):39-46.
- 355 14. Saeed M, Alhassan A, Opare-Asamoah K, Kuubiere C. A survey on obstetric fistula
356 awareness in Northern Ghana. *Eur J Exp Biol*. 2014;4(4):178-82.
- 357 15. Bashah DT, Worku AG, Mengistu MY. Do community members are aware of obstetric
358 fistula? A community based cross sectional study, Dabat district, Northwest Ethiopia.

- 359 16. Gurung S, Panthi V, Kachapati A. Awareness Regarding Vesicovaginal Fistula among
360 Women Residing in A Municipality of Kapilvastu. *Journal of Universal College of Medical*
361 *Sciences*. 2018;6(2):46-50.
- 362 17. Seteamlak A. Young Women's Knowledge on Prevention of Obstetrics Fistula and its
363 Associated Factors in Delanta District, Amhara Region, North-East Ethiopia: Harmaya
364 University; 2016.
- 365 18. Defar s. Awareness on obstetric fistula and its associated factors among reproductive age
366 women in asella town oromia region, ethiopia. 2018. Available from:
367 <http://localhost:80/xmlui/handle/123456789/14719>
- 368 19. Basheer SA, Pumpaibool T. Knowledge, attitude and maternal health care utilization
369 among married women of reproductive age towards vesicovaginal fistula in Kebbi state, Nigeria.
370 *Journal of Health Research*. 2015;29(2):93-100.
- 371 20. Azanu WK, Dassah ET, Agbeno EK, Ofori AA, Opare Addo HS. Knowledge of
372 obstetric fistula among prenatal clinic attendees and midwives in Mfantseman municipality,
373 Ghana. *International Journal of Gynecology & Obstetrics*. 2020;148:16-21.
- 374 21. Kasamba N, Kaye DK, Mbalinda SN. Community awareness about risk factors,
375 presentation and prevention and obstetric fistula in Nabitovu village, Iganga district, Uganda-
376 *BMC Pregnancy and Childbirth*-Vol. 13, 1-ISBN: 1471-2393-p. 229. 2014.
- 377 22. Banke-Thomas AO, Kouraogo SF, Siribie A, Taddese HB, Mueller JE. Knowledge of
378 obstetric fistula prevention amongst young women in urban and rural Burkina Faso: a cross-
379 sectional study. *PloS one*. 2013;8(12):e85921.
- 380 23. Tollosa DN, Kibret MA. Causes and consequences of obstetric fistula in Ethiopia: a
381 literature review. *International Journal of Medical Research & Health Sciences*. 2013;2(2):261-7.
- 382 24. Yuan F, Qian D, Huang C, Tian M, Xiang Y, He Z, et al. Analysis of awareness of health
383 knowledge among rural residents in Western China. *BMC public health*. 2015;15(1):1-8.

384

385