Additional file 6. Detailed description of studies included in this review.

Table 1. Description of studies included on cotrimoxazole preventive therapy (n= 14). *incl. papers of Okot-Chono (2009), Mugomeri (2018) which refer to both preventive therapies.

First Author (Year)	Country	Context	Study type	Study subject(s) of interest		Study aims	Data collection approach
Ansa (2014)	Ghana	urban	Comparative research	Facilities	(3 hospitals with different delivery models including one-stop-shop, partial integration and referral model)	To compare the impact of increasing integration on TB/HIV service delivery including HIV screening, ART and CPT.	Review of 590 TB patients' health records
Chan (2014)	Malawi Uganda Zimbabwe	rural, semi-urban, urban	Multi-country comparative study	Facilities	(81 health facilities, including 53 primary, 25 secondary and 3 tertiary care facilities)	To describe and compare national and inter-country delivery of training, clinical care, use of laboratories, and monitoring of ART roll-out.	Interviewer administered mixed methods questionnaire for nurses, doctors, and laboratory personnel, Review of ART registers, health records, pharmacy records
Chang (2015)	Uganda	-	Randomised trial	Patients	(442 pre-ART HIV+ adults, including 221 in the intervention group and 221 in the control group)	To assess the effect of peer support on HIV clinic attendance, preventive care use including CPT, risky sexual behaviour and ART initiation.	Survey-based outcome assessment at baseline and after one-year follow-up
Horwood (2010)	South Africa	peri-urban, rural	Cross-sectional descriptive study/ evaluation	Facilities	(6 hospital-based post-natal wards and 27 immunization units of primary health care facilities)	To evaluate the implementation of prevention of mother to child transmission of HIV services and the integration of PMTCT services with routine maternal and child health services. To compare health interventions provided at both types of health care units according to patients' health records with mothers reporting.	Review of 872 HIV+ mothers' health records, Survey-based interviews with 872 mothers regarding the services received, 26 nurses, and 27 lay counsellors regarding their work activities
Kamuhabwa (2015)	Tanzania	-	Retrospective descriptive study	Facilities	(PMTCT units at 4 public hospitals)	To assess CPT delivery in PMTCT units (including initiation, doses prescribed, duration, drug tolerance, adherence).	Review of notes of 498 infants born to HIV+ mothers recorded in PMTCT register books,
				Caregivers Providers	(321 parents/ guardians) (126 healthcare workers)	To identify challenges in the provision of CPT among children born to HIV-infected mothers.	Interviewer administered questionnaire with structured open and closed questions
Kamuhabwa (2016)	Tanzania	urban	Descriptive cross-sectional study	Patients	(353 HIV+ pregnant women)	To assess the level of implementation of CPT for malaria prevention among HIV+ pregnant women.	Review of patient health records,
				Patients	(353 HIV+ pregnant women)	To assess self-reported adherence and reasons for non-adherence.	Interviews using a structured open and closed questions questionnaire,
				Providers	(26 health providers)	To assess the level of knowledge of health providers regarding the new CPT policy.	Self-administered questionnaires,
				Providers	(18 health providers)		
				Facilities	(3 public health facilities, incl. two hospitals and one health centre)	To identify challenges encountered regarding CPT provision at the health facility level.	Focus Group Discussion,
				Facilities		To identify challenges encountered regarding the new CPT policy at the health facility level.	Facility assessment,
Louwagie (2012)	South Africa	-	Historical cohort study	Facilities	(46 health facilities with different delivery models incl. 4 semi-integrated and 42 separate facilities)	To compare access to HIV-related care, including CPT provision between non-integrated (separate) facilities and semi-integrated facilities.	Review of TB Registers, TB treatment records, electronic & paper based ART patient records of 636 TB patients from semi-integrated, and 1297 TB patients from separate facilities
*Mugomeri (2018)	Lesotho	-	Qualitative study	Providers Stake-holders	(42 public healthcare workers, 2 Ministry of Health key informants, 6 representatives of partner organizations)	To establish barriers to the implementation of IPT within the health system context of Lesotho.	Individual semi-structured interviews

Additional file 6. Detailed description of studies included in this review.

Table 1. (…continued)

First Author (Year)	Country	Context	Study type	Study subject(s) of interest		Study aims	Data collection approach
Luyirika (2013)	Uganda	-	Retrospective case study	Facilities	(public and private health facilities including 10 community clinics supported by NGO and 9 or 10? NGO partner health facilities)	To examine retrospectively how family-based care approach (which aims to integrate paediatric HIV care into adult HIV and maternal and child health services) was utilized to integrate HIV services.	Key documents review (i.e. programme strategies, guidelines, national HIV Policies, memos on CPT scale-up),
						To examine operational successes and challenges across sites and identify lessons and recommendations for a family-centred approach.	Key informant interviews with programme management, and patients' families,
						To assess changes in service uptake since the introduction of the model.	Record review using hospital management information system
Mwambete (2013)	Tanzania	urban	Serial clinical and cross-sectional resistance study	Patients	(188 HIV+ adults eligible for CPT)	To investigate incidences of faecal *E. coli* resistance to CTZ in HIV+ patients.	Stool samples for resistance profiling at the baseline visit and 4 follow-up visits,
						To compare and determine changes in CTZ resistance over time. To explore factors that may have contributed to resistance to CTZ and other commonly used antibiotics.	Patient self-reported adherence to CPT during each follow-up visit, questionnaire for initial collection of socio-demographic data
Naikoba (2017)	Uganda	rural	Cluster-randomized trial	Providers	(40 mid-level providers including 20 providers in the 5 intervention facilities and 20 providers in the 5 control facilities)	To assess the effect of one-on-one mentorship for task shifting on clinical knowledge, competence scores and facility performance.	Practical Assessment (using Case-Scenarios and clinical observation to determine mean scores for clinical knowledge and competence,
				Facilities	(10 facilities; incl. 5 intervention, 5 control facilities)	To assess health facility performance (i.e. prescription of CPT) in both arms.	Review of electronic HIV care & treatment database and TB registers
*Okot-Chono (2009)	Uganda	peri-urban, rural	Record review, qualitative study	Districts	(5 districts)	To assess the proportions of patients utilising TB-HIV collaborative services.	Review of TB registers to assess the TB-HIV services utilization (incl. CPT) among 333 TB patients,
				Patients Providers CM	(125 TB patients) (65 health providers) (70 community members)	To assess respondents knowledge, attitudes, practices and beliefs about TB-HIV collaborative services.	Standardised tools applied for 26 Focus Group Discussions, and 34 Key informant interviews,
				Providers	(? facility in-charges, ? focal persons for TB/HIV, ? district health officers, ? expert patients)	To identify reasons explaining the level of implementation of TB-HIV collaborative services.	28 In-depth interviews
Okwera (2015)	Uganda	-	Qualitative study	Patients	(n= 30 HIV+ pulmonary TB suspects)	To assess knowledge, attitudes, and practices of Cotrimoxazole use among HIV infected adults evaluated for TB.	5 Focus Group Discussions
Sibanda (2015)	Zimbabwe	-	Qualitative study	Caregivers	(n= 20 HIV+ mothers whose babies had been initiated CPT)	To explore women´s ability to maintain their babies on CPT after they had been initiated at age six weeks.	In-depth interviews

Additional file 6. Detailed description of studies included in this review.

Table 2. Description of studies included on isoniazid preventive therapy (n= 28). *incl. papers of Okot-Chono (2009), Mugomeri (2018) which refer to both preventive therapies.

First Author (Year)	Country	Context	Study type	Study subject(s) of interest		Study aims	Data collection approach
Adepoju (2020)	Nigeria	-	Retrospective cohort study	Patients	(1134 HIV+ adult attendees of 6 governmental hospitals enrolled in HIV care on IPT)	To assess IPT completion rate for 6 months IPT among PLHIV. To determine predictive factors for IPT completion.	Review of monthly prescription refill records in ART care card, IPT register, and pharmacy order form
Aisu (1995)	Uganda	urban, sub-urban	Operational Assessment	Patients	(520 HIV+ attendees of a voluntary counselling and testing centre, eligible for IPT)	To assess compliance to 6 months IPT among HIV+ individuals with latent TB.	Review of standardized forms (compliance based on attendance to follow-up visits for pill collection and pill count at each follow-up visit),
				Patients	(198 non-compliant patients)	To determine the reasons for missing appointments, to assess toxicity and determine the reasons for missing the follow-up appointment.	Interviews,
				Facility	(1 Voluntary HIV counselling and testing centre)	To assess the incremental unit cost at each step from counselling, testing and ending with completion of IPT course.	Review of expenditure reports, project budgets, UN essential drugs price list
Catalani (2014)	Kenya	rural, urban	Mixed methods assessment	Providers Stake-holders	(24 stakeholders with various background)	To explore the social context of HIV and TB care, and strategies to influence providers care practice.	Key informant interviews using an unstructured interview guide
				Facilities	(9 sites)	To explore the existing electronic medical record system and other existing recording systems.	Qualitative field notes of site observations using a semi-structured observation guide
				Providers	(9 or 10? HIV clinicians)	To evaluate a new clinical decision support system for integrating TB and HIV care and to assess clinicians perceptions of the helpfulness and accuracy of the TB reminder messages.	Mixed methods usability surveys regarding in-context usability of the support system, In-depth interviews
Durovni (2010)	Brazil	-	Preliminary findings of phased cluster-randomised trial and qualitative study	Patients	(1670 HIV+ THRio trial participants of in 29 facilities)	To assess side effects reported during the ongoing trial and to compare IPT completion prior and post training.	Analysis of Trial data set (Trial aimed at testing the effect of provider training & new IPT policy implementation on the incidence of active TB in the HIV+ clinic population),
				Providers	(10 HIV and TB clinicians)	To compare preintervention and post-intervention time to TST and time to IPT.	Interviews,
				Stake-holders	(6 administrators, 6 nurses)	To explore health providers motivations and practices concerning IPT.	2 Focus Group Discussions
Faust (2020)	Ethiopia, Nigeria, India, Angola, Brazil, China, DRC, Indonesia, Kenya, Lesotho, Liberia, Mozambique, Myanmar, South Africa, Tanzania, Thailand, Zambia, Zimbabwe	-	Survey	Countries	(18 high TB/HIV burden countries included in this systematic review)	To identify challenges experienced in TB HBCs with regard to the implementation of LTBI policies and tools.	19-questions survey with open-ended and predefined response options via email

Additional file 6. Detailed description of studies included in this review.

Table 2. (…continued)

First Author (Year)	Country	Context	Study type	Study subject(s) of interest		Study aims	Data collection approach
Gust (2011)	Botswana	-	Sub-study of the Botswana IPT prevention trial	Patients Patients	(42 HIV+ trial participants) (83 HIV+ participants lost to follow-up, 127 HIV+ non-adherent, 252 HIV+ completers)	To elicit information about participants' thoughts and opinions about the trial and the reasons for non-adherence. To identify associations between loss to follow-up/ non-adherence and selected risk factors.	Individual interviews, Focus Group Discussion, Survey instrument with primarily close-ended questions (developed through Interviews & Focus Group Discussion)
Huerga (2016)	Kenya, Swaziland	urban, rural	Two Prospective cohort studies, Operational assessment	Patients Facilities	(n= 550 HIV+ adults in Kenya; and 696 HIV+ adults in Swaziland) (1 urban private TB/HIV clinic in Kenya, two rural primary care clinics in Swaziland)	To assess the acceptability of TST, acceptability to return for TST reading, and initiation of 36 months IPT among PLHIV. To assess the operational feasibility of TST (i.e. time spent on tuberculin skin testing and reading, staff involved, working space adjustments).	Cohort study data set, Interviews with key personnel, direct observation, review of clinic registers
Jacobson (2017)	South Africa	rural	Qualitative study	Patients	(n= 30 HIV+ adults, including 17 IPT completers and 13 IPT defaulters)	To explore knowledge and attitudes of HIV+ adults towards TB, IPT, and HIV, and to explore reasons for initiating IPT and completing/ defaulting treatment.	Individual semi-structured interviews
Jarrett (2019)	South Africa	peri-urban	Multi-method assessment	Providers Patients	(10 nurses working in HIV care, able to prescribe ART, who had participated in a nurse-centred intervention) (10 HIV+ adults, who received HIV care at the intervention clinic, and had participated in a patient-centred intervention)	To examine nurse perceptions of clinical mentorship for promoting IPT uptake. To examine continued barriers to change. To assess adoption of the nurse-centred intervention. To assess patient perceptions of in-queue health education for promoting IPT uptake. To examine continued barriers to change. To assess the fidelity to the patient-centred intervention.	Face-to-face in-depths interviews using semi-structured interview guides, Records of nurse attendance on workshops (training) or in-room consultations (mentoring), Face-to-face in-depth interviews using semi-structured interview guides, Records on patient attendance on in-queue education sessions
Khan (2014)	South Africa	-	Diagnostic performance evaluation	Patients	(737 HIV+ adults in two HIV clinics)	To determine the diagnostic performance of the symptom-based TB screening questionnaire recommended by WHO for people living with HIV in resource-limited settings among adults off and on ART. To evaluate diagnostic performance of a screening strategy using chest radiology and screening questionnaire.	TB screening questionnaire, Sputum specimens for microbiologic testing (smear microscopy, mycobacterial culture), Patient information abstracted from medical charts, Chest radiology
Lai (2019)	Ethiopia	-	Cross-sectional study	Providers	(106 TB-HIV service providers who worked at least 2 months in TB-HIV care and treatment service provision at one of 40 health facilities)	To examine clinician barriers to implementing IPT for people living with HIV, and to assess factors associated with high IPT coverage at the facility level.	Interviewer-administered questionnaire
Lester (2010)	South Africa	-	Qualitative methods study	Patients Providers Providers	(20 HIV+ adults attending for ART appointments) (22 clinic health care providers incl. doctors, nurses, counsellors) (9 doctors and 2 nurses)	To describe operational barriers to IPT delivery and uptake from a provider and user perspective, and to evaluate attitudes of clinic staff and patients to IPT.	42 In-depth interviews, 42 In-depth interviews, 1 Focus group discussion with clinic staff

Additional file 6. Detailed description of studies included in this review.

Table 2. (…continued).

First Author (Year)	Country	Context	Study type	Study subject(s) of interest		Study aims	Data collection approach
Little (2018)	Malawi	rural	Sub-study of the CHEPETSA trial	Patients	(974 adult participants newly diagnosed with HIV, initiated on IPT, including 732 IPT completers, 242 non-completers)	To evaluate predictors of IPT completion in individuals newly diagnosed with HIV (i.e. the relationship between IPT completion and individual and clinic-level predictors, including concomitant receipt of ART, age, pregnancy status, IPT side effects, and alcohol use).	Review of each participant's study file (adherence based on the number of INH pills dispensed at each visit) Study team-administered questionnaires including questions about TB symptoms, possible IPT-related adverse events, ART status, pregnancy, and the number of INH pills dispensed).
McRobie (2017)	Uganda	-	Facility-level policy implementation assessment	Facilities	(24 facilities)	To assess the implementation of national policies about HIV testing, treatment and retention in 24 health facilities (policy and practice comparison).	Review of national HIV policies, 24 structured health facility surveys
				Stake-holders	(7 Key stakeholders, incl. policymakers, implementers, researchers)	To explore site-specific implementation gaps and differences between sites.	7 Key informant interviews
Meribe (2020)	Nigeria	-	Assessment of a provider-focused intervention to increase IPT initiation and completion	Facilities	(27 primary and secondary level of care military clinics and hospitals)	To describe the IPT scale-up intervention and results of the activities implemented at military healthcare facilities.	Review of routinely collected program data retrieved from the district health information system (DHIS-2), which were entered at the facility level from paper registers into DHIS-2, Quality assessments conducted at the sites during quarterly site visits
Mindachew (2011)	Ethiopia	urban	Analytical cross-sectional study	Patients	(319 HIV+ adults taking IPT for at least one month) at 4 hospitals	To determine self-reported adherence to IPT within seven days recall period among HIV+ adults and to assess factors associated with adherence to IPT.	Interviewer administered pre-tested structured questionnaire, including self-reported adherence to IPT
*Mugomeri (2018)	Lesotho	-	Qualitative study	Providers Stake-holders	(42 public healthcare workers, 2 Ministry of Health key informants, 6 representatives of partner organizations)	To establish barriers to the implementation of IPT within the health system context of Lesotho.	Individual semi-structured interviews
Mugomeri 2019	Lesotho	-	Retrospective cohort study	Patients	4122 HIV+ patients enrolled in HIV care, on ART at eight district hospitals	To assess the IPT initiation rate and retention of PLHIV on IPT, and to assess predictors of IPT initiation and interruption.	Review of the paper-based ART and IPT registers of HIV+ hospital attendees enrolled in HIV care between 2004 and 2016
Munseri (2008)	Tanzania	-	Sub-study of the TB vaccine trial	Patients	(568 HIV+ adult vaccine trial participants who were offered IPT)	To assess factors related to completion of IPT among HIV+ adults.	Review of study records (i.e. the number of patients offered and accepted IPT, completion, non-completion),
				Patients	(n= 109 HIV+ completers, 8 non-completers)	To assess knowledge, attitudes and beliefs regarding IPT.	Interviews using a standard questionnaire
Ngamvithayapong (1997)	Thailand	-	Prospective cohort study	Patients	(n= 412 HIV+ adults initiating IPT, including blood donors, outpatients, female commercial sex workers, anonymous clinic clients)	To determine the completion rate and level of adherence to nine months IPT among HIV+ individuals in northern Thailand.	Adherence Assessment (based on monthly clinic attendance for IPT collection and pill count),
				Patients	(72 HIV+ adults, incl. 50 defaulter and 22 who missed clinic appointments)	To determine the reasons for non-adherence.	Interviews,
				Patients	(n= 28 completers)	To determine motivation and methods used in order to adhere to IPT.	5 Focus Group Discussions

Additional file 6. Detailed description of studies included in this review.

Table 2. (…continued).

First Author (Year)	Country	Context	Study type	Study subject(s) of interest		Study aims	Data collection approach
*Okot-Chono (2009)	Uganda	peri-urban, rural	Record review, qualitative study	Districts	(5 districts)	To assess barriers to implementation of TB-HIV collaborative services.	Review of TB registers to assess the TB-HIV services utilization (incl. CPT) among 333 TB patients,
				Patients Providers CM	(125 TB patients) (65 health providers) (70 community members)	To assess respondents knowledge, attitudes, practices and beliefs about TB-HIV collaborative services.	Standardised tools applied for 26 Focus Group Discussions, and 34 Key informant interviews,
				Providers	(? facility in-charges, ? focal persons for TB/HIV, ? district health officers, ? expert patients)	To identify reasons explaining the level of implementation of TB/HIV collaborative services.	28 In-depth interviews
Reddy (2020)	India	60% rural	Mixed-methods study	Patients	(4020 newly registered PLHIV started on ART in 2 districts)	To assess the implementation of IPT among PLHIV, who were newly initiated on ART between 01/2017 and 06/2018, including [%] eligible for, [%] who started IPT, and among those initiated, [%] who completed IPT. To assess factors associated with initiation and non-completion of IPT.	Review of routinely collected program data,
				Providers	(22 Healthcare providers involved in IPT implementation)	Reasons for non-initiation and non-completion from healthcare providers' perspective.	Face-to-face in-depth interviews,
				Patients	(8 HIV+ adult patients, including 5 IPT completers, and 3 interrupters)	Reasons for non-initiation and non-completion from a patients' perspective.	Face-to-face in-depth interviews
Rowe (2005)	South Africa	rural	Record review, qualitative study	Patients	(n= 87 HIV+ adult clinic attendees who initiated 6-month IPT)	To describe the study population of HIV+ individuals attending the hospital-based HIV clinic that initiated IPT.	Review of patient records (i.e. patients eligible, initiated, ongoing collection of medication, side effects),
				Patients Providers	(18 HIV+ adult clinic attendees, incl. 6 eligible for IPT, 6 IPT interrupters and 6 IPT completers) (2 health care workers who routinely administered IPT)	To explore HIV+ patients and health workers' perspectives regarding the adherence to IPT and to derive lessons for improving access to care amongst HIV+ individuals in resource-poor settings.	In-depth interviews applying open-ended questions
Selehelo (2019)	South Africa	-	Qualitative study	Patients	(14 HIV+ community health centre attendees on IPT for at least 6 weeks)	To explore and describe the experiences of PLHIV regarding IPT provision to derive strategies to improve IPT uptake in a community health centre (CHC).	Face-to-face in-depth interviews
Szakacs (2006)	South Africa	sub-urban	Assessment of adherence to IPT, Cross-sectional study	Patients	(301 HIV+ adult outpatients receiving IPT at two hospitals)	To assess adherence to IPT (using in-house prepared INH urine test strips) among HIV+ adults and to identify predictors of positive urine test results.	Once off adherence assessment (urine-based INH metabolite testing) and interviewer-administered questionnaire (developed from the review of previous studies)

Additional file 6. Detailed description of studies included in this review.

Table 2. (…continued).

First Author (Year)	Country	Context	Study type	Study subject(s) of interest		Study aims	Data collection approach
Tram (2019)	Uganda	rural	Cross-sectional study (sub-study of SEARCH HIV test and treat trial)	Patients	(305 adult HIV patients on ART who started IPT in 5 government-sponsored clinics)	To evaluate IPT completion rates among trial participants receiving IPT as part of a patient-centred, multicomponent HIV care model, and patients receiving IPT as routine non-differentiated HIV care in the same health centre.	Review of IPT treatment registers to determine IPT completion (based on receipt of 6 months of IPT),
				Patients	(180 of the 305 adult HIV patients, including 94 non-differentiated care attendees, and 86 differentiated care attendees)	To assess individual, interpersonal, and structural predictors of IPT completion, as well as underlying potential differences in IPT completion by care model.	Nested (interviewer-administered) survey on barriers to IPT completion
Van Ginderdeuren (2019)	South Africa	urban	Before-and-after study)	Facilities	(n= 3 primary care health facilities)	To compare IPT and TST uptake at three primary care health facilities before and after the intervention.	Record review (IPT uptake and TST placement based on routine clinic registers in paper/ electronic format; INH prescription based on pharmacy records),
				Facilities	(n= 2 primary care health facilities)	To determine fidelity to the 2014 South African IPT guidelines at two primary care health facilities, following the implementation of the intervention.	Review of paper medical files and electronic laboratory databases to compare HCWs decisions made with decisions expected,
				Providers	(25 health care workers involved in IPT prescription	To identify remaining barriers for IPT and TST implementation perceived by HCWs.	Structured questionnaire survey
Wambiya (2018)	Kenya	urban (cities)	Qualitative study	Providers	(n= 18 healthcare providers working in three HIV clinics)	To explore factors influencing the acceptability of IPT among healthcare providers in selected HIV clinics in Nairobi.	Individual semi-structured in-depth interviews