

A Direct Capture Method for Purification and Detection of Viral Nucleic Acid Enables Epidemiological Surveillance of SARS-CoV-2

Subhanjan Mondal^{1,*}, Nathan Feirer¹, Michael Brockman, Melanie A. Preston, Sarah J. Teter, Dongping Ma, Said A. Goueli, Sameer Moorji, Brigitta Saul, James J. Cali

Promega Corporation, 5430 E Cheryl Pkwy, Fitchburg, WI, 53711.

Supplementary Information

<u>Oligo</u>	<u>Sequence</u>	<u>Reference</u>
2019-nCoV_N1-F	5'-GACCCCAAATCAGCGAAAT-3'	US CDC
2019-nCoV_N1-R	5'-TCTGGTACTGCCAGTTGAATCTG-3'	US CDC
2019-nCoV_N1-P	5'-56-FAMACCCCGCAT-ZEN-TACGTTTGGTGGACC-IABkFQ-3'	US CDC
2019-nCoV_N2-F	5'-TTACAAACATTGGCCGCAAA-IABkFQ-3'	US CDC
2019-nCoV_N2-R	5'-GCGCGACATTCCGAAGAA-3'	US CDC
2019-nCoV_N2-P	5'-56-FAM/ACAATTTGC-ZEN-CCCCAGCGCTTCAG-IABkFQ-3'	US CDC
E_Sarbeco_F1 Fwd	5'-ACAGGTACGTTAATAGTTAATAGCGT-3'	Corman <i>et al.</i> ¹
E_Sarbeco_R2 Rev	5'-ATATTGCAGCAGTACGCACACA-3'	Corman <i>et al.</i> ¹
E_Sarbeco_P1	5'-56-FAM-ACACTAGCC-ZEN-ATCCTTACTGCGCTTCG-IABkFQ-3'	Corman <i>et al.</i> ¹
PMMV FP	5'-GAGTGGTTTGACCTTAACGTTGA-3'	Kitajima <i>et al.</i> ²
PMMV RP	5'-TTGTGCGTTGCAATGCAAGT-3'	Kitajima <i>et al.</i> ²
PMMV P	5'-Quasar 670-CCTACCGAAGCAAATG-BHQ1-3'	Kitajima <i>et al.</i> ²
MS2 F1	5'-TCCTAAAAGATGGAAACCCGATT-3'	Zambenedetti <i>et al.</i> ³
MS2 R1	5'-GGCCGGCGTCTATTAGTAGATG-3'	Zambenedetti <i>et al.</i> ³
MS2 P1	5'- FAM-CCTCAGCAATCGCAGCAAACCTCCG-BHQ1-3'	Zambenedetti <i>et al.</i> ³

CC169	5'-CATTCCGGATACTGCGATTTTAAGTG-3'	This Paper
NF001	5'-CGCCAAAAGCACTCTGATTGAC-3'	This Paper
NF002	5'-GCTTCCCCGACTTCTTTTCTGA-3'	This Paper
NF003	5'-CCAACCCTATTTTCATTCTTCGCC-3'	This Paper
NF004	5'-CTGGAAGATGGAAGCGTTTTGC-3'	This Paper
DS121-Org-560	5'- CAL Fluor Orange 560-CGCCCCCAGAAGCAATTTCTGTGTAAG-BHQ1-3'	This Paper

Table S1: Primers and probes used in this study. Oligos were ordered from either IDT (Coralville, IA) or Biosearch (Novato, CA)

Pathogen	N1	N2	E
SARS-CoV-2	+	+	+
OC43	-	-	-
229E	-	-	-
HKU1	-	-	-
NL63	-	-	-
Influenza A	-	-	-
RSV	-	-	-
<i>Legionella pneumophila</i>	-	-	-
<i>Pseudomonas aeruginosa</i>	-	-	-

Table S2: The SARS-CoV-2 RT-qPCR detection kit described in the manuscript was tested for specificity of nucleic acid amplification using all three SARS-CoV-2 gene targets. Purified genomic DNA or RNA was purchased from commercial sources (American Type Culture Collection, ATCC) and amplification was performed using protocol described in the methods section. Plus sign indicates presence of RT-qPCR signal from indicated pathogen.

References

- 1 Corman, V. M. *et al.* Detection of 2019 novel coronavirus (2019-nCoV) by real-time RT-PCR. *Euro Surveill* **25**, doi:10.2807/1560-7917.ES.2020.25.3.2000045 (2020).
- 2 Kitajima, M., Sassi, H. P. & Torrey, J. R. Pepper mild mottle virus as a water quality indicator. *npj Clean Water* **1**, 19, doi:10.1038/s41545-018-0019-5 (2018).
- 3 Zambenedetti, M. R. *et al.* Internal control for real-time polymerase chain reaction based on MS2 bacteriophage for RNA viruses diagnostics. *Mem Inst Oswaldo Cruz* **112**, 339-347, doi:10.1590/0074-02760160380 (2017).