Research article

COVID-19 Pandemic and Prevalence of Self-care Practices among the Future Physicians: A Bangladesh Study

Fatema Johora¹, Asma Akter Abbasy², Fatiha Tasmin Jeenia³, Mithun Chandro Bhowmik⁴, Mohsena Aktar⁵, Nargis Akhter Choudhury⁶, Priyanka Moitra⁷, Jannatul Ferdoush^{8*}

Author Affiliations

¹Associate Professor, Department of Pharmacology & Therapeutics, Army Medical College Bogura, Bogura, Bangladesh

²Associate Professor, Department of Pharmacology & Therapeutics, Brahmanbaria Medical College, Brahmanbaria, Bangladesh

³ Assistant Professor, Department of Pharmacology & Therapeutics, Chattogram International Medical College, Chattogram, Bangladesh

⁴ Lecturer, Department of Pharmacology & Therapeutics, Rangpur Medical College, Rangpur, Bangladesh

⁵Assistant Professor, Department of Pharmacology & Therapeutics, Cumilla Medical College, Cumilla

⁶Assistant Professor, Department of Pharmacology & Therapeutics, Sheikh Hasina Medical College, Habiganj

⁷Lecturer, Department of Pharmacology & Therapeutics, Colonel Malek Medical College, Manikganj, Bangladesh

⁸Associate. Professor, Department of Pharmacology & Therapeutics, BGC Trust Medical College, Chattogram, Bangladesh

Abstract

Background: Since December 2019, the novel coronavirus, SARS-CoV-2, has garnered global attention due to its rapid transmission, which has infected more than twenty nine million people worldwide. World is facing enormous stress and anxiety as there is no effective medicine or vaccine to treat or prevent COVID-19 till date. Experts are recommending self-care like social distancing, respiratory etiquette, hand washing, using face mask to prevent corona virus infection.

Materials and methods: This descriptive cross-sectional study was designed to assess the prevalence of self-care practice among the undergraduate medical students (4th year) of 14 medical colleges of Bangladesh during COVID-19 pandemic. A structured questionnaire survey linked in the google form was used as study instrument and was distributed among

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

study population through email, messenger, whatsapp and other social media during the month of October 2020. Total 916 students were participated in the study.

Results: 79.8% of students reported self-care practice in study period. 44.98% of students went outside once in a week. 90.5%, 70.96% and 52.62% of respondents always used face mask, followed 20 seconds hand washing principle and maintained social distancing. Face masks (97.8%), sanitizers (76.7%) and gloves (71.9%) are most common items purchased as protective measures. Most of the students (76.9%) follow their hobbies as a coping strategy to overcome psychological stress, while 6% of students took professional help.

Conclusion: Suboptimal practice of self-care was found among the undergraduate medical students of Bangladesh.

INTRODUCTION

The outbreak of corona virus disease (COVID-19) becomes a global crisis. In December 2019, few cases of atypical pneumonia were reported in Wuhan, China and later identified as novel coronavirus infection caused by Severe acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2).^[1, 2] On January 30, 2020 WHO declared it's as a global health emergency and later as pandemic on March 11, 2020. By 17 September 2020, total 216 countries, areas or territories are affected with case, 29679284 are already infected and 936521 died because of COVID-19 pandemic.^[3, 4] In Bangladesh, first case of COVID-19 was detected on 8 March 2020 and rapidly spread out throughout the country.^[5] The COVID-19 causes a plethora of clinical manifestations, and the severity and outcomes may vary, depending on the underlying co-morbidities like diabetes, heart diseases, hypertension, COPD, bronchial asthma as well as age, sex, and geographic locations of the patients.^[6] The rising rates of cases are putting healthcare system in jeopardy.^[7, 8] Several therapeutic agents have been used for the treatment of COVID-19. However, there is no effective antiviral drug or vaccine to combat COVID-19 pandemic.^[9, 10] As it's a highly contagious disease, non-pharmaceutical interventions (NPIs) remain the primary strategy to control COVID-19. The NPIs include public health control measures such as quarantine, travel restrictions, isolation, social distancing, and infection prevention and control to reduce transmission of the disease. Several impact assessment studies have shown that public health interventions are associated with the reduced transmission of COVID-19.^[11, 12, 13] Prevention of infection control is one of the pivotal measures in the absence of an effective vaccine or treatment.^[14] World health organization has been repeatedly emphasizing this strategy through self-care practice.^[15]

Self-care is the ability of individuals, families and communities to promote health, prevent disease, maintain health and to cope with illness and disability with or without the support of a healthcare provider. It encompasses several issues including hygiene, nutrition, lifestyle, environmental, socio-economic factors and self medication. Promotion of self care is a means to empower individuals, families and communities for informed health decision-making. It has the potential of improving the efficiency of health systems and contributing towards health equity.^[16, 17] In a humanitarian or crisis situation like Covid-19 pandemic, self-care could play an important role to improve health-related outcome.^[18] Practicing of responsible self-care is crucial in decreasing the burden on healthcare systems through self-management of mild symptoms as well as implementation of preventive measures particularly during this time of crisis.^[19] Physical distancing, good respiratory hygiene and hand washing are important examples of self-care actions those has been promoted by WHO every day to protect individual from COVID-19, and there are many other areas in which self care can contribute to maintain physical and mental wellbeing during the coronavirus disease pandemic.^[17] Education of individual person is important to get fruitful effects of self-care practice as irresponsible practice can bring numerous potential hazards.^[19, 20] From the beginning of COVID-19 pandemic, world has been observing a pile of myths in mass media and social media.^[21, 22] Practice of self-care is influenced by several factors like education, gender, socioeconomic status and availability of medicine.^[23] But there is no evidence regarding self-care practice among the future physicians.

Hence, the present study will be carried out with the attempt to find out the prevalence of self-care practice among the undergraduate medical students of Bangladesh during Covid-19 pandemic. Finding of this research will be helpful to develop an educational intervention program on self-care practice will be integrated in order to improve rational use of medicine among the future physicians.

Material & Methods

The objective of the study was to describe pattern of self-care practice among the undergraduate medical students of Bangladesh during COVID-19 pandemic.

Study Design and Population

A descriptive cross-sectional study was designed to meet the study objective. The study population comprised of 4th year students of fourteen medical colleges of Bangladesh including government (Armed Forces Medical College, Cumilla Medical College, Colonel Malek Medical College, Manikganj and Rangpur Medical College) and non-government medical colleges (Army Medical College Bogura, Army Medical College Chattogram, BGC Trust Medical College, Brahmanbaria Medical College, Chattogram International Medical College, Jalalabad Ragib-Rabeya Medical College, Sylhet, Jashore Medical College, Khawja Yunus Ali Medical College, Sirajganj, Medical College for Women, Dhaka and US-Bangla Medical College, Narayanganj) in October 2020. Total 916 students were participated in the study.

Study Instrument

A structured questionnaire was used for data collection and questionnaire was validated before survey.

Procedure

Ethical approval was taken from the Institutional Review Board (IRB) of BGC Trust Medical College, Chittagong. Permission was taken from college authorities and informed consent was taken from the participants of the Structured Questionnaire Survey. Researchers explained the nature and purpose of the survey to the students during a virtual class. This self-administered questionnaire was linked in google form and was distributed among study population through email, messenger, whatsapp and other social media who gave consent. To assure the quality, students filled and submitted the questionnaire quickly during end of class. Later, this web-based questionnaire was sent to students who were absent in the class through email. A reminder mail or message was given on 7th day and 15th day of the primary one. The response generated by the students was received through google drive, and it did not accept double response from same participant. To maintain confidentiality, responses were anonymous.

Statistical analysis

Data was compiled, presented and analyzed using Microsoft Excel 2007, and was expressed as percentage.

Results

Nine hundreds and sixteen respondents were covered during the study period, of which 326 (35.59%) were males and 590 (64.41%) were females. Total 305 (33.23%) students responded that anyone in family including himself was diagnosed as COVID-19 and among them 242 (26.42%) were RT-PCR positive (**Table I**). 731 (79.8%) students reported self-care practice in study period. 412 (44.98%) students went outside once in a week. 829 (90.5%) respondents always used facemask while go outside. 650 (70.96%) students always followed

20 sec hand washing principle, while 482 (52.62%) students always maintained social distancing (Table II).

Table I: Demographic data

Variables	Frequency	Percentage
Gender	Male	326
	Female	590
Diagnosed case of COVID-19 in family	305	33.23%
RT-PCR positive in family	242	26.42%

Table II: Responses of medical students about different aspects of self-care practice in last 06 months

Prevalence of self-care practice	Frequency	Percentage
Yes	731	79.8%
No	94	10.26%
Not sure	91	9.93%
Rate of going outside	Frequency	Percentage
Everyday	46	5.02%
2-3 days a week	142	15.5%
Once a week	412	44.98%
Once in 2 weeks	156	17.03%
Once a month	160	17.47%
Usage of facemask while going outside	Frequency	Percentage
Always	829	90.5%
Sometimes	55	6.0%
Occasional	26	2.84%
Never	6	0.66%
Following of 20 seconds hand washing with soap-water principle	Frequency	Percentage
Always	650	70.96%
Sometimes	203	22.16%
Occasional	50	5.46%
Never	13	1.42%
Maintenance of social distancing	Frequency	Percentage
Always	482	52.62
Sometimes	333	36.35
Occasional	79	8.62
Never	22	2.4

Figure 1 showed that facemasks (97.8%) and sterilizers, sanitizers (76.7%) are most common items purchased as protective measures, followed by gloves (71.9%).

Figure 1: Items purchased as protective measures

Figure 2 showed that most of the students (76.9%) follow their hobbies to keep up mental health status, only 55 (6%) students took professional help.

Figure 2: Measures taken for psychological and emotional well-being

226 (25%) students participated online course on coronavirus developed by Directorate General Of Health Service (DGHS) (**Figure 3**).

Figure 3: Participating online course on Coronavirus disease

Discussion

For last several months, world has been observing Covid-19 pandemic with enormous stress and anxiety. Still now, there is no effective medicine or vaccine to combat COVID-19. Lots of misconception are going on about preventive measures to control this highly contagious microorganisms. Current study was conducted in this context to explore self-care practices among undergraduate medical students of Bangladesh.

Social distancing, good respiratory hygiene and hand washing have been recommended by experts as self-care practice to prevent corona virus infection. 79.8% students responded about practicing self-care practice in last 6 months but it was expected that everyone could practice self-care as they are privileged with knowledge of medicine as well as information technology, may be a certain proportion of students were not familiar with the term 'self-care'. Most of the students followed 'stay home principle' to avoid chance of infection and this finding was similar to another study conducted among medical students.^[24]

Although there were some controversies at the beginning, current best evidence suggests that facemask is beneficial for protection of both healthcare professionals and the public as wearing of mask reduces the transmissibility per contact by reducing transmission of infected droplets along with personal protection for the mask wearer. It is considered as most effective way to control spread of virus when compliance is high among larger population.^[25, 26, 27] But use of masks is an evolving and cultural phenomenon, varies widely from country to country.^[28, 29] In our study, most of the participants reported using of face mask always while going outside that was similar to two recent studies conducted in Bangladesh,^[30, 31] poor practices were observed in studies conducted on pharmacy and medical students,^[24, 32] probably cultural variation was the reason as those studies were conducted in Egypt and Jordan.^[28, 29] Hand hygiene is a key to prevent the spread of SARS-Cov-2. 20 seconds hand washing with soap water is considered as one of the most cost-effective way to mitigate COVID-19.^[33] Social distancing measures are other important tools to reduce the morbidity and mortality of

COVID-19 to some degrees, and it seems to be crucial to control the pandemic, especially in the absence of definitive treatments and vaccines.^[34, 35] Suboptimal practices were found among undergraduate medical students of Bangladesh and this was contrary to similar studies^[24, 31,32]

The extended practice of self-care among individuals has contributed to better health and more sustainable healthcare systems, and COVID-19 pandemic has prompted this trend. Throughout the pandemic, many self-care products have been recommended to help treat the symptoms of coronavirus.^[15, 18] In current study, masks, gloves, sanitizers, pulse oxymetry, thermometers were bought as protective measures, and these buying behaviors reflect the extended practice of self-care. From the very beginning of pandemic, people started to buy daily necessities as well as medicine and equipments^[36, 37], and study showed relationship of higher level of education with increased buying behavior.^[29]

COVID-19 pandemic has caused major disruption of education systems worldwide including abrupt shut down of medical schools. Sudden changes in curricular delivery, loss of peer interaction, social isolation, uncertainty, financial stress all are negatively affecting mental well-being of medical students.^[38] Most of the students resorted to following their hobbies, religious works, meditation, where few of them seek professional help to cope up the stress. And these kind of activities were used as coping strategies among medical students.^[38] Directorate General of Health Services (DGHS) introduced an online course to orient and provide useful information to healthcare professionals as well as general population on the COVID-19 with the aim to provide optimum services to COVID-19 patients while ensuring one's own protection and safety.^[39] Poor participation of medical students in this course might be because of their less familiarity and acceptance with distance learning process.^[40]

Undergraduate medical students are the 'future physicians', so it was expected that they would adhere completely with self-care practice to prevent transmission of coronavirus but suboptimal practice of self-care was found in this study.

Conclusion

Self-care practices as usage of face masks, hand hygiene, social distancing, taking care of individual's mental health are crucial measures to mitigate mortality and morbidity of COVID-19. In current study, suboptimal practices of these preventive measures were found among the undergraduate medical students of Bangladesh. Educational intervention on self-care practice might be helpful in improving current practices.

References

1. Zhu N, Zhang D, Wang W, Li X, Yang B, Song J, et al. A Novel Coronavirus from Patients with Pneumonia in China, 2019. *N Engl J Med*. 2020; 382(8):727-33
2. Zhou P, Yang XL, Wang XG, Hu B, Zhang L, Zhang W, et al. A pneumonia outbreak associated with a new coronavirus of probable bat origin. *Nature*. 2020; 579(7798): 270-3.
3. World Health Organization (WHO). Coronavirus disease (COVID-19) Weekly Epidemiological Update and Weekly Operational Update. World Health Organization, Geneva, Switzerland. 2020. Available on: <https://www.who.int/emergencies/diseases/novel-coronavirus-2019/situation-reports/> [Access on: 17/09/2020]
4. World Health Organization (WHO). Coronavirus disease (COVID-19) pandemic. World Health Organization, Geneva, Switzerland. 2020. Available on: <https://www.who.int/emergencies/diseases/novel-coronavirus-2019> [Access on: 17/09/2020]
5. World Health Organization (WHO). Coronavirus disease (COVID-2019) Bangladesh situation reports. World Health Organization, Geneva, Switzerland. 2020. Available on: [https://www.who.int/bangladesh/emergencies/coronavirus-disease-\(covid-19\)-update/coronavirus-disease-\(covid-2019\)-bangladesh-situation-reports](https://www.who.int/bangladesh/emergencies/coronavirus-disease-(covid-19)-update/coronavirus-disease-(covid-2019)-bangladesh-situation-reports) [Access on: 17/09/2020]
6. Center for Disease Control (CDC). Assessing Risk Factors for Severe COVID-19 Illness. CDC, Atlanta, USA. 2020. Available at: <https://www.cdc.gov/coronavirus/2019-ncov/covid-data/investigations-discovery/assessingrisk-factors.html> [Access on: 17/09/2020]
7. Blumenthal D, Fowler EJ, Abrahams M, Collins SR. Covid-19 — Implications for the Health Care System. *New England Journal of Medicine*, 2020. Available at: https://www.nejm.org/doi/full/10.1056/NEJMs2021088?query=recirc_curatedRelated_article [Access on: 17/09/2020]
8. World Health Organization (WHO). COVID-19 significantly impacts health services for noncommunicable diseases. World Health Organization, Geneva, Switzerland. 2020. **Available at:** <https://www.who.int/news-room/detail/01-06-2020-covid-19-significantly-impacts-health-services-for-noncommunicable-diseases> [Access on: 17/09/2020]
9. Cennimo DJ, Bergman SJ. Coronavirus Disease 2019 (COVID-19). *Medscape*, 2020. Available at: <https://emedicine.medscape.com/article/2500114-overview> [Access on: 17/09/2020]
10. Baser A, Das D, Faiz MA. Coronavirus Disease 2019 (COVID-19) – An Overview *Journal of Bangladesh College of Physicians and Surgeons*. 2020; 38: 84-90
11. Anderson RM, Heesterbeek H, Klinkenberg D, Hollingsworth TD. How will country-based mitigation measures influence the course of the COVID-19 epidemic? *Lancet*. 2020; 3 95(10228):931-4.
12. Cowling BJ, Ali ST, Ng TWY, Tsang TK, Li JCM, Fong MW, et al. Impact assessment of non-pharmaceutical interventions against coronavirus disease 2019 and influenza in Hong Kong: an observational study. *Lancet Public Health*. 2020;5(5):e279-e88.
13. Pan A, Liu L, Wang C, Guo H, Hao X, Wang Q, et al. Association of Public Health Interventions With the Epidemiology of the COVID-19 Outbreak in Wuhan, China. *JAMA*. 2020
14. Pan A, Liu L, Wang C, Guo H, Hao X, Wang Q, et al. Association of Public Health Interventions With the Epidemiology of the COVID-19 Outbreak in Wuhan, China. *JAMA*. 2020
15. World Health Organization (WHO). Self care during COVID-19. World Health Organization, Geneva, Switzerland. 2020. Available at: <https://www.who.int/news-room/photo-story/photo-story-detail/self-care-during-covid-19> [Access on: 17/09/2020]
16. World Health Organization. Self care for health. WHO Regional Office for South-East Asia. 2014. Available at: <https://apps.who.int/iris/handle/10665/205887> [Access on: 17/09/2020]
17. World Health Organization. Self-care in the context of primary health care. WHO Regional Office for South-East Asia. 2009. **Available at:** <https://apps.who.int/iris/handle/10665/206352> [Access on: 17/09/2020]
18. Global Self-Care Federation. COVID-19 statements. Global Self-Care Federation, Switzerland. 2020. Available at: <https://www.selfcarefederation.org/covid-19> [Access on: 17/09/2020]

19. World Health Organization. The Role of the pharmacist in self-care and self-medication: report of the 4th WHO Consultative Group on the Role of the Pharmacist, The Hague, The Netherlands, 26-28 August 1998. World Health Organization Geneva, Switzerland. 1998. **Available at:** <https://apps.who.int/iris/handle/10665/65860> [Access on: 17/09/2020]
20. Hughes CM, McElnay JC, Fleming GF. Benefits and risks of self-medication. *Drug Saf* 2001; 24:1027-37
21. World Health Organization (WHO). Addressing myths and rumours. World Health Organization Geneva, Switzerland. 1998. **Available at:** [https://www.who.int/bangladesh/emergencies/coronavirus-disease-\(covid-19\)-update/addressing-myths-and-rumours](https://www.who.int/bangladesh/emergencies/coronavirus-disease-(covid-19)-update/addressing-myths-and-rumours) [Access on: 17/09/2020]
22. Frenkel S, Alba D, Zhong R. Surge of virus misinformation stumps Facebook and Twitter. *The New York Times*, 2020. Available at: <https://www.nytimes.com/2020/03/08/technology/coronavirus-s-misinformation-social-media.html>. [Access on: 17/09/2020]
23. Figueiras A, Caamano F, Gestal OJJ. Sociodemographic factors related to self-medication in Spain. *Eur J Epidemiol*. 2000; 16:19-26
24. Khasawneh AI, Humeidan AA, Alsulaiman JW, Bloukh S, Ramadan M, Al-Shatanawi TN, et al. Medical Students and COVID-19: Knowledge, Attitudes, and Precautionary Measures. A Descriptive Study From Jordan. *Front Public Health*. 2020; 29:253. doi: 10.3389/fpubh.2020.00253.
25. Chu DK, Akl EA, Duda S, Solo K, Yaacoub S, Schünemann HJ; COVID-19 Systematic Urgent Review Group Effort (SURGE) study authors. Physical distancing, face masks, and eye protection to prevent person-to-person transmission of SARS-CoV-2 and COVID-19: a systematic review and meta-analysis. *Lancet*. 2020 Jun 27;395(10242):1973-1987. doi: 10.1016/S0140-6736(20)31142-9.
26. Center for Disease Control (CDC). Scientific brief: Community use of cloth masks to control the spread of SARS-CoV-2. CDC, Atlanta, USA. 2020. Available at: <https://www.cdc.gov/coronavirus/2019-ncov/more/masking-science-sars-cov2.html>
27. Li T, Liu Y, Li M, Qian X, Dai SY. Mask or no mask for COVID-19: A public health and market study. *PLOS ONE* 15(8): e0237691. <https://doi.org/10.1371/journal.pone.0237691>
28. Zhong B, Luo W, Li H, Zhang Q, Liu X, Li W, Li Y. Knowledge, attitudes, and practices towards COVID-19 among Chinese residents during the rapid rise period of the COVID-19 outbreak: a quick online cross-sectional survey. *Int J Biol Sci*. 2020; 16(10):1745–1752. doi: 10.7150/ijbs.45221
29. Clements JM. Knowledge and Behaviors Toward COVID-19 Among US Residents During the Early Days of the Pandemic: Cross-Sectional Online Questionnaire. *JMIR Public Health Surveill*. 2020 May 8;6(2):e19161. doi: 10.2196/19161. PMID: 32369759; PMCID: PMC7212816.
30. Hossain MA, Jahid MIK, Hossain KMA, Walton LM, Uddin Z, Haque MO, et al. (2020) Knowledge, attitudes, and fear of COVID-19 during the Rapid Rise Period in Bangladesh. *PLOS ONE*. 2020 15(9): e0239646. <https://doi.org/10.1371/journal.pone.0239646>
31. Ferdous MZ, Islam MS, Sikder MT, Mosaddek ASM, Zegarar-Valdivia JA, Gozal D, et al. (2020) Knowledge, attitude, and practice regarding COVID-19 outbreak in Bangladesh: An online-based cross-sectional study. *PLOS ONE* 15(10): e0239254. <https://doi.org/10.1371/journal.pone.0239254>
32. Hamza MS, Badary OA, Elmazar MM. Cross-Sectional Study on Awareness and Knowledge of COVID-19 Among Senior pharmacy Students. *J Community Health*. 2020; 1-8. doi:10.1007/s10900-020-00859-z
33. World Health Organization. Hand Hygiene for All Global Initiative. World Health Organization, Geneva, Switzerland. 2020. Available at: https://www.who.int/water_sanitation_health/sanitation-waste/sanitation/hand-hygiene-for-all/en/ [Access on: 17/09/2020]
34. Newbold, S.C., Finnoff, D., Thunström, L, Ashwort M, Shogren JF. Effects of Physical Distancing to Control COVID-19 on Public Health, the Economy, and the Environment. *Environ Resource Econ* **2020; 76**: 705–729. <https://doi.org/10.1007/s10640-020-00440-1>
35. Alimohamadi Y, Holakouie-Naieni K, Sepandi M, Taghdir M. Effect of Social Distancing on COVID-19 Incidence and Mortality in Iran Since February 20 to May 13, 2020: An Interrupted Time Series Analysis. *Risk Manag Healthc Policy*. 2020;13:1695-1700 <https://doi.org/10.2147/RMHP.S265079>
36. Haque M, Islam S, Iqbal S, Urmi UL, Kamal ZM, Rahman A, et al. Availability and price changes of potential medicines and equipment for the prevention and treatment of COVID-19 among pharmacy and drug store in Bangladesh: findings and implication. *Bangladesh J Medical Science*. 2020; 19: 36-50. <https://doi.org/10.3329/bjms.v19i0.48106>
37. Zhou YR. The global effort to tackle the coronavirus face mask shortage. *Prevention Web, UN office for Disaster Risk Reduction*. 2020. Available at: https://www.preventionweb.net/news/view/70908?fbclid=IwAR2gYKTW970bCQYQ_GJjgeOaCUuYuX8VNUhGapC5NoenQG5QTQEazLMGWxY

38. Lyons Z, Wilcox H, Leung L, Dearsley O. COVID-19 and the mental well-being of Australian medical students: impact, concerns and coping strategies used. *Australas Psychiatry*. 2020 Aug 10:1039856220947945. doi: 10.1177/1039856220947945. Epub ahead of print. PMID: 32772729; PMCID: PMC7424607.
39. Directorate General of Health Services (DGHS). Coronavirus Disease (COVID-19). DGHS, Dhaka, Bangladesh. 2020. Available at: <http://muktopaath.gov.bd/course-details/236> [Access on: 17/09/2020]
40. Khanom M, Hoque A, Sharif P, Sabuj M, Hossain MA. How were the Online Classes in Undergraduate Medical Teaching during COVID Pandemic? Students' Views of a Non-Government Medical College in Bangladesh. *BJME*.2020;11(2):1-3.