VCH SCHOOL STAFF AND STUDENT CASES AND CLUSTERS
MeSH: SARS-CoV-2, Communicable Disease Control, Schools, Epidemiology

SARS-CoV-2 transmission in K-12 schools in the Vancouver Coastal Health Region: a descriptive epidemiologic study

Diana Bark MD¹, Nalin Dhillon MSc², Martin St-Jean MSc², Brooke Kinniburgh MPH², Geoff McKee MD MPH^{1,3}, Alexandra Choi MD MHSc⁴

1. School of Population and Public Health, University of British Columbia, Vancouver BC
2. Population Health Surveillance Unit, Vancouver Coastal Health, Vancouver BC
3. Population and Public Health, British Columbia Centre for Disease Control, Vancouver BC
4. Office of the Chief Medical Health Officer, Vancouver Coastal Health, Vancouver BC

Address for correspondence: alexandra.choi1@vch.ca

Acknowledgements: We thank Amina Moustaqim-Barrette MPH (Vancouver Coastal Health) and Moe Zandy MSc (British Columbia Centre for Disease Control) for their contributions to data compilation and visualization.

Disclaimers: The views expressed in the article are the authors' own and not an official position of Vancouver Coastal Health or the British Columbia Centre for Disease Control.

Disclosures: AC is a Medical Health Officer and School Medical Officer for Vancouver Coastal Health. GM is the Medical Director for Population and Public Health at the BC Centre for Disease Control. Neither the study authors nor their institutions received payment or services at any time from any third parties for any aspects of the submitted work.

Data-Sharing Statement: The study data are not available for use by other researchers, other than that which are presented in the manuscript.

Ethical considerations: In consultation with UBC Office of Research Ethics, this study did not require institutional research ethics review according to Article 2.5 of the Tri-Council Policy Statement (TCPS2), as it was completed as part of routine public health surveillance and quality assurance/quality improvement.

VCH SCHOOL STAFF AND STUDENT CASES AND CLUSTERS MeSH: SARS-CoV-2, Communicable Disease Control, Schools, Epidemiology

Abstract

Background: There is an urgent need to assess the role of schools in the spread of SARS-CoV-2 in Canada to inform public health measures. We describe the epidemiology of SARS-CoV-2 infection in students and staff in the Vancouver Coastal Health (VCH) region in the first three months of the 2020/2021 academic year, and examine the extent of transmission in schools.

Methods: This descriptive epidemiologic study using contact tracing data included all SARS-CoV-2 cases reported to VCH between September 10 and December 18, 2020 who worked in or attended K-12 schools in-person. Case and cluster characteristics were described.

Results: There were 699 school staff and student cases during the study period, for an incidence of 55 cases per 10,000 population, compared to 73 per 10,000 population in all VCH residents. Among VCH resident staff and student cases, 53% were linked to a household case/cluster, <1.5% were hospitalized and there were no deaths. Out of 699 cases present at school, 26 clusters with school-based transmission resulted in 55 secondary cases. Staff members accounted for 54% of index cases (14/26) while comprising 14% of the school population. Among clusters, 88% had fewer than 4 secondary cases.

Interpretation: COVID-19 incidence in the school population was lower than that of the general population. There were no deaths and severe disease was rare. School-based transmissions of SARS-CoV-2 were uncommon and clusters were small. Our results support the growing body of evidence that schools do not play a major role in the spread of SARS-CoV-2.

Introduction

Early in the pandemic, schools closed across Canada to help reduce the circulation of SARS-CoV-2 and decrease risks associated with infection among staff and students(1). Evidence has since suggested that school closures and decreased in-person learning have had negative impacts on the mental health, wellbeing, educational attainment, social connection, healthcare access, and health behaviors of students, families, and staff(1-12). Combined with high socioeconomic tolls(13-15), these unintended

VCH SCHOOL STAFF AND STUDENT CASES AND CLUSTERS MeSH: SARS-CoV-2, Communicable Disease Control, Schools, Epidemiology

consequences have compelled consideration of the efficacy of school closures in limiting morbidity and mortality associated with the COVID-19 pandemic.

Evidence now indicates that age is a risk factor for severe disease and children typically experience a mild course of illness(16-22). There is also a growing body of literature suggesting children's in-person school attendance may not be a significant driver of transmission; contact-tracing and test-based studies from other jurisdictions have identified little or no transmission in the school setting(23-34). Internationally, studies illustrate that school re-opening has not impacted numbers of paediatric cases(23,35,36) and that the risk of SARS-CoV-2 infection has not increased for educational staff working with children(30,36,37). There is limited peer-reviewed data in Canada regarding risk of SARS-COV-2 transmission in schools, although Ottawa Public Health has reported low rates of school-based transmission(38). Given that transmission risk is influenced by community-level epidemiology, public health measures, health system infrastructure, and other jurisdictional factors, this study contributes needed evidence within the Canadian context. We aimed to describe cases who attended kindergarten to grade 12 (K-12) schools in-person, assess the extent of SARS-CoV-2 transmission in schools, and characterize clusters where school-based transmission was observed.

Methods

Setting

During the study period, while mass gatherings remained prohibited, schools and businesses were open with safety plans in place, and masks were encouraged but not required. Travel and gatherings in private residences were restricted gradually starting in November 2020. There were no stay-at-home requirements(39). Symptomatic testing was widely available and contact tracing was conducted for all positive cases(40).

VCH SCHOOL STAFF AND STUDENT CASES AND CLUSTERS MeSH: SARS-CoV-2, Communicable Disease Control, Schools, Epidemiology

Vancouver Coastal Health (VCH) serves over 1.1 million people(41) across urban, rural, and remote geographies. During the study period, 17,742 staff and 105,905 students worked in or attended school in-person at a total of 358 schools (104 independent and 254 public) in the VCH region. Schools in BC reopened following summer holidays on September 10, 2020 and closed for winter break after December 18, 2020. Prior to reopening, schools implemented COVID-19 safety plans developed with support from Public Health, which included public health measures (e.g. testing and contact tracing protocols), environmental measures (e.g. enhanced cleaning and disinfection), administrative measures (e.g. scheduling and work practices, health and wellness policies, and cohorting), personal measures (e.g. staying home when sick, physical distancing, hand hygiene and respiratory etiquette), and personal protective equipment. Non-medical mask use was supported, but not required(42).

Public Health Investigation and Response

SARS-CoV-2 Nucleic Acid Amplification testing (NAAT) was available for anyone with symptoms, and advised for students or staff with fever or new symptoms persisting >24 hours. Tests were generally processed within 24 hours. Positive tests were automatically reported to VCH's Office of the Chief Medical Health Officer, which investigated cases within 24 hours and identified contacts(43). Individual risk assessments were performed for all contacts, integrating cases' symptoms, ages of cases and contacts, nature and duration of contact, setting (e.g. indoor/outdoor), and presence/absence of known SARS-CoV-2 transmission. Factors constituting high-risk exposures included direct contact with infectious body fluids (e.g., via sharing vapes, being coughed or sneezed upon) and prolonged (>15 minutes) contact with a symptomatic case face to face or at close range; thus, close contacts were identified among classmates. Factors constituting low-risk exposures included the absence of interactions or interactions outdoors of limited duration and at a distance (>2m). Individuals with high-risk exposures were advised to self-isolate for 14 days and were monitored by Public Health for emergence of symptoms during their incubation period. Symptomatic individuals were directed to seek testing and few declined. Those with lower risk exposures were advised to self-monitor for symptoms and present for testing accordingly. Asymptomatic

VCH SCHOOL STAFF AND STUDENT CASES AND CLUSTERS MeSH: SARS-CoV-2, Communicable Disease Control, Schools, Epidemiology

individuals were tested at the discretion of the Medical Health Officer. Mass asymptomatic testing was undertaken in outbreak settings where uncontrolled spread was observed; this did not occur in any schools during the study period, and there were no school closures due to SARS-CoV-2 transmission.

Data sources

We obtained data from VCH's electronic COVID-19 case and contact management platform, capturing contact tracing interview data including demographics, school name, grade, role (staff/student), attendance dates, comorbidities, symptoms, and epidemiologic linkage to other cases within and outside of school. Hospitalizations and ICU admissions to VCH hospitals were determined using individual data linkages to acute care data. Provincial Health Services Authority provided hospitalization data for BC Children's Hospital. Districts confirmed denominators of staff and students attending in-person during the study period. Duplicate case entries were discarded based on unique identifiers. Cases with missing or incorrect locations were reclassified based on city of residence or postal code.

Study design

We extracted all lab-confirmed and probable cases aged 5 and older at time of case report between September 10 and December 18, 2020 who reported working in or attending a K-12 school within VCH during their incubation or infectious periods. We excluded cases who exclusively worked in or attended a daycare or post-secondary institution, attended school online, or were homeschooled. School staff and student cases were categorized as either school aged (aged 5-17) or aged 18+. We described case characteristics including age, sex, location, linkage to known cases/clusters, comorbidities, symptoms at time of initial interview, status (recovered/removed from isolation or deceased), and hospitalization/ICU admission.

To identify potential school transmission events, we analyzed all clusters where 2 or more cases were reported within a 14-day period in a school. Cases who did not interact with staff or students in the

VCH SCHOOL STAFF AND STUDENT CASES AND CLUSTERS MeSH: SARS-CoV-2, Communicable Disease Control, Schools, Epidemiology

learning setting (e.g., building maintenance staff only present after school hours) were excluded. A chart review of the cluster cases and their epidemiologically linked cases was conducted to verify the likely source of SARS-CoV-2 infection. A case was determined to be a contact's likely source if they had prolonged contact during the case's infectious period and the contact's incubation period. Sources were categorized as household, social, school, or other. Index and secondary cases were determined by symptom onset date. If there was no contact with a positive case, this was recorded. All cases with multiple potential exposures were reviewed by DB and AC. The source of acquisition was attributed to the school if the non-school source had an unclear symptom onset date and no clear source of acquisition themselves. Finally, secondary cases' close contacts who became lab-confirmed or probable cases were deemed tertiary cases. The primary outcome was the number of school-based transmissions.

All analyses were conducted via statistical computing software R Studio(44). P-values were based on Pearson's chi-squared and independent t-tests for categorical and continuous variables respectively.

Results

Description of cases in schools

In the four weeks prior to school reopening, weekly incidence of SARS-CoV-2 infection ranged 0.8-3.2 per 10,000 population in VCH based on 2019 population estimates(41), compared to 2.4-2.8 per 10,000 population in the four weeks following reopening. Weekly incidence peaked November 15-21 among VCH residents at 9.4 per 10,000 population and among K-12 staff and students at 6.6 per 10,000 population. Between September 10 and December 18, 8,755 cases among VCH residents were reported (73 per 10,000 population), of which 674 (8%) were K-12 staff or students. With an additional 25 staff and student cases who resided outside the VCH region, there were 699 cases among 123,646 staff and students working in or attending school in-person (55 per 10,000 population) in 270 unique VCH region K-12 schools during the same period. Characteristics of the 674 VCH resident staff and student cases are

VCH SCHOOL STAFF AND STUDENT CASES AND CLUSTERS

MeSH: SARS-CoV-2, Communicable Disease Control, Schools, Epidemiology

described in Table 1. Notably, adults (18+) more frequently reported comorbidities and were less frequently linked to a confirmed case/cluster compared to children (5-17 years) (p-values <0.01).

Figures 1A and B. 7-day average of new SARS-CoV-2 cases among VCH residents, reported between August 1 and December 18, 2020. Please note the different axis used in Figure 1b to compare trends between school and community cases

Figure 1A

VCH SCHOOL STAFF AND STUDENT CASES AND CLUSTERS

MeSH: SARS-CoV-2, Communicable Disease Control, Schools, Epidemiology

Figure 1B

Table 1. Characteristics of 674 student and staff SARS-CoV-2 cases residing in the VCH region reported between September 10 and December 18, 2020^a

Variable	18+ years, <i>N</i> =157 <i>n</i> (%)	5-17 years, <i>N</i> =517 <i>n</i> (%)	Total, <i>N</i> =674 <i>n</i> (%)
Age at time of report			
Median [Range]	39 [18-73]	12 [5-17]	14 [5-73]
Aged ≥65 years at time of report	2 (0.4)	NA	2 (0.3)
Sex			
Female	104 (66)	224 (43)	328 (49)

VCH SCHOOL STAFF AND STUDENT CASES AND CLUSTERS
MeSH: SARS-CoV-2, Communicable Disease Control, Schools, Epidemiology

Male	53 (33)	292 (57)	345 (51)
Health Service Delivery Area			
Coastal Rural	6 (3.8)	33 (6.4)	39 (5.8)
Coastal Urban	29 (19)	107 (21)	136 (20)
Richmond	29 (19)	99 (19)	128 (19)
Vancouver	93 (59)	278 (54)	371 (55)
Linked to confirmed case or cluster			
No	62 (40)	114 (22)	176 (26)
Yes, non-household contact	50 (32)	94 (18)	144 (21)
Yes, household contact	45 (29)	309 (60)	354 (53)
Case status			
Recovered/Removed from isolation	157 (100)	517 (100)	674 (100)
Deceased	0	0	0
Ever Hospitalized	-	-	<10 (1.5)
Ever Admitted to ICU	-	-	<10 (<1.5)
One or more comorbidities	36 (23)	37 (7.2)	73 (11)
Comorbidity types ^b			
Cancer	2 (1.3)	0 (0)	2 (0.3)
Cardiac	5 (3.2)	1 (0.2)	6 (0.9)
Diabetes	7 (4.5)	3 (0.6)	10 (1.5)
Immunocompromised	3 (1.9)	8 (1.5)	11 (1.6)
Pregnant	3 (1.9)	0 (0)	3 (0.4)
Respiratory	2 (1.3)	6 (1.2)	0 (0)
Smoking/vaping	27 (17)	22 (4.3)	49 (7.3)

VCH SCHOOL STAFF AND STUDENT CASES AND CLUSTERS

MeSH: SARS-CoV-2, Communicable Disease Control, Schools, Epidemiology

One or more COVID symptoms	155 (99)	501 (97)	656(97)
----------------------------	----------	----------	---------

^aData accurate as of January 12, 2021. Excludes 25 cases who resided outside of VCH region but worked in/attended school in-person in the VCH region. ^bComorbidity types were not mutually exclusive.

Clusters of SARS-CoV-2 infection in schools

There were 71 school clusters totaling 251 cases during the study period. No school-based transmission was suspected in 45 clusters (63%), with all 142 cases within these clusters likely acquiring SARS-CoV-2 outside of school. Within the 26 clusters containing at least one school-acquired case, a further 28 cases likely acquired infection outside of school. Of the 170 cases (19 staff and 151 students) temporally clustered but without evidence of school-based transmission, 107 (63%) were attributable to household contacts, 17 (10%) to social networks, 4 (2.4%) to sports teams, and 2 (1.2%) to work outside of school. 40 students and staff (24%) had no known contact with another case inside or outside of school.

Cases with evidence of school-based transmission

Out of 699 cases present at school, transmission stemming from 26 index cases resulted in 55 secondary cases at school. Two hundred and thirty-seven close contacts were identified from these 55 secondary cases, and 10 became tertiary cases. Staff accounted for 54% of index cases (14/26) and 25% of secondary cases (14/55). Staff cases (n=28) included 19 teachers, 6 support workers, and 3 principals/office staff members. Of the 81 index and secondary cases, 47 (58%) worked in or attended a K-7 school, 25 (31%) a grade 8-12 school, and 9 (11%) a K-12 school.

The characteristics of clusters with school-based transmission are presented in Table 2. Among 26 total clusters, 13 (50%) had only 1 secondary case, 10 (38%) had 2-3 secondary cases, and 3 (12%) had 4 or more secondary cases. All clusters with 4 or more secondary cases had an index staff case and occurred in grades K-7. The median size of clusters was 3 (range 2-8) when the index case was a staff member compared to 2 (range 2-3) when the index case was a student. There were 8 (31%) clusters involving

VCH SCHOOL STAFF AND STUDENT CASES AND CLUSTERS
MeSH: SARS-CoV-2, Communicable Disease Control, Schools, Epidemiology

students only, all among Grades 4-12, and 4 clusters (15%) involving staff members only. Of the 14 clusters including both staff and students, 9 (64%) occurred in grades K-7 and 5 (36%) in grades 8-12.

School-based activities where transmission may have occurred are presented in Table 3. Evidence of transmission was most commonly found in a shared classroom, but no specific class type was more frequently identified. For some cases, there were multiple exposures settings within the school, notably through socializing with a classmate.

Table 2. Characteristics of 26 clusters with evidence of school-based transmission

Variable	Grades K-7 <i>N=13 clusters</i>	Grades 8-12 <i>N=9 clusters</i>	Staff only <i>N=4 clusters</i>	All grades <i>N=26 clusters</i>
Number of cases per cluster				
Median [Range]	2 [2-8]	3 [2-4]	2.5 [2-4]	2.5 [2-8]
Role of index cases <i>N=26 cases</i> n (%)				
Staff	7 (54)	3 (33)	4 (100)	14 (54)
Student	6 (46)	6 (67)	0 (0)	12 (46)
Role of secondary cases <i>N=55 cases</i> n (%)				
Staff	4 (12)	3 (20)	7 (100)	14 (25)
Student	29 (88)	12 (80)	0 (0)	41 (75)

VCH SCHOOL STAFF AND STUDENT CASES AND CLUSTERS
MeSH: SARS-CoV-2, Communicable Disease Control, Schools, Epidemiology

Table 3. School settings where transmission may have occurred for 55 school-acquired SARS-CoV-2 cases^a

Exposure setting	Grades K-7 <i>N=35 exposure settings</i>	Grades 8-12 <i>N=17 exposure settings</i>	Staff only <i>N=7 exposure settings</i>	All grades n (%) <i>N=59 exposure settings</i>
Class	28 (26 students, 2 staff)	11 (10 students, 1 staff)	2 (2 staff)	41 (70)
Office/Meeting/Staff room	2 (2 staff)	1 (1 staff)	3 (3 staff)	6 (10)
Socializing at school	3 (3 students)	2 (2 students)	0	5 (8)
Unspecified interaction	0	1 (1 student)	2 (2 staff)	3 (5)
Field trip	0	2 (1 student, 1 staff)	0	2 (3)
Cohort ^b	2 (2 students)	0	0	2 (3)

^aThere were multiple potential exposure settings for some cases; an assessment of which within-school exposure setting was most likely responsible for transmission was not included as part of this study.

^bA cohort is a group of students and staff who remain together throughout a school term, the composition of which is consistent for all activities that occur in schools.

Interpretation

During the study period 8,755 cases of SARS-CoV-2 infection were reported among VCH residents (73 per 10,000 population) compared to 699 among VCH K-12 student and staff (55 per 10,000 population).

While 251 cases were temporally clustered with another case in the same school within a 14-day period, 170 (68%) of these cases had no evidence of school-based transmission. We identified 26 clusters with

evidence of school-based transmission resulting in 55 school-acquired cases (0.04% of 123,646

individuals working in or attending school) in 25/358 (7.0%) K-12 schools in the VCH region. Cluster

sizes were small, with 50% leading to 1 secondary case and 88% having fewer than 5 cases total; largest

clusters were observed in elementary schools. The index case was a staff member in over half of the

clusters (54%), although staff only constituted 35% (28/81) of all cases linked to school-based

VCH SCHOOL STAFF AND STUDENT CASES AND CLUSTERS

MeSH: SARS-CoV-2, Communicable Disease Control, Schools, Epidemiology

transmission and 14% of the overall school community. A shared classroom accounted for most transmissions; no particular class/activity was noticeably higher risk.

Our findings support the growing body of evidence that schools are likely not a significant driver of SARS-CoV-2 transmission(23,35,36). In concordance with other studies, we found that a household case was the predominant source of infection for school-aged children(30,32), and that school-based acquisitions were uncommon(29,30,32,33). Individuals with COVID-19 seldom infected others at school; out of 699 student and staff cases, we identified school-based transmissions stemming from 26 index cases (3.9%). The ratio of secondary cases to total primary cases was 0.09 in the school setting, compared to 0.27 observed in Luxembourg(30), 0.08 in Germany(32) and 0.02 in South Korea(23). While we did not systematically track the number of school contacts per index case, the secondary attack rate is likely to be very low given the low number of identified secondary cases, in concordance with prior studies across several countries(24,25,28,30). As observed elsewhere, where school-based transmission occurred, the number of secondary cases was limited(23,31,45), although one large high school outbreak was described in Israel(46) in the context of crowding and a concurrent heat wave. Interestingly, we observed a larger number of secondary cases when the index case was an adult, similar to the experience in England(31) and Singapore(26). Finally, our findings reinforced the observation that numerous cases in a school can reflect multiple introductions from the community rather than school-based transmissions(2).

Of note, our study was conducted in the context of limited circulation of SARS-CoV-2 variants of concern, strong collaboration between the public health and education sectors, good school-based infection prevention and control measures, and rapid access to testing and contact tracing. Contact tracing within 24 hours and regular monitoring of cases and contacts may have prevented transmission to tertiary cases. Further studies will be required to evaluate the impact of individual infection prevention and control measures.

VCH SCHOOL STAFF AND STUDENT CASES AND CLUSTERS MeSH: SARS-CoV-2, Communicable Disease Control, Schools, Epidemiology

Limitations

Our study was population-based and avoided bias due to cluster selection. However, our analysis was limited by reliance on routinely collected data and by potential recall bias, although case/proxy report was supplemented with information from school administrators as needed. While Medical Health Officers could request asymptomatic testing for the purposes of public health investigation, universal testing was not required. Transmissions to undetected (e.g. asymptomatic and not tested) secondary and tertiary cases were therefore not examined, which could disproportionately affect groups more likely to experience asymptomatic infection. However, studies that employed asymptomatic testing of school contacts reported low levels of transmission(24,25,28). Class sizes were not measured; thus, any association with school-based transmissions was not assessed. The potential for misclassification of the likely source of acquisition for cases with multiple exposures was minimized via independent review of these cases by two members of the research team. This study was limited to descriptive statistics and causal inferences cannot be drawn.

Conclusion

In conclusion, from September 10 to December 18, 2020 we detected minimal clusters and low rates of secondary transmission at schools. In-person school attendance may not expose students and staff to higher risks than those experienced in the community when infection prevention and control measures are in place, and adequate case and contact management capacity is available. Caution should be taken in generalizing these findings due to differences in underlying epidemiology, public health measures, health system infrastructure, and culture, among other factors that may differ by jurisdictions. However, our findings suggest that schools may be able to safely remain open in settings with moderate community transmission (i.e., weekly incidence of 10-100 cases per 100,000 as observed in this study) with infection prevention and control measures in place and sufficient public health capacity for rapid testing and contact tracing. Acknowledging harms associated with decreased school attendance, policy makers should

VCH SCHOOL STAFF AND STUDENT CASES AND CLUSTERS MeSH: SARS-CoV-2, Communicable Disease Control, Schools, Epidemiology

focus on the implementation of measures to safely operate in-person learning, and consider school closures and widespread quarantine of students and staff with great caution.

References

- (1) UNESCO. Education: From disruption to recovery. 2021; Available at: <https://en.unesco.org/covid19/educationresponse>. Accessed January 21, 2021.
- (2) European Centre for Disease Prevention and Control. COVID-19 in children and the role of school settings in transmission - first update. European Centre for Disease Prevention and Control 2020 December.
- (3) Dove N, Wong J, Gustafson R, Corneil T. Impact of School Closures on Learning, Child and Family Well-Being During the COVID-19 Pandemic. BC Centre for Disease Control & BC Children's Hospital 2020 September.
- (4) Xie X, Xue Q, Zhou Y, Zhu K, Liu Q, Zhang J, et al. Mental health status among children in home confinement during the coronavirus disease 2019 outbreak in Hubei Province, China. JAMA pediatrics 2020 September;174(9):898-900.
- (5) BC Centre for Disease Control. COVID-19 Survey and Dashboard. 2020; Available at: <http://www.bccdc.ca/health-info/diseases-conditions/covid-19/covid-19-survey>. Accessed January 21, 2021.
- (6) SickKids. Preliminary findings shared from COVID-19 child mental health research study . 2020; Available at: <https://www.sickkids.ca/en/news/archive/2020/covid19-mental-health-study-findings/>. Accessed January 21, 2021.
- (7) Moore SA, Faulkner G, Rhodes RE, Brussoni M, Chulak-Bozzer T, Ferguson LJ, et al. Impact of the COVID-19 virus outbreak on movement and play behaviours of Canadian children and youth: a national survey. International Journal of Behavioral Nutrition and Physical Activity 2020 December;17(1):1-11.
- (8) UNICEF. Averting a lost COVID generation: a six-point plan to respond, recover and reimagine a post-pandemic world for every child. UNICEF 2020 November 20.
- (9) Borkowski A, Correa JSO, Bundy DA, Burbano C, Hayashi C, Lloyd-Evans E, et al. COVID-19: Missing More Than a Classroom. The impact of school closures on children's nutrition. Innocenti Working Paper 2021-01. 2021 January.
- (10) Masonbrink AR, Hurley E. Advocating for children during the COVID-19 school closures. Pediatrics 2020 September 1;146(3).
- (11) Patrick SW, Henkhaus LE, Zickafoose JS, Lovell K, Halvorson A, Loch S, et al. Well-being of parents and children during the COVID-19 pandemic: a national survey. Pediatrics 2020 October;146(4).
- (12) Kim LE, Asbury K. 'Like a rug had been pulled from under you': The impact of COVID-19 on teachers in England during the first six weeks of the UK lockdown. Br J Educ Psychol 2020 December;90(4):1062-1083.
- (13) Statistics Canada. Impacts of COVID-19 on Canadian families and children. 2020; Available at: <https://www150.statcan.gc.ca/n1/daily-quotidien/200709/dq200709a-eng.htm>. Accessed January 21, 2021.
- (14) Hanushek EA, Woessmann L. The economic impacts of learning losses. OECD 2020.
- (15) Keogh-Brown MR, Jensen HT, Edmunds WJ, Smith RD. The impact of Covid-19, associated behaviours and policies on the UK economy: A computable general equilibrium model. SSM-population health 2020 December;12:100651.
- (16) Bailey LC, Razzaghi H, Burrows EK, Bunnell HT, Camacho PE, Christakis DA, et al. Assessment of 135 794 Pediatric Patients Tested for Severe Acute Respiratory Syndrome Coronavirus 2 Across the United States. JAMA pediatrics 2020 November 23.

VCH SCHOOL STAFF AND STUDENT CASES AND CLUSTERS

MeSH: SARS-CoV-2, Communicable Disease Control, Schools, Epidemiology

- (17) Leeb RT, Price S, Sliwa S, Kimball A, Szucs L, Caruso E, et al. COVID-19 trends among school-aged children—United States, March 1–September 19, 2020. *Morb Mortal Weekly Rep* 2020 October 2;69(39):1410.
- (18) Ding Y, Yan H, Guo W. Clinical characteristics of children with COVID-19: a meta-analysis. *Frontiers in pediatrics* 2020 July 3;8:431.
- (19) Ludvigsson JF. Children are unlikely to be the main drivers of the COVID-19 pandemic—a systematic review. *Acta Paediatrica* 2020 August;109(8):1525-1530.
- (20) Dong Y, Mo X, Hu Y, Qi X, Jiang F, Jiang Z, et al. Epidemiology of COVID-19 among children in China. *Pediatrics* 2020 June 1;145(6).
- (21) Götzinger F, Santiago-García B, Noguera-Julían A, Lanasa M, Lancella L, Carducci FIC, et al. COVID-19 in children and adolescents in Europe: a multinational, multicentre cohort study. *The Lancet Child & Adolescent Health* 2020 September 1;4(9):653-661.
- (22) Liguoro I, Pilotto C, Bonanni M, Ferrari ME, Pusiol A, Nocerino A, et al. SARS-COV-2 infection in children and newborns: a systematic review. *Eur J Pediatr* 2020 July;179:1029-1046.
- (23) Yoon Y, Kim K, Park H, Kim S, Kim Y. Stepwise School Opening and an Impact on the Epidemiology of COVID-19 in the Children. *J Korean Med Sci* 2020 November;35(46).
- (24) Brandal LT, Ofitserova TS, Meijerink H, Rykkvin R, Lund HM, Hungnes O, et al. Minimal transmission of SARS-CoV-2 from paediatric COVID-19 cases in primary schools, Norway, August to November 2020. *Eurosurveillance* 2021 January;26(1):2002011.
- (25) Macartney K, Quinn HE, Pillsbury AJ, Koirala A, Deng L, Winkler N, et al. Transmission of SARS-CoV-2 in Australian educational settings: a prospective cohort study. *The Lancet Child & Adolescent Health* 2020 November 1;4(11):807-816.
- (26) Yung CF, Kam K, Nadua KD, Chong CY, Tan NWH, Li J, et al. Novel coronavirus 2019 transmission risk in educational settings. *Clinical Infectious Diseases* 2020 June 25.
- (27) Heavey L, Casey G, Kelly C, Kelly D, McDarby G. No evidence of secondary transmission of COVID-19 from children attending school in Ireland, 2020. *Eurosurveillance* 2020 May 28;25(21):2000903.
- (28) Larosa E, Djuric O, Cassinadri M, Cilloni S, Bisaccia E, Vicentini M, et al. Secondary transmission of COVID-19 in preschool and school settings in northern Italy after their reopening in September 2020: a population-based study. *Eurosurveillance* 2020 December 10;25(49):2001911.
- (29) Zimmerman KO, Akinboyo IC, Brookhart MA, Boutzoukas AE, McGann K, Smith MJ, et al. Incidence and secondary transmission of SARS-CoV-2 infections in schools. *Pediatrics* 2021.
- (30) Mossong J, Mombaerts L, Veiber L, Pastore J, LeCoroller G, Schnell M, et al. SARS-CoV-2 Transmission in Educational Settings During an Early Summer Epidemic Wave in Luxembourg. Available at SSRN 3696896 2020 October 26.
- (31) Ismail SA, Saliba V, Bernal JL, Ramsay ME, Ladhani SN. SARS-CoV-2 infection and transmission in educational settings: a prospective, cross-sectional analysis of infection clusters and outbreaks in England. *The Lancet Infectious Diseases* 2020 December 8.
- (32) Ehrhardt J, Ekinci A, Krehl H, Meincke M, Finci I, Klein J, et al. Transmission of SARS-CoV-2 in children aged 0 to 19 years in childcare facilities and schools after their reopening in May 2020, Baden-Württemberg, Germany. *Eurosurveillance* 2020 September 10;25(36):2001587.
- (33) Falk A, Benda A, Falk P, Steffen S, Wallace Z, Høeg TB. COVID-19 Cases and Transmission in 17 K–12 Schools—Wood County, Wisconsin, August 31–November 29, 2020. *Morb Mortal Weekly Rep* 2021 January 29;70(4):136.
- (34) Danis K, Epaulard O, Bénet T, Gaymard A, Campoy S, Botelho-Nevers E, et al. Cluster of coronavirus disease 2019 (COVID-19) in the French Alps, February 2020. *Clinical Infectious Diseases* 2020 July 28;71(15):825-832.
- (35) Somekh I, Shohat T, Boker LK, Simões EA, Somekh E. Reopening Schools and the Dynamics of SARS-CoV-2 Infections in Israel: A Nationwide Study. *Clinical Infectious Diseases* 2021.
- (36) Public Health Agency of Sweden. Covid-19 in schoolchildren – A comparison between Finland and Sweden. Public Health Agency of Sweden 2020 July 7.

VCH SCHOOL STAFF AND STUDENT CASES AND CLUSTERS

MeSH: SARS-CoV-2, Communicable Disease Control, Schools, Epidemiology

- (37) National Collaborating Centre for Methods and Tools. Living Rapid Review Update 11: What is the specific role of daycares and schools in COVID-19 transmission? . National Collaborating Centre for Methods and Tools 2020 December 18.
- (38) Ottawa Public Health. Special Focus: COVID-19 in Schools. 2021 February. Available at: https://www.ottawapublichealth.ca/en/reports-research-and-statistics/resources/Documents/SpecialReport_COVIDinSchools_EN_final_2021-02-01.pdf.
- (39) Government of British Columbia. COVID-19 (Novel Coronavirus) – Orders and notices. 2021 April 28. Available at: <https://www2.gov.bc.ca/gov/content/health/about-bc-s-health-care-system/office-of-the-provincial-health-officer/current-health-topics/covid-19-novel-coronavirus#archived-orders>.
- (40) BC Centre for Disease Control. Phases of COVID-19 testing in BC. 2021. Available at: <http://www.bccdc.ca/health-info/diseases-conditions/covid-19/testing/phases-of-covid-19-testing-in-bc>.
- (41) BC Stats. British Columbia - Population Estimates. 2020; Available at: <https://bcstats.shinyapps.io/popApp/>. Accessed Jan 21, 2021.
- (42) BC Centre for Disease Control. COVID-19 Public Health Guidance for K-12 School Settings. 2020 September 11.
- (43) BC Centre for Disease Control. Interim Guidance: Public Health Management of cases and contacts associated with novel coronavirus (COVID-19) in the community. 2020 September.
- (44) R Core Team. R: A language and environment for statistical computing. 2020.
- (45) Fong MW, Cowling BJ, Leung GM, Wu P. Letter to the editor: COVID-19 cases among school-aged children and school-based measures in Hong Kong, July 2020. *Eurosurveillance* 2020;25(37):2001671.
- (46) Stein-Zamir C, Abramson N, Shoob H, Libal E, Bitan M, Cardash T, et al. A large COVID-19 outbreak in a high school 10 days after schools' reopening, Israel, May 2020. *Eurosurveillance* 2020 July 23;25(29):2001352.
- (47) BC Centre for Disease Control. COVID-19 Case Report Form. 2020 October 29.

VCH SCHOOL STAFF AND STUDENT CASES AND CLUSTERS

MeSH: SARS-CoV-2, Communicable Disease Control, Schools, Epidemiology

Appendix

The study protocol and BC Centre for Disease Control COVID-19 Public Health Guidance for K-12

School Settings document are available upon request by emailing the corresponding author.

Table A1. Data dictionary

Term	Definition
Lab confirmed case	A person with laboratory confirmation of infection with the virus that causes COVID-19 performed at a community, hospital or reference laboratory (NML or a provincial public health laboratory) running a validated assay. This consists of detection of at least one specific gene target by a NAAT assay (e.g. real-time PCR or nucleic acid sequencing).(47)
Probable-lab case	A person (who has had a laboratory test) with fever (over 38 degrees Celsius) or new onset of (or exacerbation of chronic) cough AND who meets the COVID-19 exposure criteria and in whom a laboratory diagnosis of COVID-19 is inconclusive. Inconclusive is defined as an indeterminate test on a single or multiple real-time PCR target(s) without sequencing confirmation or a positive test with an assay that has limited performance data available.(47)
Probable epi-linked case	A person (who has not had a laboratory test) with fever (over 38 degrees Celsius) or new onset of (or exacerbation of chronic) cough AND either close contact with a confirmed case of COVID-19 or lived in or worked in a closed facility known to be experiencing an outbreak of COVID-19 (e.g., long-term care facility, prison).(47)
COVID-19 Symptoms	Any one of: abdominal pain, arthralgia, chills, coma, confusion, conjunctivitis, cough, diarrhea, discoloration of toes or fingers, dizziness, fatigue, fever, headache, loss of appetite loss of sense of smell/taste, myalgia, nasal congestion, nausea, pharyngitis, rash, rhinorrhea, shortness of breath/breathing difficulty, or vomiting.(47)
Recovered / Removed from isolation	Self-isolation for a person diagnosed with COVID-19 has been discontinued based on criteria outlined by Interim Guidance(43,47).
Deceased	Death from any cause occurring in a person diagnosed with COVID-19 with no period of complete recovery between illness and death.(47)
Ever Hospitalized	Any person admitted to a hospital for at least an overnight stay, or with a prolongation of hospitalization, for reasons directly or indirectly related to their COVID-19 infection, and with no period of complete recovery between illness and admission.(47)
Ever Admitted to ICU	Any person admitted to an intensive care unit (ICU) for at least an overnight stay, or with a prolongation of ICU stay, for reasons directly or indirectly related to their COVID-19 infection, and with no period of complete recovery between illness and admission.(47)

VCH SCHOOL STAFF AND STUDENT CASES AND CLUSTERS
MeSH: SARS-CoV-2, Communicable Disease Control, Schools, Epidemiology

Cancer	Malignancy or cancer diagnosed in the last 5 years.
Cardiac	Chronic cardiac disease, excluding hypertension.
Diabetes	Client self-report at time of case interview.
Immunocompromised	Client self-report at time of case interview.
Pregnant	Client self-report at time of case interview.
Respiratory	Other chronic respiratory / pulmonary condition, excluding asthma.
Smoking/vaping	Client self-report at time of case interview.
Linked to confirmed case: Yes, non-household contact	Client report close contact with a probable or confirmed case of COVID-19 in a non-household setting within 14 days prior to illness onset.
Linked to confirmed case: Yes, household contact	Client reported close contact with a probable or confirmed case of COVID-19 in a household setting within 14 days prior to illness onset.
Linked to confirmed case: No	Client did not report close contact with a probable or confirmed case of COVID-19 within 14 days prior to illness onset.
Staff	Case reporting working in a K-12 school within the learning setting in-person, or directly interacting with those within the learning setting, including teachers, support workers, principals/vice principals and office staff members.
Student	Case (or proxy) reporting attending a K-12 school in-person.
Cohort	A group of students and staff who remain together throughout a school term, the composition of which is consistent for all activities that occur in schools (42). While classes are the primary form of grouping where students spend the majority of their time, classes within a shared cohort may interact with each other for activities like physical education or music.

VCH SCHOOL STAFF AND STUDENT CASES AND CLUSTERS
MeSH: SARS-CoV-2, Communicable Disease Control, Schools, Epidemiology

Figure A1. Study population selection flow diagram (including non-VCH resident staff and student cases)

Figure A2. School clusters identification flow diagram